

# Gröbner Bases — Homework 1

Philip Warton

January 9, 2022

## Problem 1

Let  $S = \{x - y, z\} \subseteq \mathbb{R}[x, y, z]$ . Describe the variety  $V(S)$  both using set notation and geometrically.

---

### Solution

We know that the variety  $V(S)$  is the set of points such that every function in  $S$  is equal to 0 at said point. That is, in set notation,

$$\begin{aligned} V(S) &= \{(x, y, z) \in \mathbb{R}^3 \mid x - y = 0 = z\} \\ &= \{(x, y, z) \in \mathbb{R}^3 \mid x = y \text{ and } z = 0\}. \end{aligned}$$

We can also just write this as all points in  $\mathbb{R}^3$  of the form  $(\alpha, \alpha, 0)$  where  $\alpha \in \mathbb{R}$ . Geometrically, this consists of a straight line passing through the origin that lies in the  $x$ - $y$  plane that is represented by the equation  $x = y$ .

## Problem 2

Let  $K$  be a field. Recall that for a set  $S \subseteq K^n$ , we define

$$I(S) = \{f \in K[x_1, \dots, x_n] \mid f(\vec{a}) = 0 \text{ for all } \vec{a} \in S\}$$

Prove that  $I(S)$  is an ideal of the ring  $K[x_1, \dots, x_n]$ .

---

### Solution

*Proof.* Recall that a subring  $I$  of a ring  $R$  is an ideal if for every  $x \in R, y \in I$ , we can say  $xy \in I$ . In our case specifically, let  $S \subseteq K^n$ , let

$$f \in K[x_1, \dots, x_n],$$

and let

$$g \in I(S).$$

Choose any  $\vec{a} \in S$  to be arbitrary. Then

$$\begin{aligned} (fg)(\vec{a}) &= f(\vec{a})g(\vec{a}) \\ &= f(\vec{a}) \cdot 0 \\ &= 0 \end{aligned}$$

Since  $(fg)(\vec{a}) = 0$  for any  $\vec{a} \in S$ , it follows that  $fg \in I(S)$  and it must be true that  $I(S)$  is an ideal. □