Assignment -1 in LaTeX

Mathew M Philip

EE22BTECH11211

Problem 10.13.3.21:

Two dice are thrown together. Find the probability that the product of the numbers on the top of the dice is

1) 6
2) 12
3) 7

Solution:

X = Outcome of the first dice

Y = Outcome of the second dice

Z = XY

$$
\Pr(X) = \begin{cases} 0 & x < 1 \\ \frac{1}{6} & 1 \le x \le 6 \\ 0 & x > 6 \end{cases} \tag{1}
$$

The pmf of Y is also same as above.

$$
E(X) = \sum_{X_i=1}^{X_6} X_i \Pr(X_i) = \frac{1}{6} \times \left(\sum_{X=1}^{6} X \right) \tag{2}
$$

$$
= \frac{1}{6} \times \frac{6 \times 7}{2} = \frac{7}{2} \tag{3}
$$

Similarly

$$
E(Y) = \frac{7}{2} \tag{4}
$$

Since X,Y are independent random variables

$$
E(XY) = E(X) \times E(Y) \tag{5}
$$

$$
= \frac{7}{2} \times \frac{7}{2} \tag{6}
$$

$$
= \frac{49}{4} \tag{7}
$$

$$
F_Y(y) = \begin{cases} 0 & y < 1 \\ \frac{y}{6} & 1 \le y \le 6 \\ 1 & y > 6 \end{cases} \tag{8}
$$

$$
F_Z(n) = \sum_{k=1}^{6} \Pr(X = k) F_Y\left(\left(\frac{n}{k}\right) | k\right) \tag{9}
$$

$$
= \sum_{k=1}^{6} \frac{1}{6} \times F_Y\left(\left(\frac{n}{k}\right) | k\right) \tag{10}
$$

$$
= \frac{1}{6} \times \sum_{k=1}^{6} F_Y\left(\left(\frac{n}{k}\right) | k\right) \tag{11}
$$

where $\Pr(X)$ denotes pmf of random variable X $F_Y(y)$ denotes cdf of random variable Y

$$
F\left(\frac{n}{k}\right) = \begin{cases} 1 & k < \frac{n}{6} \\ \frac{\left[\frac{n}{k}\right]}{6} & k \ge \frac{n}{6} \cap \frac{n}{k} \notin I \\ \frac{\left(\frac{n}{k}\right)}{6} & k \ge \frac{n}{6} \cap \frac{n}{k} \in I \end{cases} \tag{12}
$$

where $[x]$ denotes the greatest integer less than or equal to x

$$
F_Z(n) = \frac{1}{6} \times \sum_{k=1}^{6} F_Y\left(\frac{n}{k}\right) \tag{13}
$$

$$
= \frac{1}{6} \times \left\{ \sum_{k=1}^{\left[\frac{n}{6}\right]} F_Y\left(\frac{n}{k}\right) + \sum_{k=\left[\frac{n}{6}\right]+1}^{6} F_Y\left(\frac{n}{k}\right) \right\} \tag{14}
$$

$$
= \frac{1}{6} \times \left\{ \sum_{k=1}^{\left[\frac{n}{6}\right]} 1 + \sum_{k=\left[\frac{n}{6}\right]+1}^{6} F_Y\left(\frac{n}{k}\right) \right\} \tag{15}
$$

$$
= \frac{1}{6} \times \left\{ \left[\frac{n}{6}\right] + \sum_{k=\left[\frac{n}{6}\right]+1}^{6} F_Y\left(\frac{n}{k}\right) \right\} \tag{16}
$$

$$
\tag{17}
$$

See Fig. 3

$$
\Pr(Z = n) = F_z(n) - F_z(n-1) \tag{18}
$$

See Fig. 3

X	2	3	4	6
Y	6	4	3	2
Pr(Z=12)	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$

TABLE 2: Cases for which product of outcome = 12

No. of events for product to be $12 = 4$

$$\Pr(Z = 12) = F_z(12) - F_z(11) \qquad (22)$$

$$= 4 \times \frac{1}{36} = \frac{1}{9} \qquad (23)$$

3) Product = 7

$Z = 7$ for $(X, Y) = \{\}$

There are no events for which product of outcome = 7

$$\Pr(Z = 7) = 0 \qquad (24)$$

Fig. 3: Plot of cummulative Distribution function

Fig. 3: Plot of Probability Mass Function

Since random variables X,Y are independent

$$\Pr(X, Y) = \Pr(X) \times \Pr(Y) = \frac{1}{6} \times \frac{1}{6} = \frac{1}{36} \qquad (19)$$

1) Product = 6

X	1	2	3	6
Y	6	3	2	1
Pr(Z=6)	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$

TABLE 1: Cases for which product of outcome = 6

No. of events for product to be $6 = 4$

$$\Pr(Z = 6) = F_z(6) - F_z(5) \qquad (20)$$

$$= 4 \times \frac{1}{36} = \frac{1}{9} \qquad (21)$$

2) Product = 12