

Probability Assignment

Gautam Singh

Abstract—This document contains the solution to Question 17 of Exercise 1 in Chapter 13 of the class 12 NCERT textbook.

- 1) If A and B are events such that

$$\Pr(A|B) = \Pr(B|A) \quad (1)$$

then

- a) $A \subset B$ but $A \neq B$
- b) $A = B$
- c) $A \cap B = \phi$
- d) $\Pr(A) = \Pr(B)$

Solution: Using Bayes' Rule,

$$\Pr(A, B) = \Pr(A) \Pr(B|A) \quad (2)$$

$$= \Pr(B) \Pr(A|B) \quad (3)$$

Using (1) in (2) and (3),

$$\Pr(A) = \Pr(B) \quad (4)$$

We consider the options one by one.

- a) If $A \subset B$ and $A \neq B$, then we can write $B = A + C$, where $AC = 0$ and $C \neq 0$. Thus,

$$\Pr(B) = \Pr(A + C) \quad (5)$$

$$= \Pr(A) + \Pr(C) - \Pr(AC) \quad (6)$$

$$= \Pr(A) + \Pr(C) > \Pr(A) \quad (7)$$

However, (7) contradicts (4).

- b) We give a counterexample to show this is wrong. Consider A as the event that an even number shows on rolling a fair die and B as the event that a prime number shows on rolling a fair die. The joint pmf is shown in Table 1. Clearly,

$$\Pr(A|B) = \frac{\Pr(AB)}{\Pr(B)} = \frac{\frac{1}{6}}{\frac{1}{6} + \frac{1}{3}} = \frac{1}{2} \quad (8)$$

$$\Pr(B|A) = \frac{\Pr(AB)}{\Pr(A)} = \frac{\frac{1}{6}}{\frac{1}{6} + \frac{1}{3}} = \frac{1}{2} \quad (9)$$

- c) The same example as before provides the required counterexample, as $\Pr(AB) = \frac{1}{6}$.

- d) This is the correct answer, as discussed above.

	A	\bar{A}
B	$\frac{1}{6}$	$\frac{1}{3}$
\bar{B}	$\frac{1}{3}$	$\frac{1}{6}$

TABLE 1: Joint pmf for events A and B .