

“Sparknotes” for *Linear Algebra* by Peter Lax

Muthu Chidambaram

Last Updated: May 23, 2019

Contents

1	Fundamentals	3
1.1	Exercise 1	3
1.2	Exercise 2	3
1.3	Exercise 3	3
1.4	Exercise 4	3
1.5	Exercise 5	3
1.6	Exercise 6	3
1.7	Exercise 7	3
1.8	Exercise 8	3
1.9	Exercise 9	3
1.10	Exercise 10	3
1.11	Exercise 11	3
1.12	Exercise 12	4
1.13	Exercise 13	4
1.14	Exercise 14	4
1.15	Exercise 15	4
1.16	Exercise 16	4
1.17	Exercise 17	4
2	Duality	5
2.1	Exercise 1	5
2.2	Exercise 2	5
3	Linear Mappings	6
3.1	Exercise 1	6
3.2	Exercise 2	6
3.3	Exercise 3	6
3.4	Exercise 4	6
3.5	Exercise 5	6
3.6	Exercise 6	7
3.7	Exercise 7	7

About

“A modern mathematical proof is not very different from a modern machine, or a modern test setup: the simple fundamental principles are hidden and almost invisible under a mass of technical details.”

- Hermann Weyl

These notes contain short summaries of (my) proof ideas for exercises and some theorems from the book *Linear Algebra* by Peter Lax. I have tried to make the summaries as brief as possible, sometimes only one line or one equation. My hope is that the summaries will give enough information to reconstruct a full proof without bogging the reader down with details. In many cases, I am sure that I inadvertently sacrificed clarity in an attempt to obtain brevity, and would greatly appreciate any feedback.

Also, I like when people include (what they presume to be) relevant quotes in their notes, so I have to ask you to forgive my haughtiness in starting these notes with a quote from Hermann Weyl.

1 Fundamentals

1.1 Exercise 1

$$x + z = x = x + z' \implies z = z'.$$

1.2 Exercise 2

$$0x + x = (0 + 1)x = x.$$

1.3 Exercise 3

Coefficients can be represented as row vectors.

1.4 Exercise 4

Function can be represented as row vector by letting $a_i = f(s_i)$ for each $s_i \in S$.

1.5 Exercise 5

Follows from exercises 3 and 4.

1.6 Exercise 6

$$y_1 + z_1 + y_2 + z_2 = (y_1 + y_2) + (z_1 + z_2) \text{ and } k(y_1 + z_1) = ky_1 + kz_1.$$

1.7 Exercise 7

$$a \in Y \cap Z \implies ka \in Y, ka \in Z \implies ka \in Y \cap Z.$$

1.8 Exercise 8

$$k0 = 0, 0 + 0 = 0.$$

1.9 Exercise 9

If S contains x_i then it must contain kx_i .

1.10 Exercise 10

If $x_i = 0$, k_i can be anything.

1.11 Exercise 11

$$x = \sum_{i=1}^m \sum_{j=1}^{\dim Y_i} y_j^{(i)}.$$

1.12 Exercise 12

Complete basis for W to U and V . Use W basis vectors and additional U and V basis vectors to get $\dim X = \dim U - \dim W + \dim V - \dim W + \dim W$.

1.13 Exercise 13

Send i^{th} basis vector to e_i , where e_i is vector of all zeroes except a one in the i^{th} place. Can permute mapping to get different isomorphisms.

1.14 Exercise 14

$$x_1 - x_2 + x_2 - x_3 = x_1 - x_3.$$

1.15 Exercise 15

$$x' = x + z_x, y' = y + z_y \implies x' + y' = x + y + (z_x + z_y).$$

1.16 Exercise 16

$$x \in X_1 \oplus X_2 \implies x = (x_1, x_2) = (x_1, 0) + (0, x_2).$$

1.17 Exercise 17

Construct a basis for X from Y : $y_1, \dots, y_j, x_{j+1}, \dots, x_n$.
Then $X/Y = \text{span}\{x_{j+1}, \dots, x_n\}$.

2 Duality

Theorem 1

$$x = \sum_{i=1}^n a_i x_i \implies k_i(x) = a_i.$$

2.1 Exercise 1

$$l_1, l_2 \in Y^\perp \implies l_1(y) + l_2(y) = 0 = (l_1 + l_2)(y).$$

2.2 Exercise 2

$$\forall \xi \in Y^{\perp\perp} \implies \forall l \in Y^\perp, \xi(l) = 0 = l(y) \forall y \in Y.$$

3 Linear Mappings

3.1 Exercise 1

- (a) $x \in X \implies x = \sum_{i=1}^n k_i x_i \implies T(x) = \sum_{i=1}^n k_i T(x_i) \in U$.
(b) $T(x), T(y) \in U \implies T(x+y) \in U \implies x+y \in X$.

Theorem 1

$$x \in X, y \in N_T \implies T(x+y) = T(x) + T(y) = T(x).$$

3.2 Exercise 2

- (a) Differentiation constant and sum rules imply linearity, and multiplication by s is distributive. Take $p(s) = 1$ to see that $ST \neq TS$.
(b) Rotation by 90 degrees amounts to swapping and negating coordinates, which is linear. Take $p = (1, 1, 0)$ to see that $ST \neq TS$.

3.3 Exercise 3

- (i) $T^{-1}(T(a+b)) = T^{-1}(T(a)+T(b)) = a+b = T^{-1}(T(a)) + T^{-1}(T(b))$.
(ii) Composition of isomorphisms is an isomorphism, hence ST is invertible.

3.4 Exercise 4

- (i) Let $T : X \rightarrow U$, $S : U \rightarrow V$ and $l_v \in V'$. Then $(ST)'(l_v) = l_v(ST) = (l_v S)T = (S'l_v)T = T'S'l_v$, since $S'l_v \in U'$.
(ii) Follows from linearity of transpose (definition).
(iii) Let $T : X \rightarrow U$ be an isomorphism. Then $l_x = l_u T \implies l_x T^{-1} = l_u$ for $l_u \in U'$, $l_x \in X'$.

3.5 Exercise 5

$T''(l_{x'}) = l_{x'} T'$ where $l_{x'} \in X''$ and $l_{x'} T' \in U''$. Since we can identify elements in X'' and U'' with elements in X and U respectively, we have that T'' assigns elements of U to X .

Theorem 2'

Since $T' : U' \rightarrow X'$ we have $l_u \in N_{T'} \implies T'(l_u) = l_u T = 0$. $N_{T'}^\perp$ consists of elements $l_{u'} | l_{u'}(l_u) = 0$. From $l_u T x = 0$ we have that each $l_{u'}$ is identified with a $u \in R_T$.

3.6 Exercise 6

The first two elements of x are already 0 after applying P , so $P^2 = P$. Linearity follows from linearity of vector addition.

3.7 Exercise 7

P is linear since function addition is linear. $P^2 f = \frac{f(x)+f(-x)}{4} + \frac{f(x)+f(-x)}{4} = Pf$.