



Problem 2.3.9.

$$a) M = P_{23} E_{21} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ -1 & 1 & 0 \end{bmatrix}$$

$$b) M = E_{31} P_{23} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ -1 & 1 & 0 \end{bmatrix}$$

we need  $E_{31}$  to exchange row 3 after the  $P_{23}$ .

Problem 2.3.12.

$$\begin{aligned} \text{Sol. } & \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} & \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 1 & 3 & 4 \\ 1 & 4 & 6 \end{bmatrix} \\ & = \begin{bmatrix} 7 & 8 & 9 \\ 4 & 5 & 6 \\ 1 & 2 & 3 \end{bmatrix} \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} & = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & -2 \\ 0 & 2 & -3 \end{bmatrix} \\ & = \begin{bmatrix} 9 & 8 & 7 \\ 6 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix} \end{aligned}$$

Problem 2.3.17

$$\text{Sol. } \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 3 & 9 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 4 \\ 8 \\ 14 \end{bmatrix}$$

$$\text{It gives } \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 3 & 9 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 4 \\ 8 \\ 14 \end{bmatrix}$$

and  $a=2, b=1, c=1$  is the solution.

Problem 2.3.28.

~~Sol.~~ Since  $AB=I$ ,  $ABC=IC=C$ 

Proof.

$$\Rightarrow ACBC=C$$

$$\Rightarrow AI=C \text{ (Since } BC=I)$$

$$\Rightarrow A=C \text{ QED.}$$

Problem 2.4.6.

$$\text{Sol. When } A = \begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix}, B = \begin{bmatrix} 1 & 0 \\ 3 & 0 \end{bmatrix}$$

$$(A+B)^2 = \left( \begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 3 & 0 \end{bmatrix} \right)^2 = \begin{bmatrix} 2 & 2 \\ 3 & 0 \end{bmatrix} \begin{bmatrix} 2 & 2 \\ 3 & 0 \end{bmatrix} = \begin{bmatrix} 10 & 4 \\ 6 & 6 \end{bmatrix}$$

$$\begin{aligned} A^2 + 2AB + B^2 &= \begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix}^2 + 2 \begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 3 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 3 & 0 \end{bmatrix}^2 \\ &= \begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 14 & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 3 & 0 \end{bmatrix} = \begin{bmatrix} 16 & 2 \\ 3 & 0 \end{bmatrix} \end{aligned}$$

$$\text{So } (A+B)^2 \neq A^2 + 2AB + B^2$$

$$(A+B)^2 = A^2 + AB + BA + B^2$$



班级: 计23 姓名: 郑东森 编号: 2022010799 科目: Linear Algebra 第2页

Problem 2.4.15.

Sol. (a) True (Let  $A$  be a  $m \times n$  matrix, since  $A^2$  is defined,  $n=m$ ,  $A$  has to be square)

(b) False (Let  $A$  be a  $2 \times 3$  matrix and  $B$  be a  $3 \times 2$  matrix. Then  $AB$  and  $BA$  are both defined and  $A$  and  $B$  are not square)

(c) True (If  $AB$  and  $BA$  are both defined then  $A$  has to be a  $m \times n$  matrix and  $B$  has to be a  $n \times m$  matrix. then  $AB$  has to be a  $m \times m$  matrix,  $BA$  has to be a  $n \times n$  matrix.  $AB$  and  $BA$  are both square).

(d) False (If  $B=0$ ,  $A$  is unnecessary to be  $I$ )

Problem 2.4.18.

Sol. (a)  $\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 2 \\ 1 & 2 & 3 \end{bmatrix}$  (b)  $\begin{bmatrix} 1 & -1 & 1 \\ -1 & 1 & -1 \\ 1 & -1 & 1 \end{bmatrix}$  (c)  $\begin{bmatrix} 1 & \frac{1}{2} & \frac{1}{3} \\ 2 & 1 & \frac{2}{3} \\ 3 & \frac{3}{2} & 1 \end{bmatrix}$

Problem 2.4.21

Sol.  $A^2 = \begin{bmatrix} 0 & 2 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 2 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 4 & 0 \\ 0 & 0 & 0 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$

$A^3 = \begin{bmatrix} 0 & 2 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 4 & 0 \\ 0 & 0 & 0 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 8 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$  (Since  $A^3 = A(AA^2)$ )

$A^4 = \begin{bmatrix} 0 & 0 & 0 & 8 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 2 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$

~~$A^2 v = \begin{bmatrix} 0 & 2 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ t \end{bmatrix} = \begin{bmatrix} 2y \\ 2z \\ 2t \\ 0 \end{bmatrix}$~~

$A^2 v = \begin{bmatrix} 0 & 0 & 4 & 0 \\ 0 & 0 & 0 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} y \\ y \\ z \\ t \end{bmatrix} = \begin{bmatrix} 4z \\ 4t \\ 0 \\ 0 \end{bmatrix}$

$A^3 v = \begin{bmatrix} 0 & 0 & 0 & 8 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ t \end{bmatrix} = \begin{bmatrix} 8t \\ 0 \\ 0 \\ 0 \end{bmatrix}$   $A^4 v = 0$ .

Problem 2.4.26.

$$AB = \begin{bmatrix} 1 & 0 \\ 2 & 4 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 3 & 3 & 0 \\ 1 & 2 & 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix} \begin{bmatrix} 3 & 3 & 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 4 \\ 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 3 & 0 \\ 6 & 6 & 0 \\ 6 & 6 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 & 0 \\ 4 & 8 & 4 \\ 1 & 2 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 3 & 0 \\ 10 & 14 & 4 \\ 7 & 8 & 1 \end{bmatrix}$$





班级: 计23 姓名: 郭东林 编号: 202200799 科目: Linear Algebra 第 3 页

Problem 2.4.32.

Sol.  $AX = A \cdot [x_1 \ x_2 \ x_3] = [Ax_1 \ Ax_2 \ Ax_3] = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

Graded Problems.

Problem 1.

Sol.  $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix}$

$$\begin{bmatrix} 0 & a \\ 0 & c \end{bmatrix} = \begin{bmatrix} c & d \\ 0 & 0 \end{bmatrix}$$

so we have  $a=d, c=0$

$A = \begin{bmatrix} a & b \\ 0 & a \end{bmatrix}$  and  $\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$  commute

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$\begin{bmatrix} b & 0 \\ d & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ a & b \end{bmatrix}$$

so we have  $b=0, a=d$

$A = \begin{bmatrix} a & 0 \\ c & a \end{bmatrix}$  and  $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$  commute.

Let  $\begin{bmatrix} a & 0 \\ c & a \end{bmatrix} = \begin{bmatrix} a & b \\ 0 & a \end{bmatrix}$

we have  $c=b=0$

so  $\begin{bmatrix} a & 0 \\ 0 & a \end{bmatrix}$  commute with both  $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$

Problem 2.

$$\begin{bmatrix} -2 & -1 & 0 & 0 \\ -1 & 2 & -1 & -1 \\ 0 & -1 & 2 & 0 \\ 0 & -1 & 0 & 2 \end{bmatrix} \begin{bmatrix} -1 & 0 & 0 & 1 \\ 0 & 1 & -1 & 0 \\ 0 & -1 & 1 & 0 \\ 1 & 0 & 0 & -1 \end{bmatrix} = \begin{bmatrix} -2 & -1 & 1 & 2 \\ 0 & 3 & -3 & 0 \\ 0 & -3 & 3 & 0 \\ 2 & -1 & 1 & -2 \end{bmatrix}$$

$$\begin{aligned} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 \\ 1 & 4 & 9 & 16 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ -2 & 2 \\ 1 & 1 \\ -1 & 0 \end{bmatrix} &= \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \begin{bmatrix} 1 & -1 \end{bmatrix} + \begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix} \begin{bmatrix} -2 & 2 \end{bmatrix} + \begin{bmatrix} 1 \\ 3 \\ 9 \end{bmatrix} \begin{bmatrix} 1 & 1 \end{bmatrix} + \begin{bmatrix} 1 \\ 4 \\ 16 \end{bmatrix} \begin{bmatrix} -1 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 1 & -1 \\ 1 & -1 \\ 1 & -1 \end{bmatrix} + \begin{bmatrix} -2 & 2 \\ -4 & 4 \\ -8 & 8 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 3 & 3 \\ 9 & 9 \end{bmatrix} + \begin{bmatrix} -1 & 0 \\ -4 & 0 \\ -16 & 0 \end{bmatrix} \\ &= \begin{bmatrix} -1 & 2 \\ -4 & 6 \\ -14 & 16 \end{bmatrix} \end{aligned}$$