

ETC3550

Applied forecasting for business and economics

Ch3. The forecasters' toolbox

OTexts.org/fpp3/

Outline

- 1 A tidy forecasting workflow
- 2 Some simple forecasting methods
- 3 The workflow in action
- 4 Transformations
- 5 Distributional forecasts

Outline


- 1 A tidy forecasting workflow
- 2 Some simple forecasting methods
- 3 The workflow in action
- 4 Transformations
- 5 Distributional forecasts

A tidy forecasting workflow

The process of producing forecasts can be split up into a few fundamental steps.

- 1 Preparing data
- 2 Data visualisation
- 3 Specifying a model
- 4 Model estimation
- 5 Accuracy & performance evaluation
- 6 Producing forecasts


A tidy forecasting workflow



Outline

- 1 A tidy forecasting workflow
- 2 Some simple forecasting methods
- 3 The workflow in action
- 4 Transformations
- 5 Distributional forecasts

Some simple forecasting methods



How would you forecast these series?

Some simple forecasting methods



How would you forecast these series?

Some simple forecasting methods

Facebook closing stock price in 2018




How would you forecast these series?

Some simple forecasting methods

MEAN(y): Average method


- Forecast of all future values is equal to mean of historical data $\{y_1, \dots, y_T\}$.
- Forecasts: $\hat{y}_{T+h|T} = \bar{y} = (y_1 + \dots + y_T)/T$



Some simple forecasting methods

NAIVE(y): Naïve method


- Forecasts equal to last observed value.
- Forecasts: $\hat{y}_{T+h|T} = y_T$.
- Consequence of efficient market hypothesis.



Some simple forecasting methods

SNAIVE($y \sim \text{lag}(m)$): Seasonal naïve method

- Forecasts equal to last value from same season.
- Forecasts: $\hat{y}_{T+h|T} = y_{T+h-m(k+1)}$, where m = seasonal period and k is the integer part of $(h - 1)/m$.



Some simple forecasting methods

RW(y ~ drift()): Drift method


- Forecasts equal to last value plus average change.
- Forecasts:

$$\begin{aligned}\hat{y}_{T+h|T} &= y_T + \frac{h}{T-1} \sum_{t=2}^T (y_t - y_{t-1}) \\ &= y_T + \frac{h}{T-1} (y_T - y_1).\end{aligned}$$

- Equivalent to extrapolating a line drawn between first and last observations.

Some simple forecasting methods

Drift method




Outline

- 1 A tidy forecasting workflow
- 2 Some simple forecasting methods
- 3 The workflow in action
- 4 Transformations
- 5 Distributional forecasts

Data preparation and visualisation

```
# Set training data from 1992 to 2007  
train <- aus_production %>%  
  filter(between(year(Quarter), 1992, 2007))  
train %>% autoplot(Beer)
```



Model estimation

The `model()` function trains models to data.

```
# Fit the models
beer_fit <- train %>%
  model(
    Mean = MEAN(Beer),
    Naïve = NAIVE(Beer),
    Seasonal naïve = SNAIVE(Beer),
    Drift = RW(Beer ~ drift())
  )
```

Model estimation

```
beer_fit
```

```
## # A mable: 1 x 4
```

```
##   Mean                Naïve   Seasonal naïve Drift
```

```
##   <model>             <model> <model>             <model>
```

```
## 1 <MEAN,  $\mu=435.375$ > <NAIVE> <SNAIVE>             <RW w/ drift>
```

A mable is a model table, each cell corresponds to a fitted model.

Producing forecasts


```
beer_fc <- beer_fit %>%  
  forecast(h = 11)
```

```
## # A fable: 44 x 4 [1Q]  
## # Key:      .model [4]  
##   .model Quarter Beer .distribution  
##   <chr>      <qtr> <dbl> <dist>  
## 1 Mean      2008 Q1  435. N(435, 1964)  
## 2 Mean      2008 Q2  435. N(435, 1964)  
## 3 Mean      2008 Q3  435. N(435, 1964)  
## 4 Mean      2008 Q4  435. N(435, 1964)  
## # ... with 40 more rows
```

A fable is a forecast table with point forecasts and distributions.

Visualising forecasts

```
beer_fc %>%  
  autoplot(train, level = NULL) +  
  ggtitle("Forecasts for quarterly beer production") +  
  xlab("Year") + ylab("Megalitres") +  
  guides(colour=guide_legend(title="Forecast"))
```



Facebook closing stock price

```
# Extract training data
fb_stock <- gafa_stock %>%
  group_by(Symbol) %>%
  mutate(trading_day = row_number()) %>%
  update_tsibble(index=trading_day, regular=TRUE) %>%
  filter(Symbol == "FB",
         between(Date, ymd("2018-01-01"), ymd("2018-09-01")))

# Specify, estimate and forecast
fb_stock %>%
  model(
    Mean = MEAN(Close),
    Naïve = NAIVE(Close),
    Drift = RW(Close ~ drift())
  ) %>%
  forecast(h=42) %>%
  autoplot(fb_stock, level = NULL) +
  ggtitle("Facebook closing stock price (daily ending Sep 2018)") +
  xlab("Day") + ylab("") +
  guides(colour=guide_legend(title="Forecast"))
```

Facebook closing stock price



Your turn

- Produce forecasts from the appropriate method for Amazon closing price (`gafa_stock`) and Australian takeaway food turnover (`aus_retail`).
- Plot the results using `autoplot()`.

Outline

- 1 A tidy forecasting workflow
- 2 Some simple forecasting methods
- 3 The workflow in action
- 4 Transformations**
- 5 Distributional forecasts

Variance stabilization

If the data show different variation at different levels of the series, then a transformation can be useful.

Variance stabilization

If the data show different variation at different levels of the series, then a transformation can be useful.

Denote original observations as y_1, \dots, y_n and transformed observations as w_1, \dots, w_n .

Variance stabilization

If the data show different variation at different levels of the series, then a transformation can be useful.

Denote original observations as y_1, \dots, y_n and transformed observations as w_1, \dots, w_n .

Mathematical transformations for stabilizing variation

Square root	$w_t = \sqrt{y_t}$	↓
Cube root	$w_t = \sqrt[3]{y_t}$	Increasing
Logarithm	$w_t = \log(y_t)$	strength

Variance stabilization

If the data show different variation at different levels of the series, then a transformation can be useful.

Denote original observations as y_1, \dots, y_n and transformed observations as w_1, \dots, w_n .


Mathematical transformations for stabilizing variation

Square root	$w_t = \sqrt{y_t}$	↓
Cube root	$w_t = \sqrt[3]{y_t}$	Increasing
Logarithm	$w_t = \log(y_t)$	strength

Logarithms, in particular, are useful because they are more interpretable: changes in a log value are **relative (percent) changes on the original scale**.


Variance stabilization

```
food <- aus_retail %>%  
  filter(Industry == "Food retailing") %>%  
  summarise(Turnover = sum(Turnover))
```




Variance stabilization

```
food %>% autoplot(sqrt(Turnover)) +  
  labs(y = "Square root turnover")
```




Variance stabilization

```
food %>% autoplot(Turnover^(1/3)) +  
  labs(y = "Cube root turnover")
```




Variance stabilization

```
food %>% autoplot(log(Turnover)) +  
  labs(y = "Log turnover")
```



Variance stabilization

```
food %>% autoplot(-1/Turnover) +  
  labs(y = "Inverse turnover")
```



Box-Cox transformations

Each of these transformations is close to a member of the family of **Box-Cox transformations**:

$$w_t = \begin{cases} \log(y_t), & \lambda = 0; \\ (y_t^\lambda - 1)/\lambda, & \lambda \neq 0. \end{cases}$$

Box-Cox transformations

Each of these transformations is close to a member of the family of **Box-Cox transformations**:


$$w_t = \begin{cases} \log(y_t), & \lambda = 0; \\ (y_t^\lambda - 1)/\lambda, & \lambda \neq 0. \end{cases}$$

- $\lambda = 1$: (No substantive transformation)
- $\lambda = \frac{1}{2}$: (Square root plus linear transformation)
- $\lambda = 0$: (Natural logarithm)
- $\lambda = -1$: (Inverse plus 1)

Box-Cox transformations

Box-Cox transformations

```
food %>% autoplot(box_cox(Turnover, 1/3)) +  
  labs(y = "Box-Cox transformed turnover")
```



Box-Cox transformations

- y_t^λ for λ close to zero behaves like logs.
- If some $y_t = 0$, then must have $\lambda > 0$
- if some $y_t < 0$, no power transformation is possible unless all y_t adjusted by **adding a constant to all values**.
- Simple values of λ are easier to explain.
- Results are relatively insensitive to λ .
- Often no transformation ($\lambda = 1$) needed.
- Transformation can have very large effect on PI.
- Choosing $\lambda = 0$ is a simple way to force forecasts to be positive

Box-Cox transformations

```
food %>%
```

```
  features(Turnover, features = guerrero)
```

```
## # A tibble: 1 x 1
```

```
##   lambda_guerrero
```

```
##               <dbl>
```

```
## 1             0.00762
```

Box-Cox transformations

```
food %>%  
  features(Turnover, features = guerrero)
```

```
## # A tibble: 1 x 1  
##   lambda_guerrero  
##             <dbl>  
## 1             0.00762
```

- This attempts to balance the seasonal fluctuations and random variation across the series.
- Always check the results.
- A low value of λ can give extremely large prediction intervals.

Back-transformation

We must reverse the transformation (or *back-transform*) to obtain forecasts on the original scale. The reverse Box-Cox transformations are given by

$$y_t = \begin{cases} \exp(w_t), & \lambda = 0; \\ (\lambda W_t + 1)^{1/\lambda}, & \lambda \neq 0. \end{cases}$$

Modelling with transformations

Transformations used in the left of the formula will be automatically back-transformed. To model log-transformed food retailing turnover, you could use:

```
fit <- food %>%  
  model(SNAIVE(log(Turnover) ~ lag("year")))
```

```
## # A mable: 1 x 1  
##   SNAIVE(log(Turnover) ~ lag("year"))  
##   <model>  
## 1 <SNAIVE>
```


Forecasting with transformations

```
fc <- fit %>%  
  forecast(h = "3 years")
```

```
## # A tibble: 36 x 4 [1M]  
## # Key:   .model [1]  
##   .model          Month Turnover .distribution  
##   <chr>          <mth>      <dbl> <dist>  
## 1 "SNAIVE(log(Turnover) ~ 2018 Oct    10736. t(N(9.3, 0.004~  
## 2 "SNAIVE(log(Turnover) ~ 2018 Nov    10806. t(N(9.3, 0.004~  
## 3 "SNAIVE(log(Turnover) ~ 2018 Dec    12581. t(N(9.4, 0.004~  
## 4 "SNAIVE(log(Turnover) ~ 2019 Jan    10738. t(N(9.3, 0.004~  
## 5 "SNAIVE(log(Turnover) ~ 2019 Feb     9856. t(N(9.2, 0.004~  
## 6 "SNAIVE(log(Turnover) ~ 2019 Mar    11215. t(N(9.3, 0.004~  
## # ... with 30 more rows
```

Forecasting with transformations

```
fc %>% autoplot(filter(food, year(Month)>2010))
```



Your turn

Find a transformation that works for the Australian gas production (`aus_production`).

Bias adjustment

- Back-transformed point forecasts are medians.
- Back-transformed PI have the correct coverage.

Bias adjustment

- Back-transformed point forecasts are medians.
- Back-transformed PI have the correct coverage.

Back-transformed means

Let X be have mean μ and variance σ^2 .

Let $f(x)$ be back-transformation function, and $Y = f(X)$.

Taylor series expansion about μ :

$$f(X) = f(\mu) + (X - \mu)f'(\mu) + \frac{1}{2}(X - \mu)^2f''(\mu).$$

Bias adjustment

- Back-transformed point forecasts are medians.
- Back-transformed PI have the correct coverage.

Back-transformed means

Let X be have mean μ and variance σ^2 .

Let $f(x)$ be back-transformation function, and $Y = f(X)$.

Taylor series expansion about μ :

$$f(X) = f(\mu) + (X - \mu)f'(\mu) + \frac{1}{2}(X - \mu)^2f''(\mu).$$

$$E[Y] = E[f(X)] = f(\mu) + \frac{1}{2}\sigma^2f''(\mu)$$

Bias adjustment

Box-Cox back-transformation:

$$y_t = \begin{cases} \exp(w_t) & \lambda = 0; \\ (\lambda W_t + 1)^{1/\lambda} & \lambda \neq 0. \end{cases}$$

$$f(x) = \begin{cases} e^x & \lambda = 0; \\ (\lambda x + 1)^{1/\lambda} & \lambda \neq 0. \end{cases}$$

$$f''(x) = \begin{cases} e^x & \lambda = 0; \\ (1 - \lambda)(\lambda x + 1)^{1/\lambda - 2} & \lambda \neq 0. \end{cases}$$

Bias adjustment

Box-Cox back-transformation:

$$y_t = \begin{cases} \exp(w_t) & \lambda = 0; \\ (\lambda W_t + 1)^{1/\lambda} & \lambda \neq 0. \end{cases}$$


$$f(x) = \begin{cases} e^x & \lambda = 0; \\ (\lambda x + 1)^{1/\lambda} & \lambda \neq 0. \end{cases}$$

$$f''(x) = \begin{cases} e^x & \lambda = 0; \\ (1 - \lambda)(\lambda x + 1)^{1/\lambda - 2} & \lambda \neq 0. \end{cases}$$

$$E[Y] = \begin{cases} e^{\mu} \left[1 + \frac{\sigma^2}{2} \right] & \lambda = 0; \\ (\lambda \mu + 1)^{1/\lambda} \left[1 + \frac{\sigma^2(1-\lambda)}{2(\lambda \mu + 1)^2} \right] & \lambda \neq 0. \end{cases}$$

Bias adjustment

```
eggs <- as_tsibble(fma::eggs)
fit <- eggs %>% model(RW(log(value) ~ drift()))
fc <- fit %>% forecast(h=50)
fc_biased <- fit %>% forecast(h=50, bias_adjust = FALSE)
eggs %>% autoplot(value) +
  autolayer(fc_biased, series="Simple back transformation", level=80) +
  autolayer(fc, series="Bias adjusted", level=NULL) +
  guides(colour=guide_legend(title="Forecast"))
```



Outline

- 1 A tidy forecasting workflow
- 2 Some simple forecasting methods
- 3 The workflow in action
- 4 Transformations
- 5 Distributional forecasts

Forecast distributions

- A forecast $\hat{y}_{T+h|T}$ is (usually) the mean of the conditional distribution $y_{T+h} \mid y_1, \dots, y_T$.
- Most time series models produce normally distributed forecasts.
- The forecast distribution describes the probability of observing any future value.

Forecast distributions

Assuming residuals are normal, uncorrelated, $\text{sd} = \hat{\sigma}$:

Mean: $\hat{y}_{T+h|T} \sim N(\bar{y}, (1 + 1/T)\hat{\sigma}^2)$

Naïve: $\hat{y}_{T+h|T} \sim N(y_T, h\hat{\sigma}^2)$

Seasonal naïve: $\hat{y}_{T+h|T} \sim N(y_{T+h-m(k+1)}, (k+1)\hat{\sigma}^2)$

Drift: $\hat{y}_{T+h|T} \sim N(y_T + \frac{h}{T-1}(y_T - y_1), h\frac{T+h}{T}\hat{\sigma}^2)$

where k is the integer part of $(h - 1)/m$.

Note that when $h = 1$ and T is large, these all give the same approximate forecast variance: $\hat{\sigma}^2$.

Prediction intervals

- A prediction interval gives a region within which we expect y_{T+h} to lie with a specified probability.
- Assuming forecast errors are normally distributed, then a 95% PI is

$$\hat{y}_{T+h|T} \pm 1.96\hat{\sigma}_h$$

where $\hat{\sigma}_h$ is the st dev of the h -step distribution.

- When $h = 1$, $\hat{\sigma}_h$ can be estimated from the residuals.

Prediction intervals

```
fit <- fb_stock %>% model(NAIVE(Close))  
forecast(fit)
```

```
## # A fable: 2 x 5 [1]  
## # Key:      Symbol, .model [1]  
##   Symbol .model   trading_day Close .distribution  
##   <fct>  <chr>          <int>  <dbl> <dist>  
## 1 FB      NAIVE(Cl~      3693   176. N(176, 21)  
## 2 FB      NAIVE(Cl~      3694   176. N(176, 42)
```


Prediction intervals

```
res_sd <- sqrt(mean(augment(fit)$resid^2, na.rm = TRUE))  
last(fb_stock$Close) + 1.96 * res_sd * c(-1,1)
```

```
## [1] 166.7196 184.7404
```

```
forecast(fit, h = 1) %>%  
  transmute(interval = hilo(.distribution, level = 95))
```

```
## # A tsibble: 1 x 4 [?]  
## # Key:      Symbol, .model [1]  
##   Symbol .model      trading_day      interval  
##   <fct>  <chr>          <int>          <hilo>  
## 1 FB     NAIVE(Close)      3693 [166.7198, 184.7402]95
```

Prediction intervals

- Point forecasts are often useless without a measure of uncertainty (such as prediction intervals).
- Prediction intervals require a stochastic model (with random errors, etc).
- Multi-step forecasts for time series require a more sophisticated approach (with PI getting wider as the forecast horizon increases).

Prediction intervals

- Computed automatically from the forecast distribution.
- Use `level` argument to control coverage.
- Check residual assumptions before believing them (we will see this next class).
- Usually too narrow due to unaccounted uncertainty.