



ETC3550: Applied forecasting for business and economics

Ch12. Some practical forecasting
issues

OTexts.org/fpp2/

Outline

- 1 Models for different frequencies
- 2 Ensuring forecasts stay within limits
- 3 Forecast combinations
- 4 Missing values
- 5 Outliers

Models for different frequencies

Models for annual data

- ETS, ARIMA, Dynamic regression

Models for different frequencies

Models for annual data

- ETS, ARIMA, Dynamic regression

Models for quarterly data

- ETS, ARIMA/SARIMA, Dynamic regression, Dynamic harmonic regression, STL+ETS, STL+ARIMA

Models for different frequencies

Models for annual data

- ETS, ARIMA, Dynamic regression

Models for quarterly data

- ETS, ARIMA/SARIMA, Dynamic regression, Dynamic harmonic regression, STL+ETS, STL+ARIMA

Models for monthly data

- ETS, ARIMA/SARIMA, Dynamic regression, Dynamic harmonic regression, STL+ETS, STL+ARIMA

Models for different frequencies

Models for weekly data

- ARIMA/SARIMA, Dynamic regression, Dynamic harmonic regression, STL+ETS, STL+ARIMA, TBATS

Models for different frequencies

Models for weekly data

- ARIMA/SARIMA, Dynamic regression, Dynamic harmonic regression, STL+ETS, STL+ARIMA, TBATS


Models for daily, hourly and other sub-daily data

- ARIMA/SARIMA, Dynamic regression, Dynamic harmonic regression, STL+ETS, STL+ARIMA, TBATS

Outline

- 1 Models for different frequencies
- 2 Ensuring forecasts stay within limits
- 3 Forecast combinations
- 4 Missing values
- 5 Outliers

Positive forecasts




Outline

- 1 Models for different frequencies
- 2 Ensuring forecasts stay within limits
- 3 Forecast combinations
- 4 Missing values
- 5 Outliers

Forecast combinations

The results have been virtually unanimous: combining multiple forecasts leads to increased forecast accuracy. ... In many cases one can make dramatic performance improvements by simply averaging the forecasts. (Clemen 1989)



Outline

- 1 Models for different frequencies
- 2 Ensuring forecasts stay within limits
- 3 Forecast combinations
- 4 **Missing values**
- 5 Outliers

Missing values

Functions which can handle missing values *

`auto.arima()`, `Arima()` * `tslm()` * `nnetar()`


Models which cannot handle missing values *

`ets()` * `stl()` * `stlf()` * `tbats()`

What to do?

- 1 Model section of data after last missing value.
- 2 Estimate missing values with `na.interp()`.


Missing values



Outline

- 1 Models for different frequencies
- 2 Ensuring forecasts stay within limits
- 3 Forecast combinations
- 4 Missing values
- 5 Outliers

Outliers



```
## $index
```

```
## [1] 770
```

```
##
```

```
## $replacements
```