



ETC3550

**Applied forecasting for
business and economics**

Ch9. ARIMA models

OTexts.org/fpp3/

Outline

- 1 Stationarity and differencing
- 2 Non-seasonal ARIMA models
- 3 Estimation and order selection
- 4 ARIMA modelling in R
- 5 Forecasting
- 6 Seasonal ARIMA models
- 7 ARIMA vs ETS

Outline

- 1 Stationarity and differencing
- 2 Non-seasonal ARIMA models
- 3 Estimation and order selection
- 4 ARIMA modelling in R
- 5 Forecasting
- 6 Seasonal ARIMA models
- 7 ARIMA vs ETS

Stationarity

Definition

If $\{y_t\}$ is a stationary time series, then for all s , the distribution of (y_t, \dots, y_{t+s}) does not depend on t .

Stationarity


Definition

If $\{y_t\}$ is a stationary time series, then for all s , the distribution of (y_t, \dots, y_{t+s}) does not depend on t .


A **stationary series** is:

- roughly horizontal
- constant variance
- no patterns predictable in the long-term

Stationary?




Stationary?




Stationary?



Stationary?




Stationary?



Stationary?



Stationary?



Stationary?



Stationarity

Definition

If $\{y_t\}$ is a stationary time series, then for all s , the distribution of (y_t, \dots, y_{t+s}) does not depend on t .

Stationarity

Definition

If $\{y_t\}$ is a stationary time series, then for all s , the distribution of (y_t, \dots, y_{t+s}) does not depend on t .

Transformations help to **stabilize the variance**.


For ARIMA modelling, we also need to **stabilize the mean**.

Non-stationarity in the mean


Identifying non-stationary series

- time plot.
- The ACF of stationary data drops to zero relatively quickly
- The ACF of non-stationary data decreases slowly.
- For non-stationary data, the value of r_1 is often large and positive.


Example: Google stock price




Example: Google stock price



Example: Google stock price



Example: Google stock price



Differencing

- Differencing helps to **stabilize the mean**.
- The differenced series is the *change* between each observation in the original series:
$$y'_t = y_t - y_{t-1}.$$
- The differenced series will have only $T - 1$ values since it is not possible to calculate a difference y'_1 for the first observation.

Second-order differencing

Occasionally the differenced data will not appear stationary and it may be necessary to difference the data a second time:

Second-order differencing

Occasionally the differenced data will not appear stationary and it may be necessary to difference the data a second time:

$$\begin{aligned}y_t'' &= y_t' - y_{t-1}' \\&= (y_t - y_{t-1}) - (y_{t-1} - y_{t-2}) \\&= y_t - 2y_{t-1} + y_{t-2}.\end{aligned}$$

Second-order differencing

Occasionally the differenced data will not appear stationary and it may be necessary to difference the data a second time:

$$\begin{aligned}y_t'' &= y_t' - y_{t-1}' \\&= (y_t - y_{t-1}) - (y_{t-1} - y_{t-2}) \\&= y_t - 2y_{t-1} + y_{t-2}.\end{aligned}$$

- y_t'' will have $T - 2$ values.
- In practice, it is almost never necessary to go beyond second-order differences.

Seasonal differencing

A seasonal difference is the difference between an observation and the corresponding observation from the previous year.

Seasonal differencing

A seasonal difference is the difference between an observation and the corresponding observation from the previous year.

$$y'_t = y_t - y_{t-m}$$

where m = number of seasons.

Seasonal differencing

A seasonal difference is the difference between an observation and the corresponding observation from the previous year.


$$y'_t = y_t - y_{t-m}$$

where m = number of seasons.

- For monthly data $m = 12$.
- For quarterly data $m = 4$.


Electricity production

```
usmelec %>% autoplot(  
  Generation  
)
```




Electricity production

```
usmelec %>% autoplot(  
  log(Generation)  
)
```




Electricity production

```
usmelec %>% autoplot(  
  log(Generation) %>% difference(12)  
)
```



Electricity production

```
usmelec %>% autoplot(  
  log(Generation) %>% difference(12) %>% difference()  
)
```



Electricity production

- Seasonally differenced series is closer to being stationary.
- Remaining non-stationarity can be removed with further first difference.

If $y'_t = y_t - y_{t-12}$ denotes seasonally differenced series, then twice-differenced series is

$$\begin{aligned}y_t^* &= y'_t - y'_{t-1} \\&= (y_t - y_{t-12}) - (y_{t-1} - y_{t-13}) \\&= y_t - y_{t-1} - y_{t-12} + y_{t-13} .\end{aligned}$$

Seasonal differencing

When both seasonal and first differences are applied...

Seasonal differencing

When both seasonal and first differences are applied...

- it makes no difference which is done first—the result will be the same.
- If seasonality is strong, we recommend that seasonal differencing be done first because sometimes the resulting series will be stationary and there will be no need for further first difference.

Seasonal differencing

When both seasonal and first differences are applied...

- it makes no difference which is done first—the result will be the same.
- If seasonality is strong, we recommend that seasonal differencing be done first because sometimes the resulting series will be stationary and there will be no need for further first difference.

It is important that if differencing is used, the differences are interpretable.

Interpretation of differencing

- first differences are the change between **one observation and the next**;
- seasonal differences are the change between **one year to the next**.

Interpretation of differencing

- first differences are the change between **one observation and the next**;
- seasonal differences are the change between **one year to the next**.

But taking lag 3 differences for yearly data, for example, results in a model which cannot be sensibly interpreted.

Unit root tests

Statistical tests to determine the required order of differencing.

- 1 Augmented Dickey Fuller test: null hypothesis is that the data are non-stationary and non-seasonal.
- 2 Kwiatkowski-Phillips-Schmidt-Shin (KPSS) test: null hypothesis is that the data are stationary and non-seasonal.
- 3 Other tests available for seasonal data.

KPSS test

```
google_2018 %>%  
  features(Close, unitroot_kpss)
```

```
## # A tibble: 1 x 3  
##   Symbol kpss_stat kpss_pval  
##   <chr>      <dbl>      <dbl>  
## 1 G00G      0.573      0.0252
```

KPSS test

```
google_2018 %>%  
  features(Close, unitroot_kpss)
```

```
## # A tibble: 1 x 3  
##   Symbol kpss_stat kpss_pval  
##   <chr>      <dbl>      <dbl>  
## 1 GOOG      0.573      0.0252
```

```
google_2018 %>%  
  features(Close, unitroot_ndiffs)
```

```
## # A tibble: 1 x 2  
##   Symbol ndiffs  
##   <chr>    <int>  
## 1 GOOG      1
```


Automatically selecting differences

STL decomposition: $y_t = T_t + S_t + R_t$

Seasonal strength $F_s = \max\left(0, 1 - \frac{\text{Var}(R_t)}{\text{Var}(S_t + R_t)}\right)$

If $F_s > 0.64$, do one seasonal difference.

```
usmelec %>% mutate(log_gen = log(Generation)) %>%  
  features(log_gen, list(unitroot_nsdiffs, stl_features))
```

```
## # A tibble: 1 x 8  
##   nsdiffs trend_strength seasonal_streng~ spike linearity  
##   <int>         <dbl>         <dbl>    <dbl>    <dbl>  
## 1         1         0.994         0.941 1.54e-12    5.65  
## # ... with 3 more variables: curvature <dbl>,  
## #   seasonal_peak_year <dbl>, seasonal_trough_year <dbl>
```

Automatically selecting differences

```
usmelec %>% mutate(log_gen = log(Generation)) %>%  
  features(log_gen, unitroot_nsdiffs)
```

```
## # A tibble: 1 x 1  
##   nsdiffs  
##   <int>  
## 1      1
```

```
usmelec %>% mutate(d_log_gen = difference(log(Generation), 12)) %>%  
  features(d_log_gen, unitroot_nsdiffs)
```

```
## # A tibble: 1 x 1  
##   nsdiffs  
##   <int>  
## 1      1
```

Your turn

For the `tourism` dataset, compute the total number of trips and find an appropriate differencing (after transformation if necessary) to obtain stationary data.

Backshift notation

A very useful notational device is the backward shift operator, B , which is used as follows:

$$By_t = y_{t-1}$$

Backshift notation

A very useful notational device is the backward shift operator, B , which is used as follows:

$$By_t = y_{t-1}$$

In other words, B , operating on y_t , has the effect of **shifting the data back one period.**

Backshift notation

A very useful notational device is the backward shift operator, B , which is used as follows:

$$By_t = y_{t-1}$$

In other words, B , operating on y_t , has the effect of **shifting the data back one period**. Two applications of B to y_t **shifts the data back two periods**:

$$B(By_t) = B^2y_t = y_{t-2}$$

Backshift notation

A very useful notational device is the backward shift operator, B , which is used as follows:

$$By_t = y_{t-1}$$

In other words, B , operating on y_t , has the effect of **shifting the data back one period**. Two applications of B to y_t **shifts the data back two periods**:

$$B(By_t) = B^2y_t = y_{t-2}$$

For monthly data, if we wish to shift attention to “the same month last year”, then B^{12} is used, and the notation is $B^{12}y_t = y_{t-12}$.

Backshift notation

The backward shift operator is convenient for describing the process of *differencing*.

Backshift notation

The backward shift operator is convenient for describing the process of *differencing*. A first difference can be written as

$$y'_t = y_t - y_{t-1} = y_t - By_t = (1 - B)y_t$$

Backshift notation

The backward shift operator is convenient for describing the process of *differencing*. A first difference can be written as

$$y'_t = y_t - y_{t-1} = y_t - By_t = (1 - B)y_t$$

Note that a first difference is represented by $(1 - B)$.

Backshift notation

The backward shift operator is convenient for describing the process of *differencing*. A first difference can be written as

$$y'_t = y_t - y_{t-1} = y_t - By_t = (1 - B)y_t$$

Note that a first difference is represented by $(1 - B)$.

Similarly, if second-order differences (i.e., first differences of first differences) have to be computed, then:

$$y''_t = y_t - 2y_{t-1} + y_{t-2} = (1 - B)^2 y_t$$

Backshift notation

- Second-order difference is denoted $(1 - B)^2$.
- *Second-order difference* is not the same as a *second difference*, which would be denoted $1 - B^2$;
- In general, a d th-order difference can be written as

$$(1 - B)^d y_t$$

- A seasonal difference followed by a first difference can be written as

$$(1 - B)(1 - B^m)y_t$$

Backshift notation

The “backshift” notation is convenient because the terms can be multiplied together to see the combined effect.

$$\begin{aligned}(1 - B)(1 - B^m)y_t &= (1 - B - B^m + B^{m+1})y_t \\ &= y_t - y_{t-1} - y_{t-m} + y_{t-m-1}.\end{aligned}$$

Backshift notation

The “backshift” notation is convenient because the terms can be multiplied together to see the combined effect.

$$\begin{aligned}(1 - B)(1 - B^m)y_t &= (1 - B - B^m + B^{m+1})y_t \\ &= y_t - y_{t-1} - y_{t-m} + y_{t-m-1}.\end{aligned}$$

For monthly data, $m = 12$ and we obtain the same result as earlier.

Outline

- 1 Stationarity and differencing
- 2 Non-seasonal ARIMA models
- 3 Estimation and order selection
- 4 ARIMA modelling in R
- 5 Forecasting
- 6 Seasonal ARIMA models
- 7 ARIMA vs ETS


Autoregressive models

Autoregressive (AR) models:


$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \cdots + \phi_p y_{t-p} + \varepsilon_t,$$

where ε_t is white noise. This is a multiple regression with **lagged values** of y_t as predictors.

AR(1)




AR(2)



AR(1) model

$$y_t = 2 - 0.8y_{t-1} + \varepsilon_t$$

$$\varepsilon_t \sim N(0, 1), \quad T = 100.$$



AR(1) model


$$y_t = c + \phi_1 y_{t-1} + \varepsilon_t$$

- When $\phi_1 = 0$, y_t is **equivalent to WN**
- When $\phi_1 = 1$ and $c = 0$, y_t is **equivalent to a RW**
- When $\phi_1 = 1$ and $c \neq 0$, y_t is **equivalent to a RW with drift**
- When $\phi_1 < 0$, y_t tends to **oscillate between positive and negative values.**

AR(2) model

$$y_t = 8 + 1.3y_{t-1} - 0.7y_{t-2} + \varepsilon_t$$

$$\varepsilon_t \sim N(0, 1), \quad T = 100.$$



Stationarity conditions

We normally restrict autoregressive models to stationary data, and then some constraints on the values of the parameters are required.

General condition for stationarity

Complex roots of $1 - \phi_1 z - \phi_2 z^2 - \dots - \phi_p z^p$ lie outside the unit circle on the complex plane.

Stationarity conditions

We normally restrict autoregressive models to stationary data, and then some constraints on the values of the parameters are required.

General condition for stationarity

Complex roots of $1 - \phi_1 z - \phi_2 z^2 - \dots - \phi_p z^p$ lie outside the unit circle on the complex plane.



- For $p = 1$: $-1 < \phi_1 < 1$.
- For $p = 2$:
 $-1 < \phi_2 < 1$ $\phi_2 + \phi_1 < 1$ $\phi_2 - \phi_1 < 1$.
- More complicated conditions hold for $p \geq 3$.
- Estimation software takes care of this.

Moving Average (MA) models

Moving Average (MA) models:

$$y_t = c + \varepsilon_t + \theta_1 \varepsilon_{t-1} + \theta_2 \varepsilon_{t-2} + \cdots + \theta_q \varepsilon_{t-q},$$


where ε_t is white noise. This is a multiple regression with **past errors** as predictors. *Don't confuse this with moving average smoothing!*



MA(1) model

$$y_t = 20 + \varepsilon_t + 0.8\varepsilon_{t-1}$$


$$\varepsilon_t \sim N(0, 1), \quad T = 100.$$



MA(2) model

$$y_t = \varepsilon_t - \varepsilon_{t-1} + 0.8\varepsilon_{t-2}$$

$$\varepsilon_t \sim N(0, 1), \quad T = 100.$$



MA(∞) models

It is possible to write any stationary AR(p) process as an MA(∞) process.

Example: AR(1)

$$\begin{aligned}y_t &= \phi_1 y_{t-1} + \varepsilon_t \\&= \phi_1(\phi_1 y_{t-2} + \varepsilon_{t-1}) + \varepsilon_t \\&= \phi_1^2 y_{t-2} + \phi_1 \varepsilon_{t-1} + \varepsilon_t \\&= \phi_1^3 y_{t-3} + \phi_1^2 \varepsilon_{t-2} + \phi_1 \varepsilon_{t-1} + \varepsilon_t \\&\dots\end{aligned}$$

MA(∞) models

It is possible to write any stationary AR(p) process as an MA(∞) process.

Example: AR(1)

$$\begin{aligned}y_t &= \phi_1 y_{t-1} + \varepsilon_t \\&= \phi_1(\phi_1 y_{t-2} + \varepsilon_{t-1}) + \varepsilon_t \\&= \phi_1^2 y_{t-2} + \phi_1 \varepsilon_{t-1} + \varepsilon_t \\&= \phi_1^3 y_{t-3} + \phi_1^2 \varepsilon_{t-2} + \phi_1 \varepsilon_{t-1} + \varepsilon_t \\&\dots\end{aligned}$$

Provided $-1 < \phi_1 < 1$:

$$y_t = \varepsilon_t + \phi_1 \varepsilon_{t-1} + \phi_1^2 \varepsilon_{t-2} + \phi_1^3 \varepsilon_{t-3} + \dots$$

Invertibility

- Any $MA(q)$ process can be written as an $AR(\infty)$ process if we impose some constraints on the MA parameters.
- Then the MA model is called “invertible”.
- Invertible models have some mathematical properties that make them easier to use in practice.
- Invertibility of an ARIMA model is equivalent to forecastability of an ETS model.

Invertibility

General condition for invertibility

Complex roots of $1 + \theta_1 z + \theta_2 z^2 + \dots + \theta_q z^q$ lie outside the unit circle on the complex plane.

Invertibility

General condition for invertibility

Complex roots of $1 + \theta_1 z + \theta_2 z^2 + \dots + \theta_q z^q$ lie outside the unit circle on the complex plane.

- For $q = 1$: $-1 < \theta_1 < 1$.
- For $q = 2$:
 $-1 < \theta_2 < 1 \quad \theta_2 + \theta_1 > -1 \quad \theta_1 - \theta_2 < 1$.
- More complicated conditions hold for $q \geq 3$.
- Estimation software takes care of this.

ARIMA models

Autoregressive Moving Average models:

$$y_t = c + \phi_1 y_{t-1} + \cdots + \phi_p y_{t-p} \\ + \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q} + \varepsilon_t.$$

ARIMA models

Autoregressive Moving Average models:

$$y_t = c + \phi_1 y_{t-1} + \cdots + \phi_p y_{t-p} \\ + \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q} + \varepsilon_t.$$

- Predictors include both **lagged values of y_t and lagged errors.**
- Conditions on coefficients ensure stationarity.
- Conditions on coefficients ensure invertibility.

ARIMA models

Autoregressive Moving Average models:

$$y_t = c + \phi_1 y_{t-1} + \cdots + \phi_p y_{t-p} \\ + \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q} + \varepsilon_t.$$

- Predictors include both **lagged values of y_t and lagged errors.**
- Conditions on coefficients ensure stationarity.
- Conditions on coefficients ensure invertibility.

Autoregressive Integrated Moving Average models

- Combine ARMA model with **differencing.**
- $(1 - B)^d y_t$ follows an ARMA model.

Autoregressive Integrated Moving Average models

ARIMA(p, d, q) model

AR: p = order of the autoregressive part

I: d = degree of first differencing involved

MA: q = order of the moving average part.

- White noise model: ARIMA(0,0,0)
- Random walk: ARIMA(0,1,0) with no constant
- Random walk with drift: ARIMA(0,1,0) with const.
- AR(p): ARIMA($p,0,0$)
- MA(q): ARIMA(0,0, q)

Backshift notation for ARIMA

■ ARMA model:

$$y_t = c + \phi_1 B y_t + \dots + \phi_p B^p y_t + \varepsilon_t + \theta_1 B \varepsilon_t + \dots + \theta_q B^q \varepsilon_t$$

or $(1 - \phi_1 B - \dots - \phi_p B^p) y_t = c + (1 + \theta_1 B + \dots + \theta_q B^q) \varepsilon_t$

■ ARIMA(1,1,1) model:

$$\begin{array}{ccccc} (1 - \phi_1 B) & (1 - B) y_t & = & c + (1 + \theta_1 B) \varepsilon_t \\ \uparrow & \uparrow & & \uparrow \\ \text{AR}(1) & \text{First} & & \text{MA}(1) \\ & \text{difference} & & \end{array}$$

Backshift notation for ARIMA

■ ARMA model:

$$y_t = c + \phi_1 B y_t + \dots + \phi_p B^p y_t + \varepsilon_t + \theta_1 B \varepsilon_t + \dots + \theta_q B^q \varepsilon_t$$

or $(1 - \phi_1 B - \dots - \phi_p B^p) y_t = c + (1 + \theta_1 B + \dots + \theta_q B^q) \varepsilon_t$

■ ARIMA(1,1,1) model:

$$\begin{array}{ccccc} (1 - \phi_1 B) & (1 - B) y_t & = & c + (1 + \theta_1 B) \varepsilon_t \\ \uparrow & \uparrow & & \uparrow \\ \text{AR}(1) & \text{First} & & \text{MA}(1) \\ & \text{difference} & & \end{array}$$

Written out:

$$y_t = c + y_{t-1} + \phi_1 y_{t-1} - \phi_1 y_{t-2} + \theta_1 \varepsilon_{t-1} + \varepsilon_t$$

R model

Intercept form

$$(1 - \phi_1 B - \dots - \phi_p B^p) y'_t = c + (1 + \theta_1 B + \dots + \theta_q B^q) \varepsilon_t$$


Mean form

$$(1 - \phi_1 B - \dots - \phi_p B^p)(y'_t - \mu) = (1 + \theta_1 B + \dots + \theta_q B^q) \varepsilon_t$$

- $y'_t = (1 - B)^d y_t$
- μ is the mean of y'_t .
- $c = \mu(1 - \phi_1 - \dots - \phi_p)$.
- R uses mean form
- fable uses intercept form

Australian household expenditure

```
us_change <- read_csv(  
  "https://otexts.com/fpp3/extrfiles/us_change.csv") %>%  
  mutate(Time = yearquarter(Time)) %>%  
  as_tsibble(index = Time)
```



US personal consumption

```
fit <- us_change %>% model(arima = ARIMA(Consumption ~ PDQ(0,0,0)))  
report(fit)
```

```
## Series: Consumption  
## Model: ARIMA(1,0,3) w/ mean  
##  
## Coefficients:  
##          ar1          ma1          ma2          ma3    constant  
##          0.5885   -0.3528   0.0846   0.1739       0.3067  
## s.e.    0.1541    0.1658   0.0818   0.0843       0.0383  
##  
## sigma^2 estimated as 0.3499:  log likelihood=-164.8  
## AIC=341.6   AICc=342.1   BIC=361
```

US personal consumption

```
fit <- us_change %>% model(arima = ARIMA(Consumption ~ PDQ(0,0,0)))  
report(fit)
```

```
## Series: Consumption  
## Model: ARIMA(1,0,3) w/ mean  
##  
## Coefficients:  
##          ar1          ma1          ma2          ma3    constant  
##          0.5885   -0.3528   0.0846   0.1739     0.3067  
## s.e.    0.1541    0.1658   0.0818   0.0843     0.0383  
##  
## sigma^2 estimated as 0.3499:  log likelihood=-164.8  
## AIC=341.6   AICc=342.1   BIC=361
```


ARIMA(1,0,3) model:

$$y_t = 0.307 + 0.589y_{t-1} - 0.353\varepsilon_{t-1} + 0.0846\varepsilon_{t-2} + 0.174\varepsilon_{t-3} + \varepsilon_t,$$

where ε_t is white noise with a standard deviation of $0.592 = \sqrt{0.350}$.

US personal consumption

```
fit %>% forecast(h=10) %>%  
  autoplot(slice(us_change, (n()-80):n()))
```



Understanding ARIMA models

- If $c = 0$ and $d = 0$, the long-term forecasts will go to zero.
- If $c = 0$ and $d = 1$, the long-term forecasts will go to a non-zero constant.
- If $c = 0$ and $d = 2$, the long-term forecasts will follow a straight line.
- If $c \neq 0$ and $d = 0$, the long-term forecasts will go to the mean of the data.
- If $c \neq 0$ and $d = 1$, the long-term forecasts will follow a straight line.
- If $c \neq 0$ and $d = 2$, the long-term forecasts will follow a quadratic trend.

Understanding ARIMA models

Forecast variance and d

- The higher the value of d , the more rapidly the prediction intervals increase in size.
- For $d = 0$, the long-term forecast standard deviation will go to the standard deviation of the historical data.

Cyclic behaviour

- For cyclic forecasts, $p \geq 2$ and some restrictions on coefficients are required.
- If $p = 2$, we need $\phi_1^2 + 4\phi_2 < 0$. Then average cycle of length

$$(2\pi) / \left[\arccos(-\phi_1(1 - \phi_2)/(4\phi_2)) \right] .$$

Outline

- 1 Stationarity and differencing
- 2 Non-seasonal ARIMA models
- 3 Estimation and order selection
- 4 ARIMA modelling in R
- 5 Forecasting
- 6 Seasonal ARIMA models
- 7 ARIMA vs ETS

Maximum likelihood estimation

Having identified the model order, we need to estimate the parameters $c, \phi_1, \dots, \phi_p, \theta_1, \dots, \theta_q$.

Maximum likelihood estimation

Having identified the model order, we need to estimate the parameters $c, \phi_1, \dots, \phi_p, \theta_1, \dots, \theta_q$.

- MLE is very similar to least squares estimation obtained by minimizing

$$\sum_{t=1}^T e_t^2$$

- The `ARIMA()` model allows CLS or MLE estimation.
- Non-linear optimization must be used in either case.
- Different software will give different estimates.

Partial autocorrelations

Partial autocorrelations measure relationship between y_t and y_{t-k} , when the effects of other time lags — $1, 2, 3, \dots, k - 1$ — are removed.

Partial autocorrelations

Partial autocorrelations measure relationship between y_t and y_{t-k} , when the effects of other time lags — $1, 2, 3, \dots, k-1$ — are removed.

α_k = k th partial autocorrelation coefficient
= equal to the estimate of ϕ_k in regression:

$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \dots + \phi_k y_{t-k}.$$

Partial autocorrelations

Partial autocorrelations measure relationship between y_t and y_{t-k} , when the effects of other time lags — $1, 2, 3, \dots, k-1$ — are removed.

α_k = k th partial autocorrelation coefficient
= equal to the estimate of ϕ_k in regression:



$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \dots + \phi_k y_{t-k}.$$

- Varying number of terms on RHS gives α_k for different values of k .
- There are more efficient ways of calculating α_k .
- $\alpha_1 = \rho_1$
- same critical values of $\pm 1.96/\sqrt{T}$ as for ACF.

Example: Mink trapping




Example: Mink trapping



Example: Mink trapping

```
mink %>% gg_tsdisplay(value)
```



ACF and PACF interpretation

AR(1)

$$\begin{aligned}\rho_k &= \phi_1^k && \text{for } k = 1, 2, \dots; \\ \alpha_1 &= \phi_1 && \alpha_k = 0 \quad \text{for } k = 2, 3, \dots\end{aligned}$$

So we have an AR(1) model when

- autocorrelations exponentially decay
- there is a single significant partial autocorrelation.

ACF and PACF interpretation

AR(p)

- ACF dies out in an exponential or damped sine-wave manner
- PACF has all zero spikes beyond the p th spike

So we have an AR(p) model when

- the ACF is exponentially decaying or sinusoidal
- there is a significant spike at lag p in PACF, but none beyond p

ACF and PACF interpretation

MA(1)

$$\begin{aligned}\rho_1 &= \theta_1 & \rho_k &= 0 & \text{for } k = 2, 3, \dots; \\ \alpha_k &= -(-\theta_1)^k\end{aligned}$$

So we have an MA(1) model when

- the PACF is exponentially decaying and
- there is a single significant spike in ACF

ACF and PACF interpretation

MA(q)

- PACF dies out in an exponential or damped sine-wave manner
- ACF has all zero spikes beyond the q th spike

So we have an MA(q) model when

- the PACF is exponentially decaying or sinusoidal
- there is a significant spike at lag q in ACF, but none beyond q

Information criteria

Akaike's Information Criterion (AIC):

$$\text{AIC} = -2 \log(L) + 2(p + q + k + 1),$$

where L is the likelihood of the data,

$k = 1$ if $c \neq 0$ and $k = 0$ if $c = 0$.

Information criteria

Akaike's Information Criterion (AIC):

$$\text{AIC} = -2 \log(L) + 2(p + q + k + 1),$$

where L is the likelihood of the data,

$k = 1$ if $c \neq 0$ and $k = 0$ if $c = 0$.

Corrected AIC:

$$\text{AICc} = \text{AIC} + \frac{2(p+q+k+1)(p+q+k+2)}{T-p-q-k-2}.$$

Information criteria

Akaike's Information Criterion (AIC):

$$\text{AIC} = -2 \log(L) + 2(p + q + k + 1),$$

where L is the likelihood of the data,

$k = 1$ if $c \neq 0$ and $k = 0$ if $c = 0$.

Corrected AIC:

$$\text{AICc} = \text{AIC} + \frac{2(p+q+k+1)(p+q+k+2)}{T-p-q-k-2}.$$

Bayesian Information Criterion:

$$\text{BIC} = \text{AIC} + [\log(T) - 2](p + q + k - 1).$$

Information criteria

Akaike's Information Criterion (AIC):

$$\text{AIC} = -2 \log(L) + 2(p + q + k + 1),$$

where L is the likelihood of the data,

$k = 1$ if $c \neq 0$ and $k = 0$ if $c = 0$.

Corrected AIC:

$$\text{AICc} = \text{AIC} + \frac{2(p+q+k+1)(p+q+k+2)}{T-p-q-k-2}.$$

Bayesian Information Criterion:

$$\text{BIC} = \text{AIC} + [\log(T) - 2](p + q + k - 1).$$

Good models are obtained by minimizing either the AIC, AICc or BIC. Our preference is to use the AICc.

Outline

- 1 Stationarity and differencing
- 2 Non-seasonal ARIMA models
- 3 Estimation and order selection
- 4 ARIMA modelling in R**
- 5 Forecasting
- 6 Seasonal ARIMA models
- 7 ARIMA vs ETS

How does ARIMA() work?

A non-seasonal ARIMA process

$$\phi(B)(1 - B)^d y_t = c + \theta(B)\varepsilon_t$$

Need to select appropriate orders: p, q, d

Hyndman and Khandakar (JSS, 2008) algorithm:

- Select no. differences d and D via KPSS test and seasonal strength measure.
- Select p, q by minimising AICc.
- Use stepwise search to traverse model space.

How does ARIMA() work?

$$\text{AICc} = -2 \log(L) + 2(p + q + k + 1) \left[1 + \frac{(p+q+k+2)}{T-p-q-k-2} \right] .$$

where L is the maximised likelihood fitted to the *differenced* data,
 $k = 1$ if $c \neq 0$ and $k = 0$ otherwise.

How does ARIMA() work?

$$\text{AICc} = -2 \log(L) + 2(p + q + k + 1) \left[1 + \frac{(p+q+k+2)}{T-p-q-k-2} \right].$$

where L is the maximised likelihood fitted to the *differenced* data,
 $k = 1$ if $c \neq 0$ and $k = 0$ otherwise.

Step1: Select current model (with smallest AICc) from:

ARIMA(2, d , 2)

ARIMA(0, d , 0)

ARIMA(1, d , 0)

ARIMA(0, d , 1)

How does ARIMA() work?

$$\text{AICc} = -2 \log(L) + 2(p + q + k + 1) \left[1 + \frac{(p+q+k+2)}{T-p-q-k-2} \right].$$

where L is the maximised likelihood fitted to the *differenced* data,
 $k = 1$ if $c \neq 0$ and $k = 0$ otherwise.

Step1: Select current model (with smallest AICc) from:

ARIMA(2, d , 2)

ARIMA(0, d , 0)

ARIMA(1, d , 0)

ARIMA(0, d , 1)

Step 2: Consider variations of current model:


- vary one of p , q , from current model by ± 1 ;
- p , q both vary from current model by ± 1 ;
- Include/exclude c from current model.

Model with lowest AICc becomes current model.

Repeat Step 2 until no lower AICc can be found.


Choosing your own model

```
web_usage <- as_tsibble(WWWusage)
web_usage %>% gg_tsdisplay(value)
```



Choosing your own model

```
web_usage %>% gg_tsdisplay(difference(value))
```



Choosing your own model

```
fit <- web_usage %>% model(  
  arima = ARIMA(value ~ pdq(3, 1, 0)))  
report(fit)
```

```
## Series: value  
## Model: ARIMA(3,1,0)  
##  
## Coefficients:  
##          ar1          ar2          ar3  
##          1.151    -0.6612    0.3407  
## s.e.    0.095     0.1353    0.0941  
##  
## sigma^2 estimated as 9.656:  log likelihood=-252  
## AIC=512    AICc=512.4    BIC=522.4
```

Choosing your own model

```
web_usage %>% model(ARIMA(value ~ pdq(d=1))) %>% report()
```


```
## Series: value
## Model: ARIMA(1,1,1)
##
## Coefficients:
##           ar1      ma1
##      0.6504  0.5256
## s.e.  0.0842  0.0896
##
## sigma^2 estimated as 9.995:  log likelihood=-254.2
## AIC=514.3   AICc=514.5   BIC=522.1
```

Choosing your own model

```
web_usage %>%  
  model(ARIMA(value ~ pdq(d=1), stepwise = FALSE,  
    approximation = FALSE)) %>% report()  
  
## Series: value  
## Model: ARIMA(3,1,0)  
##  
## Coefficients:  
##          ar1          ar2          ar3  
##          1.151    -0.6612    0.3407  
## s.e.    0.095     0.1353    0.0941  
##  
## sigma^2 estimated as 9.656:  log likelihood=-252  
## AIC=512    AICc=512.4    BIC=522.4
```

Choosing your own model

```
augment(fit) %>%  
  gg_tsdisplay(.resid, plot_type = "hist")
```



Choosing your own model

```
augment(fit) %>%  
  features(.resid, ljung_box, lag = 10, dof = 3)
```

```
## # A tibble: 1 x 3  
##   .model lb_stat lb_pval  
##   <chr>    <dbl>    <dbl>  
## 1 arima      4.49    0.722
```

Choosing your own model

```
fit %>% forecast(h = 10) %>%  
  autoplot(web_usage)
```




Modelling procedure with ARIMA

- 1 Plot the data. Identify any unusual observations.
- 2 If necessary, transform the data (using a Box-Cox transformation) to stabilize the variance.
- 3 If the data are non-stationary: take first differences of the data until the data are stationary.
- 4 Examine the ACF/PACF: Is an $AR(p)$ or $MA(q)$ model appropriate?
- 5 Try your chosen model(s), and use the AICc to search for a better model.
- 6 Check the residuals from your chosen model by plotting the ACF of the residuals, and doing a portmanteau test of the residuals. If they do not look like white noise, try a modified model.
- 7 Once the residuals look like white noise, calculate forecasts.

Automatic modelling procedure with ARIMA


- 1 Plot the data. Identify any unusual observations.
- 2 If necessary, transform the data (using a Box-Cox transformation) to stabilize the variance.
- 3 Use ARIMA to automatically select a model.
- 4
- 5
- 6 Check the residuals from your chosen model by plotting the ACF of the residuals, and doing a portmanteau test of the residuals. If they do not look like white noise, try a modified model.
- 7 Once the residuals look like white noise, calculate forecasts.

Modelling procedure



Seasonally adjusted electrical equipment

```
elecequip <- as_tsibble(fpp2::elecequip)
dcmp <- elecequip %>%
  STL(value ~ season(window = "periodic"))
dcmp %>% as_tsibble %>%
  autoplot(seas_adjust) + xlab("Year") +
  ylab("Seasonally adjusted new orders index")
```




Seasonally adjusted electrical equipment

- 1 Time plot shows sudden changes, particularly big drop in 2008/2009 due to global economic environment. Otherwise nothing unusual and no need for data adjustments.
- 2 No evidence of changing variance, so no Box-Cox transformation.
- 3 Data are clearly non-stationary, so we take first differences.

Seasonally adjusted electrical equipment

```
dcmp %>% gg_tsdisplay(difference(seas_adjust))
```



Seasonally adjusted electrical equipment

- 4 PACF is suggestive of AR(3). So initial candidate model is ARIMA(3,1,0). No other obvious candidates.
- 5 Fit ARIMA(3,1,0) model along with variations: ARIMA(4,1,0), ARIMA(2,1,0), ARIMA(3,1,1), etc. ARIMA(3,1,1) has smallest AICc value.

Seasonally adjusted electrical equipment

```
fit <- dcmp %>%  
  model(  
    arima = ARIMA(seas_adjust ~ pdq(3,1,1) + PDQ(0,0,0))  
  )  
report(fit)
```

```
## Series: seas_adjust
```

```
## Model: ARIMA(3,1,1)
```

```
##
```

```
## Coefficients:
```

```
##          ar1          ar2          ar3          ma1
```

```
##          0.0044   0.0916   0.3698   -0.3921
```

```
## s.e.    0.2201   0.0984   0.0669   0.2426
```

```
##
```


```
## sigma^2 estimated as 9.577: log likelihood=-492.7
```

```
## AIC=995.4   AICc=995.7   BIC=1012
```


Seasonally adjusted electrical equipment

6

ACF plot of residuals from ARIMA(3,1,1) model look like white noise.




Seasonally adjusted electrical equipment

```
## # A tibble: 1 x 3
##   .model lb_stat lb_pval
##   <chr>    <dbl>    <dbl>
## 1 arima      24.0    0.241
```

Seasonally adjusted electrical equipment

```
fit %>% forecast %>% autoplot(dcmp)
```



Outline

- 1 Stationarity and differencing
- 2 Non-seasonal ARIMA models
- 3 Estimation and order selection
- 4 ARIMA modelling in R
- 5 Forecasting
- 6 Seasonal ARIMA models
- 7 ARIMA vs ETS

Point forecasts

- 1 Rearrange ARIMA equation so y_t is on LHS.
- 2 Rewrite equation by replacing t by $T + h$.
- 3 On RHS, replace future observations by their forecasts, future errors by zero, and past errors by corresponding residuals.

Start with $h = 1$. Repeat for $h = 2, 3, \dots$

Point forecasts

ARIMA(3,1,1) forecasts: Step 1

$$(1 - \phi_1 B - \phi_2 B^2 - \phi_3 B^3)(1 - B)y_t = (1 + \theta_1 B)\varepsilon_t,$$

Point forecasts

ARIMA(3,1,1) forecasts: Step 1

$$(1 - \phi_1 B - \phi_2 B^2 - \phi_3 B^3)(1 - B)y_t = (1 + \theta_1 B)\varepsilon_t,$$

$$\begin{aligned} \left[1 - (1 + \phi_1)B + (\phi_1 - \phi_2)B^2 + (\phi_2 - \phi_3)B^3 + \phi_3 B^4 \right] y_t \\ = (1 + \theta_1 B)\varepsilon_t, \end{aligned}$$

Point forecasts

ARIMA(3,1,1) forecasts: Step 1

$$(1 - \phi_1 B - \phi_2 B^2 - \phi_3 B^3)(1 - B)y_t = (1 + \theta_1 B)\varepsilon_t,$$

$$\begin{aligned} \left[1 - (1 + \phi_1)B + (\phi_1 - \phi_2)B^2 + (\phi_2 - \phi_3)B^3 + \phi_3 B^4 \right] y_t \\ = (1 + \theta_1 B)\varepsilon_t, \end{aligned}$$

$$\begin{aligned} y_t - (1 + \phi_1)y_{t-1} + (\phi_1 - \phi_2)y_{t-2} + (\phi_2 - \phi_3)y_{t-3} \\ + \phi_3 y_{t-4} = \varepsilon_t + \theta_1 \varepsilon_{t-1}. \end{aligned}$$

Point forecasts

ARIMA(3,1,1) forecasts: Step 1

$$(1 - \phi_1 B - \phi_2 B^2 - \phi_3 B^3)(1 - B)y_t = (1 + \theta_1 B)\varepsilon_t,$$

$$\begin{aligned} \left[1 - (1 + \phi_1)B + (\phi_1 - \phi_2)B^2 + (\phi_2 - \phi_3)B^3 + \phi_3 B^4 \right] y_t \\ = (1 + \theta_1 B)\varepsilon_t, \end{aligned}$$

$$\begin{aligned} y_t - (1 + \phi_1)y_{t-1} + (\phi_1 - \phi_2)y_{t-2} + (\phi_2 - \phi_3)y_{t-3} \\ + \phi_3 y_{t-4} = \varepsilon_t + \theta_1 \varepsilon_{t-1}. \end{aligned}$$

$$\begin{aligned} y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} \\ - \phi_3 y_{t-4} + \varepsilon_t + \theta_1 \varepsilon_{t-1}. \end{aligned}$$

Point forecasts (h=1)

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} - \phi_3y_{t-4} + \varepsilon_t + \theta_1\varepsilon_{t-1}.$$

Point forecasts (h=1)

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} \\ - \phi_3y_{t-4} + \varepsilon_t + \theta_1\varepsilon_{t-1}.$$

ARIMA(3,1,1) forecasts: Step 2

$$y_{T+1} = (1 + \phi_1)y_T - (\phi_1 - \phi_2)y_{T-1} - (\phi_2 - \phi_3)y_{T-2} \\ - \phi_3y_{T-3} + \varepsilon_{T+1} + \theta_1\varepsilon_T.$$

Point forecasts (h=1)

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} \\ - \phi_3y_{t-4} + \varepsilon_t + \theta_1\varepsilon_{t-1}.$$

ARIMA(3,1,1) forecasts: Step 2

$$y_{T+1} = (1 + \phi_1)y_T - (\phi_1 - \phi_2)y_{T-1} - (\phi_2 - \phi_3)y_{T-2} \\ - \phi_3y_{T-3} + \varepsilon_{T+1} + \theta_1\varepsilon_T.$$

ARIMA(3,1,1) forecasts: Step 3

$$\hat{y}_{T+1|T} = (1 + \phi_1)y_T - (\phi_1 - \phi_2)y_{T-1} - (\phi_2 - \phi_3)y_{T-2} \\ - \phi_3y_{T-3} + \theta_1e_T.$$

Point forecasts (h=2)

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} \\ - \phi_3y_{t-4} + \varepsilon_t + \theta_1\varepsilon_{t-1}.$$

Point forecasts ($h=2$)

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} \\ - \phi_3y_{t-4} + \varepsilon_t + \theta_1\varepsilon_{t-1}.$$

ARIMA(3,1,1) forecasts: Step 2

$$y_{T+2} = (1 + \phi_1)y_{T+1} - (\phi_1 - \phi_2)y_T - (\phi_2 - \phi_3)y_{T-1} \\ - \phi_3y_{T-2} + \varepsilon_{T+2} + \theta_1\varepsilon_{T+1}.$$

Point forecasts (h=2)

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} \\ - \phi_3y_{t-4} + \varepsilon_t + \theta_1\varepsilon_{t-1}.$$

ARIMA(3,1,1) forecasts: Step 2

$$y_{T+2} = (1 + \phi_1)y_{T+1} - (\phi_1 - \phi_2)y_T - (\phi_2 - \phi_3)y_{T-1} \\ - \phi_3y_{T-2} + \varepsilon_{T+2} + \theta_1\varepsilon_{T+1}.$$

ARIMA(3,1,1) forecasts: Step 3

$$\hat{y}_{T+2|T} = (1 + \phi_1)\hat{y}_{T+1|T} - (\phi_1 - \phi_2)y_T - (\phi_2 - \phi_3)y_{T-1} \\ - \phi_3y_{T-2}.$$

Prediction intervals

95% prediction interval

$$\hat{y}_{T+h|T} \pm 1.96\sqrt{v_{T+h|T}}$$

where $v_{T+h|T}$ is estimated forecast variance.

Prediction intervals

95% prediction interval

$$\hat{y}_{T+h|T} \pm 1.96\sqrt{v_{T+h|T}}$$

where $v_{T+h|T}$ is estimated forecast variance.

- $v_{T+1|T} = \hat{\sigma}^2$ for all ARIMA models regardless of parameters and orders.
- Multi-step prediction intervals for ARIMA(0,0,q):

$$y_t = \varepsilon_t + \sum_{i=1}^q \theta_i \varepsilon_{t-i}.$$

$$v_{T|T+h} = \hat{\sigma}^2 \left[1 + \sum_{i=1}^{h-1} \theta_i^2 \right], \quad \text{for } h = 2, 3, \dots$$

Prediction intervals

95% prediction interval

$$\hat{y}_{T+h|T} \pm 1.96\sqrt{v_{T+h|T}}$$

where $v_{T+h|T}$ is estimated forecast variance.

- Multi-step prediction intervals for ARIMA(0,0,q):

$$y_t = \varepsilon_t + \sum_{i=1}^q \theta_i \varepsilon_{t-i}.$$

$$v_{T|T+h} = \hat{\sigma}^2 \left[1 + \sum_{i=1}^{h-1} \theta_i^2 \right], \quad \text{for } h = 2, 3, \dots$$

Prediction intervals

95% prediction interval

$$\hat{y}_{T+h|T} \pm 1.96\sqrt{v_{T+h|T}}$$

where $v_{T+h|T}$ is estimated forecast variance.

- Multi-step prediction intervals for ARIMA(0,0,q):

$$y_t = \varepsilon_t + \sum_{i=1}^q \theta_i \varepsilon_{t-i}.$$

$$v_{T|T+h} = \hat{\sigma}^2 \left[1 + \sum_{i=1}^{h-1} \theta_i^2 \right], \quad \text{for } h = 2, 3, \dots$$

- AR(1): Rewrite as MA(∞) and use above result.
- Other models beyond scope of this subject.

Prediction intervals

- Prediction intervals **increase in size with forecast horizon.**
- Prediction intervals can be difficult to calculate by hand
- Calculations assume residuals are **uncorrelated** and **normally distributed.**
- Prediction intervals tend to be too narrow.
 - ▶ the uncertainty in the parameter estimates has not been accounted for.
 - ▶ the ARIMA model assumes historical patterns will not change during the forecast period.
 - ▶ the ARIMA model assumes uncorrelated future errors 99

Your turn

For the United States GDP data (from `global_economy`):

- if necessary, find a suitable Box-Cox transformation for the data;
- fit a suitable ARIMA model to the transformed data;
- check the residual diagnostics;
- produce forecasts of your fitted model. Do the forecasts look reasonable?

Outline

- 1 Stationarity and differencing
- 2 Non-seasonal ARIMA models
- 3 Estimation and order selection
- 4 ARIMA modelling in R
- 5 Forecasting
- 6 Seasonal ARIMA models
- 7 ARIMA vs ETS

Seasonal ARIMA models

ARIMA	$\underbrace{(p, d, q)}$	$\underbrace{(P, D, Q)_m}$
	↑	↑
	Non-seasonal part of the model	Seasonal part of of the model

where m = number of observations per year.

Seasonal ARIMA models


E.g., $\text{ARIMA}(1, 1, 1)(1, 1, 1)_4$ model (without constant)

Seasonal ARIMA models

E.g., ARIMA(1, 1, 1)(1, 1, 1)₄ model (without constant)
$$(1 - \phi_1 B)(1 - \Phi_1 B^4)(1 - B)(1 - B^4)y_t = (1 + \theta_1 B)(1 + \Theta_1 B^4)\varepsilon_t.$$

Seasonal ARIMA models

E.g., $\text{ARIMA}(1, 1, 1)(1, 1, 1)_4$ model (without constant)
 $(1 - \phi_1 B)(1 - \Phi_1 B^4)(1 - B)(1 - B^4)y_t = (1 + \theta_1 B)(1 + \Theta_1 B^4)\varepsilon_t.$



Seasonal ARIMA models

E.g., ARIMA(1, 1, 1)(1, 1, 1)₄ model (without constant)
 $(1 - \phi_1 B)(1 - \Phi_1 B^4)(1 - B)(1 - B^4)y_t = (1 + \theta_1 B)(1 + \Theta_1 B^4)\varepsilon_t.$

All the factors can be multiplied out and the general model written as follows:

$$\begin{aligned} y_t = & (1 + \phi_1)y_{t-1} - \phi_1 y_{t-2} + (1 + \Phi_1)y_{t-4} \\ & - (1 + \phi_1 + \Phi_1 + \phi_1 \Phi_1)y_{t-5} + (\phi_1 + \phi_1 \Phi_1)y_{t-6} \\ & - \Phi_1 y_{t-8} + (\Phi_1 + \phi_1 \Phi_1)y_{t-9} - \phi_1 \Phi_1 y_{t-10} \\ & + \varepsilon_t + \theta_1 \varepsilon_{t-1} + \Theta_1 \varepsilon_{t-4} + \theta_1 \Theta_1 \varepsilon_{t-5}. \end{aligned}$$

Common ARIMA models

The US Census Bureau uses the following models most often:

ARIMA(0,1,1)(0,1,1) _m	with log transformation
ARIMA(0,1,2)(0,1,1) _m	with log transformation
ARIMA(2,1,0)(0,1,1) _m	with log transformation
ARIMA(0,2,2)(0,1,1) _m	with log transformation
ARIMA(2,1,2)(0,1,1) _m	with no transformation

Seasonal ARIMA models

The seasonal part of an AR or MA model will be seen in the seasonal lags of the PACF and ACF.

ARIMA(0,0,0)(0,0,1)₁₂ will show:


- a spike at lag 12 in the ACF but no other significant spikes.
- The PACF will show exponential decay in the seasonal lags; that is, at lags 12, 24, 36,

ARIMA(0,0,0)(1,0,0)₁₂ will show:

- exponential decay in the seasonal lags of the ACF
- a single significant spike at lag 12 in the PACF.


European quarterly retail trade

```
eu_retail %>% autoplot(value) +  
  xlab("Year") + ylab("Retail index")
```




European quarterly retail trade

```
eu_retail %>% gg_tsdisplay(  
  value %>% difference(4))
```



European quarterly retail trade

```
eu_retail %>% gg_tsdisplay(  
  value %>% difference(4) %>% difference(1))
```




European quarterly retail trade

- $d = 1$ and $D = 1$ seems necessary.
- Significant spike at lag 1 in ACF suggests non-seasonal MA(1) component.
- Significant spike at lag 4 in ACF suggests seasonal MA(1) component.
- Initial candidate model: $\text{ARIMA}(0,1,1)(0,1,1)_4$.
- We could also have started with $\text{ARIMA}(1,1,0)(1,1,0)_4$.

European quarterly retail trade

```
fit <- eu_retail %>%  
  model(arima = ARIMA(value ~ pdq(0,1,1) + PDQ(0,1,1)))  
augment(fit) %>% gg_tsdisplay(.resid, plot_type = "hist")
```



European quarterly retail trade

```
augment(fit) %>%  
  features(.resid, ljung_box, lag = 8, dof = 2)
```

```
## # A tibble: 1 x 3  
##   .model lb_stat lb_pval  
##   <chr>    <dbl>    <dbl>  
## 1 arima      10.7    0.0997
```

European quarterly retail trade

- ACF and PACF of residuals show significant spikes at lag 2, and maybe lag 3.
- AICc of $\text{ARIMA}(0,1,1)(0,1,1)_4$ model is 75.72
- AICc of $\text{ARIMA}(0,1,2)(0,1,1)_4$ model is 74.27.
- AICc of $\text{ARIMA}(0,1,3)(0,1,1)_4$ model is 68.39.
- AICc of $\text{ARIMA}(0,1,4)(0,1,1)_4$ model is 70.73.

European quarterly retail trade


```
fit <- eu_retail %>%  
  model(  
    arima013011 = ARIMA(value ~ pdq(0,1,3) + PDQ(0,1,1))  
  )  
report(fit)
```

```
## Series: value  
## Model: ARIMA(0,1,3)(0,1,1)[4]  
##  
## Coefficients:  
##          ma1      ma2      ma3      sma1  
##      0.2630  0.3694  0.4200 -0.6636  
## s.e.  0.1237  0.1255  0.1294  0.1545  
##  
## sigma^2 estimated as 0.156: log likelihood=-28.63  
## AIC=67.26   AICc=68.39   BIC=77.65
```

European quarterly retail trade

```
augment(fit) %>%
```

```
gg_tsdisplay(.resid, plot_type = "hist")
```




European quarterly retail trade

```
augment(fit) %>%  
  features(.resid, lbjung_box, lag = 8, dof = 4)
```

```
## # A tibble: 1 x 3  
##   .model      lb_stat lb_pval  
##   <chr>      <dbl>   <dbl>  
## 1 arima013011 0.511    0.972
```

European quarterly retail trade

```
fit %>% forecast(h = "3 years") %>%  
  autoplot(eu_retail)
```



European quarterly retail trade

```
eu_retail %>% model(ARIMA(value)) %>% report()
```

```
## Series: value
```

```
## Model: ARIMA(0,1,3)(0,1,1)[4]
```

```
##
```

```
## Coefficients:
```

```
##          ma1          ma2          ma3          sma1
```

```
##          0.2630  0.3694  0.4200  -0.6636
```

```
## s.e.    0.1237  0.1255  0.1294   0.1545
```

```
##
```

```
## sigma^2 estimated as 0.156: log likelihood=-28.63
```

```
## AIC=67.26   AICc=68.39   BIC=77.65
```

European quarterly retail trade

```
eu_retail %>% model(ARIMA(value, stepwise = FALSE,  
  approximation = FALSE)) %>% report()
```

```
## Series: value
```

```
## Model: ARIMA(0,1,3)(0,1,1)[4]
```

```
##
```

```
## Coefficients:
```

```
##          ma1          ma2          ma3          sma1
```

```
##          0.2630  0.3694  0.4200  -0.6636
```


```
## s.e.  0.1237  0.1255  0.1294  0.1545
```

```
##
```


```
## sigma^2 estimated as 0.156: log likelihood=-28.63
```

```
## AIC=67.26  AICc=68.39  BIC=77.65
```

Corticosteroid drug sales



Corticosteroid drug sales



Corticosteroid drug sales

- Choose $D = 1$ and $d = 0$.
- Spikes in PACF at lags 12 and 24 suggest seasonal AR(2) term.
- Spikes in PACF suggests possible non-seasonal AR(3) term.
- Initial candidate model: $\text{ARIMA}(3,0,0)(2,1,0)_{12}$.

Corticosteroid drug sales

.model	AICc
ARIMA(3,0,1)(0,1,2)[12]	-485.5
ARIMA(3,0,1)(1,1,1)[12]	-484.3
ARIMA(3,0,1)(0,1,1)[12]	-483.7
ARIMA(3,0,1)(2,1,0)[12]	-476.3
ARIMA(3,0,0)(2,1,0)[12]	-475.1
ARIMA(3,0,2)(2,1,0)[12]	-474.9
ARIMA(3,0,1)(1,1,0)[12]	-463.4

Corticosteroid drug sales

```
fit <- h02 %>%  
  model(best = ARIMA(log(Cost) ~ 0 + pdq(3,0,1) + PDQ(0,1,2)))  
report(fit)
```

```
## Series: Cost
```

```
## Model: ARIMA(3,0,1)(0,1,2)[12]
```

```
## Transformation: log(.x)
```

```
##
```

```
## Coefficients:
```

	ar1	ar2	ar3	ma1	sma1	sma2
	-0.1602	0.5481	0.5678	0.3826	-0.5222	-0.1769
s.e.	0.1636	0.0878	0.0942	0.1895	0.0861	0.0872

```
##
```


```
## sigma^2 estimated as 0.004289: log likelihood=250.1
```

```
## AIC=-486.1 AICc=-485.5 BIC=-463.3
```

Corticosteroid drug sales

```
augment(fit) %>%
```

```
gg_tsdisplay(.resid, lag_max=36, plot_type = "hist")
```



Corticosteroid drug sales

```
augment(fit) %>%  
  features(.resid, ljung_box, lag = 36, dof = 6)
```

```
## # A tibble: 1 x 3  
##   .model lb_stat lb_pval  
##   <chr>    <dbl>    <dbl>  
## 1 best      50.5    0.0109
```


Corticosteroid drug sales

```
fit <- h02 %>% model(auto = ARIMA(log(Cost)))  
report(fit)
```

```
## Series: Cost  
## Model: ARIMA(2,1,0)(0,1,1)[12]  
## Transformation: log(.x)  
##  
## Coefficients:  
##          ar1      ar2      sma1  
##      -0.8491  -0.4207  -0.6401  
## s.e.   0.0712   0.0714   0.0694  
##  
## sigma^2 estimated as 0.004399:  log likelihood=245.4  
## AIC=-482.8   AICc=-482.6   BIC=-469.8
```

Corticosteroid drug sales

```
augment(fit) %>%  
  gg_tsdisplay(.resid, lag_max = 36, plot_type = "hist")
```



Corticosteroid drug sales

```
augment(fit) %>%  
  features(.resid, ljung_box, lag = 36, dof = 5)
```

```
## # A tibble: 1 x 3  
##   .model lb_stat lb_pval  
##   <chr>    <dbl>    <dbl>  
## 1 auto      57.5  0.00260
```


Corticosteroid drug sales

```
fit <- h02 %>%  
  model(best = ARIMA(log(Cost), stepwise = FALSE,  
    approximation = FALSE,  
    order_constraint = p + q + P + Q <= 9))  
report(fit)
```

```
## Series: Cost  
## Model: ARIMA(4,1,1)(2,1,2)[12]  
## Transformation: log(.x)  
##  
## Coefficients:  
##          ar1      ar2      ar3      ar4      ma1      sar1  
##      -0.0426  0.2097  0.2016 -0.2273 -0.7423  0.6213  
## s.e.   0.2167  0.1814  0.1144  0.0810  0.2075  0.2421  
##          sar2      sma1      sma2  
##      -0.3832 -1.2018  0.4958  
## s.e.   0.1185  0.2492  0.2136  
##  
## sigma^2 estimated as 0.004061: log likelihood=254.3  
## AIC=-488.6 AICc=-487.4 BIC=-456.1
```

Corticosteroid drug sales

```
augment(fit) %>%  
  gg_tsdisplay(.resid, lag_max = 36, plot_type = "hist")
```



Corticosteroid drug sales

```
augment(fit) %>%  
  features(.resid, ljung_box, lag = 36, dof = 9)
```

```
## # A tibble: 1 x 3  
##   .model lb_stat lb_pval  
##   <chr>    <dbl>    <dbl>  
## 1 best      35.1     0.136
```

Corticosteroid drug sales

Training data: July 1991 to June 2006

Test data: July 2006–June 2008

```
fit <- h02 %>%  
  filter_index(~ "2006 Jun") %>%  
  model(  
    ARIMA(log(Cost) ~ pdq(3, 0, 0) + PDQ(2, 1, 0)),  
    ARIMA(log(Cost) ~ pdq(3, 0, 1) + PDQ(2, 1, 0)),  
    ARIMA(log(Cost) ~ pdq(3, 0, 2) + PDQ(2, 1, 0)),  
    ARIMA(log(Cost) ~ pdq(3, 0, 1) + PDQ(1, 1, 0))  
    # ... #  
  )  
  
fit %>%  
  forecast(h = "2 years") %>%  
  accuracy(h02 %>% filter_index("2006 Jul" ~ .))
```


Corticosteroid drug sales


.model	RMSE
ARIMA(3,0,1)(1,1,1)[12]	61878
ARIMA(3,0,1)(0,1,2)[12]	62142
ARIMA(2,1,4)(0,1,1)[12]	62708
ARIMA(2,1,3)(0,1,1)[12]	62855
ARIMA(3,0,1)(0,1,1)[12]	62947
ARIMA(3,0,2)(0,1,1)[12]	62968
ARIMA(4,1,1)(2,1,2)[12]	63114
ARIMA(3,0,3)(0,1,1)[12]	63284
ARIMA(2,1,5)(0,1,1)[12]	63610
ARIMA(3,0,2)(2,1,0)[12]	65146
ARIMA(3,0,1)(2,1,0)[12]	65270
ARIMA(3,0,1)(1,1,0)[12]	66644
ARIMA(3,0,2)(2,1,0)[12]	66811

Corticosteroid drug sales

- Models with lowest AICc values tend to give slightly better results than the other models.
- AICc comparisons must have the same orders of differencing. But RMSE test set comparisons can involve any models.
- Use the best model available, even if it does not pass all tests.

Corticosteroid drug sales

```
fit <- h02 %>%  
  model(ARIMA(Cost ~ 0 + pdq(3,0,1) + PDQ(0,1,2)))  
fit %>% forecast %>% autoplot(h02) +  
  ylab("H02 Expenditure ($AUD)") + xlab("Year")
```



Outline

- 1 Stationarity and differencing
- 2 Non-seasonal ARIMA models
- 3 Estimation and order selection
- 4 ARIMA modelling in R
- 5 Forecasting
- 6 Seasonal ARIMA models
- 7 ARIMA vs ETS

ARIMA vs ETS

- Myth that ARIMA models are more general than exponential smoothing.
- Linear exponential smoothing models all special cases of ARIMA models.
- Non-linear exponential smoothing models have no equivalent ARIMA counterparts.
- Many ARIMA models have no exponential smoothing counterparts.
- ETS models all non-stationary. Models with seasonality or non-damped trend (or both) have two unit roots; all other models have one unit root.

Equivalences

ETS model	ARIMA model	Parameters
ETS(A,N,N)	ARIMA(0,1,1)	$\theta_1 = \alpha - 1$
ETS(A,A,N)	ARIMA(0,2,2)	$\theta_1 = \alpha + \beta - 2$ $\theta_2 = 1 - \alpha$
ETS(A,A,N)	ARIMA(1,1,2)	$\phi_1 = \phi$ $\theta_1 = \alpha + \phi\beta - 1 - \phi$ $\theta_2 = (1 - \alpha)\phi$
ETS(A,N,A)	ARIMA(0,0,m)(0,1,0) _m	
ETS(A,A,A)	ARIMA(0,1,m + 1)(0,1,0) _m	
ETS(A,A,A)	ARIMA(1,0,m + 1)(0,1,0) _m	

Your turn

For the `fma::condmilk` series:

- Do the data need transforming? If so, find a suitable transformation.
- Are the data stationary? If not, find an appropriate differencing which yields stationary data.
- Identify a couple of ARIMA models that might be useful in describing the time series.
- Which of your models is the best according to their AIC values?
- Estimate the parameters of your best model and do diagnostic testing on the residuals. Do the residuals resemble white noise? If not, try to find another ARIMA model which fits better.
- Forecast the next 24 months of data using your preferred model.