


Essentials of Mathematical Methods:

Foundations, Principles, and Algorithms

Yuguang Yang

version 3.0



God used beautiful mathematics in creating the world. –Paul Dirac

*Dedicated to
those who appreciate the power of mathematical methods
and enjoy learning it.*

License statement

You are free to redistribute the material in any medium or format under the following terms:

- **Attribution:** You must give appropriate credit, provide a link to the license, and indicate if changes were made. You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.
- **NonCommercial:** You may not use the material for commercial purposes.

- * The licensor cannot revoke these freedoms as long as you follow the license terms.
- * This license is created via creative commons (<https://creativecommons.org>)
- * If you have any questions regarding the license, please contact the author.

Preface

Objective

Today, mathematical methods, models, and computational algorithms are playing increasingly significant roles in addressing major challenges arising from scientific research and technological developments. Although many novel methods and algorithms, such as deep learning and artificial intelligence, are emerging and reshaping various areas at an unprecedented pace, their core ideas and working mechanisms are inherently related to and deeply rooted in some essential mathematical foundations and principles. By performing an in-depth survey on the underlying foundations, principles, and algorithms, this book aims to navigate the vast landscape of mathematical methods widely used in diverse scientific and engineering domains.

This book starts with a survey of mathematical foundations, including essential concepts and theorems in real analysis, linear algebra, and related fundamentals. Then it examines a broad spectrum of applied mathematical methods, ranging from traditional ones such as optimizations and dynamical system modeling, to state-of-the-art such as machine learning, deep learning, and reinforcement learning. The emphasis is placed on methods for stochastic and dynamical system modeling, optimal decision-making, and statistical learning. For each topic, this book organizes fundamental definitions, theorems, methods, and algorithms in a logical and illuminating way.

Features and Highlights

- Comprehensive, essential, and self-contained.
- Concepts, theorems, and discussions are developed to suit real-world applications.
- Key references and resources are provided on each topic.
- Comparisons and discussions on similar definitions and theorems.
- An evolving book with regular updates on [Github](#).

Acknowledgment

This book evolved from my study notes during my PhD studies at the Johns Hopkins University (JHU). I want to thank the following professors at JHU for their courses and valuable discussion: Daniel Robinson, Teresa Lebar, Andrea Prosperetti, Gregory Chirikjian, Michael Kahadan, James C. Spall, Marin Kobilarov, Suchi Saria, Michael Dimitz, Sean Sun, Ari Turner, Gregory Eyink, Amitabh Basu, John Miller and David Audley. I also want to thank Rachael Zhang for her editorial assistance.

Yuguang Yang, Fall 2019
yangyutu123@gmail.com

Notations

- \mathbb{R} : real numbers.
- \mathbb{R}_+ : nonnegative real numbers.
- \mathbb{R}_{++} : positive real numbers.
- $\bar{\mathbb{R}}$: extended real numbers.
- \mathbb{C} : complex numbers.
- \mathbb{F} : real or complex numbers.
- \mathbb{Q} : rational numbers.
- \mathbb{Z} : integer numbers.
- \mathbb{P} : positive numbers.
- \mathcal{P}_n : polynomial of degree of n .
- \mathbb{N} : natural numbers.
- $\mathcal{R}(A)$: the range of matrix A .
- $\mathcal{N}(A)$: the null space of matrix A .
- V : vector space.
- $\det(A)$: the determinant of matrix A .
- $\text{rank}(A)$: the rank of matrix A .
- $\|\cdot\|_2$: Euclidean 2 norm of a vector of a matrix.
- $\|\cdot\|_F$: Frobenius norm of a matrix.
- $\rho(A)$: the spectral radius of matrix A .
- $\text{Tr}(A)$: the trace of matrix A .
- $L^2[a, b]$: Lebesgue integrable function on $[a, b]$.
- $L^1[a, b]$: Lebesgue integrable function on $[a, b]$.
- $N(0, 1)$: standard Gaussian distribution.
- $N(\mu, \sigma^2)$: Gaussian distribution with mean μ and variance σ^2 .
- $MN(\mu, \Sigma)$: multivariate Gaussian distribution with mean vector μ and covariance matrix Σ .
- $\mathbf{1}(x), I(x)$ indicator function.
- $E[X], \mathbb{E}[X], \mathbb{E}[X]$ expectation of random variable X .
- $\text{Var}[X]$ variance of random variable X .

CONTENTS

i mathematical foundations

- 1 SETS, SEQUENCES, AND SERIES 2
- 2 METRIC SPACE 36
- 3 ADVANCED CALCULUS 58
- 4 LINEAR ALGEBRA AND MATRIX ANALYSIS 138
- 5 BASIC FUNCTIONAL ANALYSIS 268

ii mathematical optimization methods

- 6 UNCONSTRAINED NONLINEAR OPTIMIZATION 315
- 7 CONSTRAINED NONLINEAR OPTIMIZATION 359
- 8 LINEAR OPTIMIZATION 405
- 9 CONVEX ANALYSIS AND CONVEX OPTIMIZATION 428
- 10 BASIC GAME THEORY 477

iii classical statistical methods

- 11 PROBABILITY THEORY 499
- 12 STATISTICAL DISTRIBUTIONS 603
- 13 STATISTICAL ESTIMATION THEORY 657
- 14 MULTIVARIATE STATISTICAL METHODS 716
- 15 LINEAR REGRESSION ANALYSIS 806
- 16 MONTE CARLO METHODS 900

iv dynamics modeling methods

- 17 MODELS AND ESTIMATION IN LINEAR SYSTEMS 936
- 18 STOCHASTIC PROCESS 1010

- 19 STOCHASTIC CALCULUS 1047
- 20 MARKOV CHAIN AND RANDOM WALK 1095
- 21 TIME SERIES ANALYSIS 1144

v statistical learning methods

- 22 SUPERVISED LEARNING PRINCIPLES 1223
- 23 LINEAR MODELS FOR REGRESSION 1268
- 24 LINEAR MODELS FOR CLASSIFICATION 1287
- 25 GENERATIVE MODELS 1338
- 26 K NEAREST NEIGHBORS 1360
- 27 TREE METHODS 1366
- 28 ENSEMBLE AND BOOSTING METHODS 1390
- 29 UNSUPERVISED STATISTICAL LEARNING 1420
- 30 NEURAL NETWORK AND DEEP LEARNING 1483

vi optimal control and reinforcement learning methods

- 31 CLASSICAL OPTIMAL CONTROL THEORY 1593
- 32 REINFORCEMENT LEARNING 1611

vii appendix

- A SUPPLEMENTAL MATHEMATICAL FACTS 1691
- Alphabetical Index 1721

LIST OF ALGORITHMS

| | | |
|----|---|-----|
| 1 | A generic line search algorithm | 324 |
| 2 | Backtracking-Armijo line search algorithm | 335 |
| 3 | Steepest decent Backtracking-Armijo line search algorithm | 338 |
| 4 | Modified Newton Backtracking-Armijo line search algorithm | 338 |
| 5 | Quasi Newton with Wolfe line search algorithm | 338 |
| 6 | A generic trust-region algorithm | 340 |
| 7 | A linear conjugate algorithm | 349 |
| 8 | Iteratively reweighted least squares for p norm least square | 352 |
| 9 | Gauss-Newton method for nonlinear least-square algorithm | 354 |
| 10 | Levenberg-Marquardt method for nonlinear least-square algorithm | 355 |
| 11 | Newton method for root finding | 356 |
| 12 | Primal active-set method for strictly convex quadratic programming | 378 |
| 13 | First order gradient projection algorithm | 381 |
| 14 | The Simplex algorithm (non-degenerate system) | 419 |
| 15 | Primal-dual long-step path-following algorithm | 424 |
| 16 | A generic subgradient algorithm | 463 |
| 17 | Gradient projection algorithm with constant step size | 470 |
| 18 | Gradient projection algorithm with adaptive size | 471 |
| 19 | Proximal gradient algorithm | 472 |
| 20 | Iterative Shrinkage-Thresholding Algorithm with constant step size for L_1 optimization | 474 |
| 21 | Iterative reweighed estimation for multivariate normal distribution | 723 |
| 22 | Alternating sparse canonical correlation analysis algorithm | 743 |
| 23 | EM algorithm for least square with nonconstant variance | 869 |
| 24 | Accept-Reject algorithm for random number generation. | 904 |

| | | |
|----|--|------|
| 25 | Importance sampling for Monte Carlo integration. | 913 |
| 26 | MCMC Metropolis-Hasting algorithm | 917 |
| 27 | MCMC Gibbs sampling algorithm | 919 |
| 28 | Recursive linear least square estimation of dynamical systems. | 1001 |
| 29 | Recursive nonlinear least square of dynamical systems | 1003 |
| 30 | Kalman filtering | 1007 |
| 31 | Extended Kalman filter | 1008 |
| 32 | Coordinate descent for Lasso regression. | 1279 |
| 33 | Iteratively reweighted least squares for logistic regression | 1293 |
| 34 | Perceptron learning algorithm | 1323 |
| 35 | Soft margin SVM algorithm | 1332 |
| 36 | Multinomial Naive Bayes classification | 1347 |
| 37 | KNN classification and regression algorithm | 1361 |
| 38 | A generic decision tree generation algorithm | 1374 |
| 39 | ID3 classification decision tree algorithm | 1379 |
| 40 | A regression tree growth algorithm | 1387 |
| 41 | A basic bagging algorithm | 1396 |
| 42 | Generic Adaboost classifier algorithm | 1402 |
| 43 | Adaboost regressor algorithm | 1405 |
| 44 | A generic additive model algorithm | 1407 |
| 45 | Generic gradient boosting algorithm | 1410 |
| 46 | Gradient tree boosting algorithm | 1411 |
| 47 | XGBoost algorithm | 1417 |
| 48 | Iterative reweighted least square PCA with outliers algorithm | 1432 |
| 49 | Random sample consensus PCA with outliers algorithm | 1432 |
| 50 | Orthogonal Matching Pursuit | 1434 |
| 51 | K-SVD for dictionary learning. | 1436 |
| 52 | Online dictionary learning | 1437 |
| 53 | Stochastic gradient descent for matrix factorization based recommender systems. | 1447 |

| | | |
|----|--|------|
| 54 | Isomap algorithm | 1456 |
| 55 | Kernel PCA algorithm | 1457 |
| 56 | Laplacian Eigenmap algorithm | 1462 |
| 57 | Diffusion map algorithm | 1465 |
| 58 | K-means algorithm. | 1470 |
| 59 | DBSCAN algorithm | 1473 |
| 60 | General spectral clustering algorithm | 1475 |
| 61 | Gaussian mixture model EM algorithm | 1479 |
| 62 | Full batch gradient descent algorithm | 1499 |
| 63 | Minibatch stochastic gradient descent algorithm | 1500 |
| 64 | Adam stochastic gradient descent algorithm. | 1505 |
| 65 | Solve forward backward stochastic differential equation via deep learning. . | 1527 |
| 66 | Minibatch stochastic gradient descent training of GAN. | 1580 |
| 67 | Minibatch stochastic gradient descent training of conditional GAN. | 1587 |
| 68 | Minibatch stochastic gradient descent training of Wasserstein GAN. | 1589 |
| 69 | The policy iteration algorithm for MDP | 1618 |
| 70 | Value iteration algorithm for a finite state MDP | 1620 |
| 71 | First-visit MC value function estimation | 1627 |
| 72 | MC-based reinforcement learning control | 1629 |
| 73 | TD(o) estimation of a value function. | 1630 |
| 74 | SARSA learning | 1631 |
| 75 | Q-learning algorithm | 1632 |
| 76 | TD(n) estimation of a value function. | 1635 |
| 77 | n -step SARSA learning | 1636 |
| 78 | A generic batch policy-gradient algorithm (REINFORCE) | 1647 |
| 79 | A basic Monte-Carlo policy-gradient algorithm | 1648 |
| 80 | Actor-Critic policy-gradient method | 1650 |
| 81 | Policy-gradient method with a value function baseline | 1655 |
| 82 | Neural Fitted Q Iteration (NFQ) | 1660 |
| 83 | Deep Q-learning with experience replay | 1662 |

| | | |
|----|--|------|
| 84 | Asynchronous Deep Q-Learning for each thread | 1667 |
| 85 | Deep Q-learning with universal value function approximator | 1668 |
| 86 | Deep deterministic policy gradient algorithm (DDPG) | 1671 |
| 87 | Twin-delayed deep deterministic policy gradient (TD3) | 1673 |
| 88 | Trust Region Policy Optimization | 1675 |
| 89 | Proximal Policy Optimization | 1678 |
| 90 | Soft Actor-Critic (SAC) policy optimization | 1681 |
| 91 | Isotropic multivariate Gaussian evolution strategies for reinforcement learning | 1683 |
| 92 | Isotropic multivariate Gaussian parallelized evolution strategies for reinforcement learning | 1683 |
| 93 | Q learning with prioritized experience replay | 1685 |
| 94 | Deep Q-learning with hindsight experience replay | 1686 |

LIST OF FIGURES

| | |
|---------------|---|
| Figure 3.1.1 | An example 3D curve generated by $z = t, x = t \cos(t), y = t \sin(t)$. 69 |
| Figure 3.1.2 | An example smooth surface generated by $z = x \exp(-2x^2 - y^2)$ 72 |
| Figure 4.9.1 | Demonstration of SVD for matrices of different shapes. The dashed lines highlight the compact form SVD. 207 |
| Figure 4.12.1 | Illustration of different quadratic forms. 228 |
| Figure 6.1.1 | Demonstration of local minimizer (red, green, and blue), strict local minimizer (red and blue), and global minimizer (blue). 318 |
| Figure 6.1.2 | A complex objective function in unconstrained optimization. 319 |
| Figure 6.1.3 | Illustration of different cases in unconstrained quadratic optimization. 323 |
| Figure 6.2.1 | Drawbacks of steepest gradient descent. 326 |
| Figure 6.2.2 | Demonstration on the step choices on the iterative algorithm. (a) Large step size. (b) Small step size. 331 |
| Figure 6.2.3 | Armijo sufficient decrease condition. 335 |
| Figure 6.3.1 | Demonstration of the dogleg path as an approximation to the exact solution path in the trust-region subproblem. 343 |
| Figure 6.4.1 | Demonstration of coordinate descent procedures when A is diagonal and non-diagonal. 346 |
| Figure 7.1.1 | Demonstration of KKT condition at a local minimal for $f(x_1, x_2) = x_1^2 + x_2^2$ under constraint $x_1 + x_2 = 1$. 363 |
| Figure 7.2.1 | Demonstration of KKT condition at a local minimal for $f(x_1, x_2) = x_1^2 + x_2^2$ under constraint $x_1 + x_2 \geq 1$. 373 |
| Figure 8.2.1 | The geometry of linear programming. (a) The feasible region is an open space extending to infinity if A is not full column rank. (b-d) Example feasible regions if $\text{rank}(A) = n, m \geq n$. Red arrows are direction of $-c$. When moving along $-c$ in the feasible region, the objective function will decrease. 409 |
| Figure 8.2.2 | Demonstration on multiple minimizers forming a convex set. 411 |
| Figure 8.3.1 | Overview of geometry approach to linear programming. 413 |
| Figure 9.1.1 | Example 2D affine hull and 3D affine hull. 430 |
| Figure 9.1.2 | Affine subspace and linear subspace. 432 |
| Figure 9.2.1 | (left) A convex set. (right) A non-convex set. 436 |

- Figure 9.2.2 (left) The affine hull of two points in a plane is a line passing through them. (right) The convex hull of two points in a plane is a line segment containing them. [438](#)
- Figure 9.2.3 An illustration of separating hyperplane theorem for two convex bodies. [440](#)
- Figure 9.2.4 An illustration of Farkas' lemma. (left) When b lies outside the cone (that is, $Ax = b, x \geq 0$ has no solution), there exists a hyperplane, characterized by normal vector y , separating b and the cone. (right) When b lies inside the cone (that is, $Ax = b, x \geq 0$ has a solution), there does not exist a hyperplane, characterized by normal vector y , separating b and the cone. [443](#)
- Figure 9.2.5 An illustration of Farkas' lemma variant where the cone is open set. (left) When b lies outside the cone (that is, $Ax = b, x > 0$ has no solution), there exists a hyperplane, characterized by normal vector y , separating b and the cone. (right) When b lies inside the cone (that is, $Ax = b, x > 0$ has a solution), there does not exist a hyperplane, characterized by normal vector y , separating b and the cone. [443](#)
- Figure 9.3.1 Demonstration of convex functions. (a) A convex function satisfying $f(\alpha x_1 + (1 - \alpha)x_2) \leq \alpha f(x_1) + (1 - \alpha)f(x_2)$. (b) A non-convex function where the green points does not satisfy the relation. [445](#)
- Figure 9.3.2 The epigraph (green area) of a convex function. [447](#)
- Figure 9.3.3 Illustration of linear underestimator. [450](#)
- Figure 9.5.1 Demonstration of optimality condition $\nabla f(x^*)^T(x - x^*) \geq 0, \forall x \in X$ when x^* lies on the boundary of X . [460](#)
- Figure 11.1.1 An illustration of measurable functions. [506](#)
- Figure 11.11.1 Visualization of central limit theorem. Samples are drawn from uniform and lognormal distributions. Sample means \bar{X}_n converge to normal distribution in distribution when n is large. [576](#)
- Figure 12.1.1 Comparison of Laplace distribution and normal distribution. [613](#)
- Figure 12.1.2 Density of $LN(0,1)$ and $LN(0,0.5)$. Note the positive skewness. [620](#)
- Figure 12.1.3 Density of $NLN(0,1)$ and $NLN(0,0.5)$. Note the negative skewness. [621](#)
- Figure 12.2.1 Distributions with left-skewness (black) and right-skewness (red). [642](#)
- Figure 12.2.2 Distributions with zero excess Kurtosis (Normal distribution, black), positive excess kurtosis (Laplace distribution, red), and negative excess Kurtosis (Uniform distribution, blue). [643](#)
- Figure 12.2.3 Percentile points at $\alpha = 0.1, 0.2, \dots, 0.9$ for a standard normal distribution. [645](#)

| | |
|---------------|--|
| Figure 12.2.4 | QQ plot of different sample distributions against standard normal distribution, including standard normal $N(0, 1)$, shifted-scaled normal $N(50, 5)$, Student's t with degree 1, and lognormal $LN(0, 1)$. Red solid lines are the fitted linear lines. 647 |
| Figure 13.1.1 | Statistical estimation and inference scheme. 660 |
| Figure 13.1.2 | An example of biased estimator with smaller variance than unbiased estimator 665 |
| Figure 13.1.3 | Visualization of log-likelihood function for normal distributed samples. 673 |
| Figure 13.6.1 | Demonstration for rejection regions for upper-tailed one-sided hypothesis (a), two-sided hypothesis(b), and lower-tailed one-sided hypothesis (c). 700 |
| Figure 13.7.1 | QQ plot with different sample distributions, including normal, Laplace ($b = 4$), Uniform $U([-1, 1])$, Lognormal $LN(0, 1)$. Red solid lines are the fitted linear lines. 709 |
| Figure 14.2.1 | Principal components for 2D samples. 727 |
| Figure 14.2.2 | PCA eigenface analysis. 735 |
| Figure 14.2.3 | PCA eigen-digit analysis for MNIST dataset. 736 |
| Figure 14.2.4 | Demonstration of interest rate curve dynamics. 738 |
| Figure 14.2.5 | Demonstration of first three dominating PCA factor in the swap rate curve daily change. 738 |
| Figure 14.4.1 | Gaussian copula with different correlations. 760 |
| Figure 14.4.2 | Student T copula with different correlations. 764 |
| Figure 14.4.3 | Generated correlated default time via Gaussian copula with different correlations. The hazard rate for both parties is $h(t) = 0.01$. 781 |
| Figure 14.5.1 | Correlation structure for the Gaussian copula implied by the factor model. 792 |
| Figure 14.5.2 | Factor model using (a) external factors or (b) internal factors. 793 |
| Figure 14.5.3 | Scatter plot of AAPL return vs. market excess return, SMB excess return and HML excess return. 797 |
| Figure 14.6.1 | A fully connected graphical model representing the joint distribution $P(X_1, \dots, X_N)$. 800 |
| Figure 14.6.2 | Graphical model examples. 802 |
| Figure 14.6.3 | d-separation examples. 804 |
| Figure 15.1.1 | Demonstration of simple linear regression model $y = \beta_1 x + \beta_0 + \epsilon$ and multiple linear regression model $y = \beta_1 x_1 + \beta_2 x_2 + \beta_0 + \epsilon$. Scatter points are observed data. The solid line in the left and the plane in the right are the mean responses. 809 |
| Figure 15.3.1 | Demo of heteroskedasticity in linear regression. The noises are larger at larger x values. 867 |

| | |
|---------------|--|
| Figure 15.3.2 | Demonstrations on linear regression with auto-correlated error. Observations are generated by $y_i = x_i + \epsilon_i, \epsilon_i = \rho\epsilon_{i-1} + z, z \sim N(0, 1)$. 872 |
| Figure 15.3.3 | Illustration of an outlier, a high-leverage point, and a influential point. Left subfigure shows a red-colored outlier, which does not have high leverage and large influence on the regression result. Middle subfigure shows a red-colored high-leverage point, which is not an outlier or influential point due to its weak influence on the regression result. Right subfigure shows an influential point that is both an outlier and a high-leverage point. 877 |
| Figure 15.3.4 | Visual scatter and box plots to identify outliers. 878 |
| Figure 15.3.5 | Different function choice for M-estimation linear regression 883 |
| Figure 15.3.6 | Common linear regression diagnosis plots 886 |
| Figure 15.4.1 | Diagnosis plots for a toy linear regression example 889 |
| Figure 15.4.2 | Diagnosis plots for the Boston Housing example 891 |
| Figure 16.4.1 | Brownian motion interpolation Demo. 925 |
| Figure 18.1.1 | An illustration of a random walk mapping a sample point, ω , to a trajectory parameterized by time, where red trajectory sample point HHT, and blue trajectory has sample point THT. 1014 |
| Figure 18.3.1 | Sample trajectories of Brownian motion process. 1019 |
| Figure 18.4.1 | Variance of X_t in a Brownian bridge 1029 |
| Figure 18.5.1 | A typical realized trajectory from the Poisson process with jumps at T_1, T_2 , and T_3 . 1034 |
| Figure 19.4.1 | The variance function $Var[X(t)]$ for Brownian motion (red) and OU process(black) with $a = 0.5, \sigma = 1$. 1081 |
| Figure 19.4.2 | Representative trajectories from three OU processes with different k . k has the unit of inverse year. Mean level $\mu = 50$ and volatility $\sigma = 20$. 1083 |
| Figure 20.1.1 | Example Markov chains. Arrows and numbers are transition directions and probabilities. 1098 |
| Figure 20.1.2 | Markov chain diagram for a random walk on the state space \mathbb{Z} . 1099 |
| Figure 20.2.1 | Demonstration accessibility in a Markov chain. In chain (a), states A and B are accessible to each other or they can communicate. In chain (b), state A can access B but B cannot access A. 1101 |
| Figure 20.2.2 | Demonstration of partitioning state space by communicating classes. Green and orange states belong to different communicating classes. Note that a communicating class can consist of only one state. 1102 |

| | |
|---------------|---|
| Figure 20.2.3 | Classification of communicating classes into recurrent class and state space by communicating classes. Green states form a communicating class belonging to the transient class. Orange states form a communicating class belonging to the recurrent/closed/adsorbing class. 1109 |
| Figure 20.2.4 | Example periodic and aperiodic Markov chains. 1111 |
| Figure 21.1.1 | Example time series including a white noise process (upper left), a seasonable time series with periodicity 20, the US new privately owned housing [source] , and the US GDP time series [source] . 1146 |
| Figure 21.1.2 | Demonstration on using STL to decompose an example CO2 concentration time series. 1151 |
| Figure 21.2.1 | Example trajectories of AR(1) models ($X_t = aX_{t-1} + Z_t$) with different choices of a . 1156 |
| Figure 21.2.2 | Example trajectories of MA(1) models (upper) and MA(2) models (lower). 1163 |
| Figure 21.2.3 | Representative simulated trajectories for AR(1) process with coefficient 1, which forms a unit-root process, and with coefficient -1. 1172 |
| Figure 21.2.4 | The ACF and PACF correlogram for a white noise process. 1179 |
| Figure 21.2.5 | The ACF and PACF correlogram for an AR(1) process with coefficient 0.8. 1179 |
| Figure 21.2.6 | The ACF and PACF correlogram for an MA(1) process. 1180 |
| Figure 21.2.7 | The ACF and PACF correlogram for an ARMA(1,1) process. 1180 |
| Figure 21.2.8 | Diagnosis plot of residuals for AR(2) model estimation. 1189 |
| Figure 21.2.9 | Diagnosis plot of residuals for AR(1) model fitted to time series generated by AR(2) ground truth model. 1190 |
| Figure 21.4.1 | Stock index SP500 daily return between 2014 and 2019. 1203 |
| Figure 21.4.2 | Simulated representative trajectories from ARCH(1) model with coefficients $a = 0.9$ and $a = 0.5$. 1205 |
| Figure 22.1.1 | Scheme of a supervised learning task. Training samples are fed into a learning system to obtain an optimized model, which will be further used in a prediction system for regression and classification tasks. 1225 |
| Figure 22.1.2 | Input feature data type examples. 1225 |
| Figure 22.2.1 | A simple regression problem illustrating underfitting and overfitting. Solid lines are models, and points are samples. 1229 |
| Figure 22.2.2 | The commonly observed phenomenon of overfitting and underfitting in machine learning. 1230 |
| Figure 22.2.3 | The performance of different types of models as training proceeds. 1233 |

| | |
|---------------|---|
| Figure 22.3.1 | Regression loss functions: MSE Loss, MAE Loss, Huber loss ($\delta = 0.1, 1, 3$), and Log-Cosh Loss. 1237 |
| Figure 22.3.2 | Common classification loss functions. 1239 |
| Figure 22.3.3 | Scheme for ROC curves diagram. A and B demote ROC curves of different model. 1244 |
| Figure 22.4.1 | Scheme for cross-validation error calculation procedure. 1247 |
| Figure 22.4.2 | Hyperparameter search via turning regularization parameter λ . At large λ , heavy regularization causes underfitting; at small λ , insufficient regularization causes overfitting. 1249 |
| Figure 22.5.1 | Left: A sample of nine real-world time series reveals a diverse range of temporal patterns [4, 5]. Right: Examples of different classes of methods for quantifying the different types of structure, such as those seen in time series on the left: (i) distribution (the distribution of values in the time series, regardless of their sequential ordering); (ii) autocorrelation properties (how values of a time series are correlated to themselves through time); (iii) stationarity (how statistical properties change across a recording); (iv) entropy (measures of complexity or predictability of the time series quantified using information theory); and (v) nonlinear time-series analysis (methods that quantify nonlinear properties of the dynamics).[9] 1258 |
| Figure 23.1.1 | Correlation among features and the label MEDV. 1271 |
| Figure 23.1.2 | Pair plot among features and the label. 1272 |
| Figure 23.2.1 | Comparison different penalization: Lasso $L1$, elastic net, and Ridge $L2$. Orange regions are admissible set for parameter β . Contours are objective function value as a function of parameter β . Black solid circles are the minimizers $\hat{\beta}$ when there are no penalties, and red solid circles are the minimizers when penalties are applied. 1281 |
| Figure 23.2.2 | Penalized regression path in a toy regression problem. 1282 |
| Figure 23.3.1 | Linear regression with non-linear high order terms. The samples are generated via $y = 2x_1 + x_2 - 0.8x_1x_2 + 0.5x_1^2 + \epsilon$, where ϵ is noise. 1285 |
| Figure 23.3.2 | Linear model enhancement flowchart. 1285 |
| Figure 24.1.1 | Logistic regression for classification on the Iris data set. 1290 |
| Figure 24.1.2 | Pair plot analysis results on South Africa heart disease problem. 1298 |
| Figure 24.1.3 | $L1$ regularization path for South Africa heart disease problem. 1299 |
| Figure 24.1.4 | Logistic regression result with $L1$ penalty. 1301 |
| Figure 24.1.5 | Balanced accuracy score vs. inverse regularization strength in credit card fraud detection problem. 1302 |
| Figure 24.1.6 | Logistic regression coefficients corresponding to each class. 1303 |

| | |
|---------------|--|
| Figure 24.2.1 | Geometry of decision boundary in linear Gaussian discriminant model. 1307 |
| Figure 24.2.2 | Comparison of Gaussian LDA and Gaussian QDA on binary classification. Decision boundary of LDA is simply a line in 2D input space and unable to discriminate difficult cases. Gaussian discrimination can have richer decision boundary geometry. LDA using polynomial features is a special case of GDA. 1310 |
| Figure 24.2.3 | The decision boundary geometry of LDA and GDA can be understood via their decision functions 1311 |
| Figure 24.3.1 | The linear discriminant w that maximizes the separability for 2D sample points belonging to two classes. 1313 |
| Figure 24.3.2 | Fisher linear discriminate will fail to achieve class separability for complex data structures. 1317 |
| Figure 24.4.1 | Scheme of a hyperplane. 1322 |
| Figure 24.4.2 | Binary classification using the Perceptron learning algorithm. The hyperplane learned separates the two clusters. 1324 |
| Figure 24.5.1 | Left: existence of multiple separating hyperplanes in 2D binary classification problem. Right: hyperplanes with maximum margin. 1326 |
| Figure 24.5.2 | SVM classification with different regularization strength. Small C tends to emphasize the margin and ignore the outliers in the training data, while large C may tend to overfit the training data. 1329 |
| Figure 24.5.3 | SVM classification using Gaussian kernel. The original problem cannot be separated by linear kernel. 1334 |
| Figure 24.5.4 | Comparison of classification loss functions. 1336 |
| Figure 25.2.1 | Histogram of the features group by class label (fraud vs. genuine). 1351 |
| Figure 25.2.2 | Feature density similarity and predictive performance of model 1352 |
| Figure 26.2.1 | Binary classification via KNN algorithm with different choices $K = 1, 3, 5, 7$. Scattered points are training examples classified into two different classes (red and blue). Colored regions are corresponding decision boundaries. 1365 |
| Figure 27.2.1 | Demonstration of decision trees. 1373 |
| Figure 27.2.2 | Different types of impurity measure. Entropy function, Gini function and classification error function. 1376 |
| Figure 27.2.3 | Different splitting strategy when variables taking more than two discrete values. 1377 |
| Figure 27.2.4 | The visualization of the decision tree classifier for Iris data set. The tree grows until all examples are classified correctly. The splitting criterion is Gini impurity. 1382 |

| | |
|---------------|---|
| Figure 27.2.5 | The visualization of the decision tree classifier for Iris data set. The tree grows until examples in each node is smaller than 10. The splitting criterion is Gini impurity. 1383 |
| Figure 27.2.6 | The visualization of the decision tree classifier for Iris data set. The tree grows until examples in each node is smaller than 10. The splitting criterion is entropy and information gain. 1383 |
| Figure 27.3.1 | Demonstration of a tree and input space partitioning. 1386 |
| Figure 27.3.2 | 2D input space partitions cannot be represented by a regression tree. 1386 |
| Figure 27.3.3 | Regression tree demonstration in a toy example. 1388 |
| Figure 27.3.4 | Variable importance from regression tree in the Boston Housing Pricing problem. 1389 |
| Figure 28.1.1 | The correctness probability of a majority vote is greater than the correctness probability of individual votes when individual accuracy probability is greater than 0.5. 1394 |
| Figure 28.3.1 | Illustration of adaptive boosting where sample weights are adjusted iteratively based on the classification error. 1401 |
| Figure 29.1.1 | Demonstration of SVD for matrices of two different shapes. The dashed lines highlight the compact form SVD. 1423 |
| Figure 29.1.2 | Principal components for 2D samples. 1427 |
| Figure 29.2.1 | Singular value spectrum of LSA on 20-news-group text data. 1442 |
| Figure 29.2.2 | A simplified illustration of the latent factor approach, which characterizes both users and movies using two axes—male versus female and serious versus escapist. [9] 1447 |
| Figure 29.3.1 | Triangle and Tetrahydron reconstructed from distance matrix by MDS method. 1454 |
| Figure 29.3.2 | Application of MDS, based on Euclidean distance, to Swiss Roll data set cannot fully reveal of the global structure. 1455 |
| Figure 29.3.3 | Isomap analysis of MNIST dataset. 1466 |
| Figure 29.3.4 | Isomap analysis of digit '5' in MNIST dataset 1467 |
| Figure 29.4.1 | Demonstration of k-means clustering on a data set with two blobs. 1469 |
| Figure 29.4.2 | K-means performance can be affected by a number of factors, including incorrect number of clusters/blobs, non-spherical clusters/blobs, clusters/blobs with unequal variance, and bad initial cluster centers. 1471 |
| Figure 29.4.3 | Clustering comparison between Kmeans and Kmeans++. 1472 |
| Figure 29.4.4 | DBSCAN demo with different ϵ . 1474 |
| Figure 29.4.5 | Spectral clustering demo. 1476 |
| Figure 29.4.6 | Clustering comparison between K-means and GMM. 1480 |
| Figure 29.4.7 | K-means application to image segmentation. 1481 |

| | |
|---------------|--|
| Figure 30.1.1 | Scheme of an artificial neuron. 1487 |
| Figure 30.1.2 | Common activation functions in artificial neural networks. 1488 |
| Figure 30.1.3 | Scheme for an artificial neural network. 1489 |
| Figure 30.1.4 | A four-layer feed-forward neural network. 1489 |
| Figure 30.1.5 | A three-layer feed-forward neural network with three output units. 1490 |
| Figure 30.1.6 | A four-layer feed-forward neural network. 1492 |
| Figure 30.2.1 | An example saddle point at $(0,0,0)$, which locally minimizes the x direction but maximizes the y direction.. 1499 |
| Figure 30.2.2 | SGD without momentum and with momentum. SGD with momentum can accumulate gradient/velocity in horizontal direction and move faster towards the minimum located at the center. 1503 |
| Figure 30.3.1 | Dropout technique for a simple feedforward neural networks. The original network (left) and the neural network after some neurons being dropped out (right). 1511 |
| Figure 30.4.1 | Neural network architecture for polynomial regression. 1513 |
| Figure 30.4.2 | Polynomial regression with degree $d = 1, 3, 6, 10$. 1514 |
| Figure 30.4.3 | Visualization of first layer weight for a one-layer linear multi-class classification neural network 1515 |
| Figure 30.4.4 | Example images from the Fashion MNIST dataset. 1516 |
| Figure 30.4.5 | The confusion matrix from fashion MNIST classification results. 1517 |
| Figure 30.4.6 | Classification results for a set of randomly selected samples. 1518 |
| Figure 30.4.7 | (a) Embedding layer maps large, sparse one-hot vectors to short, dense vectors. (b) Example of low dimensional embeddings that capture semantic meanings. 1519 |
| Figure 30.4.8 | (a) The Skip-gram architecture that predicts surrounding words given the central word. (b) The CBOW architecture that predicts the central word given its surrounding context words. The one-hot vector has size V ; the dense vector has length $D \ll V$. Also note that no nonlinearity activation is applied between input and hidden layer. 1520 |
| Figure 30.4.9 | A feed-forward neural network architecture for sentiment analysis. 1524 |
| Figure 30.5.1 | Comparison of receptive fields in fully-connected layer and local-connected layer in CNN. Credit 1528 |
| Figure 30.5.2 | Demo for one kernel 'convoluting' with an input image. 1529 |
| Figure 30.5.3 | Pooling layer demo. 1530 |
| Figure 30.5.4 | A typical CNN architecture for image classification tasks. 1531 |
| Figure 30.5.5 | Scheme for LeNet [24] 1531 |
| Figure 30.5.6 | Architecture of AlexNet. 1532 |

| | |
|----------------|--|
| Figure 30.5.7 | A typical VGG architecture: VGG-19 scheme. 1532 |
| Figure 30.5.8 | The depth of the configurations increases from the left (A) to the right (E), as more layers are added (the added layers are shown in bold). The convolutional layer parameters are denoted as “conv-receptive field size-number of channels” [26] 1533 |
| Figure 30.5.9 | Training error (left) and testing error (right) on CIFAR-10 with 20-layer and 56-layer vanilla CNN networks. The deeper network has both higher training error and testing error.[27] 1534 |
| Figure 30.5.10 | Scheme for a residual block. 1534 |
| Figure 30.5.11 | Scheme of a 18 layer ResNet 1535 |
| Figure 30.6.1 | Example images from CIFAR10 image dataset. 1536 |
| Figure 30.6.2 | Visualization of convolution layer. 1538 |
| Figure 30.6.3 | Grad-CAM method applied to understand image classification. Middle and right are class-discriminative localization map superimposed onto the original image. 1539 |
| Figure 30.6.4 | A CNN based autoencoder. An autoencoder consists of an encoder that transforms high-dimensional image into a low-dimensional code and a decoder that unfolds the code to reconstruct the image. 1541 |
| Figure 30.6.5 | Comparison of reconstruction performance on random samples from MNIST data set. Top row is the original data. Middle row is autoencoder result based on a 49 dimensional code. Bottom row is PCA result based on a 50 dimensional code. 1541 |
| Figure 30.6.6 | Denoising autoencoders applied to remove the noise in the MNIST dataset. 1542 |
| Figure 30.6.7 | Demonstration of neural style transfer with Van Gogh painting style. 1544 |
| Figure 30.6.8 | Demonstration of neural style transfer with Picasso painting style. 1545 |
| Figure 30.6.9 | Application of CNN in deep reinforcement learning in Q learning approach and Actor-Critic approach. Two streams of sensory inputs are fed to the neural network, including a pixel image of the robot’s neighborhood fed into a convolutional layer and the target’s position fed into a fully connected layer. 1547 |
| Figure 30.6.10 | Demonstration of a CNN filter applying to a sentence to produce a one-dimensional feature vector. 1548 |
| Figure 30.6.11 | CNN for sentence classification proposed in [43]. 1549 |
| Figure 30.7.1 | Scheme of recurrent units in a neural network (left). Recurrent neural network can be unrolled (right) 1551 |
| Figure 30.7.2 | Scheme for backpropogation through time in a simple RNN. Red arrows are backpropogation directions. 1552 |

| | | |
|---------------|--|------|
| Figure 30.7.3 | Scheme of an LSTM cell. Modified from [45, p. 149]. | 1554 |
| Figure 30.7.4 | Scheme of a GRU cell. Modified from [45, p. 152]. | 1557 |
| Figure 30.7.5 | Different types of units in RNNs: (a) Vanilla RNN cell. (b) LSTM cell. (c) GRU cell. Credit | 1558 |
| Figure 30.7.6 | Typical RNN connection to the output layer. | 1559 |
| Figure 30.7.7 | Scheme for stacked RNN. | 1560 |
| Figure 30.7.8 | Scheme for bidirectional RNN. | 1560 |
| Figure 30.8.1 | RNN architecture for time series prediction . (a) In the training phase, RNN are updated by minimizing the next step prediction error. (b) In the prediction phase, trained RNN is used to sequentially predict next step state value based on preceding predicted state value. | 1562 |
| Figure 30.8.2 | RNN one-step forward and multiple forward prediction performance for Sine time series. | 1563 |
| Figure 30.8.3 | RNN architecture for time series prediction with covariates. (a) In the training phase, RNN are updated by minimizing the next step prediction error. (b) In the prediction phase, trained RNN is used to sequentially predict next step state value based on preceding predicted state value. Note that covariate time series are assumed available for all time steps. | 1564 |
| Figure 30.8.4 | DeepAR model architecture for time series prediction. Outputs are the parameters characterizing the condition distribution of x_{t+1} conditioned on histories of x_t and z_t . (a) In the training phase, RNN are updated by minimizing the negative log likelihood function. (b) In the prediction phase, a predicted \hat{x}_{t+1} are sampled from predicted conditional distribution and then used to predict next step conditional distribution. | 1565 |
| Figure 30.8.5 | A RNN architecture for MNIST image recognition. | 1567 |
| Figure 30.8.6 | RNN architecture for sentiment analysis. | 1568 |
| Figure 30.8.7 | A RNN for character-level word classification. | 1569 |
| Figure 30.8.8 | A RNN for character-level language model. During the training session, one-hot coded characters are directly fed into RNN and predict the next character in the word. | 1570 |
| Figure 30.8.9 | A RNN for character-level language model. During the word generation session, the network starts with a user input character and continues the generation process with predicted character from the previous step. | 1571 |
| Figure 30.9.1 | Seq2seq modeling for language transformation. The input and output sequences might have different lengths, and not in synchrony. | 1573 |

- Figure 30.9.2 The Encoder-decoder architecture for seq2seq language modeling. The input sequence is fed into the encoder RNN and terminated by an explicit <EOS> symbol. Then the decoder RNN starts with the context vector and the final prediction of the encoder to generate an output sequence until an explicit <EOS> symbol is produced. [1574](#)
- Figure 30.9.3 The Encoder-decoder architecture with attention mechanism for seq2seq modeling. During the encoding phase, all hidden states, rather than the final one, are saved to construct different context vectors via linear combination for the decoding stage. During the decoding phase, relevant context vectors are constructed and fed into the each hidden states in the decoder. [1576](#)
- Figure 30.9.4 A bidirectional RNN encoder system with attention mechanism. [\[51\]](#) [1577](#)
- Figure 30.10.1 Scheme for a canonical GAN. The discriminator is trained to distinguish between real and fake image, while the generator is trained to generate realistic image to 'fool' the discriminator. The generator usually uses a decoder-like neural network structure that generate a high-dimensional data from a sample point in the low-dimensional latent space. [1579](#)
- Figure 30.10.2 Generated image samples from a GAN consisting of feed-forward networks. [1581](#)
- Figure 30.10.3 DCGAN generator network architecture. A 100 dimensional uniform noise is passing through a series of fractionally-strided convolutions then is converted into a final image. Notably, no fully connected or pooling layers are used [\[56\]](#) [1584](#)
- Figure 30.10.4 Understanding DCGAN. Each row represents the image generated as we interpolate a random point z in the latent space. Images show smooth transitions from one bedroom and another bedroom.[\[56\]](#) [1584](#)
- Figure 30.10.5 Scheme for a conditional GAN. The discriminator is trained to distinguish between real and fake image given external label information, while the generator is trained to generate realistic image to 'fool' the discriminator given external label information. The generator usually uses a decoder-like neural network structure that generate a high-dimensional data from a sample point in the low-dimensional latent space. [1586](#)
- Figure 32.1.1 Policy iteration involves iteratively carrying out policy evaluation and policy improvement procedures. [1614](#)

- Figure 32.2.1 One core component in reinforcement learning is agent environment interaction. The agent takes actions based on observations on the environment and a decision-making module that maps observations to action. The environment model updates system state and provides rewards according to the action [1622](#)
- Figure 32.2.2 Scheme of the Atari game *breakout*. [1623](#)
- Figure 32.2.3 Policy evaluation and policy improvement framework in the context reinforcement learning. [1624](#)
- Figure 32.3.1 Neural network parameterization for a Gaussian policy (a) and (b) a generic stochastic policy. [1652](#)
- Figure 32.4.1 A typical feed-forward neural network used in NFQ to approximate the Q function. [1659](#)
- Figure 32.4.2 The network architecture for canonical deep Q learning. The network takes a state or an observation, denoted by s , as the input, and outputs multiple values corresponding $Q(s, a)$. The number of outputs [1661](#)
- Figure 32.4.3 A typical single stream Q -network (top) in deep Q learning and a dueling Q -network (bottom). The dueling network has two streams to separately estimate (scalar) state-value and the advantage function for each action; the green output module synthesize the Q value from two streams. Both networks output Q -values for each action. [\[10\]](#). [1664](#)
- Figure 32.4.4 In a DQRN, recurrent layers are usually placed on the last layer before output. Unlike canonical DQN, we need to feed sequence of observation into the network and finally output Q values. Above scheme only unfolds to two steps.[\[11\]](#) [1665](#)
- Figure 32.4.5 An example architecture that implements the asynchronous reinforcement learning paradigm. Multiple agents interact with multiple instances of environments in parallel. Agents collect experiences to train a globally shared network that learns control policies. [1666](#)
- Figure 32.4.6 A comparison between a typical deep Q learning network (left) and a typical universal value function approximator network (right). [1668](#)
- Figure 32.4.7 A typical deep neural network architecture for DDPG reinforcement learning. The actor network outputs control policy; the critic network outputs estimate of Q function. Through back-prorogation, the critic network improves estimation accuracy and the actor network improves policy. [1670](#)
- Figure 32.5.1 Illustration of planning curriculum on a swiss roll manifold. Red targets can generating along the path connecting from an intended start point to the target goal position. [1687](#)

Figure 32.5.2 One representative trajectory on the curved surface via a control policy learned via curriculum learning on a low-dimensional manifold. [1688](#)

LIST OF TABLES

| | | |
|--------------|---|------|
| Table 13.7.1 | Test on mean with known variance σ^2 | 708 |
| Table 13.7.2 | Test on mean with unknown variance σ^2 | 710 |
| Table 13.7.3 | Test on variance | 710 |
| Table 13.7.4 | Test on variance comparison between two samples | 711 |
| Table 14.2.1 | Eigenvectors and eigenvalues for swap rate daily change | 739 |
| Table 14.5.1 | statistics on Fama-French 3 factors from July 1963 to Dec. 1991. | 796 |
| Table 14.5.2 | AAPL stock return modeled by the Fama-French 3 factor model. | 798 |
| Table 20.2.1 | Summary of Markov chain state property[3, p. 140] | 1113 |
| Table 21.2.1 | Summary of PACF and ACF for AR, MA, and ARMA processes. | 1179 |
| Table 23.1.1 | Linear regression results. | 1273 |
| Table 24.1.1 | Logistic regression results on South Africa heart disease problem. | 1299 |
| Table 29.2.1 | Most frequent words in the top 8 topics | 1443 |
| Table 29.2.2 | Rating matrix or utility matrix, where each row is a user's ratings for different movies. SW ₁ , SW ₂ are Star wars episodes; HP ₁ and HP ₂ are Harry Potter episodes; TW is Twilight; BM is Batman. | 1445 |
| Table 32.2.1 | Estimating cumulative rewards $G_t^{(n)}$ of different steps n as the target for value function V . $G^{(1)}$ corresponds to temporal-difference TD(0) and $G^{(\infty)}$ corresponds to Monte-Carlo estimation. If the process terminates at K and $K < n$, then we use $G_t^{(n)} = G_t^{(K)}$. Trajectories are generated under policy π . | 1634 |
| Table A.9.1 | Closed Newton-Cotes Formula | 1711 |

CONTENTS

i mathematical foundations

| | | |
|---------|---|----|
| 1 | SETS, SEQUENCES, AND SERIES | 2 |
| 1.1 | Sets | 4 |
| 1.1.1 | Definitions and basic properties | 4 |
| 1.1.2 | DeMorgan's Law | 5 |
| 1.1.3 | Set equivalence and partition | 5 |
| 1.1.4 | Countability | 6 |
| 1.2 | Functions | 8 |
| 1.2.1 | Basic concepts | 8 |
| 1.2.2 | Inverse image vs. inverse function | 8 |
| 1.2.3 | Set operations in function mapping | 9 |
| 1.2.4 | Parameter change of function | 9 |
| 1.3 | Real numbers | 10 |
| 1.3.1 | Rational numbers | 10 |
| 1.3.2 | Dense subset | 10 |
| 1.3.3 | Axiom of completeness | 11 |
| 1.4 | Sequence in \mathbb{R} | 14 |
| 1.4.1 | Basics | 14 |
| 1.4.2 | Cauchy criterion | 16 |
| 1.4.3 | Sequence characterization of a dense subset | 17 |
| 1.5 | Monotone sequence | 19 |
| 1.5.1 | Fundamentals | 19 |
| 1.5.2 | Applications | 19 |
| 1.6 | Subsequence and limits | 23 |
| 1.6.1 | Subsequence | 23 |
| 1.6.2 | Bolzano-Weierstrass theorem | 23 |
| 1.6.3 | Subsequence limits | 24 |
| 1.7 | Infinite series | 27 |
| 1.7.1 | Fundamental results | 27 |
| 1.7.2 | Tests for convergence | 28 |
| 1.7.3 | Inequalities and l_2 series | 30 |
| 1.7.3.1 | Holder's and Minkowski's inequality | 30 |
| 1.7.3.2 | Cauchy-Schwarz inequality | 31 |
| 1.7.4 | Alternating series | 32 |

| | | |
|---------|--|----|
| 1.8 | Notes on bibliography | 34 |
| 2 | METRIC SPACE | 36 |
| 2.1 | Metric space | 38 |
| 2.1.1 | Definitions | 38 |
| 2.1.2 | metric space vs. normed (vector) space vs. Banach space | 40 |
| 2.2 | Sequences in metric space | 41 |
| 2.3 | Closed sets & open sets in metric space | 43 |
| 2.3.1 | Closed set | 43 |
| 2.3.2 | Open sets | 43 |
| 2.3.3 | Further characterization and properties | 44 |
| 2.3.4 | Open and closed sets in \mathbb{R}^n | 46 |
| 2.4 | Compact sets | 48 |
| 2.4.1 | Basic concepts | 48 |
| 2.4.1.1 | closed set vs. compact set | 49 |
| 2.4.2 | Compact sets in \mathbb{R}^N | 49 |
| 2.4.3 | The Heine-Borel Theorem and boundedness of continuous function | 50 |
| 2.5 | Completeness of metric space | 51 |
| 2.5.1 | Sequence and completeness | 51 |
| 2.5.2 | Completeness of \mathbb{R}^n | 51 |
| 2.6 | Topology space | 53 |
| 2.6.1 | Definitions | 53 |
| 2.6.2 | Continuous function, Homeomorphism in topological space | 54 |
| 2.6.3 | Subspaces of topological space | 54 |
| 2.7 | Notes on bibliography | 56 |
| 3 | ADVANCED CALCULUS | 58 |
| 3.1 | Continuous functions | 61 |
| 3.1.1 | Continuous function on \mathbb{R} | 61 |
| 3.1.1.1 | Basics | 61 |
| 3.1.1.2 | Continuity and inverse | 64 |
| 3.1.2 | Continuous function in metric space | 65 |
| 3.1.3 | Boundedness and extreme value theorems | 66 |
| 3.1.4 | More on extreme values | 68 |
| 3.1.5 | Curves and surfaces | 68 |
| 3.1.5.1 | Curvature | 71 |
| 3.1.5.2 | Surfaces | 71 |
| 3.2 | Uniform continuity | 74 |
| 3.2.1 | Uniform continuity on real line | 74 |
| 3.2.1.1 | Concepts | 74 |
| 3.2.1.2 | Lipschitz continuity | 76 |
| 3.2.2 | Uniform continuity on metric space | 79 |

| | | |
|---------|--|-----|
| 3.2.3 | Locally and globally Lipschitz continuous | 80 |
| 3.3 | Differentiation | 82 |
| 3.3.1 | Differential function concept | 82 |
| 3.3.2 | Differential rules | 83 |
| 3.3.3 | Mean value theorem | 85 |
| 3.4 | Function sequence and series | 88 |
| 3.4.1 | Pointwise convergence, uniform convergence | 88 |
| 3.4.2 | Properties of uniform convergence | 89 |
| 3.4.2.1 | Uniform convergence preserve continuity | 89 |
| 3.4.2.2 | Exchange limits and integration | 90 |
| 3.4.2.3 | Exchange limits and differential | 90 |
| 3.4.2.4 | Linearity of uniform convergence | 90 |
| 3.5 | Power series | 92 |
| 3.5.1 | Fundamentals | 92 |
| 3.5.2 | Term-by-term operation | 94 |
| 3.5.3 | Power series and analytic function | 94 |
| 3.5.4 | Approximation by polynomials | 95 |
| 3.6 | Taylor polynomial and Taylor series | 97 |
| 3.6.1 | Taylor polynomial and approximation | 97 |
| 3.6.2 | Taylor series and Taylor's theorem | 99 |
| 3.6.3 | Common Taylor series | 101 |
| 3.6.4 | Useful approximations | 103 |
| 3.7 | Riemann Integral Theory | 105 |
| 3.7.1 | Construction of Riemann integral | 105 |
| 3.7.2 | Riemann integrability | 106 |
| 3.7.2.1 | Basics | 106 |
| 3.7.2.2 | Lebesgue characterization of integrability | 107 |
| 3.7.2.3 | limits and integrability | 107 |
| 3.7.2.4 | Algebraic properties | 108 |
| 3.7.3 | First Fundamental Theorem of Calculus | 109 |
| 3.7.4 | Second Fundamental Theorem of Calculus | 109 |
| 3.7.4.1 | Fundamentals | 109 |
| 3.7.4.2 | Differentiating definite integrals | 112 |
| 3.7.4.3 | Application to differential equation | 113 |
| 3.7.5 | Essential theorems | 114 |
| 3.7.6 | Integration rules | 115 |
| 3.7.7 | Improper Riemann integrals | 115 |
| 3.8 | Basic measure theory | 118 |
| 3.8.1 | Measurable space | 118 |
| 3.8.1.1 | σ algebra | 118 |
| 3.8.1.2 | Measurable space and positive measure | 119 |

| | | | |
|------|---------|--|-----|
| | 3.8.1.3 | Borel algebra and Lebesgue measure | 119 |
| | 3.8.2 | Measurable functions and properties | 121 |
| | 3.8.2.1 | Measurable function and measurability | 121 |
| | 3.8.2.2 | Properties | 122 |
| | 3.8.3 | Convergence of measurable functions | 124 |
| | 3.8.4 | Almost everywhere convergence | 124 |
| 3.9 | | Lebesgue integral | 126 |
| | 3.9.1 | Simple function and its Lebesgue integral | 126 |
| | 3.9.2 | Lebesgue integral of measurable functions | 128 |
| | 3.9.2.1 | Integral of non-negative functions | 128 |
| | 3.9.2.2 | Integral of general functions | 130 |
| | 3.9.3 | Riemann vs. Lebesgue integrals | 131 |
| | 3.9.4 | Convergence theorems | 131 |
| | 3.9.4.1 | Applications | 133 |
| 3.10 | | Notes on bibliography | 136 |
| 4 | | LINEAR ALGEBRA AND MATRIX ANALYSIS | 138 |
| 4.1 | | Theory for system of linear equations | 143 |
| | 4.1.1 | Overview | 143 |
| | 4.1.2 | Homogeneous systems | 143 |
| | 4.1.3 | Non-homogeneous systems | 144 |
| | 4.1.4 | Overdetermined vs. underdetermined systems | 145 |
| | 4.1.5 | Solution methods | 146 |
| | 4.1.6 | Error bounds in numerical solutions | 151 |
| | 4.1.6.1 | Condition number | 151 |
| | 4.1.6.2 | Error bounds | 151 |
| 4.2 | | Vector space theory | 153 |
| | 4.2.1 | Vector space | 153 |
| | 4.2.2 | Subspace | 154 |
| | 4.2.3 | Sum and direct sum | 155 |
| | 4.2.4 | Basis and dimensions | 157 |
| | 4.2.5 | Complex vector space vs. real vector space | 159 |
| 4.3 | | Linear maps & linear operators | 161 |
| | 4.3.1 | Basic concepts of linear maps | 161 |
| | 4.3.2 | Fundamental theorem of linear maps | 162 |
| | 4.3.3 | Isomorphism | 164 |
| | 4.3.4 | Coordinate map properties | 166 |
| | 4.3.5 | Change of basis and similarity | 167 |
| | 4.3.5.1 | Change of basis for coordinate vector | 167 |
| | 4.3.5.2 | Change of basis for linear maps | 167 |
| | 4.3.6 | Linear maps and matrices | 167 |
| | 4.3.6.1 | Similarity | 169 |

| | | |
|---------|--|-----|
| 4.4 | Fundamental theorems of ranks and linear algebra | 170 |
| 4.4.1 | Basics of ranks | 170 |
| 4.4.2 | Fundamental theorem of ranks | 171 |
| 4.4.3 | Fundamental theorem of linear algebra | 172 |
| 4.5 | Complementary subspaces and projections | 174 |
| 4.5.1 | General complementary subspaces | 174 |
| 4.5.2 | Orthogonal complementary spaces and projections | 176 |
| 4.5.3 | Decomposition of orthogonal projectors | 180 |
| 4.6 | Orthonormal basis and projections | 183 |
| 4.6.1 | Gram-Schmidt Procedure | 183 |
| 4.6.2 | Orthogonal-triangular decomposition | 183 |
| 4.6.3 | Orthonormal basis for linear operators | 184 |
| 4.6.4 | Riesz representation theorem | 185 |
| 4.7 | Eigenvectors and eigenvalues of Matrices: general theory | 186 |
| 4.7.1 | Existence and properties of eigenvalues | 186 |
| 4.7.2 | Properties of eigenvectors | 188 |
| 4.7.3 | Right and left eigenvectors | 189 |
| 4.7.4 | Diagonalizable matrices | 190 |
| 4.8 | Eigenvalue and eigenvectors of matrices: case studies | 193 |
| 4.8.1 | Real diagonalizable matrix | 193 |
| 4.8.2 | Real symmetric matrix | 194 |
| 4.8.2.1 | Spectral properties | 194 |
| 4.8.2.2 | Rayleigh quotients | 196 |
| 4.8.2.3 | Pointcare inequality | 199 |
| 4.8.3 | Hermitian matrix | 200 |
| 4.8.4 | Matrix congruence | 202 |
| 4.8.5 | Complex symmetric matrix | 203 |
| 4.8.6 | Unitary, orthonormal & rotation matrix | 203 |
| 4.9 | Singular Value Decomposition theory | 205 |
| 4.9.1 | SVD fundamentals | 205 |
| 4.9.2 | SVD and matrix norm | 207 |
| 4.9.3 | SVD vs. eigendecomposition | 208 |
| 4.9.4 | SVD low rank approximation | 209 |
| 4.9.4.1 | Frobenius norm low rank approximation | 209 |
| 4.9.4.2 | Two-norm low rank approximation | 211 |
| 4.10 | Generalized eigenvectors and Jordan normal forms | 213 |
| 4.10.1 | Generalized eigenvectors | 213 |
| 4.10.2 | Upper triangle matrix and nilpotent matrix | 216 |
| 4.10.3 | Jordan normal forms | 218 |
| 4.11 | Matrix factorization | 222 |
| 4.11.1 | Orthogonal-triangular decomposition | 222 |

| | | |
|----------|--|-----|
| 4.11.2 | LU decomposition | 223 |
| 4.11.3 | Cholesky decomposition | 223 |
| 4.12 | Positive definite matrices and quadratic forms | 225 |
| 4.12.1 | Quadratic forms | 225 |
| 4.12.2 | Real symmetric non-negative definite matrix | 226 |
| 4.12.2.1 | Characterization | 226 |
| 4.12.2.2 | Decomposition and transformation | 229 |
| 4.12.2.3 | Matrix square root | 230 |
| 4.12.2.4 | Maximization of quadratic forms | 231 |
| 4.12.2.5 | Gramian matrix | 233 |
| 4.12.3 | Completing the square | 234 |
| 4.13 | Matrix norm and spectral estimation | 235 |
| 4.13.1 | Basics | 235 |
| 4.13.2 | Singularity from matrix norm and spectral radius | 236 |
| 4.13.3 | Gerschgorin theorem | 237 |
| 4.13.4 | Irreducible matrix and stronger results | 238 |
| 4.14 | Pseudoinverse of matrix | 239 |
| 4.14.1 | Pseudoinverse for full rank system | 239 |
| 4.14.2 | Pseudoinverse for general matrix | 241 |
| 4.14.3 | Application in linear systems | 243 |
| 4.15 | Multilinear forms | 246 |
| 4.15.1 | Bilinear forms | 246 |
| 4.15.2 | Multilinear forms | 247 |
| 4.16 | Determinant | 250 |
| 4.16.1 | Basic properties | 250 |
| 4.16.2 | Vandermonde matrix and determinant | 256 |
| 4.17 | Numerical iteration analysis | 258 |
| 4.17.1 | Numerical linear equation solution | 258 |
| 4.17.1.1 | Goals and general principles | 258 |
| 4.17.1.2 | Jacobi algorithm | 258 |
| 4.17.1.3 | Gauss Seidel algorithm | 259 |
| 4.17.2 | Power method for eigen-decomposition | 259 |
| 4.18 | Appendix: supplemental results for polynomials | 262 |
| 4.18.1 | Basics | 262 |
| 4.18.2 | Factorization of polynomial over \mathbb{C} | 263 |
| 4.18.3 | Factorization of polynomial over \mathbb{R} | 264 |
| 4.19 | Notes on bibliography | 266 |
| 5 | BASIC FUNCTIONAL ANALYSIS | 268 |
| 5.1 | Normed vector space | 270 |
| 5.1.1 | Basic properties | 270 |
| 5.1.2 | Equivalence of norms | 272 |

| | | |
|---------|--|-----|
| 5.2 | Contraction mapping and fixed point theorems | 274 |
| 5.2.1 | Complete normed space (Banach space) | 274 |
| 5.2.2 | Contraction mapping | 275 |
| 5.2.3 | Banach fixed point theorem | 276 |
| 5.2.4 | Applications in root finding | 277 |
| 5.2.5 | Application to numerical linear equations | 277 |
| 5.2.6 | Applications to integral and differential equations | 278 |
| 5.3 | Inner product space and Hilbert space | 281 |
| 5.3.1 | Inner product space (pre-Hilbert space) and Hilbert space | 281 |
| 5.3.1.1 | Foundations | 281 |
| 5.3.2 | Hilbert spaces | 283 |
| 5.3.2.1 | Basics | 283 |
| 5.3.3 | Orthogonal decomposition of Hilbert spaces | 284 |
| 5.3.3.1 | Orthogonality | 284 |
| 5.3.4 | Projection and orthogonal decomposition | 285 |
| 5.4 | Approximations in Hilbert space | 288 |
| 5.4.1 | Approximation via projection | 288 |
| 5.4.2 | Application examples | 289 |
| 5.4.2.1 | Orthogonal projection and normal equations in \mathbb{R}^n | 289 |
| 5.4.2.2 | Approximation by continuous polynomials | 291 |
| 5.4.2.3 | Legendre polynomial via Gram-Schmidt process | 292 |
| 5.5 | Orthonormal systems | 293 |
| 5.5.1 | Basic definitions | 293 |
| 5.5.2 | Gram-Schmidt process | 293 |
| 5.5.3 | Properties of orthonormal systems | 293 |
| 5.5.4 | Orthonormal expansion in Hilbert space | 295 |
| 5.5.5 | Complete orthonormal system | 296 |
| 5.5.5.1 | Weierstrass approximation theorem for polynomials | 298 |
| 5.5.5.2 | Examples of complete orthonormal function set | 298 |
| 5.6 | Theory for trigonometric Fourier Series | 300 |
| 5.6.1 | Basic definitions | 300 |
| 5.6.2 | Completeness of Fourier series | 302 |
| 5.6.3 | Complex representation | 303 |
| 5.7 | Fourier transform | 305 |
| 5.7.1 | Definitions and basic concepts | 305 |
| 5.7.2 | Convolution theorem | 307 |
| 5.7.3 | Fourier transform and Fourier series | 308 |
| 5.7.4 | Discrete Fourier transform | 309 |
| 5.7.4.1 | Properties | 309 |
| 5.8 | Notes on bibliography | 312 |

ii mathematical optimization methods

| | | |
|---------|---|-----|
| 6 | UNCONSTRAINED NONLINEAR OPTIMIZATION | 315 |
| 6.1 | Optimality conditions | 317 |
| 6.1.1 | Optimality concepts | 317 |
| 6.1.2 | Necessary and sufficient conditions | 319 |
| 6.1.3 | Special case: unconstrained quadratic programming | 322 |
| 6.2 | Line search method | 324 |
| 6.2.1 | A generic algorithm | 324 |
| 6.2.2 | Theory and computation of descent directions | 325 |
| 6.2.2.1 | Gradient descent direction and properties | 325 |
| 6.2.2.2 | Curvature-modified descent direction | 326 |
| 6.2.2.3 | Quasi-Newton method | 328 |
| 6.2.2.4 | Subspace optimization in quadratic forms | 330 |
| 6.2.3 | Theory and computation of step length | 331 |
| 6.2.3.1 | Overview | 331 |
| 6.2.3.2 | Lipschitz bounded convex functions | 331 |
| 6.2.3.3 | Backtracking-Armijo step size search | 334 |
| 6.2.3.4 | Wolfe condition | 336 |
| 6.2.4 | Complete algorithms | 337 |
| 6.3 | Trust region method | 339 |
| 6.3.1 | Motivation and the framework | 339 |
| 6.3.2 | Cauchy point method | 340 |
| 6.3.3 | Exact solution method | 342 |
| 6.3.4 | Approximate method | 342 |
| 6.4 | Conjugate gradient method | 345 |
| 6.4.1 | Motivating problems | 345 |
| 6.4.2 | Theory conjugate direction | 346 |
| 6.4.3 | Linear conjugate gradient algorithm | 347 |
| 6.5 | Least square problems | 350 |
| 6.5.1 | Linear least square theory and algorithm | 350 |
| 6.5.1.1 | Linear least square problems | 350 |
| 6.5.1.2 | SVD methods | 351 |
| 6.5.1.3 | Extension to L^p norm optimization | 351 |
| 6.5.2 | nonlinear least square problem | 352 |
| 6.5.3 | Line search Gauss-Newton method | 353 |
| 6.5.4 | Trust region method | 355 |
| 6.5.5 | Application: roots for nonlinear equation | 355 |
| 6.6 | Notes on bibliography | 357 |
| 7 | CONSTRAINED NONLINEAR OPTIMIZATION | 359 |
| 7.1 | Quadratic optimization I: equality constraints | 361 |
| 7.1.1 | Problem formulation | 361 |

| | | |
|---------|---|-----|
| 7.1.2 | Optimality condition | 361 |
| 7.1.2.1 | General case | 361 |
| 7.1.2.2 | Positive semi-definitive quadratic programming | 364 |
| 7.1.3 | Solving KKT systems | 365 |
| 7.1.3.1 | Factorization approach | 365 |
| 7.1.3.2 | Range space approach | 366 |
| 7.1.4 | Linear least square with linear constraints | 366 |
| 7.1.4.1 | Least norm problem | 366 |
| 7.1.5 | Application: Markovitz Portfolio Optimization Model | 368 |
| 7.2 | Quadratic optimization II: inequality constraints | 371 |
| 7.2.1 | Problem formulation | 371 |
| 7.2.2 | Optimality conditions | 371 |
| 7.2.2.1 | Pure inequality case | 371 |
| 7.2.2.2 | General constrained optimization | 373 |
| 7.2.2.3 | Positive semi-definitive quadratic programming | 374 |
| 7.2.3 | Primal active-set method | 375 |
| 7.2.4 | Gradient projection method | 380 |
| 7.2.5 | Dual convex quadratic programming | 381 |
| 7.3 | General equality constrained optimization | 383 |
| 7.3.1 | Feasible path and optimality | 383 |
| 7.3.2 | Constraint qualification and Lagrange theory | 384 |
| 7.3.3 | Second order condition | 387 |
| 7.4 | General inequality constrained optimization | 391 |
| 7.4.1 | Feasible path and optimality | 391 |
| 7.4.2 | Constraint qualifications and KKT conditions | 392 |
| 7.4.3 | Second order conditions | 396 |
| 7.5 | Envelope theorem and sensitive analysis | 400 |
| 7.6 | Notes on bibliography | 403 |
| 8 | LINEAR OPTIMIZATION | 405 |
| 8.1 | Equality constrained linear programming | 406 |
| 8.2 | Inequality constrained linear programming | 408 |
| 8.2.1 | Linear optimization with inequality constraints | 408 |
| 8.2.2 | Geometry of linear programming | 408 |
| 8.2.3 | Optimality property and condition | 410 |
| 8.2.4 | Standard form of linear programming | 411 |
| 8.2.5 | Application examples | 412 |
| 8.3 | Linear programming geometry and simplex algorithm | 413 |
| 8.3.1 | Geometrical approach to linear programming | 413 |
| 8.3.1.1 | Overview | 413 |
| 8.3.1.2 | Vertex and optimality | 413 |
| 8.3.1.3 | Descent direction at a vertex | 417 |

| | | |
|---------|--|-----|
| 8.3.1.4 | Stepping along a descent direction | 418 |
| 8.3.2 | The simplex algorithm | 419 |
| 8.4 | Interior point method | 420 |
| 8.4.1 | Optimality condition | 420 |
| 8.4.2 | Newton step and perturbed system | 422 |
| 8.4.3 | Algorithms | 424 |
| 8.5 | Notes on bibliography | 426 |
| 9 | CONVEX ANALYSIS AND CONVEX OPTIMIZATION | 428 |
| 9.1 | Affine sets | 430 |
| 9.1.1 | Basic concepts | 430 |
| 9.1.2 | Affine independence and dimensions | 432 |
| 9.2 | Convex sets and properties | 436 |
| 9.2.1 | Concepts of convex sets | 436 |
| 9.2.2 | Projection theorems | 438 |
| 9.2.3 | Separation theorems | 439 |
| 9.2.3.1 | Separating hyperplane theorem | 439 |
| 9.2.3.2 | Farka's lemma | 441 |
| 9.3 | Convex functions | 445 |
| 9.3.1 | Basic concepts | 445 |
| 9.3.2 | Connection to convex set | 447 |
| 9.3.3 | Strongly convex functions | 447 |
| 9.3.4 | Operations preserve convexity | 448 |
| 9.3.5 | Convexity and derivatives | 449 |
| 9.3.6 | Subgradient | 451 |
| 9.4 | Duality theory | 453 |
| 9.5 | Convex optimization and optimality conditions | 457 |
| 9.5.1 | Local optimality vs. global optimality | 457 |
| 9.5.2 | Unconstrained optimization optimality conditions | 458 |
| 9.5.3 | Constrained optimization optimality conditions | 458 |
| 9.6 | Subgradient methods | 463 |
| 9.6.1 | A generic algorithm for unconstrained problem | 463 |
| 9.6.2 | Convergence under Lipschitz smoothness | 465 |
| 9.6.3 | Projected gradient methods | 468 |
| 9.6.3.1 | Foundations | 468 |
| 9.6.3.2 | Algorithms | 470 |
| 9.6.4 | Proximal gradient methods | 471 |
| 9.6.4.1 | Foundations | 471 |
| 9.6.4.2 | Algorithms | 472 |
| 9.6.4.3 | Case study: sparsity regularization problem | 473 |
| 9.7 | Notes on bibliography | 475 |
| 10 | BASIC GAME THEORY | 477 |

| | | |
|----------|---|-----|
| 10.1 | Static normal form game | 478 |
| 10.1.1 | Normal form game concepts | 478 |
| 10.1.2 | Pure strategy and equilibrium | 478 |
| 10.1.2.1 | Solution concepts | 478 |
| 10.1.3 | Mixed strategy and equilibrium | 482 |
| 10.1.4 | Pareto optimality | 484 |
| 10.2 | Zero-sum matrix game | 485 |
| 10.2.1 | Fundamentals | 485 |
| 10.2.2 | Optimal strategy and Nash equilibrium | 486 |
| 10.2.3 | Saddle points as solutions | 488 |
| 10.2.4 | Maxmin strategies and Nash equilibrium | 489 |
| 10.2.5 | Linear programming approach to optimal strategy | 492 |
| 10.3 | Notes on bibliography | 496 |

iii classical statistical methods

| | | |
|----------|--|-----|
| 11 | PROBABILITY THEORY | 499 |
| 11.1 | σ algebra | 501 |
| 11.1.1 | σ algebra concepts | 501 |
| 11.1.2 | Generation of sigma algebra | 501 |
| 11.1.3 | Partition of sample space | 502 |
| 11.1.4 | Filtration & information | 502 |
| 11.1.5 | Borel σ algebra | 503 |
| 11.1.6 | Measurable set and measurable space | 504 |
| 11.2 | Probability space | 507 |
| 11.2.1 | Event, sample point and sample space | 507 |
| 11.2.2 | Probability space | 507 |
| 11.2.3 | Properties of probability measure | 509 |
| 11.2.4 | Conditional probability | 510 |
| 11.2.4.1 | Basics | 510 |
| 11.2.4.2 | Bayes' theorem | 511 |
| 11.2.4.3 | Independence of events and sigma algebra | 512 |
| 11.3 | Measurable map and random variable | 515 |
| 11.3.1 | Random variable | 515 |
| 11.3.2 | σ algebra of random variables | 516 |
| 11.3.3 | Independence of random variables | 517 |
| 11.4 | Distributions of random variables | 519 |
| 11.4.1 | Basic concepts | 519 |
| 11.4.1.1 | Probability mass function | 519 |
| 11.4.1.2 | Distributions on \mathbb{R}^n | 519 |
| 11.4.1.3 | Probability density function | 520 |
| 11.4.1.4 | Conditional distributions | 521 |

| | | |
|----------|---|-----|
| 11.4.1.5 | Bayes law | 522 |
| 11.4.2 | Independence | 523 |
| 11.4.3 | Conditional independence | 525 |
| 11.4.4 | Transformations | 525 |
| 11.4.4.1 | Transformation for univariate distribution | 525 |
| 11.4.4.2 | Location-scale transformation | 526 |
| 11.4.4.3 | Transformation for multivariate distribution | 527 |
| 11.5 | Expectation, variance, and covariance | 531 |
| 11.5.1 | Expectation | 531 |
| 11.5.2 | Expectation in the Lebesgue framework | 532 |
| 11.5.3 | Properties of expectation | 534 |
| 11.5.4 | Variance and covariance | 534 |
| 11.5.5 | Conditional variance | 535 |
| 11.5.6 | Delta method | 536 |
| 11.6 | Moment generating functions and characteristic functions | 538 |
| 11.6.1 | Moment generating function | 538 |
| 11.6.2 | Characteristic function | 541 |
| 11.6.3 | Joint moment generating functions for random vectors | 542 |
| 11.6.4 | Cumulants | 543 |
| 11.7 | Conditional expectation | 545 |
| 11.7.1 | General intuitions | 545 |
| 11.7.2 | Formal definitions | 545 |
| 11.7.3 | Different versions of conditional expectation | 547 |
| 11.7.3.1 | Conditioning on an event | 547 |
| 11.7.3.2 | Conditioning on a discrete random variable as a new random variable | 547 |
| 11.7.3.3 | Condition on random variable vs. event vs σ algebra | 548 |
| 11.7.4 | Properties | 548 |
| 11.7.4.1 | Linearity | 548 |
| 11.7.4.2 | Taking out what is known | 549 |
| 11.7.4.3 | Law of total expectation | 549 |
| 11.7.4.4 | Law of iterated expectations | 551 |
| 11.7.4.5 | Conditioning on independent random variable/ σ algebra | 551 |
| 11.7.4.6 | Least Square minimizing property | 552 |
| 11.8 | The Hilbert space of random variables | 553 |
| 11.8.1 | Definitions | 553 |
| 11.8.2 | Subspaces, projections, and approximations | 553 |
| 11.8.3 | Connection to conditional expectation | 558 |
| 11.9 | Probability inequalities | 561 |
| 11.9.1 | Chebychev inequalities | 561 |

| | | |
|----------|--|-----|
| 11.9.2 | Jensen's inequality | 562 |
| 11.9.3 | Holder's, Minkowski, and Cauchy-Schwarz inequalities | 563 |
| 11.9.4 | Popoviciu's inequality for variance | 565 |
| 11.10 | Convergence of random variables | 567 |
| 11.10.1 | Different levels of equivalence among random variables | 567 |
| 11.10.2 | Convergence almost surely | 567 |
| 11.10.3 | Convergence in probability | 568 |
| 11.10.4 | Mean square convergence | 570 |
| 11.10.5 | Convergence in distribution | 570 |
| 11.11 | Law of Large Number and Central Limit theorem | 573 |
| 11.11.1 | Law of Large Numbers | 573 |
| 11.11.2 | Central limit theorem | 575 |
| 11.12 | Finite sampling models | 578 |
| 11.12.1 | Counting principles | 578 |
| 11.12.2 | Matching problem | 581 |
| 11.12.3 | Birthday problem | 583 |
| 11.12.4 | Coupon collection problem | 584 |
| 11.12.5 | Balls into bins model | 585 |
| 11.13 | Order statistics | 588 |
| 11.14 | Information theory | 592 |
| 11.14.1 | Concept of entropy | 592 |
| 11.14.2 | Entropy maximizing distributions | 593 |
| 11.14.3 | KL divergence | 597 |
| 11.14.4 | Conditional entropy and mutual information | 598 |
| 11.14.5 | Cross-entropy | 599 |
| 11.15 | Notes on bibliography | 601 |
| 12 | STATISTICAL DISTRIBUTIONS | 603 |
| 12.1 | Common distributions and properties | 604 |
| 12.1.1 | Overview | 604 |
| 12.1.2 | Bernoulli distribution | 604 |
| 12.1.3 | Poisson distribution | 604 |
| 12.1.4 | Geometric distribution | 606 |
| 12.1.5 | Binomial distribution | 607 |
| 12.1.6 | Normal distribution | 609 |
| 12.1.7 | Half-normal distribution | 612 |
| 12.1.8 | Laplace distribution | 612 |
| 12.1.9 | Multivariate Gaussian/normal distribution | 613 |
| 12.1.9.1 | Basic definitions | 613 |
| 12.1.9.2 | Affine transformation and its consequences | 615 |
| 12.1.9.3 | Marginal and conditional distribution | 616 |
| 12.1.9.4 | Box Muller transformation | 618 |

| | | |
|-----------|--|-----|
| 12.1.10 | Lognormal distribution | 619 |
| 12.1.10.1 | Univariate lognormal distribution | 619 |
| 12.1.10.2 | Extension to univariate lognormal distribution | 620 |
| 12.1.10.3 | Multivariate lognormal distribution | 622 |
| 12.1.11 | Exponential distribution | 623 |
| 12.1.12 | Gamma distribution | 624 |
| 12.1.13 | Hypergeometric distribution | 626 |
| 12.1.14 | Beta distribution | 626 |
| 12.1.15 | Multinomial distribution | 629 |
| 12.1.16 | Dirichlet distribution | 630 |
| 12.1.17 | χ^2 -distribution | 631 |
| 12.1.17.1 | Basic properties | 631 |
| 12.1.17.2 | Noncentral chi-squared distribution | 633 |
| 12.1.18 | Wishart distribution | 633 |
| 12.1.19 | t -distribution | 634 |
| 12.1.19.1 | Standard t distribution | 634 |
| 12.1.19.2 | classical t distribution | 635 |
| 12.1.19.3 | Multivariate t distribution | 635 |
| 12.1.20 | F -distribution | 636 |
| 12.1.21 | Empirical distributions | 637 |
| 12.1.22 | Heavy-tailed distributions | 638 |
| 12.1.22.1 | Basic characterization | 638 |
| 12.1.22.2 | Pareto and power distribution | 638 |
| 12.1.22.3 | Student t distribution family | 639 |
| 12.1.22.4 | Gaussian mixture distributions | 640 |
| 12.2 | Characterizing distributions | 642 |
| 12.2.1 | Skewness and kurtosis | 642 |
| 12.2.2 | Percentiles and quantiles | 644 |
| 12.2.2.1 | Basics | 644 |
| 12.2.2.2 | Cornish-Fisher expansion | 646 |
| 12.3 | Moment matching approximation methods | 648 |
| 12.4 | Gaussian quadratic forms | 650 |
| 12.4.0.1 | Quadratic forms and chi-square distribution | 650 |
| 12.4.0.2 | Applications | 653 |
| 12.5 | Notes on bibliography | 656 |
| 13 | STATISTICAL ESTIMATION THEORY | 657 |
| 13.1 | Parameter estimators | 660 |
| 13.1.1 | Overview | 660 |
| 13.1.2 | Statistic and Estimator | 661 |
| 13.1.2.1 | Statistic | 661 |
| 13.1.2.2 | Estimator properties | 662 |

| | | |
|----------|--|-----|
| 13.1.2.3 | Variance-bias decomposition | 664 |
| 13.1.2.4 | Consistence | 666 |
| 13.1.2.5 | Efficiency | 668 |
| 13.1.2.6 | Robust statistics | 669 |
| 13.1.3 | Method of moments | 670 |
| 13.1.4 | Maximum likelihood estimation | 671 |
| 13.1.4.1 | Basic concepts | 671 |
| 13.1.4.2 | MLE examples | 673 |
| 13.1.4.3 | Bias and consistence of MLE | 676 |
| 13.2 | Information and efficiency | 679 |
| 13.2.1 | Fisher information | 679 |
| 13.2.2 | Cramer-Rao lower bound | 683 |
| 13.2.2.1 | Preliminary: information inequality | 683 |
| 13.2.2.2 | Cramer-Rao lower bound: univariate case | 684 |
| 13.2.2.3 | Cramer-Rao lower bound: multivariate case | 685 |
| 13.2.3 | Efficient estimators | 687 |
| 13.2.4 | Asymptotic normality and efficiency of MLE | 688 |
| 13.3 | Sufficiency and data reduction | 689 |
| 13.3.1 | Sufficient estimators | 689 |
| 13.3.2 | Factorization theorem | 690 |
| 13.4 | Bayesian estimation theory | 693 |
| 13.4.1 | Overview | 693 |
| 13.4.2 | Basics | 693 |
| 13.5 | Bootstrap method | 696 |
| 13.6 | Hypothesis testing theory | 698 |
| 13.6.1 | Basics | 698 |
| 13.6.2 | Characterizing errors and power | 701 |
| 13.6.3 | Power of a statistical test | 702 |
| 13.6.4 | Common statistical tests | 704 |
| 13.6.4.1 | Chi-square goodness-of-fit test | 704 |
| 13.6.4.2 | Chi-square test for statistical independence | 706 |
| 13.6.4.3 | Kolmogorov-Smirnov goodness-of-fit test | 707 |
| 13.7 | Hypothesis testing on normal distributions | 708 |
| 13.7.1 | Normality test | 708 |
| 13.7.2 | Sample mean with known variance | 708 |
| 13.7.3 | Sample mean with unknown variance | 710 |
| 13.7.4 | Variance test | 710 |
| 13.7.5 | Variance comparison test | 711 |
| 13.7.6 | Person correlation t test | 711 |
| 13.7.7 | Two sample tests | 712 |
| 13.7.7.1 | Two-sample z test | 712 |

| | | |
|----------|---|-----|
| 13.7.7.2 | Two-sample t test | 712 |
| 13.7.7.3 | Paired Data | 713 |
| 13.7.8 | Interval estimation for normal distribution | 713 |
| 13.8 | Notes on bibliography | 715 |
| 14 | MULTIVARIATE STATISTICAL METHODS | 716 |
| 14.1 | Multivariate data and distribution | 718 |
| 14.1.1 | Sample statistics | 718 |
| 14.1.2 | Multivariate Gaussian distribution | 719 |
| 14.1.3 | Estimation methods | 721 |
| 14.1.3.1 | Maximum likelihood estimation | 721 |
| 14.1.3.2 | Weighted estimation | 723 |
| 14.2 | Principal component analysis (PCA) | 725 |
| 14.2.1 | Statistical fundamentals of PCA | 725 |
| 14.2.1.1 | PCA for random vectors | 725 |
| 14.2.1.2 | Sample principal components | 726 |
| 14.2.2 | Geometric fundamentals of PCA | 729 |
| 14.2.2.1 | Optimization approach | 729 |
| 14.2.2.2 | Properties | 731 |
| 14.2.3 | Probabilistic PCA | 732 |
| 14.2.4 | Applications | 734 |
| 14.2.4.1 | Eigenfaces and eigendigits | 734 |
| 14.2.4.2 | Interest rate curve dynamics modeling | 736 |
| 14.3 | Canonical correlation analysis | 740 |
| 14.3.1 | Basics | 740 |
| 14.3.2 | Sparse CCA | 742 |
| 14.4 | Copulas and dependence modeling | 744 |
| 14.4.1 | Definitions and properties | 744 |
| 14.4.2 | Copulas and distributions | 747 |
| 14.4.2.1 | Fundamentals | 747 |
| 14.4.2.2 | Survival copula | 754 |
| 14.4.2.3 | Partial differential and conditional distribution | 755 |
| 14.4.3 | Common copula functions | 759 |
| 14.4.3.1 | Gaussian copula | 759 |
| 14.4.3.2 | t copula | 764 |
| 14.4.3.3 | Common copula functions: other copula | 764 |
| 14.4.4 | Dependence and copula | 765 |
| 14.4.4.1 | Linear correlations | 765 |
| 14.4.4.2 | Rank correlations | 767 |
| 14.4.4.3 | Tail dependence | 773 |
| 14.4.5 | Estimating copula function | 775 |
| 14.4.5.1 | Empirical copula method | 775 |

| | | |
|----------|--|-----|
| 14.4.5.2 | Maximum likelihood method | 776 |
| 14.4.6 | Applications of copula | 777 |
| 14.4.6.1 | Generating correlated uniform random number | 777 |
| 14.4.6.2 | Generating general correlated random number | 779 |
| 14.4.6.3 | Multivariate distribution approximation with Gaussian copula | 783 |
| 14.5 | Covariance structure and factor analysis | 784 |
| 14.5.1 | The orthogonal factor model | 784 |
| 14.5.1.1 | Motivation and factor models | 784 |
| 14.5.1.2 | Covariance structure implied by factor model | 784 |
| 14.5.2 | Parameter estimation | 786 |
| 14.5.2.1 | Data collection and preparation | 786 |
| 14.5.2.2 | PCA method | 787 |
| 14.5.2.3 | Maximum likelihood method | 788 |
| 14.5.3 | Factor score estimation | 788 |
| 14.5.4 | Application I: Joint default modeling | 789 |
| 14.5.4.1 | Single factor model | 789 |
| 14.5.4.2 | Multiple factor model | 792 |
| 14.5.5 | Application II: factor models for stock return | 793 |
| 14.5.5.1 | Overview | 793 |
| 14.5.5.2 | The Fama-French 3 factor model | 795 |
| 14.6 | Graphical models | 799 |
| 14.6.1 | Fundamentals | 799 |
| 14.7 | Notes on Bibliography | 805 |
| 15 | LINEAR REGRESSION ANALYSIS | 806 |
| 15.1 | Linear regression analysis: basics | 808 |
| 15.1.1 | Linear regression models | 808 |
| 15.1.2 | Ordinary least square (OLS): fundamentals | 811 |
| 15.1.2.1 | Review on orthogonal projections | 811 |
| 15.1.2.2 | OLS results | 812 |
| 15.1.2.3 | OLS results with demeaned data | 817 |
| 15.1.2.4 | Gauss-Markov theorem | 820 |
| 15.1.2.5 | Variance decomposition | 821 |
| 15.1.2.6 | Residual and variance estimation | 824 |
| 15.1.3 | Ordinary least square (OLS): Additional topics | 825 |
| 15.1.3.1 | Orthogonal input and successive regression | 825 |
| 15.1.3.2 | Frisch-Waugh-Lovell(FWL) theorem and partial regression | 827 |
| 15.1.3.3 | Forecasting analysis with normality assumption | 828 |
| 15.1.4 | Hypothesis testing and analysis of variance | 830 |
| 15.1.4.1 | Distribution of coefficients | 831 |

| | | |
|----------|---|-----|
| 15.1.4.2 | t test and normality test of single coefficients | 833 |
| 15.1.4.3 | F lack-of-fit test | 835 |
| 15.1.4.4 | χ^2 test for variance | 837 |
| 15.1.5 | Maximum likelihood method with normality assumption | 838 |
| 15.1.6 | Asymptotic properties of least square solutions | 840 |
| 15.1.6.1 | Asymptotic properties of standard OLS | 840 |
| 15.1.6.2 | Asymptotic efficiency of standard OLS | 842 |
| 15.1.7 | Partial and multiple correlation | 842 |
| 15.1.7.1 | Multiple correlation coefficient, R^2 | 842 |
| 15.1.7.2 | Partial correlation coefficient | 845 |
| 15.1.8 | Generalized linear regression (GLR) | 846 |
| 15.1.8.1 | Linear regression with structural error | 846 |
| 15.1.8.2 | Generalized least square solution | 847 |
| 15.1.8.3 | Gauss-Markov theorem for GLR | 849 |
| 15.1.8.4 | Feasible GLS | 850 |
| 15.1.9 | Linear structure in joint distributions | 850 |
| 15.2 | Model specification and selection | 853 |
| 15.2.1 | Model order mis-specification | 853 |
| 15.2.1.1 | Omission of relevant regressors | 853 |
| 15.2.1.2 | Inclusion of irrelevant regressors | 854 |
| 15.2.2 | Model selection methods | 856 |
| 15.2.2.1 | Adjusted R square method | 856 |
| 15.2.2.2 | F test method | 856 |
| 15.2.2.3 | Information criterion methods | 858 |
| 15.2.2.4 | Bayesian information criterion (BIC) | 859 |
| 15.2.3 | Test for structure change | 860 |
| 15.3 | Linear regression analysis: diagnostics & solutions | 862 |
| 15.3.1 | Multi-collinearity | 862 |
| 15.3.1.1 | Detection and characterization | 862 |
| 15.3.1.2 | Regressor linear regression and variance inflation factor | 862 |
| 15.3.1.3 | Principal component linear regression (PCLR) | 865 |
| 15.3.2 | Rank deficiency and rigid regression | 865 |
| 15.3.3 | Heteroskedasticity | 867 |
| 15.3.3.1 | Test for heteroskedasticity | 867 |
| 15.3.3.2 | Heteroskedasticity robust estimator | 868 |
| 15.3.3.3 | Feasible weighted least square | 868 |
| 15.3.4 | Residual normality test | 870 |
| 15.3.4.1 | Jarque-Bera test | 870 |
| 15.3.4.2 | D'Agostino's K^2 test | 870 |
| 15.3.5 | Autocorrelation of errors | 871 |

| | | |
|----------|---|-----|
| 15.3.5.1 | Motivation and general remarks | 871 |
| 15.3.5.2 | Test of autocorrelation of errors | 872 |
| 15.3.5.3 | Models with known autocorrelation | 874 |
| 15.3.5.4 | Transformation to generalized linear regression | 875 |
| 15.3.6 | Outliers analysis and robust linear regression | 877 |
| 15.3.6.1 | Outliers and influential points | 877 |
| 15.3.6.2 | Outlier impact analysis | 878 |
| 15.3.6.3 | Robust M-estimation linear regression | 881 |
| 15.3.7 | Visual diagnosis | 884 |
| 15.4 | Linear regression case studies | 887 |
| 15.4.1 | Standard linear regression | 887 |
| 15.4.2 | Boston Housing example | 889 |
| 15.5 | Multivariate multiple linear regression (MMLR) | 892 |
| 15.5.1 | Canonical MMLR | 892 |
| 15.5.1.1 | Motivation and model | 892 |
| 15.5.1.2 | Ordinary least square solution | 893 |
| 15.5.2 | Reduced rank regression | 894 |
| 15.6 | Notes on Bibliography | 899 |
| 16 | MONTE CARLO METHODS | 900 |
| 16.1 | Generating random variables | 902 |
| 16.1.1 | Inverse transform method | 902 |
| 16.1.2 | Box-Muller method for standard normal random variable | 904 |
| 16.1.3 | Acceptance-rejection method | 904 |
| 16.1.4 | Composition approach | 905 |
| 16.1.5 | Generate dependent continuous random variables | 908 |
| 16.1.5.1 | Multivariate normal and lognormal distribution | 908 |
| 16.1.5.2 | Multivariate student t distribution | 908 |
| 16.1.5.3 | General joint distribution | 909 |
| 16.1.6 | Generate discrete random variables | 909 |
| 16.1.6.1 | Generate single discrete random variables | 909 |
| 16.1.6.2 | Generate correlated discrete random variables | 910 |
| 16.2 | Monte Carlo integration | 912 |
| 16.2.1 | Naive approach | 912 |
| 16.2.2 | Importance sampling | 913 |
| 16.3 | Markov chain Monte Carlo | 916 |
| 16.3.1 | Basics | 916 |
| 16.3.1.1 | Markov chain Monte Carlo (MCMC) | 916 |
| 16.3.2 | Metropolis-Hasting algorithm | 917 |
| 16.3.3 | Gibbs sampling | 919 |
| 16.4 | Monte Carlo for random processes | 920 |
| 16.4.1 | Simulating stochastic differential equations | 920 |

| | | |
|----------|--|-----|
| 16.4.1.1 | Simulating Brownian motion | 920 |
| 16.4.1.2 | Simulating linear arithmetic SDE | 920 |
| 16.4.1.3 | Simulating linear geometric SDE | 921 |
| 16.4.1.4 | Simulation mean-reversion(OU) process | 921 |
| 16.4.2 | Stochastic interpolation | 922 |
| 16.4.2.1 | Interpolating Gaussian processes | 922 |
| 16.4.2.2 | Interpolating one Dimensional Brownian motions | 923 |
| 16.4.2.3 | Interpolating multi-dimensional Brownian motions | 925 |
| 16.5 | Monte Carlo variance reduction | 928 |
| 16.5.1 | Antithetic sampling | 928 |
| 16.5.1.1 | Basic principles | 928 |
| 16.5.1.2 | Methods and analysis | 929 |
| 16.5.2 | Control variates | 931 |
| 16.5.2.1 | Basic principles | 931 |
| 16.5.2.2 | Multiple control variates | 933 |
| 16.6 | Notes on bibliography | 934 |

iv dynamics modeling methods

| | | |
|----------|--|-----|
| 17 | MODELS AND ESTIMATION IN LINEAR SYSTEMS | 936 |
| 17.1 | Difference equation | 937 |
| 17.1.1 | Introduction | 937 |
| 17.1.2 | Solution structure of linear difference equations | 938 |
| 17.1.3 | Solution to non-homogeneous equation | 940 |
| 17.1.4 | Linear equations with constant coefficients | 941 |
| 17.1.4.1 | Basic case | 941 |
| 17.1.4.2 | General case | 944 |
| 17.2 | Differential equations | 947 |
| 17.2.1 | Linear differential equations | 947 |
| 17.2.1.1 | Concepts | 947 |
| 17.2.1.2 | Wronskian and linear independence | 948 |
| 17.2.1.3 | General solution theory | 951 |
| 17.2.1.4 | Existence & uniqueness of solution | 954 |
| 17.2.2 | Linear homogeneous differential equations with constant coefficients | 955 |
| 17.2.2.1 | The key identity | 955 |
| 17.2.2.2 | The case of real roots | 957 |
| 17.2.2.3 | The case of complex roots | 958 |
| 17.2.2.4 | The complete solution set | 958 |
| 17.2.3 | Solution to non-homogeneous ODEs | 961 |
| 17.2.3.1 | General principles | 961 |
| 17.2.3.2 | Key identity approach | 962 |

| | | |
|----------|--|------|
| 17.2.4 | First order linear differential equation | 968 |
| 17.3 | Linear system | 970 |
| 17.3.1 | Solution space for linear homogeneous system | 970 |
| 17.3.2 | Linear independence and the Wronskian | 972 |
| 17.3.3 | The fundamental system and solution method | 974 |
| 17.3.4 | The non-homogeneous linear equation | 975 |
| 17.3.5 | Conversion of linear differential/difference equation to linear systems | 979 |
| 17.3.6 | Solution method for discrete system | 980 |
| 17.4 | Linear system with constant coefficients | 981 |
| 17.4.1 | General solutions | 981 |
| 17.4.2 | System eigenvector method: continuous-time system | 981 |
| 17.4.2.1 | Diagonalizable system | 981 |
| 17.4.2.2 | two-by-two non-diagonalizable system | 988 |
| 17.4.2.3 | Non-diagonalizable system | 989 |
| 17.4.3 | Equilibrium point | 991 |
| 17.4.3.1 | Discrete-time system | 991 |
| 17.4.3.2 | Continuous-time system | 992 |
| 17.4.4 | Stability | 993 |
| 17.4.5 | Complex eigenvalues/eigenvectors | 995 |
| 17.4.6 | Boundedness of linear systems | 995 |
| 17.4.7 | One dimensional Nonlinear dynamical system analysis | 996 |
| 17.5 | Least square estimation of constant vectors | 998 |
| 17.5.1 | linear static estimation from single measurement with no prior information | 998 |
| 17.5.2 | linear static estimation from single measurement with prior information | 999 |
| 17.5.3 | Batch and recursive least square estimation with multiple measurements | 1000 |
| 17.5.4 | Nonlinear least square estimation | 1003 |
| 17.6 | Kalman filter | 1004 |
| 17.6.1 | Preliminary: error propagation in linear systems | 1004 |
| 17.6.1.1 | Discrete-time system | 1004 |
| 17.6.1.2 | Continuous-time system | 1004 |
| 17.6.2 | Batch estimation | 1005 |
| 17.6.3 | From batch estimation to Kalman filter | 1006 |
| 17.6.4 | Extended Kalman filter for nonlinear system | 1007 |
| 17.7 | Notes on bibliography | 1009 |
| 18 | STOCHASTIC PROCESS | 1010 |
| 18.1 | Stochastic process | 1012 |
| 18.1.1 | Basic definition and concepts | 1012 |

| | | |
|----------|---|------|
| 18.1.2 | Stationarity | 1014 |
| 18.2 | Gaussian process | 1017 |
| 18.2.1 | Basic Gaussian process | 1017 |
| 18.2.2 | Stationarity | 1018 |
| 18.3 | Brownian motion (Wiener process) | 1019 |
| 18.3.1 | Definition and elementary properties | 1019 |
| 18.3.2 | Multi-dimensional Brownian motion | 1021 |
| 18.3.3 | Asymptotic behaviors | 1022 |
| 18.3.4 | The reflection principle | 1023 |
| 18.3.5 | Quadratic variation | 1024 |
| 18.3.6 | Discrete-time approximations and simulation | 1026 |
| 18.4 | Brownian motion variants | 1027 |
| 18.4.1 | Gaussian process generated by Brownian motion | 1027 |
| 18.4.2 | Brownian bridge | 1028 |
| 18.4.2.1 | Constructions | 1028 |
| 18.4.2.2 | Applications | 1032 |
| 18.4.3 | Geometric Brownian motion | 1032 |
| 18.5 | Poisson process | 1034 |
| 18.5.1 | Basics | 1034 |
| 18.5.2 | Arrival and Inter-arrival Times | 1035 |
| 18.6 | Martingale theory | 1037 |
| 18.6.1 | Preliminaries: Filtration and adapted process | 1037 |
| 18.6.1.1 | Basic concepts in filtration | 1037 |
| 18.6.1.2 | Filtration for Brownian motion | 1039 |
| 18.6.2 | Basics of martingales | 1039 |
| 18.6.3 | Martingale transformation | 1042 |
| 18.7 | Stopping time | 1043 |
| 18.7.1 | Stopping time examples | 1043 |
| 18.7.1.1 | First passage time | 1043 |
| 18.7.1.2 | Trivial stopping time | 1043 |
| 18.7.1.3 | Counter example: last exit time | 1044 |
| 18.7.2 | Wald's equation | 1044 |
| 18.7.3 | Optional stopping | 1045 |
| 18.7.4 | martingale method for first hitting time | 1045 |
| 18.8 | Notes on bibliography | 1046 |
| 19 | STOCHASTIC CALCULUS | 1047 |
| 19.1 | Ito stochastic integral | 1048 |
| 19.1.1 | Motivation | 1048 |
| 19.1.2 | Construction of Ito integral | 1048 |
| 19.1.2.1 | Ito integral of a simple process | 1048 |
| 19.1.2.2 | Ito integral of a general process | 1051 |

| | | |
|----------|---|------|
| 19.1.3 | Ito integral with deterministic integrands (Wiener integral) | 1055 |
| 19.1.3.1 | Basics | 1055 |
| 19.1.3.2 | Integration by parts | 1058 |
| 19.2 | Stochastic differential equations | 1061 |
| 19.2.1 | Ito Stochastic differential equations | 1061 |
| 19.2.2 | Ito's lemma | 1062 |
| 19.2.3 | Useful results of Ito's lemma | 1065 |
| 19.2.3.1 | Product rule and quotient rule | 1065 |
| 19.2.3.2 | Logarithm and exponential | 1066 |
| 19.2.3.3 | Ito integral by parts | 1066 |
| 19.2.3.4 | Differentiate integrals of Ito process | 1067 |
| 19.3 | Linear SDE | 1069 |
| 19.3.1 | State-independent linear arithmetic SDE | 1069 |
| 19.3.2 | State-independent linear geometric SDE | 1069 |
| 19.3.3 | Multiple dimension extension | 1071 |
| 19.3.4 | Exact SDE | 1073 |
| 19.3.5 | Calculation mean and variance from SDE | 1074 |
| 19.3.6 | Integrals of Ito SDE | 1075 |
| 19.4 | Ornstein-Uhlenbeck(OU) process | 1080 |
| 19.4.1 | OU process | 1080 |
| 19.4.1.1 | Constant coefficient OU process | 1080 |
| 19.4.1.2 | Time-dependent coefficient OU process | 1084 |
| 19.4.1.3 | Integral of OU process | 1086 |
| 19.4.2 | Exponential OU process | 1089 |
| 19.4.3 | Parameter estimation for OU process | 1091 |
| 19.4.4 | Multiple factor extension | 1091 |
| 19.5 | Notes on bibliography | 1094 |
| 20 | MARKOV CHAIN AND RANDOM WALK | 1095 |
| 20.1 | Discrete-time Markov chain | 1097 |
| 20.1.1 | The model | 1097 |
| 20.1.2 | Evolution of discrete chain | 1099 |
| 20.2 | Classification of states | 1101 |
| 20.2.1 | accessibility and communicating classes | 1101 |
| 20.2.2 | Transient and recurrent states and classes | 1103 |
| 20.2.2.1 | Transient and recurrent states | 1103 |
| 20.2.2.2 | From states to classes | 1107 |
| 20.2.2.3 | Qualitative classification of recurrent and transient classes | 1108 |
| 20.2.3 | Periodicity | 1109 |
| 20.2.4 | Positive and null recurrent | 1111 |
| 20.2.5 | Summary | 1113 |

| | | |
|----------|--|------|
| 20.3 | Absorption analysis | 1114 |
| 20.3.1 | Matrix structure for adsorption analysis | 1114 |
| 20.3.2 | Absorbing Markov chains | 1116 |
| 20.3.3 | Hitting and return analysis | 1118 |
| 20.3.4 | Examples | 1120 |
| 20.3.4.1 | Consecutive coin toss game | 1120 |
| 20.4 | Limiting behavior & distributions | 1122 |
| 20.4.1 | Preliminary: eigenvalue propoties of stochastic matrices | 1122 |
| 20.4.1.1 | Preliminary: Frobenius-Perron matrix theory | 1122 |
| 20.4.1.2 | More general situations | 1124 |
| 20.4.2 | Limiting theorem | 1126 |
| 20.4.2.1 | Limiting distribution | 1126 |
| 20.4.2.2 | Extensions via Long run return analysis | 1130 |
| 20.4.3 | Application: PageRank algorithm | 1132 |
| 20.5 | Detailed balance and spectral properties | 1134 |
| 20.6 | Random walk | 1136 |
| 20.6.1 | Basic concepts and properties | 1136 |
| 20.6.2 | Persistent random walk | 1136 |
| 20.6.3 | Asymptotic properties | 1138 |
| 20.6.4 | Gambler's ruin problems | 1138 |
| 20.7 | Notes on bibliography | 1143 |
| 21 | TIME SERIES ANALYSIS | 1144 |
| 21.1 | Overview of time series analysis | 1145 |
| 21.1.1 | Introduction to time series | 1145 |
| 21.1.2 | Stationarity | 1146 |
| 21.1.2.1 | Stationarity concept | 1146 |
| 21.1.2.2 | Rolling analysis | 1148 |
| 21.1.3 | Remove trend and seasonality | 1149 |
| 21.2 | Linear stationary process theory | 1152 |
| 21.2.1 | Preliminaries: the lag operator and polynomial | 1152 |
| 21.2.2 | Linear process | 1153 |
| 21.2.3 | Autoregressive (AR) process | 1155 |
| 21.2.3.1 | Basics | 1155 |
| 21.2.3.2 | Stationarity and invertibility condition | 1158 |
| 21.2.3.3 | Forecasting | 1159 |
| 21.2.4 | Moving average (MA) process | 1162 |
| 21.2.4.1 | Basics | 1162 |
| 21.2.4.2 | Stationarity and invertibility | 1165 |
| 21.2.4.3 | Forecasting | 1166 |
| 21.2.5 | ARMA process | 1169 |
| 21.2.5.1 | Basic properties | 1169 |

| | | |
|----------|--|------|
| 21.2.6 | Unit root AR process | 1171 |
| 21.2.6.1 | Unit root process | 1171 |
| 21.2.6.2 | Trend stationarity vs. unit root process | 1173 |
| 21.2.6.3 | Unit root test | 1173 |
| 21.2.6.4 | Forecasting | 1174 |
| 21.2.7 | Correlation analysis | 1174 |
| 21.2.7.1 | Autocorrelation statistical analysis | 1174 |
| 21.2.7.2 | Partial autocorrelation function theory | 1175 |
| 21.2.7.3 | Correlogram analysis example | 1178 |
| 21.2.8 | Model analysis and calibration | 1180 |
| 21.2.8.1 | Order selection | 1180 |
| 21.2.8.2 | Yule-Walker equations and related methods | 1181 |
| 21.2.8.3 | Linear regression approach | 1184 |
| 21.2.8.4 | Maximum likelihood estimation | 1186 |
| 21.2.8.5 | Example: a toy example | 1187 |
| 21.2.9 | Wold Representation theorem | 1190 |
| 21.3 | Extensions to multivariate time series | 1193 |
| 21.3.1 | Introduction | 1193 |
| 21.3.2 | Vector autoregressive models | 1194 |
| 21.3.2.1 | VAR(1) model | 1194 |
| 21.3.2.2 | VAR(2) model | 1196 |
| 21.3.2.3 | VAR(p) model | 1198 |
| 21.3.3 | Vector moving-average model | 1200 |
| 21.4 | Autoregressive conditional heteroscedastic model | 1203 |
| 21.4.1 | ARCH models | 1203 |
| 21.4.1.1 | The motivation and the model | 1203 |
| 21.4.1.2 | Statistical properties | 1205 |
| 21.4.1.3 | Variance forecasting | 1211 |
| 21.4.1.4 | Detect ARCH effect | 1214 |
| 21.4.1.5 | Parameter estimation | 1215 |
| 21.4.2 | GARCH models | 1215 |
| 21.4.2.1 | The model | 1215 |
| 21.4.2.2 | Connecting GARCH to ARCH | 1218 |
| 21.4.2.3 | Variance forecasting | 1218 |
| 21.5 | Notes on Bibliography | 1221 |

v statistical learning methods

| | | |
|--------|---------------------------------|------|
| 22 | SUPERVISED LEARNING PRINCIPLES | 1223 |
| 22.1 | The supervised learning problem | 1224 |
| 22.1.1 | Concepts | 1224 |
| 22.1.2 | Framework | 1226 |

| | | |
|----------|---|------|
| 22.2 | Variance bias trade-off | 1228 |
| 22.2.1 | Underfitting and Overfitting | 1228 |
| 22.2.2 | Variance and bias trade-off | 1230 |
| 22.2.3 | Examples | 1233 |
| 22.2.3.1 | Linear regression | 1233 |
| 22.2.4 | No free lunch theorem | 1234 |
| 22.3 | Model loss and evaluation | 1236 |
| 22.3.1 | Common loss functions | 1236 |
| 22.3.2 | Model evaluation metrics | 1240 |
| 22.3.2.1 | Regression metrics | 1240 |
| 22.3.2.2 | Classification metrics | 1241 |
| 22.3.2.3 | ROC and PRC metrics | 1243 |
| 22.3.2.4 | Metrics for imbalanced data | 1245 |
| 22.4 | Model selection methods | 1246 |
| 22.4.1 | The training-validation-testing idea | 1246 |
| 22.4.2 | Cross-validation | 1246 |
| 22.5 | Data and feature engineering | 1251 |
| 22.5.1 | Data preprocessing | 1251 |
| 22.5.1.1 | Data standardization | 1251 |
| 22.5.1.2 | Data normalization | 1252 |
| 22.5.1.3 | Handle categorical data | 1252 |
| 22.5.1.4 | Handle missing values | 1253 |
| 22.5.1.5 | Dimensional reduction | 1253 |
| 22.5.1.6 | Centering kernel matrix | 1253 |
| 22.5.2 | Feature engineering I: basic routines | 1254 |
| 22.5.2.1 | Nonlinear transformation | 1254 |
| 22.5.2.2 | Polynomial features | 1254 |
| 22.5.2.3 | Binning | 1254 |
| 22.5.3 | Feature engineering II: feature selection | 1255 |
| 22.5.3.1 | Filtering methods | 1255 |
| 22.5.3.2 | Recursive elimination methods | 1256 |
| 22.5.3.3 | Regularization methods | 1256 |
| 22.5.4 | Feature engineering III: feature extraction | 1256 |
| 22.5.4.1 | Text analytics | 1256 |
| 22.5.4.2 | Image | 1257 |
| 22.5.4.3 | Time series | 1258 |
| 22.5.5 | Imbalanced data | 1259 |
| 22.5.5.1 | Motivations | 1259 |
| 22.5.5.2 | Data resampling: undersampling | 1259 |
| 22.5.5.3 | Data resampling: upsampling | 1259 |
| 22.5.5.4 | Choice of loss functions, algorithms, and metrics | 1260 |

| | | |
|----------|--|------|
| 22.6 | Kernel methods | 1261 |
| 22.6.1 | Basic concepts of kernels and feature maps | 1261 |
| 22.6.2 | Mercer's theorem | 1261 |
| 22.6.3 | Common kernels | 1263 |
| 22.6.4 | Kernel trick and elementary algorithms using kernels | 1265 |
| 22.6.5 | Elementary algorithms | 1265 |
| 22.7 | Note on bibliography | 1267 |
| 23 | LINEAR MODELS FOR REGRESSION | 1268 |
| 23.1 | Standard linear regression | 1269 |
| 23.1.1 | Ordinary linear regression | 1269 |
| 23.1.2 | Application examples | 1270 |
| 23.1.2.1 | Boston housing prices | 1270 |
| 23.2 | Penalized linear regression | 1274 |
| 23.2.1 | Ridge regression | 1274 |
| 23.2.1.1 | Basics | 1274 |
| 23.2.1.2 | Dual form of ridge regression | 1277 |
| 23.2.2 | Lasso regression | 1277 |
| 23.2.3 | Elastic net | 1280 |
| 23.2.4 | Shrinkage Comparison | 1280 |
| 23.2.5 | Effective degree of freedom | 1283 |
| 23.3 | Basis function extension | 1284 |
| 23.4 | Note on bibliography | 1286 |
| 24 | LINEAR MODELS FOR CLASSIFICATION | 1287 |
| 24.1 | Logistic regression | 1289 |
| 24.1.1 | Logistic regression model | 1289 |
| 24.1.2 | Parameter estimation via maximum likelihood estimation | 1290 |
| 24.1.3 | Logistic regression with regularization | 1294 |
| 24.1.4 | Feature augmentation strategies | 1295 |
| 24.1.5 | Multinomial logistic regression | 1296 |
| 24.1.6 | Application examples | 1297 |
| 24.1.6.1 | South Africa heart disease | 1297 |
| 24.1.6.2 | Credit card fraud detection | 1300 |
| 24.1.6.3 | MNIST | 1302 |
| 24.2 | Gaussian discriminate analysis | 1304 |
| 24.2.1 | Linear Gaussian discriminant model | 1304 |
| 24.2.1.1 | The model | 1304 |
| 24.2.1.2 | Model parameter estimation | 1305 |
| 24.2.1.3 | Geometry of decision boundary | 1305 |
| 24.2.2 | Quadratic Gaussian discriminant model | 1307 |
| 24.2.2.1 | The model | 1307 |
| 24.2.2.2 | Model parameter estimation | 1309 |

| | | |
|----------|---|------|
| 24.2.3 | Application examples | 1309 |
| 24.2.3.1 | A toy example | 1309 |
| 24.3 | Fisher Linear discriminate analysis (Fisher LDA) | 1312 |
| 24.3.1 | One dimensional linear discriminant | 1312 |
| 24.3.1.1 | Basics | 1312 |
| 24.3.1.2 | Application in classification | 1314 |
| 24.3.1.3 | Possible issues | 1316 |
| 24.3.2 | Multi-dimensional linear discriminate | 1317 |
| 24.3.2.1 | Basics | 1317 |
| 24.3.2.2 | Application in classification | 1318 |
| 24.3.3 | Supervised dimensional reduction via Fisher LDA | 1320 |
| 24.4 | Separating hyperplane and Perceptron learning algorithm | 1322 |
| 24.4.1 | Basic geometry of hyperplanes | 1322 |
| 24.4.2 | The Perceptron learning algorithm | 1323 |
| 24.5 | Support vector machine classifier | 1325 |
| 24.5.1 | Motivation and formulation | 1325 |
| 24.5.2 | Optimality condition and dual form | 1326 |
| 24.5.3 | Soft margin SVM | 1328 |
| 24.5.3.1 | Basics | 1328 |
| 24.5.3.2 | Optimality condition for soft margin SVM | 1329 |
| 24.5.3.3 | Algorithm | 1331 |
| 24.5.4 | SVM with kernels | 1333 |
| 24.5.5 | A unified perspective from loss functions | 1334 |
| 24.6 | Note on bibliography | 1337 |
| 25 | GENERATIVE MODELS | 1338 |
| 25.1 | Naive Bayes classifier (NBC) | 1342 |
| 25.1.1 | Overview | 1342 |
| 25.1.2 | Binomial NBC | 1342 |
| 25.1.3 | Multinomial NBC | 1344 |
| 25.1.4 | Gaussian NBC | 1347 |
| 25.1.5 | Discussion | 1349 |
| 25.2 | Application | 1350 |
| 25.2.1 | Classifying documents using bag of words | 1350 |
| 25.2.2 | Credit card fraud prediction | 1350 |
| 25.3 | Supporting mathematical results | 1353 |
| 25.3.1 | Beta-binomial model | 1353 |
| 25.3.1.1 | The model | 1353 |
| 25.3.1.2 | Parameter inference | 1354 |
| 25.3.2 | Dirichlet-multinomial model | 1356 |
| 25.3.2.1 | The model | 1356 |
| 25.3.2.2 | Parameter inference | 1359 |

| | | |
|----------|---|------|
| 26 | K NEAREST NEIGHBORS | 1360 |
| 26.1 | Principles | 1361 |
| 26.1.1 | The algorithm | 1361 |
| 26.1.2 | Metrics and features | 1362 |
| 26.2 | Application examples | 1364 |
| 27 | TREE METHODS | 1366 |
| 27.1 | Preliminaries: entropy concepts | 1367 |
| 27.1.1 | Concept of entropy | 1367 |
| 27.1.2 | Conditional entropy and mutual information | 1368 |
| 27.2 | Classification tree | 1372 |
| 27.2.1 | Basic concepts of decision tree learning | 1372 |
| 27.2.2 | A generic tree-growth algorithm | 1373 |
| 27.2.3 | Splitting criterion | 1374 |
| 27.2.4 | Tree pruning | 1378 |
| 27.2.5 | Practical algorithms | 1378 |
| 27.2.6 | Examples | 1381 |
| 27.2.6.1 | Tree structures in Iris data classification | 1381 |
| 27.3 | Regression tree | 1384 |
| 27.3.1 | Basics | 1384 |
| 27.3.2 | Practical algorithms | 1387 |
| 27.3.3 | Examples | 1387 |
| 27.3.3.1 | A toy example | 1387 |
| 27.3.3.2 | Boston Housing prices | 1388 |
| 28 | ENSEMBLE AND BOOSTING METHODS | 1390 |
| 28.1 | Motivation and overview | 1393 |
| 28.2 | Bagging Methods | 1395 |
| 28.2.1 | A basic bagging method | 1395 |
| 28.2.2 | Tree bagging | 1396 |
| 28.2.3 | Random Forest | 1398 |
| 28.3 | Adaboost | 1401 |
| 28.3.1 | Adaboost classifier | 1401 |
| 28.3.2 | Adaboost regressor | 1405 |
| 28.3.3 | Additive model framework | 1406 |
| 28.3.3.1 | Generic additive model algorithm | 1406 |
| 28.3.3.2 | Adaboost as a special additive model | 1407 |
| 28.4 | Gradient boosting machines | 1409 |
| 28.4.1 | Fundamental | 1409 |
| 28.4.2 | Gradient boosting tree | 1410 |
| 28.5 | XGBoost | 1415 |
| 28.6 | Notes on Bibliography | 1419 |

| | | |
|----------|---|------|
| 29 | UNSUPERVISED STATISTICAL LEARNING | 1420 |
| 29.1 | Singular value decomposition (SVD) and matrix factorization | 1421 |
| 29.1.1 | SVD theory | 1421 |
| 29.1.1.1 | SVD fundamentals | 1421 |
| 29.1.1.2 | SVD and matrix norm | 1423 |
| 29.1.1.3 | SVD low rank approximation | 1424 |
| 29.1.2 | Principal component analysis (PCA) | 1426 |
| 29.1.2.1 | Statistical perspective of PCA | 1426 |
| 29.1.2.2 | Geometric fundamentals of PCA | 1429 |
| 29.1.2.3 | Robust PCA with outliers | 1431 |
| 29.1.3 | Sparse coding and dictionary learning | 1433 |
| 29.1.3.1 | Sparse coding | 1433 |
| 29.1.3.2 | Dictionary learning | 1434 |
| 29.1.3.3 | Online dictionary learning | 1436 |
| 29.1.4 | Non-negative matrix factorization | 1438 |
| 29.2 | Advanced applications of matrix factorization methods | 1440 |
| 29.2.1 | Latent semantic analysis | 1440 |
| 29.2.2 | Collaborative filtering in recommender systems | 1443 |
| 29.2.3 | Co-occurrence based word embedding | 1448 |
| 29.3 | Manifold learning | 1450 |
| 29.3.1 | Overview | 1450 |
| 29.3.2 | Preliminary: multidimensional scaling (MDS) | 1450 |
| 29.3.2.1 | Motivation | 1450 |
| 29.3.2.2 | Solution to classical MDS | 1451 |
| 29.3.3 | Isomap | 1455 |
| 29.3.4 | Kernel PCA | 1456 |
| 29.3.5 | Laplacian eigenmap | 1458 |
| 29.3.5.1 | Preliminary: graph Laplacian | 1458 |
| 29.3.5.2 | Laplacian eigenmap | 1460 |
| 29.3.6 | Diffusion map | 1463 |
| 29.3.7 | Application examples | 1466 |
| 29.3.7.1 | MNIST | 1466 |
| 29.4 | Clustering | 1468 |
| 29.4.1 | Overview | 1468 |
| 29.4.2 | K-means | 1468 |
| 29.4.2.1 | Canonical K-means | 1468 |
| 29.4.2.2 | K means++ | 1471 |
| 29.4.2.3 | Kernel K means | 1472 |
| 29.4.3 | Density-based spatial clustering of applications with noise (DB-SCAN) | 1473 |
| 29.4.4 | Spectral clustering | 1475 |

| | | |
|----------|--|------|
| 29.4.5 | Gaussian mixture models (GMM) | 1476 |
| 29.4.5.1 | Preliminaries: Expectation Maximization (EM) algorithm | 1476 |
| 29.4.5.2 | The GMM model and algorithm | 1478 |
| 29.4.6 | Application examples | 1480 |
| 29.4.6.1 | Image segmentation | 1480 |
| 29.5 | Notes on Bibliography | 1482 |
| 30 | NEURAL NETWORK AND DEEP LEARNING | 1483 |
| 30.1 | Neural network foundations | 1485 |
| 30.1.1 | From machine learning to deep learning | 1485 |
| 30.1.2 | Neurons and neural networks | 1486 |
| 30.1.2.1 | Artificial neurons | 1486 |
| 30.1.2.2 | Artificial neural networks | 1488 |
| 30.1.3 | Universal approximation | 1491 |
| 30.1.4 | Training via backpropagation | 1492 |
| 30.2 | Optimization algorithms | 1498 |
| 30.2.1 | Motivation | 1498 |
| 30.2.2 | Full Batch gradient descent | 1499 |
| 30.2.3 | Minibatch stochastic gradient descent | 1500 |
| 30.2.4 | Adaptive gradient method | 1501 |
| 30.2.4.1 | Adaptive gradient (AdaGrad) | 1501 |
| 30.2.4.2 | RMSProp & AdaDelta | 1501 |
| 30.2.5 | Momentum method | 1503 |
| 30.2.6 | Combined together: adaptive momentum (Adam) | 1504 |
| 30.3 | Training and regularization techniques | 1506 |
| 30.3.1 | Choices of activation functions | 1506 |
| 30.3.2 | Weight initialization | 1506 |
| 30.3.2.1 | Motivation | 1506 |
| 30.3.2.2 | Xvaier initialization | 1507 |
| 30.3.2.3 | He initialization | 1508 |
| 30.3.3 | Data normalization | 1508 |
| 30.3.3.1 | Initial data standardization | 1508 |
| 30.3.3.2 | Batch normalization | 1509 |
| 30.3.4 | Regularization | 1510 |
| 30.3.4.1 | L_p regularization | 1510 |
| 30.3.4.2 | Weight decay | 1510 |
| 30.3.4.3 | Early stopping | 1511 |
| 30.3.4.4 | Dropout | 1511 |
| 30.3.4.5 | Data augmentation | 1512 |
| 30.3.4.6 | Label smoothing | 1512 |
| 30.4 | Feed-forward neural network examples | 1513 |

| | | |
|----------|--|------|
| 30.4.1 | Linear regression and classification | 1513 |
| 30.4.2 | Image classification | 1515 |
| 30.4.3 | Word embedding | 1518 |
| 30.4.4 | Sentiment analysis | 1522 |
| 30.4.5 | Approximating numerical partial differential equations | 1524 |
| 30.5 | Convolutional neural networks (CNN) | 1528 |
| 30.5.1 | Foundations | 1528 |
| 30.5.2 | CNN classical architectures | 1531 |
| 30.5.2.1 | LeNet | 1531 |
| 30.5.2.2 | AlexNet | 1531 |
| 30.5.2.3 | VGG | 1532 |
| 30.5.2.4 | ResNet | 1533 |
| 30.6 | CNN application examples | 1536 |
| 30.6.1 | Image classification | 1536 |
| 30.6.2 | Visualizing CNN | 1537 |
| 30.6.2.1 | Visualizing filters | 1537 |
| 30.6.2.2 | Visualizing classification activation map | 1538 |
| 30.6.3 | Autoencoders and denoising | 1540 |
| 30.6.3.1 | Autoencoders | 1540 |
| 30.6.3.2 | Denoising autoencoder | 1541 |
| 30.6.4 | Neural style transfer | 1542 |
| 30.6.5 | Visual based deep reinforcement learning | 1545 |
| 30.6.6 | Sentence classification | 1547 |
| 30.7 | Recurrent neural networks (RNN) | 1550 |
| 30.7.1 | Recurrent units | 1550 |
| 30.7.1.1 | Simple recurrent unit (SRU) | 1550 |
| 30.7.1.2 | Simple RNN and its approximation capability | 1551 |
| 30.7.1.3 | Backpropogation through time (BPTT) | 1551 |
| 30.7.2 | Recurrent unit variants | 1553 |
| 30.7.2.1 | Long short term memory (LSTM) | 1553 |
| 30.7.2.2 | Gated Recurrent Unit (GRU) | 1556 |
| 30.7.3 | Common RNN architectures | 1557 |
| 30.8 | RNN application examples | 1561 |
| 30.8.1 | Time series prediction | 1561 |
| 30.8.1.1 | Simple RNN prediction | 1561 |
| 30.8.1.2 | Deep autoregressive (DeepAR) model | 1564 |
| 30.8.1.3 | Deep factor model | 1566 |
| 30.8.2 | MNIST classification with sequential observation | 1567 |
| 30.8.3 | Sentiment classification | 1567 |
| 30.8.4 | Character-level language modeling | 1568 |
| 30.8.4.1 | Word classification | 1568 |

| | | |
|-----------|---|------|
| 30.8.4.2 | Text generation | 1569 |
| 30.9 | Sequence-to-sequence modeling | 1572 |
| 30.9.1 | Encoder decoder model | 1572 |
| 30.9.2 | Attention mechanism | 1574 |
| 30.10 | Generative adversarial network (GAN) | 1578 |
| 30.10.1 | Canonical GAN | 1578 |
| 30.10.1.1 | Basics | 1578 |
| 30.10.1.2 | An example | 1580 |
| 30.10.1.3 | Understand training difficulties in GAN | 1581 |
| 30.10.1.4 | Deep Convolutional GAN (DCGAN) | 1583 |
| 30.10.2 | Conditional GAN | 1585 |
| 30.10.3 | Wasserstein GAN (WGAN) | 1587 |
| 30.11 | Notes on Bibliography | 1591 |

vi optimal control and reinforcement learning methods

| | | |
|----------|--|------|
| 31 | CLASSICAL OPTIMAL CONTROL THEORY | 1593 |
| 31.1 | Basic problem | 1594 |
| 31.2 | Controllability & observability | 1595 |
| 31.3 | Dynamic programming principle | 1596 |
| 31.3.1 | Principle of optimality | 1596 |
| 31.3.2 | The Hamilton-Jacobi-Bellman equation (finite horizon) | 1596 |
| 31.3.3 | The Hamilton-Jacobi-Bellman equation (infinite horizon) | 1597 |
| 31.4 | Deterministic linear quadratic control | 1600 |
| 31.4.1 | Linear quadratic control (finite horizon) | 1600 |
| 31.4.2 | Linear quadratic control(infinite horizon) | 1601 |
| 31.5 | Continuous-time stochastic optimal control | 1603 |
| 31.5.1 | HJB equation for general nonlinear systems | 1603 |
| 31.5.2 | Linear Gaussian quadratic system | 1604 |
| 31.6 | Stochastic dynamic programming | 1605 |
| 31.6.1 | Discrete-time Stochastic dynamic programming: finite horizon | 1605 |
| 31.6.2 | Discrete-time stochastic dynamic programming: infinite horizon | 1607 |
| 31.6.2.1 | Fundamentals | 1607 |
| 31.6.2.2 | Convergence analysis | 1608 |
| 31.7 | Notes on bibliography | 1610 |
| 32 | REINFORCEMENT LEARNING | 1611 |
| 32.1 | Preliminaries | 1612 |
| 32.1.1 | Notations | 1612 |
| 32.1.2 | Finite state Markov decision process | 1612 |
| 32.1.3 | Policy iteration and value iteration | 1614 |

| | | |
|----------|--|------|
| 32.1.3.1 | Policy iteration | 1614 |
| 32.1.3.2 | Value iteration | 1618 |
| 32.2 | Reinforcement learning theory | 1622 |
| 32.2.1 | Overview | 1622 |
| 32.2.2 | State-action Value function (Q function) | 1624 |
| 32.2.3 | Monte-Carlo method | 1626 |
| 32.2.3.1 | On-policy value estimation | 1626 |
| 32.2.3.2 | Off-policy value estimation | 1628 |
| 32.2.3.3 | MC-based reinforcement learning control | 1628 |
| 32.2.4 | TD(0) learning | 1629 |
| 32.2.4.1 | TD(0) for value estimation | 1629 |
| 32.2.4.2 | On-policy reinforcement learning control | 1630 |
| 32.2.4.3 | Off-policy reinforcement learning control | 1631 |
| 32.2.5 | TD(n) learning | 1633 |
| 32.2.5.1 | Motivation and concepts | 1633 |
| 32.2.5.2 | TD(n) for value estimation | 1634 |
| 32.2.5.3 | TD(n) for reinforcement learning control | 1636 |
| 32.2.6 | Standing challenges in reinforcement learning | 1636 |
| 32.2.6.1 | Curse of dimensionality | 1636 |
| 32.2.6.2 | Sample efficiency | 1637 |
| 32.2.6.3 | Exploration-exploitation dilemma | 1637 |
| 32.2.6.4 | Deadly triad | 1637 |
| 32.3 | Policy gradient learning | 1638 |
| 32.3.1 | Stochastic policy gradient fundamentals | 1638 |
| 32.3.1.1 | Preliminaries: derivative and expectation | 1638 |
| 32.3.1.2 | Theoretical framework based on finite-horizon trajectories | 1639 |
| 32.3.1.3 | Theoretical framework based on distributions* | 1642 |
| 32.3.1.4 | Estimate policy gradient and basic algorithms | 1646 |
| 32.3.1.5 | Bootstrap and Actor-Critic methods | 1648 |
| 32.3.1.6 | Common stochastic policies and their representations | 1650 |
| 32.3.2 | Advanced methods for policy gradient estimation | 1652 |
| 32.3.2.1 | Stochastic policy gradient with baseline | 1652 |
| 32.3.2.2 | Generalized advantage estimation | 1655 |
| 32.3.2.3 | Summary of stochastic gradient descent forms | 1656 |
| 32.3.3 | Deterministic policy gradient | 1657 |
| 32.4 | Algorithms zoo | 1659 |
| 32.4.1 | Neural Fitted Q Iteration (NFQ) | 1659 |
| 32.4.2 | Canonical deep Q learning | 1660 |
| 32.4.3 | DQN variants | 1662 |

| | | |
|----------|---|------|
| 32.4.3.1 | Overview | 1662 |
| 32.4.3.2 | Double Q learning | 1663 |
| 32.4.3.3 | Dueling network | 1663 |
| 32.4.3.4 | Deep Recurrent Q network (DRQN) | 1664 |
| 32.4.3.5 | Asynchronous Methods | 1664 |
| 32.4.4 | Universal value function approximator | 1666 |
| 32.4.5 | Deep deterministic policy gradient (DDPG) algorithm | 1669 |
| 32.4.6 | Twin-delayed deep deterministic policy gradient (TD3) | 1671 |
| 32.4.7 | Trust Region Policy Optimization (TRPO) | 1674 |
| 32.4.7.1 | TRPO | 1674 |
| 32.4.7.2 | Evaluating Hessian of KL-divergence | 1676 |
| 32.4.8 | Proximal Policy Optimization (PPO) | 1677 |
| 32.4.9 | Soft Actor-Critic(SAC) | 1679 |
| 32.4.9.1 | Entropy regulated reinforcement learning | 1679 |
| 32.4.9.2 | The SAC algorithm | 1680 |
| 32.4.10 | Evolution strategies | 1681 |
| 32.5 | Advanced training strategies | 1684 |
| 32.5.1 | Priority experience replay | 1684 |
| 32.5.2 | Hindsight experience generation | 1685 |
| 32.5.3 | Reverse goal generation | 1686 |
| 32.5.4 | Reverse goal generation on low-dimensional manifolds | 1687 |
| 32.5.4.1 | Key idea | 1687 |
| 32.5.4.2 | Example: navigation on a curved surface | 1688 |
| 32.6 | Notes on bibliography | 1689 |

vii appendix

| | | |
|-------|--|------|
| A | SUPPLEMENTAL MATHEMATICAL FACTS | 1691 |
| A.1 | Basic logic for proof | 1692 |
| A.2 | Some common limits | 1693 |
| A.3 | Common series summation | 1695 |
| A.4 | Some common spaces | 1696 |
| A.4.1 | Notations on continuously differentiable functions | 1697 |
| A.5 | Different modes of continuity | 1698 |
| A.5.1 | continuity vs. uniform continuity | 1699 |
| A.6 | Exchanges of limits | 1700 |
| A.6.1 | Overall remark | 1700 |
| A.6.2 | exchange limits with infinite summations | 1700 |
| A.6.3 | Exchange limits with integration and differentiation | 1700 |
| A.6.4 | Exchange differentiation with integration | 1701 |
| A.6.5 | Exchange limit and function evaluations | 1702 |
| A.7 | Useful inequalities | 1703 |

| | | |
|--------------------|---|------|
| A.7.1 | Gronwall's inequality | 1703 |
| A.7.2 | Inequality for norms | 1703 |
| A.7.3 | Young's inequality for product | 1704 |
| A.8 | Useful properties of matrix | 1705 |
| A.8.1 | Matrix derivatives | 1705 |
| A.8.2 | Matrix inversion lemma | 1705 |
| A.8.3 | Block matrix | 1706 |
| A.8.4 | Matrix trace | 1707 |
| A.8.5 | Matrix elementary operator | 1708 |
| A.8.6 | Matrix determinant | 1710 |
| A.9 | Numerical integration | 1711 |
| A.9.1 | Gaussian quadrature | 1712 |
| A.10 | Vector calculus | 1714 |
| A.11 | Numerical linear algebra computation complexity | 1715 |
| A.12 | Distributions | 1716 |
| A.13 | Common integrals | 1717 |
| A.14 | Nonlinear root finding | 1718 |
| A.14.1 | Bisection method | 1718 |
| A.14.2 | Newton method | 1718 |
| A.14.3 | Secant method | 1718 |
| A.15 | Interpolation | 1720 |
| A.15.1 | cubic interpolation | 1720 |
| Alphabetical Index | | 1721 |

Part I

MATHEMATICAL FOUNDATIONS