

Lecture 05

Geometry of Least Squares

16 September 2015

Taylor B. Arnold
Yale Statistics
STAT 312/612

The Yale University logo, featuring the word "Yale" in a blue, serif font.

Goals for today

1. Geometry of least squares
2. Projection matrix P and annihilator matrix M
3. Application to

GEOMETRY OF LEAST SQUARES

Last time, we established that the least squares solution to the model:

$$y = X\beta + \epsilon$$

Yields the solution:

$$\hat{\beta} = (X^t X)^{-1} X^t y$$

As long as the matrix $X^t X$ is invertable.

Define the column space of the matrix X as:

$$\mathcal{R}(X) = \{\theta : \theta = X\beta, \beta \in \mathbb{R}^p\} \subset \mathbb{R}^n$$

This is the space spanned by the p columns on X sitting in n -dimensional space.

Define the column space of the matrix X as:

$$\mathcal{R}(X) = \{\theta : \theta = X\beta, \beta \in \mathbb{R}^p\} \subset \mathbb{R}^n$$

This is the space spanned by the p columns on X sitting in n -dimensional space.

Notice that the least squares problem can be re-written as:

$$\hat{\theta} = \arg \min_{\theta} \{ \|y - \theta\|_2^2, \quad \text{s.t. } \theta = X\beta \}$$

Where $\hat{\beta} = X\hat{\theta}$.

Theorem 3.2 (p.g. 37, Rao & Toutenburg) The minimum, $\hat{\theta}$ is attained when $(y - \hat{\theta}) \perp \mathcal{R}(X)$. In other words, $(y - \hat{\theta})$ is perpendicular to all vectors in \mathcal{R} .

Proof. Pick a $\hat{\theta}$ in \mathcal{R} such that $(y - \hat{\theta}) \perp \mathcal{R}(X)$.

Proof. Pick a $\hat{\theta}$ in \mathcal{R} such that $(y - \hat{\theta}) \perp \mathcal{R}(X)$. This implies that $X^t(y - \hat{\theta}) = 0$. Then for all $\theta \in \mathcal{R}$:

$$\|y - \theta\|^2 = (y - \hat{\theta} + \hat{\theta} - \theta)^t (y - \hat{\theta} + \hat{\theta} - \theta)$$

Proof. Pick a $\hat{\theta}$ in \mathcal{R} such that $(y - \hat{\theta}) \perp \mathcal{R}(X)$. This implies that $X^t(y - \hat{\theta}) = 0$. Then for all $\theta \in \mathcal{R}$:

$$\begin{aligned} \|y - \theta\|^2 &= (y - \hat{\theta} + \hat{\theta} - \theta)^t (y - \hat{\theta} + \hat{\theta} - \theta) \\ &= (y - \hat{\theta})^t (y - \hat{\theta}) + (\hat{\theta} - \theta)^t (\hat{\theta} - \theta) + 2(y - \hat{\theta})^t (\hat{\theta} - \theta) \end{aligned}$$

Proof. Pick a $\hat{\theta}$ in \mathcal{R} such that $(y - \hat{\theta}) \perp \mathcal{R}(X)$. This implies that $X^t(y - \hat{\theta}) = 0$. Then for all $\theta \in \mathcal{R}$:

$$\begin{aligned} \|y - \theta\|^2 &= (y - \hat{\theta} + \hat{\theta} - \theta)^t (y - \hat{\theta} + \hat{\theta} - \theta) \\ &= (y - \hat{\theta})^t (y - \hat{\theta}) + (\hat{\theta} - \theta)^t (\hat{\theta} - \theta) + 2(y - \hat{\theta})^t (\hat{\theta} - \theta) \\ &= (y - \hat{\theta})^t (y - \hat{\theta}) + (\hat{\theta} - \theta)^t (\hat{\theta} - \theta) \end{aligned}$$

Proof. Pick a $\hat{\theta}$ in \mathcal{R} such that $(y - \hat{\theta}) \perp \mathcal{R}(X)$. This implies that $X^t(y - \hat{\theta}) = 0$. Then for all $\theta \in \mathcal{R}$:

$$\begin{aligned} \|y - \theta\|^2 &= (y - \hat{\theta} + \hat{\theta} - \theta)^t(y - \hat{\theta} + \hat{\theta} - \theta) \\ &= (y - \hat{\theta})^t(y - \hat{\theta}) + (\hat{\theta} - \theta)^t(\hat{\theta} - \theta) + 2(y - \hat{\theta})^t(\hat{\theta} - \theta) \\ &= (y - \hat{\theta})^t(y - \hat{\theta}) + (\hat{\theta} - \theta)^t(\hat{\theta} - \theta) \\ &= \|y - \hat{\theta}\|_2^2 + \|\hat{\theta} - \theta\|_2^2 \end{aligned}$$

Proof. Pick a $\hat{\theta}$ in \mathcal{R} such that $(y - \hat{\theta}) \perp \mathcal{R}(X)$. This implies that $X^t(y - \hat{\theta}) = 0$. Then for all $\theta \in \mathcal{R}$:

$$\begin{aligned} \|y - \theta\|^2 &= (y - \hat{\theta} + \hat{\theta} - \theta)^t (y - \hat{\theta} + \hat{\theta} - \theta) \\ &= (y - \hat{\theta})^t (y - \hat{\theta}) + (\hat{\theta} - \theta)^t (\hat{\theta} - \theta) + 2(y - \hat{\theta})^t (\hat{\theta} - \theta) \\ &= (y - \hat{\theta})^t (y - \hat{\theta}) + (\hat{\theta} - \theta)^t (\hat{\theta} - \theta) \\ &= \|y - \hat{\theta}\|_2^2 + \|\hat{\theta} - \theta\|_2^2 \\ &\geq \|y - \hat{\theta}\|_2^2 \end{aligned}$$

Proof: Pick a $\hat{\theta}$ in \mathcal{R} such that $(y - \hat{\theta}) \perp \mathcal{R}(X)$. This implies that $X^t(y - \hat{\theta}) = 0$. Then for all $\theta \in \mathcal{R}$:

$$\begin{aligned} \|y - \theta\|^2 &= (y - \hat{\theta} + \hat{\theta} - \theta)^t(y - \hat{\theta} + \hat{\theta} - \theta) \\ &= (y - \hat{\theta})^t(y - \hat{\theta}) + (\hat{\theta} - \theta)^t(\hat{\theta} - \theta) + 2(y - \hat{\theta})^t(\hat{\theta} - \theta) \\ &= (y - \hat{\theta})^t(y - \hat{\theta}) + (\hat{\theta} - \theta)^t(\hat{\theta} - \theta) \\ &= \|y - \hat{\theta}\|_2^2 + \|\hat{\theta} - \theta\|_2^2 \\ &\geq \|y - \hat{\theta}\|_2^2 \end{aligned}$$

So, if such a $\hat{\theta}$ exists it attains the minimum. To see that it does, write $\hat{\theta} = X\hat{\beta}$.

Proof: Pick a $\hat{\theta}$ in \mathcal{R} such that $(y - \hat{\theta}) \perp \mathcal{R}(X)$. This implies that $X^t(y - \hat{\theta}) = 0$. Then for all $\theta \in \mathcal{R}$:

$$\begin{aligned} \|y - \theta\|^2 &= (y - \hat{\theta} + \hat{\theta} - \theta)^t(y - \hat{\theta} + \hat{\theta} - \theta) \\ &= (y - \hat{\theta})^t(y - \hat{\theta}) + (\hat{\theta} - \theta)^t(\hat{\theta} - \theta) + 2(y - \hat{\theta})^t(\hat{\theta} - \theta) \\ &= (y - \hat{\theta})^t(y - \hat{\theta}) + (\hat{\theta} - \theta)^t(\hat{\theta} - \theta) \\ &= \|y - \hat{\theta}\|_2^2 + \|\hat{\theta} - \theta\|_2^2 \\ &\geq \|y - \hat{\theta}\|_2^2 \end{aligned}$$

So, if such a $\hat{\theta}$ exists it attains the minimum. To see that it does, write $\hat{\theta} = X\hat{\beta}$. Then:

$$\begin{aligned} X^t(y - \hat{\theta}) &= X^t(y - X\hat{\beta}) \\ &= X^ty - X^tX\hat{\beta} \end{aligned}$$

To see that such a $\hat{\theta}$ does exist, write $\hat{\theta} = X\hat{\beta}$.

To see that such a $\hat{\theta}$ does exist, write $\hat{\theta} = X\hat{\beta}$. Then:

$$X^t(y - \hat{\theta}) = X^t(y - X\hat{\beta})$$

To see that such a $\hat{\theta}$ does exist, write $\hat{\theta} = X\hat{\beta}$. Then:

$$\begin{aligned} X^t(y - \hat{\theta}) &= X^t(y - X\hat{\beta}) \\ &= X^ty - X^tX\hat{\beta} \end{aligned}$$

To see that such a $\hat{\theta}$ does exist, write $\hat{\theta} = X\hat{\beta}$. Then:

$$\begin{aligned} X^t(y - \hat{\theta}) &= X^t(y - X\hat{\beta}) \\ &= X^ty - X^tX\hat{\beta} \\ &= X^ty - X^tX(X^tX)^{-1}X^ty \end{aligned}$$

To see that such a $\hat{\theta}$ does exist, write $\hat{\theta} = X\hat{\beta}$. Then:

$$\begin{aligned}X^t(y - \hat{\theta}) &= X^t(y - X\hat{\beta}) \\&= X^ty - X^tX\hat{\beta} \\&= X^ty - X^tX(X^tX)^{-1}X^ty \\&= X^ty - X^ty \\&= 0\end{aligned}$$

And therefore our proposed $\hat{\theta} \in \mathcal{R}(X)$.

From this geometric interpretation of the least squares estimator, we introduce an important matrix P_X called the *projection matrix*.

$$P_X = X(X^tX)^{-1}X^t$$

I'll often drop the subscript as it should be understood that the projection is on the data matrix X .

Notice that $PX = X$:

$$\begin{aligned} PX &= X(X^tX)^{-1}X^tX \\ &= X \end{aligned}$$

Notice that $PX = X$:

$$\begin{aligned} PX &= X(X^tX)^{-1}X^tX \\ &= X \end{aligned}$$

And Py gives the fitted values \hat{y} :

$$\begin{aligned} Py &= X(X^tX)^{-1}X^tXy \\ &= X\hat{\beta} \\ &= \hat{\theta} \\ &= \hat{y} \end{aligned}$$

Do you see why the projection matrix is called the projection matrix?

Notice that $PX = X$:


$$\begin{aligned} PX &= X(X^tX)^{-1}X^tX \\ &= X \end{aligned}$$

And Py gives the fitted values \hat{y} :

$$\begin{aligned} Py &= X(X^tX)^{-1}X^tXy \\ &= X\hat{\beta} \\ &= \hat{\theta} \\ &= \hat{y} \end{aligned}$$

Do you see why the projection matrix is called the projection matrix?

The projection matrix is sometimes called the *hat matrix*. Any thoughts as to why?



A closely related matrix to P is the *annihilator matrix* M :

$$M = I_n - P$$

A closely related matrix to P is the *annihilator matrix* M :

$$M = I_n - P$$

It gets its name because $MX = 0$.

The matrix $P = X(X^tX)^{-1}X^t$ is clearly symmetric. It is also idempotent:

$$\begin{aligned} P^2 &= X(X^tX)^{-1}X^tX(X^tX)^{-1}X^t \\ &= X(X^tX)^{-1}(X^tX)(X^tX)^{-1}X^t \\ &= X(X^tX)^{-1}X^t \\ &= P \end{aligned}$$

M is also symmetric

$$\begin{aligned} M^t &= (I_n - P)^t \\ &= (I_n - P^t) \\ &= M \end{aligned}$$

M is also symmetric

$$\begin{aligned}M^t &= (I_n - P)^t \\&= (I_n - P^t) \\&= M\end{aligned}$$

And idempotent:

$$\begin{aligned}M^2 &= (I_n - P)^2 \\&= (I_n - P)(I_n - P)\end{aligned}$$

M is also symmetric

$$\begin{aligned}M^t &= (I_n - P)^t \\&= (I_n - P^t) \\&= M\end{aligned}$$

And idempotent:

$$\begin{aligned}M^2 &= (I_n - P)^2 \\&= (I_n - P)(I_n - P) \\&= I_n - 2 * P + P^2\end{aligned}$$

M is also symmetric

$$\begin{aligned}M^t &= (I_n - P)^t \\&= (I_n - P^t) \\&= M\end{aligned}$$

And idempotent:

$$\begin{aligned}M^2 &= (I_n - P)^2 \\&= (I_n - P)(I_n - P) \\&= I_n - 2 * P + P^2 \\&= I_n - 2 * P + P\end{aligned}$$

M is also symmetric

$$\begin{aligned}M^t &= (I_n - P)^t \\&= (I_n - P^t) \\&= M\end{aligned}$$

And idempotent:

$$\begin{aligned}M^2 &= (I_n - P)^2 \\&= (I_n - P)(I_n - P) \\&= I_n - 2 * P + P^2 \\&= I_n - 2 * P + P \\&= I_n - P \\&= M\end{aligned}$$

M is also symmetric

$$\begin{aligned}M^t &= (I_n - P)^t \\&= (I_n - P^t) \\&= M\end{aligned}$$

And idempotent:

$$\begin{aligned}M^2 &= (I_n - P)^2 \\&= (I_n - P)(I_n - P) \\&= I_n - 2 * P + P^2 \\&= I_n - 2 * P + P \\&= I_n - P \\&= M\end{aligned}$$

These properties both make sense given the geometric interpretation of P and M as projections; into the column space of X and the complement of the column space of X .

These properties are quite useful. Notice how we can easily rewrite the following for the residual vector $r = y - X\hat{\beta}$:

$$r = y - X\hat{\beta}$$

These properties are quite useful. Notice how we can easily rewrite the following for the residual vector $r = y - X\hat{\beta}$:

$$\begin{aligned} r &= y - X\hat{\beta} \\ &= y - Py \end{aligned}$$

These properties are quite useful. Notice how we can easily rewrite the following for the residual vector $r = y - X\hat{\beta}$:

$$\begin{aligned}r &= y - X\hat{\beta} \\&= y - Py \\&= (I_n - P)y\end{aligned}$$

These properties are quite useful. Notice how we can easily rewrite the following for the residual vector $r = y - X\hat{\beta}$:

$$\begin{aligned}r &= y - X\hat{\beta} \\&= y - Py \\&= (I_n - P)y \\&= My\end{aligned}$$

These properties are quite useful. Notice how we can easily rewrite the following for the residual vector $r = y - X\hat{\beta}$:

$$\begin{aligned} r &= y - X\hat{\beta} \\ &= y - Py \\ &= (I_n - P)y \\ &= My \\ &= M(X\beta - \epsilon) \end{aligned}$$

These properties are quite useful. Notice how we can easily rewrite the following for the residual vector $r = y - X\hat{\beta}$:

$$\begin{aligned} r &= y - X\hat{\beta} \\ &= y - Py \\ &= (I_n - P)y \\ &= My \\ &= M(X\beta - \epsilon) \\ &= M\epsilon \end{aligned}$$

These properties are quite useful. Notice how we can easily rewrite the following for the residual vector $r = y - X\hat{\beta}$:

$$\begin{aligned} r &= y - X\hat{\beta} \\ &= y - Py \\ &= (I_n - P)y \\ &= My \\ &= M(X\beta - \epsilon) \\ &= M\epsilon \end{aligned}$$

The matrices P and M not only help make the derivation easier, they also give geometric insight into what we are doing.

One particularly useful formula will be writing the squared residuals as:

$$\begin{aligned} ||r||_2^2 &= ||M\epsilon||_2^2 \\ &= \epsilon^t M^t M \epsilon \\ &= \epsilon^t M \epsilon \end{aligned}$$

One particularly useful formula will be writing the squared residuals as:

$$\begin{aligned} ||r||_2^2 &= ||M\epsilon||_2^2 \\ &= \epsilon^t M^t M \epsilon \\ &= \epsilon^t M \epsilon \end{aligned}$$

So the matrix M translates the sum of squared residuals into the sum of the square errors, which are estimated by the residuals.

APPLICATIONS