



BITS Pilani

Pilani Campus

Mathematical Foundations for Data Science

MFDS Team



BITS Pilani
Pilani Campus

DSECL ZC416, MFDS

Lecture No.11

Slides are adapted version of slides from McGraw Hill Education

Agenda




- Functions
 - Domain
 - Codomain and
 - Range
- Types of Functions
 - Injective,
 - Surjective
 - Bijective
- Inverse of functions
- Composition of functions
- Floor and Ceiling operators

Functions

Definition: Let A and B be nonempty sets. A *function* f from A to B , denoted $f: A \rightarrow B$ is an assignment of each element of A to exactly one element of B . We write $f(a) = b$ if b is the unique element of B assigned by the function f to the element a of A .

Functions are sometimes called *mappings* or *transformations*.



Functions

A function $f: A \rightarrow B$ can also be defined as a subset of $A \times B$ (a relation). This subset is restricted to be a relation where no two elements of the relation have the same first element.

Specifically, a function f from A to B contains one, and only one ordered pair (a, b) for every element $a \in A$.

$$\forall x [x \in A \rightarrow \exists y [y \in B \wedge (x, y) \in f]]$$

$$\forall x, y_1, y_2 [(x, y_1) \in f \wedge (x, y_2) \in f] \rightarrow y_1 = y_2]$$


Functions

Given a function $f: A \rightarrow B$:

- We say f maps A to B or f is a *mapping* from A to B .
- A is called the *domain* of f .
- B is called the *co-domain* of f .

If $f(a) = b$,

- then b is called the *image* of a under f .
- a is called the *preimage* of b .



The range of f is the set of all images of points in A under f . We denote it by $f(A)$.

Two functions are *equal* when they have the same domain, the same co-domain and map each element of the domain to the same element of the co-domain.

Representing Functions

Functions may be specified in different ways:

- An explicit statement of the assignment.

Students and grades example.

- A formula.


$$f(x) = x + 1$$

- A computer program.

- A Java program that when given an integer n , produces the n th Fibonacci Number

Functions

- $f(a) =$
- The image of d is ?
- The preimage(s) of z is (are) ? $\{a, c, d\}$
- The domain of f is ? A
- The co-domain of f is ? B
- The preimage of y is ? b
- $f(A) = \{y, z\}$




Question on Functions & Sets

If $f : A \rightarrow B$ and S is a subset of A , then

$$f(S) = \{f(s) | s \in S\}$$


$f\{a,b,c\}$ is ? $\{y,z\}$

$f\{c,d\}$ is ? $\{z\}$




Injection

Definition: A function f is said to be *one-to-one*, or *injective*, if and only if $f(a) = f(b)$ implies that $a = b$ for all a and b in the domain of f . A function is said to be an *injection* if it is one-to-one.




Surjection

Definition: A function f from A to B is called *onto* or *surjective*, if and only if for every element $a \in A$ there is an element $b \in B$ with $f(a) = b$. A function f is called a *surjection* if it is *onto*.




Bijection

Definition: A function f is a *one-to-one correspondence*, or a *bijection*, if it is both one-to-one and onto (surjective and injective).




Different Types of Correspondences




NOT a
Function

A has many B




General
Function

B can have many A




Injective
(not surjective)

B can't have many A



Surjective
(not injective)

Every B has some A



Bijjective
(injective, surjective)

A to B, perfectly

Definitions

Suppose that $f : A \rightarrow B$.

To show that f is injective Show that if $f(x) = f(y)$ for arbitrary $x, y \in A$ with $x \neq y$, then $x = y$.

To show that f is not injective Find particular elements $x, y \in A$ such that $x \neq y$ and $f(x) = f(y)$.

To show that f is surjective Consider an arbitrary element $y \in B$ and find an element $x \in A$ such that $f(x) = y$.

To show that f is not surjective Find a particular $y \in B$ such that $f(x) \neq y$ for all $x \in A$.

Examples

Example 1: Let f be the function from $\{a,b,c,d\}$ to $\{1,2,3\}$ defined by $f(a) = 3$, $f(b) = 2$, $f(c) = 1$, and $f(d) = 3$. Is f an onto function?

Solution: Yes, f is onto since all three elements of the codomain are images of elements in the domain. If the codomain were changed to $\{1,2,3,4\}$, f would not be onto.

Example 2: Is the function $f(x) = x^2$ from the set of integers to the set of integers onto?


Solution: No, f is not onto because there is no integer x with $x^2 = -1$, for example.

Inverse Functions


Definition: Let f be a bijection from A to B . Then the *inverse* of f , denoted f^{-1} , is the function from B to A defined as

$$f^{-1}(y) = x \text{ iff } f(x) = y$$

No inverse exists unless f is a bijection. Why?



Inverse Functions



Examples

Example 1: Let f be the function from $\{a,b,c\}$ to $\{1,2,3\}$ such that $f(a) = 2$, $f(b) = 3$, and $f(c) = 1$. Is f invertible and if so what is its inverse?

Solution: The function f is invertible because it is a one-to-one correspondence. The inverse function f^{-1} reverses the correspondence given by f , so $f^{-1}(1) = c$, $f^{-1}(2) = a$, and $f^{-1}(3) = b$.

Example 2: Let $f: \mathbf{Z} \rightarrow \mathbf{Z}$ be such that $f(x) = x + 1$. Is f invertible, and if so, what is its inverse?

Solution: The function f is invertible because it is a one-to-one correspondence. The inverse function f^{-1} reverses the correspondence so $f^{-1}(y) = y - 1$.

Contd..


Example 3: Let $f: \mathbf{R} \rightarrow \mathbf{R}$ be such that $f(x) = x^2$. Is f invertible, and if so, what is its inverse?

Solution: The function f is not invertible because it is not one-to-one .


Composition

Definition: Let $f: B \rightarrow C$, $g: A \rightarrow B$. The *composition of f with g* , denoted $f \circ g$ is the function from A to C defined by

$$f \circ g(x) = f(g(x))$$



Composition



Examples

Example 1: If $f(x) = x^2$ and $g(x) = 2x + 1$, then

$$f(g(x)) = (2x + 1)^2$$

$$g(f(x)) = 2x^2 + 1$$

Example 2: Let g be the function from the set $\{a, b, c\}$ to itself such that $g(a) = b$, $g(b) = c$, and $g(c) = a$. Let f be the function from the set $\{a, b, c\}$ to the set $\{1, 2, 3\}$ such that $f(a) = 3$, $f(b) = 2$, and $f(c) = 1$.

What is the composition of f and g , and what is the composition of g and f .

Solution: The composition $f \circ g$ is defined by

$$f \circ g(a) = f(g(a)) = f(b) = 2.$$

$$f \circ g(b) = f(g(b)) = f(c) = 1.$$

$$f \circ g(c) = f(g(c)) = f(a) = 3.$$

Note that $g \circ f$ is not defined, because the range of f is not a subset of the domain of g .

Contd..

Example 3: Let f and g be functions from the set of integers to the set of integers defined by $f(x) = 2x + 3$ and $g(x) = 3x + 2$.

What is the composition of f and g , and also the composition of g and f ?


Solution:

$$f \circ g (x) = f(g(x)) = f(3x + 2) = 2(3x + 2) + 3 = 6x + 7$$


$$g \circ f (x) = g(f(x)) = g(2x + 3) = 3(2x + 3) + 2 = 6x + 11$$

Graphs

Let f be a function from the set A to the set B . The *graph* of the function f is the set of ordered pairs $\{(a,b) \mid a \in A \text{ and } f(a) = b\}$.



Graph of $f(n) = 2n + 1$
from \mathbb{Z} to \mathbb{Z}



Graph of $f(x) = x^2$
from \mathbb{Z} to \mathbb{Z}

Important Functions

The *floor* function, denoted $f(x) = \lfloor x \rfloor$

is the largest integer less than or equal to x .

The *ceiling* function, denoted $f(x) = \lceil x \rceil$


is the smallest integer greater than or equal to x

Example:


$$\lceil 3.5 \rceil = 4 \quad \lfloor 3.5 \rfloor = 3$$

$$\lceil -1.5 \rceil = -1 \quad \lfloor -1.5 \rfloor = -2$$

Floor and Ceiling Functions



(a) $y = [x]$



(b) $y = [x]$

Graph of (a) Floor and (b) Ceiling Functions

Properties

TABLE 1 Useful Properties of the Floor and Ceiling Functions.

(n is an integer, x is a real number)

(1a) $\lfloor x \rfloor = n$ if and only if $n \leq x < n + 1$

(1b) $\lceil x \rceil = n$ if and only if $n - 1 < x \leq n$

(1c) $\lfloor x \rfloor = n$ if and only if $x - 1 < n \leq x$

(1d) $\lceil x \rceil = n$ if and only if $x \leq n < x + 1$

(2) $x - 1 < \lfloor x \rfloor \leq x \leq \lceil x \rceil < x + 1$

(3a) $\lfloor -x \rfloor = -\lceil x \rceil$

(3b) $\lceil -x \rceil = -\lfloor x \rfloor$

(4a) $\lfloor x + n \rfloor = \lfloor x \rfloor + n$

(4b) $\lceil x + n \rceil = \lceil x \rceil + n$

A Classical Proof

Example: Prove that x is a real number, then

$$\lfloor 2x \rfloor = \lfloor x \rfloor + \lfloor x + 1/2 \rfloor$$

Solution: Let $x = n + \varepsilon$, where n is an integer and $0 \leq \varepsilon < 1$.

Case 1: $\varepsilon < 1/2$

- $2x = 2n + 2\varepsilon$ and $\lfloor 2x \rfloor = 2n$, since $0 \leq 2\varepsilon < 1$.
- $\lfloor x + 1/2 \rfloor = n$, since $x + 1/2 = n + (1/2 + \varepsilon)$ and $0 \leq 1/2 + \varepsilon < 1$.
- Hence, $\lfloor 2x \rfloor = 2n$ and $\lfloor x \rfloor + \lfloor x + 1/2 \rfloor = n + n = 2n$.

Case 2: $\varepsilon \geq 1/2$

- $2x = 2n + 2\varepsilon = (2n + 1) + (2\varepsilon - 1)$ and $\lfloor 2x \rfloor = 2n + 1$, since $0 \leq 2\varepsilon - 1 < 1$.
- $\lfloor x + 1/2 \rfloor = \lfloor n + (1/2 + \varepsilon) \rfloor = \lfloor n + 1 + (\varepsilon - 1/2) \rfloor = n + 1$ since $0 \leq \varepsilon - 1/2 < 1$.
- Hence, $\lfloor 2x \rfloor = 2n + 1$ and $\lfloor x \rfloor + \lfloor x + 1/2 \rfloor = n + (n + 1) = 2n + 1$.