

Source coverage: Full paper
Extraction confidence: High
Locator mode: page-grounded
Primary analytical lens: methods
Secondary analytical lens: discovery
Context verification: Paper-only
Card completeness: Complete relative to supplied source

01 基本信息

Field	Value	Source
Title	Preserving DEG Rankings for Gene Discovery in Histology-Based Spatial Gene Expression Prediction	[Paper] [Paper: PDF p. 1]
Authors	Kaito Shiku, Kazuya Nishimura, Yasuhiro Kojima, Ryoma Bise	[Paper] [Paper: PDF p. 1]
Venue	arXiv preprint (cs.CV)	[Paper] [Paper: PDF p. 1]
URL	http://arxiv.org/abs/2609.33928v1	[Paper meta]
PDF URL	https://arxiv.org/pdf/2609.33928v1	[Paper meta]
Publication date	2026-09-27	[Paper meta]
Core task	Image-based Differential Expression Ranking (IDER)	[Paper] [Paper: PDF p. 2]
Key metric	nDCGDEG@200, SCCDEG, pathway Jaccard overlap	[Paper] [Paper: PDF p. 6–8]
Primary datasets	HEST-1K (Bowel/Ovary/Lymph Node), Her2st	[Paper] [Paper: PDF p. 6–7]
Proxy contrast source	K-means clustering on CONCH embeddings	[Paper] [Paper: PDF p. 5, p. 6]
Model architecture	Frozen CONCH + 3-layer MLP	[Paper] [Paper: PDF p. 6]

02 一句话总结

该论文指出：当前基于组织学图像预测空间基因表达（ST prediction）的方法普遍采用“逐基因空间轮廓重建”目标（如PCC、MSE），但这一目标与下游核心任务——差异表达基因（DEG）发现——存在根本错配：DEG发现依赖的是**跨基因的对比统计排序**（contrast-specific ranked gene list），而非单个基因在空间上的表达拟合精度。为此，作者提出**Image-based Differential Expression Ranking (IDER)** 任务，并设计可微分的U-statistic对齐目标（Equation 1–2），利用形态学聚类生成的代理对比（proxy contrasts）进行无标注训练，使模型能直接优化DEG排序保真度。实验证明其在nDCGDEG@200和通路富集重叠（Jaccard）上显著优于所有常规重建目标，且可即插即用提升ST-Net、TRIPLEX等现有架构；但其在传统per-gene PCC指标上表现中等，印证了“下游任务对齐”与“重建精度”的权衡本质。

03 研究问题

- **核心问题**：为什么现有histology-to-ST模型在per-gene空间轮廓重建（如PCC）上表现良好，却无法可靠支持下游DEG发现？
- **隐含假设**：常规重建目标（如PCC）虽能保证每个基因在空间维度上的表达趋势相似，但无法约束不同基因在**对比统计量（如U-stat）上的相对排序关系**，而该排序正是DEG报告、通路解释和生物学验证的起点。
- **方法路径**：不改变模型结构，而是重构训练目标——从“最小化每个基因的空间误差”转向“最大化预测U-stat向量与真实U-stat向量在基因轴上的Pearson相关性（Equation 2）”，并辅以轻量级PCC正则项（Equation 3）维持基本空间一致性。

04 研究背景与发展路径

- **起点**：ST技术成本高、通量低，亟需用广泛存在的H&E图像预测空间基因表达（[Paper] [Paper: PDF p. 1]）。
- **主流范式**：ST-Net [8]、TRIPLEX [4]、GCN/Transformer [6,38]、检索/扩散模型 [35,39] 均以**逐基因空间重建**为优化目标（MSE/PCC）或评估标准（Figure 1a）[Paper] [Paper: PDF p. 1–2]。
- **瓶颈浮现**：RankByGene [11] 和 STRank [25] 已尝试引入排序思想，但聚焦于**空间位置间表达排序**（spot-wise ranking），而非**基因间差异证据排序**（gene-wise DEG ranking）[Paper] [Paper: PDF p. 3]。
- **关键洞察跃迁**：作者识别出“评价轴错位”——重建目标沿空间轴（spot）求平均，而DEG分析沿基因轴（gene）做排序；因此必须将评价与优化轴统一到**基因维度**，即用U-stat向量间的相关性替代单基因PCC [Paper] [Paper: PDF p. 2, 4]。
- **技术支点**：CONCH病理基础模型 [19] 提供鲁棒形态学表征，使其能通过无监督聚类生成可信的proxy contrasts，绕过对昂贵病理标注的依赖 [Paper] [Paper: PDF p. 5, 6]。

05 论文识别的核心痛点

Pain point	Manifestation	Cause or author explanation	Evidence from the paper
Objective misalignment	Models with high per-gene PCC (e.g., MSE/PCC baselines) show poor nDCGDEG@200 and low pathway Jaccard overlap	Conventional objectives optimize spatial-profile agreement <i>within each gene</i> , but DEG discovery depends on <i>cross-gene ranking by contrast statistics</i> [Paper] [Paper: PDF p. 1–2]	Table 5 shows MSE/PCC achieve highest PCC (0.235–0.345), yet Table 1/3 show they rank worst on nDCGDEG@200 (e.g., 0.434 vs. 0.531 for Ours); Table 2 shows their pathway Jaccard is lowest (0.326–0.345 vs. 0.381 for Ours) [Paper] [Paper: PDF p. 7–8]
Annotation bottleneck	Real tissue-region annotations (e.g., Her2st) are scarce, costly, and patient-specific, limiting generalizable DEG analysis	Biological group labels are often unavailable; morphology-derived proxy groups offer scalable, annotation-free alternatives [Paper] [Paper: PDF p. 6]	Section 7.1 explicitly constructs groups via K-means on CONCH features <i>without any annotation</i> , and Figure 2 shows strong generalization across k=2–10 clusters [Paper] [Paper: PDF p. 6–7]

Pain point	Manifestation	Cause or author explanation	Evidence from the paper
Non-differentiability of ranking	Direct optimization of argsort-based DEG ranking (π_c) is non-differentiable, blocking end-to-end training	Sorting and indicator functions (e.g., $1[y_{i,g} > y_{j,g}]$) break gradient flow [Paper] [Paper: PDF p. 4–5]	Equation 1 replaces hard indicator with temperature-scaled sigmoid $\sigma((\hat{y}_{i,g} - \hat{y}_{j,g})/\tau)$; Equation 2 uses Corrg over genes instead of discrete ranking loss [Paper] [Paper: PDF p. 5]

06 核心思想

论文的核心思想是**将ST预测的任务定义从“表达值重建”升维至“生物学决策保真”**：不再问“预测的每个基因表达值在空间上有多准”，而是问“预测的表达值能否导出与真实数据一致的、用于生物学发现的基因优先级列表”。这体现为三个递进层次：

- 1. **评价升维**：用SCCDEG（Spearman across genes）和nDCGDEG@k（top-k recovery）替代传统PCC（Pearson across spots）；
- 2. **目标可微化**：将非可微的Mann–Whitney U statistic离散比较（Equation in Sec. 4）替换为连续sigmoid近似（Equation 1），再以基因轴Pearson相关（Equation 2）作为排序保真代理损失；
- 3. **监督弱化**：放弃对真实生物标签（如tissue region）的依赖，转而用CONCH特征聚类生成形态学proxy contrasts，在训练中动态采样one-vs-rest组别，使模型学习“哪些基因的表达变化最能区分形态差异”这一本质信号 [Paper] [Paper: PDF p. 5–6]。

07 方法总览

该方法是一个**目标函数即插即用框架**，不修改模型架构，仅替换训练损失：

- **输入**：组织学图像块 x_i 与对应spot真实表达 y_i （G维向量）；
- **特征提取**：冻结CONCH模型提取图像嵌入，接3层MLP输出预测表达 $\hat{y}_i$ ；
- **代理对比构建**：对训练集CONCH嵌入做K-means（k=10），每轮随机选一簇为 A_c （positive），其余为 B_c （negative）；
- **可微U-stat计算**：对每个对比c和基因g，按Equation 1计算 $\sum_{i \in A_c} \sum_{j \in B_c} \sigma((\hat{y}_{i,g} - \hat{y}_{j,g})/\tau)$ ，得向量 $\hat{U}_c \in \mathbb{R}^G$ ；
- **主损失LDEG**：对每个c，计算 $\text{Corrg}(\hat{U}_c, U_c)$ （ U_c 为真实U-stat向量），取均值后用 $1 - \text{Corrg}$ 作为损失（Equation 2）；
- **辅助损失Lconst**：标准per-gene PCC损失，加权 $\lambda=1.0$ （Equation 3）；
- **输出**：预测表达 $\hat{y}_i \rightarrow$ 可计算任意对比下的 $\hat{U}_c \rightarrow \text{argsort} \downarrow (\hat{U}_c)$ 即为预测DEG排名。

08 核心模块拆解

Module	Function	Why needed	Input and output	Supporting evidence	Known or expected effect of removal
Morphology-derived proxy contrast generator	Generates K-means clusters on frozen CONCH embeddings to define Ac/Bc groups without biological labels	Enables training without costly pathologist annotations; provides diverse morphology-defined contrasts for robust DEG learning	Input: Training image patches x_i → Output: Cluster assignments $l_i \in \{1, \dots, K\}$	Described in Sec. 5 & 7.1; used in all experiments (e.g., $k=10$ for training, $k=2-10$ for Figure 2) [Paper] [Paper: PDF p. 5-6]	Without it, model cannot train on unannotated data (e.g., HEST-1K); would require manual labels or fail in cluster-based setting
Differentiable U-stat approximator	Replaces hard indicator $1[\hat{y}_{i,g} > \hat{y}_{j,g}]$ with sigmoid $\sigma((\hat{y}_{i,g} - \hat{y}_{j,g})/\tau)$ to compute $\hat{U}_{c,g}$	Makes Mann-Whitney U statistic differentiable for end-to-end optimization; τ controls gradient smoothness	Input: Predicted expression $\hat{y}_{i,g}, \tau$ → Output: Scalar $\hat{U}_{c,g}$	Equation 1 explicitly defined; τ is a tunable hyperparameter [Paper] [Paper: PDF p. 5]	Removal reverts to non-differentiable sorting; training fails as gradients vanish (not empirically tested but implied by design rationale)
DEG-ranking surrogate objective (LDEG)	Computes $1 - \text{Corr}(\hat{U}_c, U_c)$ across genes to align predicted and true contrast statistics	Directly optimizes preservation of gene ordering by differential evidence, not raw expression values	Input: Vectors $\hat{U}_c, U_c \in \mathbb{R}^G \rightarrow$ Output: Scalar loss LDEG	Equation 2; ablation in Table 4 shows “Ours w/o DEG ranking” (MSE on U) performs worse than full LDEG [Paper] [Paper: PDF p. 5, 8]	Removal reduces nDCGDEG@200 by ~0.08 (Ovary) and SCCDEG by ~0.10 (Table 4), proving ranking alignment is essential
Auxiliary expression-consistency regularizer (Lconst)	Standard PCC loss averaged over genes, correlating $\hat{y}_{i,g}$ and $y_{i,g}$ across spots	Prevents degeneration where $\hat{U}_{c,g}$ aligns but individual gene spatial patterns collapse; stabilizes training	Input: Matrices $\hat{Y}, Y \in \mathbb{R}^{N \times G} \rightarrow$ Output: Scalar Lconst	Equation 3; described as “lightweight” and used with $\lambda=1.0$ [Paper] [Paper: PDF p. 5]	Removal (tested implicitly via ablation “Baseline” in Table 4) yields lower SCCDEG (0.584 vs. 0.686) and much lower nDCGDEG (0.451 vs. 0.531), showing spatial consistency aids ranking

09 关键公式与符号

- **No verifiable closed-form equation beyond those provided.** The paper defines three core equations, all present in input:

- **Equation 1:** $\hat{U}_{c,g} = \sum_{i \in A_c} \sum_{j \in B_c} \sigma((\hat{y}_{i,g} - \hat{y}_{j,g})/\tau)$ — differentiable U-stat approximation; σ = sigmoid, τ = temperature, $\hat{y}_{i,g}$ = predicted expression of gene g at spot i .
- **Equation 2:** $L_{DEG} = 1 - (1/|C|) \sum_{c \in C} \text{corr}_g(\hat{U}_c, U_c)$ — DEG-ranking surrogate loss; corr_g = Pearson correlation computed *across genes* ($g=1..G$), not spots.
- **Equation 3:** $L = L_{DEG} + \lambda L_{\text{const}}$ — overall loss; L_{const} = conventional PCC loss (defined in Sec. 5 but not numbered).
- **Key symbols:**
 - $U_{c,g}$: True one-sided Mann–Whitney U statistic for gene g under contrast c (Sec. 4).
 - $\pi_c = \text{argsort}_{\downarrow g}(U_{c,g})$: Reference DEG ranking list for contrast c (Sec. 4).
 - $\text{SCCDEG}_c = \text{Spearman}_g(U_c, \hat{U}_c)$: Spearman rank correlation between true/predicted U-stat vectors (Eq. 4).
 - $\text{nDCGDEG}_{c@k}$: Normalized Discounted Cumulative Gain for top- k DEGs (Eq. 5).
 - A_c, B_c : Positive/negative spot sets in one-vs-rest contrast (Sec. 4 & 5).

10 实验设计与证据链

Experiment	Claim tested	Comparison and conditions	Result	Supported conclusion	Unsupported stronger conclusion	Source
Cluster-based DEG (Table 1, Figure 2)	IDER objective improves DEG ranking under morphology-derived contrasts	All methods trained on HEST-1K (Ovary/Lymph Node/Bowel); groups from K-means ($k=10$); evaluation on $\text{nDCGDEG}@50/100/200$ & SCCDEG	Ours achieves highest scores across all metrics and datasets (e.g., $\text{nDCGDEG}@200$: 0.531 vs. 0.453 for MSE&PCC in Ovary)	Morphology-derived proxy contrasts + IDER objective enable superior DEG ranking vs. reconstruction objectives	Not claimed: superiority generalizes to <i>all</i> ST datasets or clinical cohorts (only public benchmarks tested)	[Paper] [Paper: PDF p. 7], Table 1, Figure 2
Real annotation-based DEG (Table 3)	IDER-trained model generalizes to pathologist-annotated tissue regions without seeing labels during training	Trained on Her2st using proxy clusters; evaluated on real annotations (adipose, connective, etc.)	Ours outperforms all baselines on $\text{SCCDEG}/\text{nDCG}$ DEG for every tissue type (e.g., $\text{SCCDEG}=0.475$ for connective tissue vs. 0.317 for MSE&PCC)	Proxy contrasts capture morphology-biology associations sufficiently to transfer to real annotations	Not claimed: performance matches expert-level annotation quality; no human-in-the-loop validation	[Paper] [Paper: PDF p. 7–8], Table 3

Experiment	Claim tested	Comparison and conditions	Result	Supported conclusion	Unsupported stronger conclusion	Source
Pathway enrichment overlap (Table 2)	IDER preserves downstream functional interpretation	Top-200 DEGs from predicted/measured profiles fed to Reactome/KEGG; Jaccard index computed	Ours achieves highest Jaccard across all datasets (e.g., 0.571 for Lymph Node vs. 0.443 for NB)	Better DEG ranking → better pathway-level biological signal recovery	Not claimed: recovered pathways are experimentally validated; no wet-lab confirmation	[Paper] [Paper: PDF p. 8], Table 2
Ablation (Table 4)	Inter-gene ranking alignment (not just U-stat magnitude) is critical	“Ours w/o DEG ranking” adds MSE loss on Uc but no Corrg; “Baseline” uses only PCC	Removing ranking alignment drops nDCGDEG@200 by 0.08 (Ovary) and SCCDEG by 0.10	Optimizing <i>relative order</i> of U-stat values matters more than absolute U-stat fidelity	Not claimed: Corrg is optimal choice; other ranking losses (e.g., ListNet) were not compared	[Paper] [Paper: PDF p. 8], Table 4
Plug-in (Table 6)	IDER objective is architecture-agnostic	Applied to ST-Net and TRIPLEX on Her2st; same training protocol	Both architectures show large nDCGDEG@200 gains (e.g., ST-Net: 0.073 → 0.247)	IDER loss can be adopted as a drop-in replacement for existing ST models	Not claimed: gains scale linearly with model size; no test on foundation-scale models	[Paper] [Paper: PDF p. 9], Table 6

11 对结论的正确理解

- **核心结论成立**：IDER目标确实提升了DEG排序保真度（nDCGDEG@200）、全局排序一致性（SCCDEG）及下游通路解释力（Jaccard），且该提升独立于模型架构（plug-in results）和对比来源（proxy vs. real annotations）。
- **边界清晰**：提升是**相对的**——相比MSE/PCC等基线，而非绝对完美（e.g., nDCGDEG@200 max=1.0, Ours=0.595）；提升是**task-specific**——在per-gene PCC上逊于专用重建模型（Table 5），证明其牺牲部分重建精度换取下游任务对齐。
- **机制可信**：Ablation（Table 4）证实ranking alignment（Corrg）比U-stat magnitude fitting（MSE on U）更关键；generalization across k=2–10 clusters (Figure 2) confirms proxy contrasts encode robust morphology signals.
- **不可外推**：未证明该方法在single-cell resolution、multi-omics integration或perturbation prediction场景有效；未验证其在临床诊断级WSI或跨机构泛化能力。

12 作者明确承认的限制

Limitation	Specific manifestation	Future direction proposed by authors	Source
Dependence on morphology-derived proxy contrasts	Proxy groups may not fully capture biologically meaningful tissue states (e.g., subtle molecular subtypes)	Explore integration with weak supervision from limited expert annotations or multi-modal signals (e.g., proteomics) [Paper] [Paper: PDF p. 9]	[Paper] [Paper: PDF p. 9] (end of Sec. 7.3)
Evaluation limited to bulk-spot ST	Current setup assumes spot-level expression; does not address single-cell deconvolution or sub-spot heterogeneity	Extend IDER to single-cell resolution by defining contrasts at cellular level or leveraging cell-type priors	[Paper] [Paper: PDF p. 9] (end of Sec. 7.3)
No clinical outcome correlation	DEG ranking improvement not linked to patient survival, treatment response, or diagnostic accuracy	Future work should correlate IDER performance with clinical endpoints in retrospective cohorts	[Paper] [Paper: PDF p. 9] (end of Sec. 7.3)

13 批判性分析

[Analysis] Observation	Potential issue or alternative explanation	Why it matters	How to test it	Basis
Proxy contrasts rely on CONCH embeddings, which are trained on generic pathology tasks, not ST-specific morphology	CONCH may miss ST-relevant histological nuances (e.g., stromal-tumor interface patterns critical for DEG) → proxy groups could misalign with true biology	If proxy groups poorly reflect biological contrasts, IDER optimization may learn spurious correlations, harming generalization to real annotations	Compare IDER performance when proxy groups are generated by ST-specialized foundation models (e.g., HistoST) vs. CONCH on Her2st; analyze cluster purity w.r.t. real annotations	[Paper] [Paper: PDF p. 5–6] states use of CONCH but no ablation on embedding source; Her2st annotations exist but weren't used for proxy generation
nDCGDEG@k focuses only on top-k recovery, ignoring ranking quality in mid/lower tiers	A model could “game” nDCG by boosting top-k genes while scrambling the rest, yielding high nDCG but poor overall biological utility	Downstream analyses (e.g., pathway enrichment) depend on broader gene sets; over-optimizing top-k may neglect functionally coherent modules outside top-200	Report nDCG@500 or Kendall's tau for full ranking; correlate full $\hat{U}$ vector with pathway membership (e.g., via GSEA) instead of binary top-k relevance	[Paper] [Paper: PDF p. 6] defines nDCGDEG@k but only reports @50/100/200; Table 2 uses top-200 for pathways, suggesting sensitivity to cutoff

[Analysis] Observation	Potential issue or alternative explanation	Why it matters	How to test it	Basis
Auxiliary Lconst uses simple PCC, which is known to be sensitive to outliers and zero-inflation in ST data	Outlier spots (e.g., necrotic regions) could dominate Lconst, pulling optimization away from DEG-critical morphology	Compromised spatial consistency might degrade DEG ranking in morphologically ambiguous regions (e.g., tumor margins)	Replace Lconst with robust correlation (e.g., Spearman) or zero-inflated loss (e.g., ZINB) and measure impact on nDCGDEG@200	[Paper] [Paper: PDF p. 5] calls Lconst “lightweight” but gives no justification for PCC choice; ST data is known to be sparse/outlier-prone [24]

14 学到的知识

- **任务定义升维是关键创新点：** 将ST预测从 “reconstruction” 重新定义为 “downstream decision fidelity” (IDER)，直击领域痛点，比单纯改进模型架构更具范式意义。
- **可微排序代理的设计智慧：** 用温度控制的sigmoid近似U-stat + 基因轴Pearson相关，巧妙绕过argsort不可微困境，比listwise ranking losses（如ListMLE）更契合DEG分析的统计本质。
- **代理监督的价值被低估：** K-means on CONCH features generates surprisingly effective proxy contrasts —no ground-truth labels needed, yet generalizes to real annotations (Table 3) and varying granularities (Figure 2).
- **下游指标与上游目标必须对齐：** Table 5 proves that high PCC ≠ good DEG ranking; thus，评估必须匹配最终用途，而非沿用历史惯性指标。
- **模块化即插即用潜力：** Table 6显示IDET loss boosts ST-Net/TRIPLEX without architecture changes—为快速迭代ST模型提供新范式。

15 与既有知识的连接

- **候选连接/方法论连接：**
- 与**single-cell foundation models**： IDER’ s use of morphology-derived proxy contrasts parallels scFoundation’ s use of cell-type priors for contrastive learning; both avoid hard labels by leveraging unsupervised structure.
- 与**spatial transcriptomics**： IDER directly addresses the “spatial DEG detection” gap in ST analysis tools (e.g., SPARK, SpatialDE) by making prediction DEG-aware.
- 与**graph neural networks**： GNN-based ST models (e.g., [6,38]) could integrate IDER loss to enhance neighborhood-aware DEG ranking, as current GNNs focus on spatial smoothing, not cross-gene prioritization.
- 与**multi-omics**： IDER’ s contrast-centric view aligns with multi-omics integration frameworks (e.g., MOFA+) that seek joint latent spaces preserving contrast-specific signals across modalities.
- 与**biomedical AI**： IDER exemplifies “clinical-task-aligned AI” —prioritizing decision outputs (DEG lists) over intermediate representations (expression matrices), resonating with diagnostic AI principles.
- **弱连接/方法论连接：**

- **perturbation prediction:** IDER's contrast statistics resemble perturbation response signatures; adapting IDER to predict gene ranking shifts under drug/CRISPR perturbations is plausible but untested.
- **cell state representation:** The $\hat{U}_c$ vector per contrast acts as a morphology-conditioned cell-state signature; could be used as input to cell-state classifiers, but paper doesn't explore this.
- **cross-modal alignment:** IDER aligns histology (image) and transcriptomics (U-stat) at the *decision level* (ranking), offering an alternative to feature-level alignment (e.g., CLIP-style); however, no explicit cross-modal loss is defined.

16 研究想法

[Hypothesis] 以下研究想法为基于论文证据和局限的可检验假设，不代表论文作者已验证。

- **name:** Morphology-Guided Perturbation DEG Ranking (MorphoPert)

originating limitation/observation: Proxy contrasts in IDER are static (K-means on baseline images), but perturbation responses (e.g., drug treatment) induce dynamic morphology changes not captured by baseline clustering [Analysis in Sec. 13]. **core hypothesis:** Learning perturbation-specific proxy contrasts—by clustering histology features *after* in silico perturbation simulation or using time-series WSI—will improve prediction of perturbation-induced DEG rankings. **delta from paper:** Replaces static K-means with perturbation-conditioned contrast generation (e.g., GAN-based morphological perturbation + contrastive clustering). **initial method:** Train a conditional GAN to simulate drug-treated histology from baseline WSI; extract features from perturbed images; cluster to get Ac/Bc (treated/untreated); apply IDER loss on simulated contrasts. **validation:** Test on CMap or LINCS perturbation datasets with matched histology + RNA-seq; compare nDCGDEG@200 for predicted vs. measured perturbation DEGs. **failure modes:** GAN artifacts dominate clustering; perturbation morphology too subtle for CONCH to resolve; lack of paired perturbed histology-RNA-seq data. **innovation status:** unverified

- **name:** Graph-Aware IDER (GraphIDER)

originating limitation/observation: IDER treats spots as independent in U-stat computation ($\Sigma_i \in A_c \Sigma_j \in B_c$), ignoring spatial adjacency and tissue graph structure known to modulate DEG signals [Paper] [Paper: PDF p. 3 cites GCN works]. **core hypothesis:** Constraining U-stat comparisons to spatially adjacent spot pairs (i,j) within and across Ac/Bc will yield more biologically grounded DEG rankings, especially for locally interacting tissues (e.g., tumor-stroma interface). **delta from paper:** Modifies Equation 1 to sum only over spatially neighboring (i,j) pairs, weighted by graph adjacency matrix A_{ij} . **initial method:** Build spatial graph from Visium spot coordinates (k-NN); replace $\Sigma_i \in A_c \Sigma_j \in B_c$ with $\Sigma_i \in A_c \Sigma_j \in B_c A_{ij} \cdot \sigma((\hat{y}_i, g - \hat{y}_j, g)/\tau)$; retain $\text{Corrg}(\hat{U}_c, U_c)$ as LDEG. **validation:** Apply to Her2st with known tumor/stroma boundaries; compare nDCGDEG@200 for stroma-enriched genes (e.g., COL1A1) vs. baseline IDER. **failure modes:** Over-smoothing in dense graphs; loss of long-range contrast signals; increased computational cost. **innovation status:** unverified

- **name:** Cross-Modal Alignment via DEG Ranking (CM-ALIGN)

originating limitation/observation: IDER aligns histology→transcriptomics, but multi-omics studies need alignment across modalities (e.g., histology + proteomics + methylation); current IDER is unimodal [Paper] [Paper: PDF p. 3 mentions multi-omics briefly]. **core hypothesis:** Jointly optimizing DEG rankings

across two modalities (e.g., histology-predicted and proteomics-predicted expression) using shared contrast statistics will enforce cross-modal biological consistency. **delta from paper:** Extends IDER to dual-modality: compute $\hat{U}_{c,hist}$ and $\hat{U}_{c,prot}$ for same contrast c ; add loss term $Corrg(\hat{U}_{c,hist}, \hat{U}_{c,prot})$ to align rankings across modalities. **initial method:** Use paired histology-proteomics dataset (e.g., CPTAC); train separate predictors f_{hist} , f_{prot} ; for each contrast c , compute $\hat{U}_{c,hist}$ and $\hat{U}_{c,prot}$; minimize $1 - Corrg(\hat{U}_{c,hist}, \hat{U}_{c,prot})$ alongside individual IDER losses. **validation:** On CPTAC breast cancer cohort; assess if CM-ALIGN improves concordance of top DEGs between predicted proteomics and measured transcriptomics. **failure modes:** Modality-specific noise dominates alignment; lack of large-scale paired multi-omics ST datasets; domain gap between histology and proteomics feature spaces. **innovation status:** unverified