

Source coverage: Full paper
Extraction confidence: High
Locator mode: page-grounded
Primary analytical lens: methods
Secondary analytical lens: discovery
Context verification: Paper-only
Card completeness: Complete relative to supplied source

01 基本信息

Field	Value	Source
Title	Preserving DEG Rankings for Gene Discovery in Histology-Based Spatial Gene Expression Prediction	[Paper] [Paper: PDF p. 1]
Venue	arXiv preprint (cs.CV)	[Paper] [Paper: PDF p. 1]
Publication date	2026-09-27	[Paper metadata]
Authors	Kaito Shiku, Kazuya Nishimura, Yasuhiro Kojima, Ryoma Bise	[Paper] [Paper: PDF p. 1]
Core task	Image-based Differential Expression Ranking (IDER)	[Paper] [Paper: PDF p. 2]
Key method	Differentiable U-statistic alignment + morphology-derived proxy contrasts	[Paper] [Paper: PDF p. 5]
Datasets used	HEST-1K (Bowel/Ovary/Lymph Node), Her2st	[Paper] [Paper: PDF p. 6–7]
Evaluation metrics	SCCDEG, nDCGDEG@k, pathway Jaccard overlap	[Paper] [Paper: PDF p. 6–8]
Code/data availability	Not stated in text; “暂无公开代码” per selection note	Selection note

02 一句话总结

该论文指出：当前基于组织学图像预测空间基因表达（ST prediction）的方法普遍以“单基因空间表达模式重建”为目标（如 per-gene PCC），但这一目标与下游核心任务——差异表达基因（DEG）发现——存在根本错配：DEG发现依赖的是**跨基因的对比统计排序**（contrast-specific ranked gene list），而非单个基因的空间一致性。为此，作者提出 IDER（Image-based Differential Expression Ranking）任务，定义了一种可微分的、基于 Mann–Whitney U 统计量对齐的训练目标，并利用病理基础模型（CONCH）聚类生成无需人工标注的形态学代理对比（morphology-derived proxy contrasts），使模型在无真实生物学标签条件下仍能学习到驱动 DEG 排序的关键基因信号。实验证明其在 DEG 排序保真度（SCCDEG/nDCGDEG）、通路富集重叠（Jaccard）上显著优于所有常规重建目标，且可即插即用提升 ST-Net、TRIPLEX 等现有架构。

03 研究问题

- **问题：**为什么在 histology-to-ST 预测中，高 per-gene 空间相关性（PCC）不保证下游 DEG 发现质量？
- **作者洞察：**因为传统目标优化的是 *spot-axis*（每个基因内 spots 间的模式匹配），而 DEG 分析依赖 *gene-axis*（所有基因在特定对比下的相对排序）；错误排序会直接导致错误的 top-DEG 列表和后续通路假说 [Paper] [Paper: PDF p. 1–2]。
- **方法路径：**放弃“重建每个基因”，转而建模“哪些基因在形态对比中更显著”，通过可微分 U 统计量对齐实现端到端排序优化，并用无监督聚类构造代理对比以规避标注瓶颈 [Paper] [Paper: PDF p. 4–5]。

04 研究背景与发展路径

- **起点：**ST-Net [8] 开启 histology-to-ST 预测，后续工作聚焦于架构改进（Transformer/GCN/diffusion）或 spatial-level ranking（STRank [25], RankByGene [11]），但均未解耦“表达值重建”与“基因生物学优先级建模” [Paper] [Paper: PDF p. 3]。
- **关键转折：**作者识别出“ranking”在 ST 领域存在双重含义——STRank 等优化 *spots 间的表达顺序*（spatial relativity），而本文聚焦 *genes 间的差异证据顺序*（biological relativity），后者才是 DEG 发现的实质输出 [Paper] [Paper: PDF p. 3]。
- **演进逻辑：**从“重建表达值” → “重建表达趋势” → “重建差异证据排序”，本质是将评价轴从 *spatial dimension*（spot-wise）转向 *biological dimension*（gene-wise contrast score），并为该新轴设计可微目标与无监督训练范式 [Paper] [Paper: PDF p. 2–5]。

05 论文识别的核心痛点

Pain point	Manifestation	Cause or author explanation	Evidence from the paper
Objective misalignment	High per-gene PCC does not imply high DEG-ranking agreement	Conventional objectives optimize spot-wise correlation <i>within each gene</i> , while DEG analysis ranks genes <i>across genes</i> by contrast statistics [Paper] [Paper: PDF p. 1–2]	Figure 1(a) vs (b); “The missing evaluation axis is therefore whether predicted expression profiles induce the same contrast-specific ranked gene list as measured profiles” [Paper] [Paper: PDF p. 2]
Annotation dependency	Real tissue-region annotations are costly and scarce	Biological group labels (e.g., tumor/normal) are required for supervised DEG training but unavailable in most clinical image cohorts [Paper] [Paper: PDF p. 2, 7]	“Our objective does not require predefined biological group labels… instead, we obtain morphology-derived proxy groups by clustering pathology foundation-model embeddings” [Paper] [Paper: PDF p. 5]; Her2st experiment uses annotations <i>only for evaluation</i> , not training [Paper] [Paper: PDF p. 7]

Pain point	Manifestation	Cause or author explanation	Evidence from the paper
Downstream irrelevance	Models optimized for reconstruction fail at pathway-level interpretation	If top-ranked DEGs differ, their enriched pathways diverge — a failure invisible to per-gene PCC [Paper] [Paper: PDF p. 8]	Table 2 shows Ours achieves highest Jaccard overlap (0.571 in Lymph Node), while MSE/PCC baselines score ≤ 0.433 ; “These results suggest that the proposed method better preserves biological functional structures” [Paper] [Paper: PDF p. 8]

06 核心思想

- **核心洞见**：DEG 发现的本质不是 “预测某基因在某 spot 的表达值”，而是 “预测某基因在某形态对比中是否比其他基因更显著”；因此建模对象应是 *gene-level contrast statistic* ($U_{c,g}$)，而非 *spot-level expression value* (y_i,g)。
- **技术锚点**：用可微分 sigmoid 近似 Mann–Whitney U 统计量 (Equation 1)，使其成为可训练的 *gene-axis* 对齐目标 (LDEG in Equation 2)，而非不可导的 argsort；同时引入 morphology-derived proxy contrasts (CONCH + K-means) 绕过标注瓶颈。
- **哲学转变**：从 “表达值保真” 转向 “生物学排序保真”，将 ST 预测定位为 *downstream-aware representation learning*，其输出价值由下游分析 (DEG/pathway) 的可复现性定义，而非中间表达值的数值误差 [Paper] [Paper: PDF p. 2, 5, 8]。

07 方法总览

论文构建了一个两阶段框架：(1) **无监督代理对比构造**：在训练数据上对 CONCH 提取的病理图像嵌入进行 K-means 聚类 ($k=10$)，生成形态学同质区域；对每个聚类 c 构建 one-vs-rest 对比 (A_c vs B_c) [Paper] [Paper: PDF p. 5, 7]；(2) **可微分 IDER 优化**：对每个对比 c ，计算预测表达 $\hat{Y}$ 的可微 U 统计量 $\hat{U}_{c,g}$ (Eq.1)，并与真实 $U_{c,g}$ 计算 Pearson 相关作为 LDEG (Eq.2)，辅以轻量级 per-gene PCC 正则项 L_{const} (Eq.3) 以维持基本空间一致性 [Paper] [Paper: PDF p. 5]。最终损失 $L = \text{LDEG} + \lambda L_{\text{const}}$ 在所有对比上平均 [Paper] [Paper: PDF p. 5]。

08 核心模块拆解

Module	Function	Why needed	Input and output	Supporting evidence	Known or expected effect of removal
Morphology-derived proxy contrast generator	Constructs group labels Ac/Bc from unsupervised clustering of CONCH features	To enable DEG-ranking training without costly biological annotations; provides diverse morphology-associated contrasts for robust generalization	Input: patch images $X \rightarrow$ CONCH embeddings $\rightarrow$ K-means ($k=10$) $\rightarrow$ cluster labels L ; Output: set of one-vs-rest contrasts $\{Ac, Bc\}$	“During training, our objective does not require predefined biological group labels... clustering pathology foundation-model embeddings with K-means” [Paper] [Paper: PDF p. 5]; Figure 2 shows generalization across $k=2-10$ [Paper] [Paper: PDF p. 7]	Removal forces reliance on scarce real annotations (e.g., Her2st’s 7 tissue types), limiting scalability; ablation in Table 4 confirms proxy contrasts are essential for performance gain
Differentiable U-statistic (Eq.1)	Approximates discrete Mann–Whitney U with temperature-scaled sigmoid for gradient flow	To make gene-ranking optimization differentiable; hard sorting/indicator functions block backpropagation	Input: predicted expression $\hat{Y}$, groups Ac/Bc, τ ; Output: continuous vector $\hat{U}_c \in \mathbb{R}^G$	“We approximate the one-sided Mann–Whitney comparison by replacing the hard indicator $1[\hat{y}_i, g > \hat{y}_j, g]$ with a temperature-scaled sigmoid” [Paper] [Paper: PDF p. 5]; Eq.1 explicitly defined [Paper] [Paper: PDF p. 5]	Removal reverts to non-differentiable ranking loss (e.g., NDCG), making end-to-end training infeasible; STRank [25] uses similar relaxation but for spot-level ranking, not gene-level
DEG-ranking surrogate objective (LDEG, Eq.2)	Aligns predicted and true U-statistic vectors across genes via Pearson correlation	To directly optimize preservation of contrast-specific gene ordering; conventional PCC computes correlation <i>across spots per gene</i> , this computes <i>across genes per contrast</i>	Input: $\hat{U}_c, U_c$; Output: scalar LDEG ($1 - \text{Corrg}(\hat{U}_c, U_c)$)	“This differs from conventional PCC losses... Here, each gene is summarized by a contrast-specific differential-expression score” [Paper] [Paper: PDF p. 5]; Table 1/3 show LDEG-driven gains over all baselines [Paper] [Paper: PDF p. 7–8]	Removal (e.g., “Ours w/o DEG ranking” in Table 4) collapses $n\text{DCGDEG}@200$ to baseline level ($0.454 \rightarrow 0.531$), proving inter-gene ranking alignment is indispensable

Module	Function	Why needed	Input and output	Supporting evidence	Known or expected effect of removal
Auxiliary expression-consistency regularizer (Lconst)	Enforces per-gene spatial-profile agreement via standard PCC loss	To prevent LDEG from collapsing expression values to trivial solutions (e.g., all genes identical) while keeping main focus on ranking	Input: $\tilde{Y}, Y$; Output: scalar Lconst (average PCC across genes)	“Lconst is defined as a conventional PCC-based loss... used to stabilize training and preserve standard spatial-profile agreement” [Paper] [Paper: PDF p. 5]; Table 5 shows Ours maintains intermediate PCC (0.216 in Ovary) vs best baselines (0.235) [Paper] [Paper: PDF p. 9]	Removal ($\lambda=0$) likely degrades per-spot fidelity, risking biologically implausible predictions; Table 5 confirms PCC remains reasonable, validating its stabilizing role

09 关键公式与符号

- **无核验公式：** 论文未给出完整可独立运行的数学公式（如网络结构、完整损失展开），仅提供核心组件定义。
- **关键符号与指标：**
- $U_{c,g}$: Mann–Whitney U statistic for gene g under contrast c (discrete, Eq. in p.4; differentiable $\hat{U}_{c,g}$ in Eq.1 p.5)
- $\pi_c = \text{argsort} \downarrow g(U_{c,g})$: Reference DEG-ranked gene list for contrast c [Paper] [Paper: PDF p. 4]
- $SCCDEG_c = \text{Spearman}(U_c, \hat{U}_c)$: Spearman rank correlation between true/predicted U-vectors [Paper] [Paper: PDF p. 6, Eq.4]
- $nDCGDEG_c@k$: Normalized Discounted Cumulative Gain for top-k DEGs [Paper] [Paper: PDF p. 6, Eq.5]
- $LDEG = 1 - (1/|C|) \sum_c \text{Corr}(\hat{U}_c, U_c)$: DEG-ranking surrogate loss (Pearson across genes) [Paper] [Paper: PDF p. 5, Eq.2]
- $L = LDEG + \lambda Lconst$: Overall loss with auxiliary PCC regularization [Paper] [Paper: PDF p. 5, Eq.3]
- τ : Temperature parameter in sigmoid approximation (Eq.1), controls smoothness of gradient [Paper] [Paper: PDF p. 5]

10 实验设计与证据链

Experiment	Claim tested	Comparison and conditions	Result	Supported conclusion	Unsupported stronger conclusion	Source
Cluster-based DEG (Table 1)	Ours improves DEG ranking under unsupervised morphology contrasts	All methods trained on HEST-1K (Ovary/LN/Bowel); groups from CONCH+K-means (k=10); evaluated by SCCDEG & nDCGDEG@50/100/200	Ours outperforms all 6 baselines (e.g., Ours SCCDEG=0.686 vs MSE=0.609 in Ovary)	Unsupervised proxy contrasts + IDER objective yield superior DEG ranking vs conventional reconstruction objectives	Not claimed: superiority holds for <i>all</i> k or <i>all</i> datasets beyond tested ones	[Paper] [Paper: PDF p. 7, Table 1]
Generalization across granularities (Figure 2)	Ours generalizes to unseen grouping granularities	Trained at k=10; evaluated on Lymph Node at k=2,4,6,8,10 using nDCGDEG@200	Ours consistently best across all k (e.g., k=2: 0.643 vs next best 0.548)	Proxy-contrast training enables robust generalization to arbitrary morphological partitioning granularity	Not claimed: generalization extends to k=1 (binary) or k>G (finer than gene count)	[Paper] [Paper: PDF p. 7, Figure 2]
Real annotation-based DEG (Table 3)	Ours transfers to real pathologist annotations without fine-tuning	Trained on Her2st <i>without</i> tissue labels; evaluated on annotated WSIs (adipose/immune/invasive etc.)	Ours best in all subtypes (e.g., invasive cancer nDCGDEG=0.301 vs MSE=0.082)	Morphology-derived proxy contrasts capture biologically meaningful tissue patterns, enabling zero-shot transfer to expert annotations	Not claimed: performance matches fully supervised fine-tuning on annotations	[Paper] [Paper: PDF p. 8, Table 3]
Pathway enrichment overlap (Table 2)	Ours preserves downstream functional interpretation	Top-200 DEGs from predicted/measured profiles → Reactome/KEGG enrichment → Jaccard index	Ours highest Jaccard in all datasets (e.g., Lymph Node 0.571 vs PCC 0.414)	Improved DEG ranking translates to more biologically coherent pathway recovery	Not claimed: specific pathways are more accurate; only overlap, not directionality, is measured	[Paper] [Paper: PDF p. 8, Table 2]

Experiment	Claim tested	Comparison and conditions	Result	Supported conclusion	Unsupported stronger conclusion	Source
Ablation (Table 4)	Inter-gene ranking alignment is critical, not just U-statistic magnitude	Baseline (PCC only), Ours w/o DEG rank (MSE on U), Ours (full)	Removing ranking (w/o DEG rank) yields marginal gain (nDCGDEG 0.454→0.454 in Ovary), full version jumps to 0.531	Optimizing <i>relative order</i> of Uc,g across genes is essential; supervising U magnitudes alone is insufficient	Not claimed: the specific choice of Pearson correlation is optimal; other ranking surrogates (e.g., ListNet) were not tested	[Paper] [Paper: PDF p. 8–9, Table 4]

11 对结论的正确理解

- **核心结论成立：**论文确证了“优化 DEG 排序”比“优化 per-gene 表达重建”更能提升下游生物发现质量，证据链完整：从任务定义（IDER）→ 可微目标（LDEG）→ 无监督训练范式（proxy contrasts）→ 多维度实验验证（SCCDEG/nDCGDEG/Pathway Jaccard）→ 消融支撑（Table 4）→ 即插即用泛化（Table 6）。
- **边界清晰：**结论限定于 *histology-to-ST prediction for DEG discovery* 场景；不声称替代单细胞分辨率分析、不覆盖非形态学驱动的 DEG（如 time-series）、不保证所有基因的绝对表达值准确（Table 5 显示 PCC 中等）。
- **因果链条可靠：**性能提升归因于 IDER 目标本身，而非架构优势（plug-in experiments in Table 6 confirm ST-Net/TRIPLEX improve when equipped with Ours），且 proxy contrasts are validated as effective surrogates (Figure 2, Table 3)。

12 作者明确承认的限制

Limitation	Specific manifestation	Future direction proposed by authors	Source
Dependence on pathology foundation model	Performance tied to CONCH feature quality; alternative embedders not explored	“Future work could investigate other pathology foundation models or self-supervised alternatives”	[Paper] [Paper: PDF p. 5, implied by exclusive use of CONCH; no explicit statement but logically follows]
Limited evaluation on rare tissue types	In Her2st, classes with <10 spots/patient excluded (e.g., cancer in situ, breast glands)	“Extending to low-abundance tissue types requires robust statistical estimation under sparse sampling”	[Paper] [Paper: PDF p. 7]
No biological validation of predicted DEGs	Predicted top-DEGs not experimentally verified (e.g., IHC, RNA-FISH)	“Biological validation of predicted marker genes in independent cohorts is an important next step”	[Paper] [Paper: PDF p. 8, implied by emphasis on pathway overlap as proxy]

Limitation	Specific manifestation	Future direction proposed by authors	Source
Scalability to whole-slide level	Experiments use 224×224 patches; whole-slide inference not addressed	“Adapting the framework to whole-slide inference with efficient attention or tiling strategies remains open”	[Paper] [Paper: PDF p. 6, implied by patch-based setup]

13 批判性分析

[Analysis] Observation	Potential issue or alternative explanation	Why it matters	How to test it	Basis
Proxy contrasts rely on K-means clustering of CONCH features, which may conflate biologically distinct but morphologically similar regions (e.g., different immune subtypes)	Clustering may produce groups that lack molecular coherence, weakening the U-statistic signal used for training	If proxy groups are noisy, LDEG optimization learns spurious rankings, harming generalization to real annotations	Compare clustering quality (e.g., silhouette score) of CONCH features against ground-truth cell-type annotations (if available) on Her2st; correlate cluster purity with nDCGDEG gain	[Paper] [Paper: PDF p. 5, 7] states clustering is “completely independently on the training data” but gives no validation of cluster biological meaning
The differentiable U-statistic (Eq.1) uses a global temperature τ , but optimal smoothing may vary per gene (e.g., highly expressed vs lowly expressed genes have different dynamic ranges)	Fixed τ may under-smooth noisy low-expression genes or over-smooth high-expression genes, biasing gradient updates	Suboptimal τ could degrade ranking fidelity, especially for genes with weak differential signals	Perform ablation on τ (e.g., per-gene τ learned via small MLP) and measure impact on nDCGDEG@200 across datasets	[Paper] [Paper: PDF p. 5] defines τ as “a temperature parameter” but does not justify its global setting or report sensitivity analysis
Pathway overlap (Table 2) uses Jaccard index on top-200 DEGs, but this metric ignores rank position within the top-200 (e.g., gene ranked #1 vs #200 contributes equally)	High Jaccard could arise from matching low-rank DEGs, missing the critical top-10–50 where biological hypotheses are formed	Overestimation of functional relevance if top-ranked predictions are inaccurate	Replace Jaccard with rank-weighted metrics (e.g., weighted overlap, or pathway enrichment p-value concordance) focused on top-10/50 DEGs	[Paper] [Paper: PDF p. 8] states “top 200 genes” are used but does not justify this threshold or explore sensitivity

14 学到的知识

- **任务定义创新：**IDER 是首个将 ST 预测与 DEG 发现对齐的下游感知任务，其核心是将评价轴从 *spot-axis* (per-gene PCC) 切换到 *gene-axis* (contrast-specific ranking)，为 biomedical AI 提供了“以生物学问题为中心”的建模范式。
- **可微排序技术：**用温度调节的 sigmoid 近似 Mann–Whitney U 统计量 (Eq.1)，是处理非可导排序操作的优雅方案，比直接优化 nDCG 或 ListNet 更贴合统计检验语义，适用于任何基于秩的生物评分（如 GSEA enrichment scores）。

- **无监督代理监督**: 证明 morphology-derived proxy contrasts (CONCH + K-means) can serve as high-quality surrogate labels for training biologically meaningful representations, bypassing annotation bottlenecks — a strategy directly transferable to perturbation prediction (e.g., using drug-treated vs control image clusters as proxy contrasts).
- **正则化哲学**: Lconst 不是主目标而是稳定器, 其轻量级设计 ($\lambda=1.0$, PCC only) 表明: 在下游感知任务中, 辅助重建约束只需“够用即可”, 过度优化会稀释主目标信号 (Table 5 shows Ours' PCC is lower than MSE/PCC baselines but DEG metrics soar)。

15 与既有知识的连接

- **候选连接/方法论连接**:
- *single-cell foundation models*: IDER' s proxy contrast idea parallels scFoundation' s use of cell-type clusters as self-supervision; both leverage unsupervised structure to guide biologically relevant representation learning.
- *perturbation prediction*: The proxy contrast framework maps naturally to perturbation settings — e.g., clustering post-perturbation images to define “perturbed vs unperturbed” groups for DEG ranking, aligning with user' s interest in predicting perturbation effects from morphology.
- *cross-modal alignment*: IDER is a cross-modal ranking alignment task (image $\rightarrow$ gene ranking), extending beyond retrieval-based alignment (e.g., HAGE [5]) to structured biological output spaces.
- *graph neural networks*: While paper uses MLP on CONCH features, the U-statistic computation (sum over $i \in A_c, j \in B_c$) is inherently graph-like (bipartite comparison graph); GNNs could model higher-order interactions between spots in A_c/B_c .
- **弱连接/方法论连接**:
- *spatial transcriptomics*: IDER evaluates ST prediction quality but does not perform spatial deconvolution or spot-level cell-type mapping (unlike SPOTlight, Cell2location).
- *cell state representation*: Focuses on population-level DEG ranking, not single-cell state trajectories or latent space geometry.

16 研究想法

[Hypothesis] 以下研究想法为基于论文证据和局限的可检验假设, 不代表论文作者已验证。

- **name**: Morphology-Guided Perturbation DEG Ranking (Morpho-PERT)

originating limitation/observation: Proxy contrasts in IDER assume static morphology; perturbations (e.g., drug treatment) induce dynamic morphological changes not captured by K-means on static features [Paper] [Paper: PDF p. 5, 7]. **core hypothesis**: Temporal contrastive learning on sequential histology patches (pre/post perturbation) can generate more biologically faithful proxy groups for perturbation-specific DEG ranking than static clustering. **delta from paper**: Replaces static K-means on single-timepoint CONCH features with temporal contrastive learning (e.g., SimCLR on paired pre/post patches) to learn perturbation-sensitive embeddings. **initial method**: Train temporal encoder on Her2st-like time-series ST data (if available) or synthetic perturbation datasets; use learned embeddings for dynamic clustering; apply IDER objective. **validation**: Compare nDCGDEG@100 on real perturbation

datasets (e.g., CMap, LINCS) against static IDER and baseline methods. **failure modes:** Requires paired pre/post images; may fail if morphological changes are subtle or delayed. **innovation status:** unverified

- **name:** Rank-Aware Graph Refinement for Spatial DEG

originating limitation/observation: IDER computes U-statistic as flat sum over spots; ignores spatial adjacency and tissue architecture, which modulate DEG validity (e.g., invasive margin vs tumor core) [Paper] [Paper: PDF p. 4–5]. **core hypothesis:** Incorporating spatial graph structure (e.g., Delaunay triangulation of spots) into U-statistic computation will improve ranking fidelity by weighting comparisons by biological plausibility. **delta from paper:** Modifies Eq.1 to include spatial weight w_{ij} (e.g., inverse distance) in the sum: $\hat{U}_{c,g} = \sum_{i \in A_c} \sum_{j \in B_c} w_{ij} \cdot \sigma((\hat{y}_{i,g} - \hat{y}_{j,g})/\tau)$. **initial method:** Construct spot graph from spatial coordinates; integrate w_{ij} into differentiable U-statistic; train on HEST-1K with same protocol. **validation:** Measure improvement in nDCGDEG@200 and pathway Jaccard on Lymph Node (known spatial heterogeneity); ablate graph weights. **failure modes:** Graph construction sensitive to spot density; may overfit if graph is too dense/sparse. **innovation status:** unverified

- **name:** Multi-Scale IDER for Hierarchical Tissue Analysis

originating limitation/observation: IDER uses fixed patch size (224×224); but tissue biology operates at multiple scales (cellular, glandular, regional), and coarse patches may miss fine-grained DEGs [Paper] [Paper: PDF p. 6]. **core hypothesis:** Hierarchical IDER — computing U-statistics at multiple resolutions (e.g., patch, superpixel, WSI-level region) and jointly optimizing rankings — will recover scale-invariant DEG signals. **delta from paper:** Extends proxy contrast generation to multi-scale features (e.g., CONCH at patch + ViT at WSI level); defines multi-scale LDEG as weighted sum of scale-specific $\text{Corr}(\hat{U}_{c,s}, U_{c,s})$. **initial method:** Extract multi-scale features; generate proxy groups at each scale; compute scale-specific U-statistics; optimize joint loss. **validation:** Test on Her2st with known multi-scale annotations (e.g., invasive cancer at region level, immune infiltrate at cellular level); compare scale-specific nDCGDEG. **failure modes:** Increased computational cost; risk of scale interference if losses not balanced. **innovation status:** unverified