

Executed Command

Reference: pipelines/PIPELINE_RUN_GUIDE.md

Script: pipelines/compare_hardcoded_vs_qiskit_pipeline.py

/opt/anaconda3/bin/python3 Fermi-Hamil-JW-VQE-TROTTER-

PIPELINE/pipelines/compare_hardcoded_vs_qiskit_pipeline.py --l-values 2 --run-pipelines --enable-drive
--drive-pattern dimer_bias --drive-omega 2.0 --drive-tbar 2.0 --t-final 2.0 --num-times 21 --trotter-steps
32 --drive-amplitudes 0.0,0.2 --with-drive-amplitude-comparison-pdf --artifacts-dir
artifacts/amp_compare_smoke --skip-qpe

Hardcoded vs Qiskit Pipeline Comparison Summary

generated_utc: 2026-02-21T00:10:16.463575+00:00

all_pass: True

l_values: [2]

trajectory_comparison_basis: trotter trajectories start from
each pipeline's selected initial_state_source (default: vqe)

exact_trajectory_labels: Exact_Hardcode, Exact_Qiskit

exact_trajectory_method: python_matrix_eigendecomposition

thresholds:

```
{'doublon_trotter_max_abs_delta': 0.001,  
 'energy_trotter_max_abs_delta': 0.001,  
 'fidelity_max_abs_delta': 0.0001,  
 'ground_state_energy_abs_delta': 1e-08,  
 'n_dn_site0_trotter_max_abs_delta': 0.005,  
 'n_up_site0_trotter_max_abs_delta': 0.005}
```

hardcoded_qiskit_import_isolation:

```
{'offending_imports': [],  
 'pass': True,  
 'qiskit_imports': [{'line': 497, 'module': 'qiskit'},  
                    {'line': 498, 'module': 'qiskit.circuit.library'},  
                    {'line': 499, 'module': 'qiskit.primitives'},  
                    {'line': 500, 'module': 'qiskit.quantum_info'},  
                    {'line': 501, 'module': 'qiskit.synthesis'},  
                    {'line': 502, 'module': 'qiskit_algorithms'},  
                    {'line': 503, 'module': 'qiskit_algorithms.minimum_eigensolvers'}],  
 'qpe_adapter_range': {'end_line': 599, 'start_line': 463}}
```

Delta metric definitions:

$$\Delta F(t) = |F_{hc}(t) - F_{qk}(t)|$$
$$\Delta E_{trot}(t) = |E_{trot_hc}(t) - E_{trot_qk}(t)|$$
$$\Delta n_{up0}(t) = |n_{up0_hc}(t) - n_{up0_qk}(t)|$$


$\Delta n_{dn0}(t) = |n_{dn0_hc}(t) - n_{dn0_qk}(t)|$



$\Delta D(t) = |D_{hc}(t) - D_{qk}(t)|$

$F_{\text{pipeline}}(t)$ is the pipeline's stored trajectory fidelity value (as computed internally vs that pipeline's exact evolution).

Per-L pass flags:

L=2 pass=True metrics_json=/Users/jakestrobels/Documents/Holstein_implementation/Holstein_test/artifacts/amp_compare_smoke/hardcoded_vs_qiskit_pipeline_L2_metrics.json







QPE comparison skipped: no finite QPE energy estimates were found in per-L payloads.


Bundle L=2: Run Settings & Metrics Summary

```
L=2  t=1.0  u=4.0  dv=0.0  boundary=periodic  ordering=blocked  initial_state_source=vqe  t_final=2.0  num_times=21  suzuk.  
thresholds:  
  doublon_trotter_max_abs_delta: 1.00e-03  
  energy_trotter_max_abs_delta: 1.00e-03  
  fidelity_max_abs_delta: 1.00e-04  
  ground_state_energy_abs_delta: 1.00e-08  
  n_dn_site0_trotter_max_abs_delta: 5.00e-03  
  n_up_site0_trotter_max_abs_delta: 5.00e-03  
  
max |Δ|:  
  gs_energy: 0.00e+00  
  doublon_trotter: 1.32e-05  
  energy_trotter: 2.86e-05  
  fidelity: 1.05e-09  
  n_dn_site0_trotter: 2.79e-05  
  n_up_site0_trotter: 2.45e-05  
result: PASS
```


Bundle Page: L=2 Fidelity & Energy




L=2 Site-0 n_{up}





L=2 Site-0 n_{dn}



L=2 Doublon







When initial_state_source=vqe, Trotter $E(t=0) = \langle \psi_{\text{vqe}} | H | \psi_{\text{vqe}} \rangle = \text{VQE energy}$.
VQE energy \neq exact ground state energy unless VQE fully converged.



Bundle Delta Diagnostics L=2

$\Delta X(t) = |X_{hc}(t) - X_{qk}(t)|$, where $X_{pipeline}(t)$ is that pipeline's stored trajectory value.



Bundle metrics page L=2

Trotterization comparison uses each path's configured initial state.

Trajectory labels: Exact_Hardcode and Exact_Qiskit.

Exact trajectory method: python_matrix_eigendecomposition.

For VQE-init runs, both exact(t) and trotter(t) start from the VQE ansatz state.

Delta metric definitions:

$\Delta F(t) = |F_{hc}(t) - F_{qk}(t)|$

$\Delta E_{trot}(t) = |E_{trot_hc}(t) - E_{trot_qk}(t)|$

$\Delta n_{up0}(t) = |n_{up0_hc}(t) - n_{up0_qk}(t)|$

$\Delta n_{dn0}(t) = |n_{dn0_hc}(t) - n_{dn0_qk}(t)|$

$\Delta D(t) = |D_{hc}(t) - D_{qk}(t)|$

$F_{pipeline}(t)$ is the pipeline's stored trajectory fidelity value (as computed internally vs that pipeline's exact evolution).

ground_state_energy_abs_delta = 0.0

fidelity max/mean/final = 1.0517273718591014e-09 / 2.1139742337591575e-10 / 2.5473967379952e-10

energy_trotter max/mean/final = 2.8569770569086828e-05 / 1.1392085824032025e-05 / 3.854281204862531e-06

n_up_site0_trotter max/mean/final = 2.4453936811663635e-05 / 1.2263117297871604e-05 / 4.229340619166244e-06

n_dn_site0_trotter max/mean/final = 2.7934482115687942e-05 / 1.1312123054409349e-05 / 1.2385115255586499e-06

doublon_trotter max/mean/final = 1.3169303991367176e-05 / 5.808441969007691e-06 / 1.246908090368759e-05

checks:

```
{'doublon_trotter_max_abs_delta': True,
 'energy_trotter_max_abs_delta': True,
 'fidelity_max_abs_delta': True,
 'ground_state_energy_abs_delta': True,
 'n_dn_site0_trotter_max_abs_delta': True,
 'n_up_site0_trotter_max_abs_delta': True}
```

PASS = True