

Multiple Linear Regression

outline:

- Setup model & get LS est
↳ matrix formulation of regression
- Gauss-Markov
- Fitted values & estimate σ^2

④ Model Set Up

$$\text{SLR: } y_i = \beta_0 + \beta_1 x_i + \varepsilon_i \quad i=1, \dots, n$$

$$y_1 = \beta_0 + \beta_1 x_1 + \varepsilon_1$$

$$y_2 = \beta_0 + \beta_1 x_2 + \varepsilon_2$$

\vdots

$$y_n = \beta_0 + \beta_1 x_n + \varepsilon_n$$

Let's define

$$\underbrace{y}_{(n \times 1)} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}, \quad \underbrace{X}_{(n \times 2)} = \begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ \vdots & \vdots \\ 1 & x_n \end{bmatrix}, \quad \underbrace{\beta}_{2 \times 1} = \begin{bmatrix} \beta_0 \\ \beta_1 \end{bmatrix}, \quad \underbrace{\varepsilon}_{n \times 1} = \begin{bmatrix} \varepsilon_1 \\ \varepsilon_2 \\ \vdots \\ \varepsilon_n \end{bmatrix}$$

$$Y = X\beta + \varepsilon$$

$$Y = \begin{bmatrix} \beta_0 + \beta_1 x_1 \\ \beta_0 + \beta_1 x_2 \\ \vdots \\ \beta_0 + \beta_1 x_n \end{bmatrix} + \varepsilon$$

← this is equivalent to my system of n equations

Assumptions

① X is fixed

② $\varepsilon \sim N(0, \begin{bmatrix} \sigma^2 & 0 & 0 & \dots & 0 \\ 0 & \sigma^2 & & & \\ \vdots & & \ddots & & \\ 0 & \dots & 0 & \sigma^2 \end{bmatrix})$

$$I = \begin{bmatrix} 1 & & 0 \\ & \ddots & \\ 0 & & 1 \end{bmatrix}$$

$$= N(0, \sigma^2 I_n)$$

Aside:

if I have a random vector $Z = \begin{bmatrix} z_1 \\ z_2 \end{bmatrix}$

then $E(Z) = \begin{bmatrix} E(z_1) \\ E(z_2) \end{bmatrix} = \mu$ is the mean vector


* $\text{Var}(Z) = \begin{bmatrix} \text{Var}(z_1) & \text{cov}(z_1, z_2) \\ \text{cov}(z_2, z_1) & \text{Var}(z_2) \end{bmatrix} = \Sigma$ is the covariance matrix.

Random variable z : $\text{Var}(z) = E((x-\mu)^2)$

random vector z : $\text{Var}(z) = E(\underbrace{(x-\mu)}_{n \times 1} \underbrace{(x-\mu)^T}_{1 \times n})$

Multivariate Normal Dist

$$\underline{z} = \begin{bmatrix} z_1 \\ \vdots \\ z_n \end{bmatrix} \sim N(\underline{0}, \sigma^2 \underline{I})$$



$$\begin{aligned} f(z_1, \dots, z_n) &= \prod_{i=1}^n f(z_i) \\ &= \prod_{i=1}^n \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right) e^{-z_i^2 / 2\sigma^2} \\ &= (2\pi\sigma^2)^{-n/2} e^{-\sum_i z_i^2 / 2\sigma^2} \end{aligned}$$

In general a random vector $z \sim N(\mu, \Sigma)$

has the pdf:

$$f(z_1, \dots, z_n) = \frac{1}{(\sqrt{2\pi})^n |\Sigma|^{1/2}} e^{-\frac{(x-\mu)^T \Sigma^{-1} (x-\mu)}{2}}$$

② Parameter Estimation under Matrix Setup

In SLR the idea was to minimize

$$Q(\beta_0, \beta_1) = \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2$$

In the matrix setup, we minimize

$$\begin{aligned} Q(\beta) &= (Y - X\beta)^T (Y - X\beta) \\ &= \|Y - X\beta\|^2 \end{aligned}$$

If I want to minimize $Q(\beta)$ w.r.t. β which is a vector, we need matrix calculus.

$$\textcircled{1} \frac{\partial (\vec{\beta}^T A)}{\partial \vec{\beta}} = A$$

$$\textcircled{2} \frac{\partial (\vec{\beta}^T A \vec{\beta})}{\partial \vec{\beta}} = \underline{\underline{2A\vec{\beta}}} \quad \text{if } A \text{ is symmetric}$$

$$\textcircled{3} \frac{\partial Q}{\partial \vec{\beta}} = 0$$

$$y^T X \vec{\beta} = \langle y, X \vec{\beta} \rangle = \langle X \vec{\beta}, y \rangle$$

$$= (X \vec{\beta})^T y$$

$$= \vec{\beta}^T X^T y$$

Goal is to minimize

$$Q(\vec{\beta}) = \|y - X\vec{\beta}\|^2 = (y - X\vec{\beta})^T (y - X\vec{\beta})$$

$$= y^T y - \boxed{y^T X \vec{\beta}} - (X \vec{\beta})^T y + (X \vec{\beta})^T (X \vec{\beta})$$

$$= \underline{y^T y} - \underline{y^T X \vec{\beta}} - \underbrace{\vec{\beta}^T X^T y}_{= y^T X \vec{\beta}} + \underline{\vec{\beta}^T X^T X \vec{\beta}}$$

$$\hookrightarrow \frac{\partial Q}{\partial \vec{\beta}} = 0 - X^T y - X^T y + 2 X^T X \vec{\beta} \stackrel{!}{=} 0$$

$$= -2 X^T y + 2 X^T X \vec{\beta} = 0$$

$$\Rightarrow (X^T X) \vec{\beta} = X^T y \quad \uparrow$$

$$\Rightarrow \hat{\beta} = (X^T X)^{-1} X^T Y \quad \left. \vphantom{\hat{\beta}} \right\} \text{the Normal Eq.}$$

$$\begin{pmatrix} \hat{\beta}_0 \\ \hat{\beta}_1 \end{pmatrix} = \begin{pmatrix} \bar{Y} - \hat{\beta}_1 \bar{X} \\ \frac{SS_{XY}}{SS_X} \end{pmatrix} \quad \begin{matrix} \text{[...]} \\ \swarrow \text{SER} \end{matrix} \quad [\text{worked out in Cody Notes}]$$

Why did I do this?

Now I can add more predictors to the design matrix, X :

Multiple Linear Regression Model

$$Y_i = \beta_0 + \beta_1 X_{i1} + \dots + \beta_{p-1} X_{i(p-1)} + \varepsilon_i \quad i=1, \dots, n$$



$$Y = X\beta + \varepsilon$$

where

$$X = \begin{matrix} n \times p \\ \begin{bmatrix} 1 & X_{11} & X_{12} & \dots & X_{1(p-1)} \\ 1 & X_{21} & X_{22} & & \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & X_{n1} & X_{n2} & \dots & X_{n(p-1)} \end{bmatrix} \end{matrix} \quad \& \quad \beta = \begin{pmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_{p-1} \end{pmatrix}$$

So the LS estimate for the MLR model is

$$\text{Just } \hat{\beta} = (X^T X)^{-1} X^T Y \quad (\text{assuming that } X \text{ has full rank})$$

Fitted values

$$\hat{y} = X\hat{\beta} = X(X^T X)^{-1} X^T y$$

$H = \text{"hat matrix"}$

$$= Hy$$

