

## ✓ Congratulations! You passed!

Go to next item

Grade received 100% Latest Submission Grade 100% To pass 70% or higher

1.

1 / 1 point

Gradient descent is an algorithm for finding values of parameters  $w$  and  $b$  that minimize the cost function  $J$ .

When  $\frac{\partial J(w,b)}{\partial w}$  is a negative number (less than zero), what happens to  $w$  after one update step?

- ☐ It is not possible to tell if  $w$  will increase or decrease.
- ☒  $w$  increases.
- ☐  $w$  decreases
- ☐  $w$  stays the same

✓ Correct

The learning rate is always a positive number, so if you take  $w$  minus a negative number, you end up with a new value for  $w$  that is larger (more positive).

2.

1 / 1 point

For linear regression, what is the update step for parameter  $b$ ?

- ☐  $b = b - \alpha \frac{1}{m} \sum_{i=1}^m (f_{w,b}(x^{(i)}) - y^{(i)}) x^{(i)}$
- ☒  $b = b - \alpha \frac{1}{m} \sum_{i=1}^m (f_{w,b}(x^{(i)}) - y^{(i)})$

✓ Correct

The update step is  $b = b - \alpha \frac{\partial J(w,b)}{\partial b}$  where  $\frac{\partial J(w,b)}{\partial b}$  can be computed with this expression:  $\sum_{i=1}^m (f_{w,b}(x^{(i)}) - y^{(i)})$