

OVARIAN TUMORS

BY
MANYEPA.

INTRODUCTIONS.

- Ovary composed of three cell types.
 - Surface epithelium
 - Germ cells
 - Sex cord stroma
- Tumor can arise from any of the three or from metastasis.

SURFACE EPITHELIAL TUMORS

- Most common type of ovarian tumors
- Derived from coelomic epithelium that lines the ovary
- Two most common subtypes of surface epithelial tumors are;
 - Serous
 - Mucinous
- Both usually are cystic

Benign tumors (cystadenoma)

- Composed of a single cyst with simple, flat lining
- Most commonly arises in premenopausal women (30 – 40 years)

Malignant tumors (cystadenocarcinoma)

- Composed of complex cysts with thick, shaggy lining
- Most commonly arises in post menopausal women (60 – 70 years old)

Borderline tumors

- Features in between benign and malignant tumors
- Better prognosis than clearly malignant tumors
- But still carry metastatic potential

BRCA1 mutation carriers

- Increased risk for serous carcinoma of the ovary and fallopian tube

Other types of surface tumors

- Endometrioid
- Brenner tumor

Surface tumors presents late

- Vague abdominal symptoms (pain, fullness)
- Sign of compression (urinary frequency)
- Poor prognosis
- Epithelial carcinomas tend to spread locally especially to peritoneum.

CA-125

- Useful serum marker to monitor treatment response.
- Screening for recurrence

Germ cell tumors

- Second most common ovarian tumor (15% of cases)
- Usually occur in women of reproductive age
- Tumor subtypes mimic tissues normally produced by germ cells

Cystic teratoma

- Cystic tumor composed of fetal tissue derived from two or three embryologic layers
- Most common germ cell tumors in females
- Bilateral in 10% of cases

Benign

- Presence of immature tissue or somatic malignancy indicates malignant potential.

Struma Ovarii

- Cystic Teratoma composed primarily of thyroid tissue

Dysgerminoma

- Composed of large cells with clear cytoplasm and central nuclei
- Most common malignant germ cell tumor
- Testicular counterpart is termed seminoma
- Good prognosis, responds to radiotherapy
- Serum LDH maybe elevated.

Endodermal sinus Tumor

- Malignant tumor that mimic yolk sac
- Most common germ cell tumor in children
- Serum AFP is often elevated

Choriocarcinoma

- Composed of trophoblasts and syncytiotrophoblasts (Villi are absent)
- Small, hemorrhagic tumor with early hematogenous spread
- High B – hcG
- Poor response to Chemotherapy

Embryonal carcinoma

- Composed of large primitive cells
- Aggressive with early metastasis

Sex cord – stromal tumors

- Resemble sex cord – stroma tissue of ovary

Granulosa-theca cell tumor

- Neoplasm of granulosa and theca cells
- Often produces estrogen
- Presents with signs of estrogen excess (symptoms vary with age)

Sertoli-leydig cell tumor

- Sertoli cells from tubules
- Leydig cells contain characteristic Reinke crystals
- May produce androgen;
 - Hirsutism or virilization

Fibroma

- Benign tumor of fibroblasts
- Associated with pleural effusions and ascites

Metastasis

- Kruckenberg tumor
- Pseudomyxoma peritonei