Ex.No-1 NUMPY

AIM:

To calculate the values for the mathematical formulas using NumPy library

# INTEGRATED DEVELOPMENT ENVIRONMENT (IDE) REQUIRED:

JUPYTER NOTEBOOK

## REQUIRED LIBRARIES FOR PYTHON:

Numpy

## **PROCEDURE:**

## A) Euclidean distance

The mathematical formula for calculating the Euclidean distance between 2 points in 2D space:

$$d(p,q) = \sqrt[2]{(q_1-p_1)^2+(q_2-p_2)^2}$$

## B) Dot Product

$$u={5\brack 12},\quad v={8\brack 6}$$
 Dot product is  $u\cdot v=u_1\times v_1+u_2\times v_2$   $=5\times 8+12\times 6$   $=112$ 

## C) Solving a System of Linear Equations

A system of linear equations can be represented in matrix form as AX=B, where A is the matrix of coefficients, X is the column vector of variables, and B is the column vector of solutions. To solve for  $\mathbf{X}$ , we can use:  $\mathbf{X} = \mathbf{A}^{-1} \mathbf{B}$  assuming A is invertible.

#### **PROGRAM:**

#### A) Calculating the Euclidean Distance Between Two Points

importnumpy as np

defeuclidean\_distance(p, q):

```
returnnp.sqrt(np.sum((q - p) ** 2))
# Example usage
p = np.array([1, 2])
q = np.array([4, 6])
distance = euclidean_distance(p, q)
print("Output for Calculating the Euclidean Distance Between Two Points is: ",distance)
B) Calculating the Dot Product of Two Vectors
importnumpy as np
A = np.array([1, 3, -5])
B = np.array([4, -2, -1])
dot_product = np.dot(A, B)
print("Output for dot product of two vectors A and B is ",dot product)
C) Solving a System of Linear Equations
importnumpy as np
# Coefficients matrix A and result vector b
A = np.array([[3, 1], [1, 2]])
b = np.array([9, 8])
# Solve for x
x = np.linalg.solve(A, b)
print("Output solution of System of Linear Equations is ",x)
Output:
A)Output for Calculating the Euclidean Distance between Two Points is: 5.0. Exercise 2 –B)Output for
dot product of two vectors A and B is 3
C)Output solution of System of Linear Equations is [2. 3.]
Result:
```

The programs were run successfully