

Find Key:

You are provided with 3 numbers : input1, input2 and input3.

Each of these are 4 digits numbers within ≥ 1000 and ≤ 9999

i.e.,

$1000 \leq \text{input1} \leq 9999$

$1000 \leq \text{input2} \leq 9999$

$1000 \leq \text{input3} \leq 9999$

You are expected to find the key using below formula:

Key = Sum of Largest digits of each number + Sum of Second Largest digits of each number

For Example, input1=3521, input2=2452 input3=1352

Key = $(5+5+5) + (3+4+3) = 25$

Assuming three numbers are passed to given function and complete the function to find and return the Key

• FIND KEY(DIFFERENT TESTCASE WITH DIFFERENT MODEL-2)**Find Key:**

You are provided with 3 numbers input1,input2,input3.

Each of these are four digit numbers within the range ≥ 1000 and ≤ 9999

i.e

$1000 \leq \text{input1} \leq 9999$

$1000 \leq \text{input2} \leq 9999$

$1000 \leq \text{input3} \leq 9999$

you are expected to find the key using the below formula

Key=[smallest digit in the thousands place of all three numbers][LARGEST digit in the hundreds place of all the three numbers]

[smallest digit in the tens place of all three numbers][LARGEST digit in the units place of all three numbers]

for e.g if input1=3521,input2=2452,input3=1352,then Key=[1][5][2][2]=1522

Assuming that the 3 numbers are passed to the given function.Complete the function to find and return the key.

FIND KEY(DIFFERENT TESTCASE WITH DIFFERENT MODEL-3)

You are provided with 3 numbers input1,input2,input3.

Each of these are four digit numbers within the range ≥ 1000 and ≤ 9999

i.e

$1000 \leq \text{input1} \leq 9999$

$1000 \leq \text{input2} \leq 9999$

$1000 \leq \text{input3} \leq 9999$

you are expected to find the key using the below formula

Key=[LARGEST digit in the thousands place of all three numbers][smallest digit in the hundreds place of all the three numbers]
[LARGEST digit in the tens place of all three numbers][smallest digit in the units place of all three numbers]

for e.g if input1=3521,input2=2452,input3=1352,then Key=[3][3][5][1]=3351

Assuming that the 3 numbers are passed to the given function.Complete the function to find and return the key.

ANSWER:

• TWO DIGIT REDUCED SUBTRACTED FORM

Given a number, you are expected to find its two-digit “Reduced Subtracted Form(RSF)”

The “Reduced Subtracted Form(RSF)” of a number can be found by concatenating the difference between its adjacent digits

To find the two-digit “Reduced Subtracted Form(RSF)”,we need to continue this process till the resultant RSF is not a two digit number.

For eg if the input number is 6928, its RSF can be found by concatenating the difference between (6 and 9), (9 and 2) and (2 and 8) as shown below –

Difference between 6 and 9 is 3

Difference between 9 and 2 is 7

Difference between 2 and 8 is 6

So, the “Reduced Subtracted Form(RSF)” of 6928 = 376

The resultant RSF (376) is not a two-digit number, so we must continue finding its “Reduced Subtracted Form(RSF)”

Difference between 3 and 7 is 4

Difference between 7 and 6 is 1

So, the “Reduced Subtracted Form(RSF)” of 376 = 41

The resultant RSF (41) is two-digit number, so we have reached the “two-digit Reduced Subtracted Form”.

Therefore, the two-digit RSF of 6928 = 41

Let’s see another example

If input1 = 5271

Expected output = 21

Explanation:

RSF of 5271 = (5-2)(2-7)(7-1)=356

RSF of 356=(3-5)(5-6)=21

Note1: input1 will be always be ≥ 100

Note2:Note that while concatenating the differences, we are expected to use the absolute values(non-negative)

Note3: The input values for all test cases in this program have been designed such that their two-digits RSF will definitely result in a two-digit number

Find Password:

Detective Buckshee junior has been approached by the shantiniketan kids society for help in finding the password to the games complex. After hearing the scenario, detective Buckshee junior realizes that he will need a programmer's support. He contacts you requests your help. Please help the detective by writing a function to generate the password.

The scenario is as below-

Five numbers are available with the kids.

These numbers are either stable or unstable

A numbers is stable if each of its digit occur the same number of times, i.e the frequency of each digit in the number is the same.

For e.g:2277,4004,11,23,583835,1010 are examples of stable numbers.

Similarly,A number is unstable if the frequency of each digit in the number is NOT the same.

For eg:221,4314,101,233,58135,101 are examples of unstable numbers.

The password can be found as below-

i.e password=(Number of unstable numbers*10)+Number of stable numbers

Assuming that the fiVE numbers are passed to a function as

input1,input2,input3,input4,input5.complete the function to find and return the password.

For example:

If input1=12,input2=1313,input3=122,input4=678 and input5=898 , we see that there are

THREE stable numbers i.e 12,1313 and 678 and

TWO unstable numbers i.e 122 and 898

so,the password should be=(Number of Unstable numbers*10)+Number of stable numbers=(2*10)+3=23

Find Password:

Detective Buckshee junior has been approached by the shantiniketan kids society for help in finding the password to the games complex.After hearing the scenario,detective Buckshee junior realises that he will need a programmer's support.He contacts you requests your help. Please help the detective by writing a function to generate the password.

The scenario is as below-

Five numbers are available with the kids.

These numbers are either stable or unstable

A numbers is stable if each of its digit occur the same number of times, i.e the frequency of each

digit in the number is the same.

For e.g:2277,4004,11,23,583835,1010 are examples of stable numbers.

Similarly ,A number is unstable if the frequency of each digit in the number is NOT the same.For eg:221,4314,101,233,58135,101 are examples of unstable numbers.

The password can be found as below-

i.e password=(Number of stable numbers*10)+Number of unstable numbers

Assuming that the five numbers are passed to a function as

input1,input2,input3,input4,input5.complete the function to find and return the password.

For example:

If input1=12,input2=1313,input3=122,input4=678 and input5=898 , we see that there are THREE stable numbers i.e 12,1313 and 678 and

TWO unstable numbers i.e 122 and 898

so,the password should be=(Number of stable numbers*10)+Number of Unstable numbers=(3*10)+1=32

ANSWER:

20) Find Password:

Detective Buckshee junior has been approached by the shantiniketan kids society for help in finding the password to the games complex.After hearing the scenario,detective Buckshee junior realises that he will need a programmer's support.He contacts you requests your help. Please help the detective by writing a function to generate the password.

The scenario is as below-

Five numbers are available with the kids.

These numbers are either stable or unstable

A numbers is stable if each of its digit occur the same number of times, i.e the frequency of each digit in the number is the same.

For e.g:2277,4004,11,23,583835,1010 are examples of stable numbers.

Similarly ,A number is unstable if the frequency of each digit in the number is NOT the same.For eg:221,4314,101,233,58135,101 are examples of unstable numbers.

The password can be found as below-

i.e Password=Maximum of all stable numbers+Minimum of all Unstable numbers

[28-07-2025 14:33] Jogendra: Assuming that the five numbers are passed to a function as input1,input2,input3,input4,input5.complete the function to find and return the password.

For example:

If input1=12,input2=1313,input3=122,input4=678 and input5=898 , we see that there are THREE stable numbers i.e 12,1313 and 678 and TWO unstable numbers i.e 122 and 898 so,the password should be=Maximum of all stable numbers+Minimum of all Unstable numbers=1313+122=1435

FIND PASSWORD(DIFFERENT TESTCASE WITH MODEL-4)

Detective Buckshee junior has been approached by the shantiniketan kids society for help in finding the password to the games complex. After hearing the scenario, detective Buckshee junior realizes that he will need a programmer's support. He contacts you requests your help. Please help the detective by writing a function to generate the password.

The scenario is as below-

Five numbers are available with the kids.

These numbers are either stable or unstable

A numbers is stable if each of its digit occur the same number of times, i.e the frequency of each digit in the number is the same.

For e.g:2277,4004,11,23,583835,1010 are examples of stable numbers.

Similarly, A number is unstable if the frequency of each digit in the number is NOT the same.

For eg:221,4314,101,233,58135,101 are examples of unstable numbers.

The password can be found as below-

i.e Password=sum of all stable numbers - sum of all Unstable numbers

Assuming that the five numbers are passed to a function as input1, input2, input3, input4, input5.

Complete the function to find and return the password.

For example:

If input1=12,input2=1313,input3=122,input4=678 and input5=898 , we see that there are THREE stable numbers 12,1313 and 678 and

TWO unstable numbers 122 and 898

So, the password should be=sum of all stable numbers – sum of all Unstable numbers=983

FIND PASSWORD (DIFFERENT TESTCASE WITH MODEL – 5)

Detective Buckshee junior has been approached by the shantiniketan kids society for help in finding the password to the games complex. After hearing the scenario, detective Buckshee junior realizes that he will need a programmer's support. He contacts you requests your help. Please help the detective by writing a function to generate the password.

The scenario is as below-

Five numbers are available with the kids.

These numbers are either stable or unstable

A number is stable if each of its digit occur the same number of times, i.e the frequency of each digit in the number is the same.

For e.g:2277,4004,11,23,583835,1010 are examples of stable numbers.

Similarly, A number is unstable if the frequency of each digit in the number is NOT the same.

For eg:221,4314,101,233,58135,101 are examples of unstable numbers.

The password can be found as below-

i.e Password=Maximum of all stable numbers - Minimum of all Unstable numbers

Assuming that the five numbers are passed to a function as input1, input2, input3, input4, input5.

Complete the function to find and return the password.

For example:

If input1=12,input2=1313,input3=122,input4=678 and input5=898 , we see that there are THREE stable numbers 12,1313 and 678 and

TWO unstable numbers 122 and 898

So, the Password should be=Maximum of all stable numbers - Minimum of all Unstable numbers=1313-122=1191

FIND PASSWORD(DIFFERENT TESTCASE WITH MODEL-6)

Detective Buckshee junior has been approached by the shantiniketan kids society for help in finding the password to the games complex. After hearing the scenario, detective Buckshee junior realizes that he will need a programmer's support. He contacts you requests your help. Please help the detective by writing a function to generate the password.

[28-07-2025 14:33] Jogendra: The scenario is as below-

Five numbers are available with the kids.

These numbers are either stable or unstable

A number is stable if each of its digit occur the same number of times, i.e the frequency of each digit in the number is the same.

For e.g:2277,4004,11,23,583835,1010 are examples of stable numbers.

Similarly, A number is unstable if the frequency of each digit in the number is NOT the same.

For eg:221,4314,101,233,58135,101 are examples of unstable numbers.

The password can be found as below-

i.e Password=Maximum of all Unstable numbers - Minimum of all Unstable numbers

Assuming that the five numbers are passed to a function as input1, input2, input3, input4, input5.

Complete the function to find and return the password.

For example:

If input1=12,input2=1313,input3=122,input4=678 and input5=898 , we see that there are THREE stable numbers 12,1313 and 678 and

TWO unstable numbers 122 and 898

So, the Password should be=Maximum of all Unstable numbers - Minimum of all Unstable numbers=898-122=776

FIND PASSWORD(DIFFERENT TESTCASE WITH MODEL-7)

Detective Buckshee junior has been approached by the shantiniketan kids society for help in finding the password to the games complex. After hearing the scenario, detective Buckshee junior realizes that he will need a programmer's support. He contacts you requests your help. Please help the detective by writing a function to generate the password.

The scenario is as below-

Five numbers are available with the kids.

These numbers are either stable or unstable

A numbers is stable if each of its digit occur the same number of times, i.e the frequency of each digit in the number is the same.

For e.g:2277,4004,11,23,583835,1010 are examples of stable numbers.

Similarly, A number is unstable if the frequency of each digit in the number is NOT the same.

For eg:221,4314,101,233,58135,101 are examples of unstable numbers.

The password can be found as below-

i.e Password=sum of all Unstable numbers

Assuming that the five numbers are passed to a function as input1, input2, input3, input4, input5.

Complete the function to find and return the password.

For example:

If input1=12,input2=1313,input3=122,input4=678 and input5=898 , we see that there are

THREE stable numbers 12,1313 and 678 and

TWO unstable numbers 122 and 898

So, the Password should be=sum of all Unstable numbers=898-122=1020

FIND PASSWORD(DIFFERENT TESTCASE WITH MODEL-8)

Detective Buckshee junior has been approached by the shantiniketan kids society for help in finding the password to the games complex. After hearing the scenario, detective Buckshee junior realizes that he will need a programmer's support. He contacts you requests your help. Please help the detective by writing a function to generate the password.

The scenario is as below-

Five numbers are available with the kids.

These numbers are either stable or unstable

A numbers is stable if each of its digit occur the same number of times, i.e the frequency of each digit in the number is the same.

For e.g:2277,4004,11,23,583835,1010 are examples of stable numbers.

Similarly, A number is unstable if the frequency of each digit in the number is NOT the same.

For eg:221,4314,101,233,58135,101 are examples of unstable numbers.

The password can be found as below-

i.e Password=Maximum of all stable numbers - Minimum of all stable numbers

Assuming that the five numbers are passed to a function as input1, input2, input3, input4, input5.
Complete the function to find and return the password.

[28-07-2025 14:33] Jogendra: For example:

If input1=12,input2=1313,input3=122,input4=678 and input5=898 , we see that there are

THREE stable numbers 12,1313 and 678 and

TWO unstable numbers 122 and 898

So, the Password should be=Maximum of all stable numbers - Minimum of all stable numbers=1313-12=1301

FIND PASSWORD(DIFFERENT TESTCASE WITH MODEL-9)

Detective Buckshee junior has been approached by the shantiniketan kids society for help in finding the password to the games complex. After hearing the scenario, detective Buckshee junior realizes that he will need a programmer's support. He contacts you requests your help. Please help the detective by writing a function to generate the password.

The scenario is as below-

Five numbers are available with the kids.

These numbers are either stable or unstable

A numbers is stable if each of its digit occur the same number of times, i.e the frequency of each digit in the number is the same.

For e.g:2277,4004,11,23,583835,1010 are examples of stable numbers.

Similarly, A number is unstable if the frequency of each digit in the number is NOT the same.

For eg:221,4314,101,233,58135,101 are examples of unstable numbers.

The password can be found as below-

i.e Password=Maximum of all stable numbers + Minimum of all stable numbers

Assuming that the five numbers are passed to a function as input1, input2, input3, input4, input5.
Complete the function to find and return the password.

For example:

If input1=12,input2=1313,input3=122,input4=678 and input5=898 , we see that there are

THREE stable numbers 12,1313 and 678 and

TWO unstable numbers 122 and 898

So, the Password should be=Maximum of all stable numbers + Minimum of all stable numbers=1313 + 12=1325
