



United States Department of the Interior

FISH AND WILDLIFE SERVICE
South Florida Ecological Services Office
1339 20th Street
Vero Beach, Florida 32960

May 25, 2007



Colonel Paul L. Grosskruger
U.S. Army Corps of Engineers
701 San Marcos Boulevard, Room 372
Jacksonville, Florida 32207-8175

Service Federal Activity Code: 41420-2006-FA-1582
Date Received: July 24, 2006
Formal Consultation Initiation Date: April 23, 2007
Project: ICWW maintenance dredging
and beach placement
Applicant: U.S. Army Corps of Engineers
County: Palm Beach County

Dear Colonel Grosskruger:

This document transmits the Fish and Wildlife Service's (Service) biological opinion based on our review of the proposed maintenance dredging of the intracoastal waterway (ICWW) and beach placement activities south of Jupiter Inlet, Palm Beach County, Florida. The U.S. Army Corps of Engineers (Corps) determined on July 20, 2006, the proposed project "may affect" the threatened loggerhead sea turtle (*Caretta caretta*), the endangered leatherback sea turtle (*Dermochelys coriacea*), the endangered green sea turtle (*Chelonia mydas*), the endangered hawksbill sea turtle (*Eretmochelys imbricata*), and the endangered Kemp's ridley sea turtle (*Lepidochelys kempii*), and we concur with your determination. This Biological Opinion is provided in accordance with section 7 of the Endangered Species Act of 1973, as amended (Act) (16 U.S.C. 1531 *et seq.*).

In the letter dated July 20, 2006, the Corps also determined the proposed action "may affect, but is not likely to adversely affect," the West Indian manatee, wood stork, piping plover, and beach jacquemontia. In order to protect these species, the Corps will ensure specific construction safety precautions are implemented as outlined in the *Standard Manatee Conditions for In-Water Work* (Florida Fish and Wildlife Conservation Commission (FWC) 2005), *Draft Species Conservation Guidelines – South Florida - Wood Storks* (Service 2004), and standard migratory bird protection measures. In addition, a qualified botanist will conduct a pre-construction survey for beach jacquemontia. If any specimens are likely to be destroyed by the project, a buffer will be created around these specimens, or they will be relocated in accordance with the species' recovery plan. Based upon implementation of the above stated conditions, the Service concurs with the determination in regard to these stated listed species.

This Biological Opinion is based on information provided in the Corps' letter and biological assessment dated July 20, 2006, and telephone conversations and email correspondence with the Corps. A complete administrative record of this consultation is on file at the South Florida Ecological Services Office, Vero Beach, Florida.



CONSULTATION HISTORY

On July 24, 2006, the Service received the Corps' request for formal consultation and Biological Assessment concerning the proposed dredging of approximately 16,000 linear feet of the ICWW adjacent to Jupiter Inlet. In addition, beach compatible material removed from the ICWW is proposed for beach placement along approximately 5,500 linear feet south of the Inlet.

On November 15, 2006, the Service requested additional information from the Corps.

On December 11, 2006, the Service received the requested information from the Corps. In addition, the Corps inquired if the proposed ICWW dredging project and the Jupiter Inlet District sand trap dredging project could be combined into the same Biological Opinion.

On December 20, 2006, the Service informed the Corps via email that both dredging projects could be combined into one Biological Opinion.

On January 22, 2007, the Service received the Corps' Public Notice and request for formal consultation dated January 19, 2007, regarding the proposed dredging of Jupiter Inlet sand trap permit application [SAJ-1989-506(IP-LAO)].

On February 20, 2007, the Service received an email from the Corps requesting that the Service provide a separate Biological Opinion for each of the proposed dredging projects.

On March 8, 2007, the Service sent out an email to the Corps requesting additional information.

On March 20, 2007, the Service received the requested information from the Corps. The Service sent out another email requesting clarification of information.

On April 23, 2007, the Corps provided the requested information and the Service initiated formal consultation.

BIOLOGICAL OPINION

DESCRIPTION OF THE PROPOSED ACTION

The Corps proposes to maintenance dredge approximately 16,000 feet of the ICWW along Palm Beach County, Florida (Figure 1). The proposed action is necessary to restore the authorized channel depths in order to maintain unobstructed navigation of the ICWW and to control shoreline erosion south of the Inlet. The project area is highly dynamic with past maintenance dredging projects having taken place in 1983, 1987, 1991, 1995, and 2004. Approximately 130,000 cubic yards (cy) of material will be removed from the channel between Cuts P-1 and P-9 (Figure 1) most likely using a cutterhead dredge or a small hopper dredge as an alternative. The pipeline corridor will run along the bottom of the channel, exit south of the jetty, and then south on the beach. Surveys have determined that seagrasses are not present within the pipeline corridor. The beach compatible material will be discharged 6 to 10 feet from the end of the

discharge pipe and form a large pile where it will be graded using bulldozers along 5,500 feet of shoreline between DEP monuments R-13 and R-19 (Figure 1). Based on estimated erosional losses within the project area, considerably more than 130,000 cy of material has been lost, hence, the beach profile will not be elevated above historical heights. As documented by past dredging projects and recent sample analyses, no beach incompatible material is present. If any beach incompatible material is encountered, it will be deposited within an agency approved upland or offshore site.

The project is expected to begin February 15, 2008, and end April 30, 2008. Completion of the entire project is expected to take approximately 45 days; however, the contractor will have up to April 30, 2008, due to possible delays based on equipment malfunctions and adverse weather. The project is expected to take place during both daylight and night-time hours.

The action area is defined as all areas to be affected directly or indirectly by the action and not merely the immediate area involved in the action. The Service identifies the action area to include shoreline in Palm Beach County between DEP monuments R-13 and R-19 (1.04 miles) and the ICWW between Cuts P-1 and P-9 (3 miles).

STATUS OF THE SPECIES/CRITICAL HABITAT

Species/critical habitat description

Loggerhead Sea Turtle

The loggerhead sea turtle, listed as a threatened species on July 28, 1978 (43 Federal Register [FR] 32800), inhabits the continental shelves and estuarine environments along the margins of the Atlantic, Pacific, and Indian Oceans. Loggerhead sea turtles nest within the continental U.S. from Louisiana to Virginia. Major nesting concentrations in the U.S. are found on the coastal islands of North Carolina, South Carolina, and Georgia, and on the Atlantic and Gulf coasts of Florida (Hopkins and Richardson 1984).

No critical habitat has been designated for the loggerhead sea turtle.

Green Sea Turtle

The green sea turtle was federally listed on July 28, 1978 (43 FR 32800). Breeding populations of the green turtle in Florida and along the Pacific Coast of Mexico are listed as endangered; all other populations are listed as threatened. The green turtle has a worldwide distribution in tropical and subtropical waters. Major green turtle nesting colonies in the Atlantic occur on Ascension Island, Aves Island, Costa Rica, and Surinam. Within the U.S., green turtles nest in small numbers in the U.S. Virgin Islands and Puerto Rico, and in larger numbers along the east coast of Florida, particularly in Brevard, Indian River, St. Lucie, Martin, Palm Beach, and Broward Counties (NOAA Fisheries and Service 1991a). Nesting has also been documented along the Gulf coast of Florida on Santa Rosa Island (Okaloosa and Escambia Counties) and from Pinellas County through Collier County (FWC statewide nesting database). Green turtles have been known to nest in Georgia, but only on rare occasions (Georgia Department of Natural

Resources statewide nesting database). The green turtle also nests sporadically in North Carolina and South Carolina (North Carolina Wildlife Resources Commission statewide nesting database; South Carolina Department of Natural Resources statewide nesting database). Unconfirmed nesting of green turtles in Alabama has also been reported (Bon Secour National Wildlife Refuge nesting reports).

Critical habitat for the green sea turtle has been designated for the waters surrounding Culebra Island, Puerto Rico, and its outlying keys.

Leatherback Sea Turtle

The leatherback sea turtle, listed as an endangered species on June 2, 1970 (35 FR 8491), nests on shores of the Atlantic, Pacific and Indian Oceans. Non-breeding animals have been recorded as far north as the British Isles and the Maritime Provinces of Canada and as far south as Argentina and the Cape of Good Hope (Pritchard 1992). Nesting grounds are distributed worldwide, with the Pacific Coast of Mexico supporting the world's largest known concentration of nesting leatherbacks. The largest nesting colony in the wider Caribbean region is found in French Guiana, but nesting occurs frequently, although in lesser numbers, from Costa Rica to Columbia and in Guyana, Surinam, and Trinidad (NOAA Fisheries and Service 1992, National Research Council 1990).

The leatherback regularly nests in the U.S. in Puerto Rico, the U.S. Virgin Islands, and along the Atlantic coast of Florida as far north as Georgia (NOAA Fisheries and Service 1992). Leatherback turtles have been known to nest in Georgia, South Carolina, and North Carolina, but only on rare occasions (North Carolina Wildlife Resources Commission, South Carolina Department of Natural Resources, and Georgia Department of Natural Resources statewide nesting databases). Leatherback nesting has also been reported on the northwest coast of Florida (LeBuff 1990, FWC statewide nesting database); a false crawl (non-nesting emergence) has been observed on Sanibel Island (LeBuff 1990).

Marine and terrestrial critical habitat for the leatherback sea turtle has been designated at Sandy Point on the western end of the island of St. Croix, U.S. Virgin Islands.

Hawksbill Sea Turtle

The hawksbill sea turtle was listed as an endangered species on June 2, 1970 (35 FR 8491). The hawksbill is found in tropical and subtropical seas of the Atlantic, Pacific, and Indian Oceans. The species is widely distributed in the Caribbean Sea and western Atlantic Ocean. Within the continental U.S., hawksbill sea turtle nesting is rare and is restricted to the southeastern coast of Florida (Volusia through Dade Counties) and the Florida Keys (Monroe County) (Meylan 1992, Meylan et al. 1995). However, hawksbill tracks are difficult to differentiate from those of loggerheads and may not be recognized by surveyors. Therefore, surveys in Florida likely underestimate actual hawksbill nesting numbers (Meylan et al. 1995). In the U.S. Caribbean, hawksbill nesting occurs on beaches throughout Puerto Rico and the U.S. Virgin Islands (NOAA Fisheries and Service 1993).

Critical habitat for the hawksbill sea turtle has been designated for selected beaches and/or waters of Mona, Monito, Culebrita, and Culebra Islands, Puerto Rico.

Kemp's Ridley Sea Turtle

The Kemp's ridley sea turtle was listed as endangered on December 2, 1970 (35 FR 18320). The range of the Kemp's ridley includes the Gulf coasts of Mexico and the U.S., and the Atlantic coast of North America as far north as Nova Scotia and Newfoundland. Most Kemp's ridleys nest on the coastal beaches of the Mexican states of Tamaulipas and Veracruz, although a very small number of Kemp's ridleys nest consistently along the Texas coast (Turtle Expert Working Group 1998). In addition, rare nesting events have been reported in Florida, Alabama, South Carolina, and North Carolina. Hatchlings, after leaving the nesting beach, are believed to become entrained in eddies within the Gulf of Mexico, where they are dispersed within the Gulf and Atlantic by oceanic surface currents until they reach about 8 inches in length, at which size they enter coastal shallow water habitats (Ogren 1989). Outside of nesting, adult Kemp's ridleys are believed to spend most of their time in the Gulf of Mexico, while juveniles and subadults also regularly occur along the eastern seaboard of the U.S. (Service and NOAA Fisheries 1992). No critical habitat has been designated for the Kemp's ridley sea turtle.

Life history

Loggerhead Sea Turtle

Loggerheads are known to nest from one to seven times within a nesting season (Talbert et al. 1980, Richardson and Richardson 1982, Lenarz et al. 1981); the mean is approximately 4.1 (Murphy and Hopkins 1984). The interval between nesting events within a season varies around a mean of about 14 days (Dodd 1988). Mean clutch size varies from about 100 to 126 eggs along the southeastern United States coast (NOAA Fisheries and Service 1991b). Nesting migration intervals of years are most common in loggerheads, but the number can vary from 1 to 7 years (Dodd 1988). Age at sexual maturity is believed to be about 20 to 30 years (Turtle Expert Working Group 1998).

Green Sea Turtle

Green turtles deposit from one to nine clutches within a nesting season, but the overall average is 3.3. The mean interval between nesting events within a season is 13 days (Hirth 1997). Mean clutch size varies widely among populations. Average clutch size reported for Florida was 136 eggs in 130 clutches (Witherington and Ehrhart 1989). Only occasionally do females produce clutches in successive years. Usually 2, 3, 4, or more years intervene between breeding seasons (NOAA Fisheries and Service 1991a). Age at sexual maturity is believed to be 20 to 50 years (Hirth 1997).

Leatherback Sea Turtle

Leatherbacks nest an average of five to seven times within a nesting season, with an observed maximum of 11 (NOAA Fisheries and Service 1992). The interval between nesting events

within a season is about 10 days. Clutch size averages 80 to 85 yolked eggs, with the addition of usually a few dozen smaller, yolkless eggs, mostly laid toward the end of the clutch (Pritchard 1992). Nesting migration intervals of 2 to 3 years were observed in leatherbacks nesting on Sandy Point National Wildlife Refuge, St. Croix, U.S. Virgin Islands (McDonald and Dutton 1996). Leatherbacks are believed to reach sexual maturity in 6 to 10 years (Zug and Parham 1996).

Hawksbill Sea Turtle

Hawksbills nest on average 4.5 times per season at intervals of approximately 14 days (Corliss et al. 1989). In Florida and the U.S. Caribbean, clutch size is approximately 140 eggs, although several records exist of over 200 eggs per nest (NOAA Fisheries and Service 1993). On the basis of limited information, nesting migration intervals of 2 to 3 years appear to predominate. Hawksbills are recruited into the reef environment at about 14 inches in length and are believed to begin breeding about 30 years later. However, the time required to reach 14 inches in length is unknown and growth rates vary geographically. As a result, actual age at sexual maturity is not known.

Kemp's Ridley Sea Turtle

Nesting occurs from April into July during which time the turtles appear off the Tamaulipas and Veracruz coasts of Mexico. Precipitated by strong winds, the females swarm to mass nesting emergences, known as *arribadas* or *arribazones*, to nest during daylight hours. Clutch size averages 100 eggs (Service and NOAA Fisheries 1992). Some females breed annually and nest an average of 1 to 4 times in a season at intervals of 10 to 28 days. Age at sexual maturity is believed to be between 7 to 15 years (Turtle Expert Working Group 1998).

Population dynamics

Loggerhead Sea Turtle

Total estimated nesting in the Southeast is approximately 50,000 to 90,000 nests per year (FWC statewide nesting database 2004, Georgia Department of Natural Resources statewide nesting database 2004, South Carolina Department of Natural Resources statewide nesting database 2004, North Carolina Wildlife Resources Commission statewide nesting database 2004). In 1998, 85,988 nests were documented in Florida alone. However, in 2001, 2002, 2003, and 2004, this number dropped to 69,657, 62,905, 56,852, and 47,173, respectively. An analysis of nesting data from the Florida Index Nesting Beach Survey (INBS) Program from 1989 to 2004, a period encompassing index surveys that are more consistent and more accurate than surveys in previous years, has shown no detectable trend but, more recently (1998 through 2004), has shown evidence of a declining trend (Blair Witherington, FWC, personal communication, 2005). Given inherent annual fluctuations in nesting and the short time period over which the decline has been noted, caution is warranted in interpreting the decrease in terms of nesting trends.

From a global perspective, the southeastern U.S. nesting aggregation is of paramount importance to the survival of the species and is second in size only to that which nests on islands in the Arabian Sea off Oman (Ross 1982, Ehrhart 1989, NOAA Fisheries and Service 1991b). The status of the Oman loggerhead nesting population, reported to be the largest in the world (Ross 1979), is uncertain because of the lack of long-term standardized nesting or foraging ground

surveys and its vulnerability to increasing development pressures near major nesting beaches and threats from fisheries interactions on foraging grounds and migration routes (Earl Possardt, Service, personal communication, 2005). The loggerhead nesting aggregations in Oman, the southeastern U.S., and Australia have been estimated to account for about 88 percent of nesting worldwide (NOAA Fisheries and Service 1991b). About 80 percent of loggerhead nesting in the southeastern U.S. occurs in six Florida counties (Brevard, Indian River, St. Lucie, Martin, Palm Beach, and Broward Counties) (NOAA Fisheries and Service 1991b).

Green Sea Turtle

About 150 to 2,750 females are estimated to nest on beaches in the continental U.S. annually (FWC 2006). In the U.S. Pacific, over 90 percent of nesting throughout the Hawaiian archipelago occurs at the French Frigate Shoals, where about 200 to 700 females nest each year (NOAA Fisheries and Service 1998a). Elsewhere in the U.S. Pacific, nesting takes place at scattered locations in the Commonwealth of the Northern Marianas, Guam, and American Samoa. In the western Pacific, the largest green turtle nesting group in the world occurs on Raine Island, Australia, where thousands of females nest nightly in an average nesting season (Limpus et al. 1993). In the Indian Ocean, major nesting beaches occur in Oman where 30,000 females are reported to nest annually (Ross and Barwani 1995).

Leatherback Sea Turtle

Recent estimates of global nesting populations indicate 26,000 to 43,000 nesting females annually (Spotila et al. 1996). The largest nesting populations at present occur in the western Atlantic in French Guiana (4,500 to 7,500 females nesting/year) and Colombia (estimated several thousand nests annually), and in the western Pacific in West Papua (formerly Irian Jaya) and Indonesia (about 600 to 650 females nesting/year). In the United States, small nesting populations occur on the Florida east coast (100 females/year) (FWC 2006), Sandy Point, U.S. Virgin Islands (50 to 190 females/year) (Alexander et al. 2002), and Puerto Rico (30 to 90 females/year).

Hawksbill Sea Turtle

About 15,000 females are estimated to nest each year throughout the world with the Caribbean accounting for 20 to 30 percent of the world's hawksbill population. Only five regional populations remain with more than 1,000 females nesting annually (Seychelles, Mexico, Indonesia, and two in Australia). Mexico is now the most important region for hawksbills in the Caribbean with 3,000 nests per year (Meylan 1999). Other significant, but smaller populations in the Caribbean still occur in Martinique, Jamaica, Guatemala, Nicaragua, Grenada, Dominican Republic, Turks and Caicos Islands, Cuba, Puerto Rico, and U.S. Virgin Islands. In the U.S. Caribbean, about 150 to 500 nests per year are laid on Mona Island, Puerto Rico, and 70 to 130 nests per year on Buck Island Reef National Monument, U.S. Virgin Islands. In the U.S. Pacific, hawksbills nest only on main island beaches in Hawaii, primarily along the east coast of the island of Hawaii. Hawksbill nesting has also been documented in American Samoa and Guam (NOAA Fisheries and Service 1998b).

Kemp's Ridley Sea Turtle

The 40,000 nesting females estimated from a single mass nesting emergence in 1947 reflected a much larger total number of nesting turtles in that year than exists today (Carr 1963, Hildebrand 1963). However, nesting in Mexico has been steadily increasing in recent years - from 702 nests in 1985 to over 10,000 nests in 2005 (Service 2005). Despite protection for the nests, turtles have been and continue to be lost to incidental catch by shrimp trawls (Service and NOAA Fisheries 1992).

Status and distribution

Loggerhead Sea Turtle

Genetic research involving analysis of mitochondrial DNA has identified five different loggerhead subpopulations/nesting aggregations in the western North Atlantic: (1) the Northern Subpopulation occurring from North Carolina to around Cape Canaveral, Florida (about 29° N.); (2) South Florida Subpopulation occurring from about 29° N. on Florida's east coast to Sarasota on Florida's west coast; (3) Dry Tortugas, Florida, Subpopulation, (4) Northwest Florida Subpopulation occurring at Eglin Air Force Base and the beaches near Panama City; and (5) Yucatán Subpopulation occurring on the eastern Yucatán Peninsula, Mexico (Bowen 1994, 1995, Bowen et al. 1993, Encalada et al. 1998, Pearce 2001). These data indicate gene flow between these five regions is very low. If nesting females are extirpated from one of these regions, regional dispersal will not be sufficient to replenish the depleted nesting subpopulation.

The Northern Subpopulation has declined substantially since the early 1970s. Recent estimates of loggerhead nesting trends from standardized daily beach surveys showed significant declines ranging from 1.5 to 2.0 percent annually (Mark Dodd, Georgia Department of Natural Resources, personal communication, 2005). Nest totals from aerial surveys conducted by the South Carolina Department of Natural Resources showed a 3.3 percent annual decline in nesting since 1980. Overall, there is strong statistical evidence to suggest the Northern Subpopulation has sustained a long-term decline.

Data from all beaches where nesting activity has been recorded indicate the South Florida Subpopulation has shown significant increases over the last 25 years. However, an analysis of nesting data from the Florida INBS Program from 1989 to 2002, a period encompassing index surveys that are more consistent and more accurate than surveys in previous years, has shown no detectable trend and, more recently (1998 through 2002), has shown evidence of a declining trend (Blair Witherington, FWC, personal communication, 2003). Given inherent annual fluctuations in nesting and the short time period over which the decline has been noted, caution is warranted in interpreting the decrease in terms of nesting trends.

A near census of the Florida Panhandle Subpopulation undertaken from 1989 to 2002 reveals a mean of 1,028 nests per year, which equates to about 251 females nesting per year (FWC 2006). Evaluation of long-term nesting trends for the Florida Panhandle Subpopulation is difficult because of changed and expanded beach coverage. Although there are now 8 years (1997 to 2004) of INBS data for the Florida Panhandle Subpopulation, the time series is too short to detect a trend (Blair Witherington, FWC, personal communication, 2005).

A near census of the Dry Tortugas Subpopulation undertaken from 1995 to 2001 reveals a mean of 213 nests per year, which equates to about 50 females nesting per year (FWC 2006). The trend data for the Dry Tortugas Subpopulation are from beaches that were not part of the State of Florida's INBS program prior to 2004, but have moderately good monitoring consistency. There are 7 continuous years (1995 to 2001) of data for this Subpopulation, but the time series is too short to detect a trend (Blair Witherington, FWC, personal communication, 2005).

Nesting surveys in the Yucatán Subpopulations have been too irregular to date to allow for a meaningful trend analysis (Turtle Expert Working Group 1998, 2000).

Threats include incidental take from channel dredging and commercial trawling, longline, and gill net fisheries; loss or degradation of nesting habitat from coastal development and beach armoring; disorientation of hatchlings by beachfront lighting; excessive nest predation by native and non-native predators; degradation of foraging habitat; marine pollution and debris; watercraft strikes; and disease. There is particular concern about the extensive incidental take of juvenile loggerheads in the eastern Atlantic by longline fishing vessels from several countries.

Green Sea Turtle

Total population estimates for the green turtle are unavailable, and trends based on nesting data are difficult to assess because of large annual fluctuations in numbers of nesting females. For instance, in Florida, where the majority of green turtle nesting in the southeastern U.S. occurs, estimates range from 150 to 2,750 females nesting annually (FWC 2006). Populations in Surinam, and Tortuguero, Costa Rica, may be stable, but there is insufficient data for other areas to confirm a trend.

A major factor contributing to the green turtle's decline worldwide is commercial harvest for eggs and food. Fibropapillomatosis, a disease of sea turtles characterized by the development of multiple tumors on the skin and internal organs, is also a mortality factor and has seriously impacted green turtle populations in Florida, Hawaii, and other parts of the world. The tumors interfere with swimming, eating, breathing, vision, and reproduction, and turtles with heavy tumor burdens may die. Other threats include loss or degradation of nesting habitat from coastal development and beach armoring; disorientation of hatchlings by beachfront lighting; excessive nest predation by native and non-native predators; degradation of foraging habitat; marine pollution and debris; watercraft strikes; and incidental take from channel dredging and commercial fishing operations.

Leatherback Sea Turtle

Declines in leatherback nesting have occurred over the last 2 decades along the Pacific coasts of Mexico and Costa Rica. The Mexican leatherback nesting population, once considered to be the world's largest leatherback nesting population (historically estimated to be 65 percent of the worldwide population), is now less than 1 percent of its estimated size in 1980. Spotila et al. (1996) estimated the number of leatherback sea turtles nesting on 28 beaches throughout the world from the literature and from communications with investigators studying those beaches.

The estimated worldwide population of leatherbacks in 1995 was about 34,500 females on these beaches with a lower limit of about 26,200 and an upper limit of about 42,900. This is less than one third the 1980 estimate of 115,000. Leatherbacks are rare in the Indian Ocean and in very low numbers in the western Pacific Ocean. The largest population is in the western Atlantic. Using an age-based demographic model, Spotila et al. (1996) determined leatherback populations in the Indian Ocean and western Pacific Ocean cannot withstand even moderate levels of adult mortality and even the Atlantic populations are being exploited at a rate that cannot be sustained. They concluded leatherbacks are on the road to extinction and further population declines can be expected unless we take action to reduce adult mortality and increase survival of eggs and hatchlings.

The crash of the Pacific leatherback population is believed primarily to be the result of exploitation by humans for the eggs and meat, as well as incidental take in numerous commercial fisheries of the Pacific. Other factors threatening leatherbacks globally include loss or degradation of nesting habitat from coastal development; disorientation of hatchlings by beachfront lighting; excessive nest predation by native and non-native predators; degradation of foraging habitat; marine pollution and debris; and watercraft strikes.

Hawksbill Sea Turtle

The hawksbill sea turtle has experienced global population declines of 80 percent or more during the past century and continued declines are projected (Meylan and Donnelly 1999). Most populations are declining, depleted, or remnants of larger aggregations. Hawksbills were previously abundant, as evidenced by high-density nesting at a few remaining sites and by trade statistics. The decline of this species is primarily due to human exploitation for tortoiseshell. While the legal hawksbill shell trade ended when Japan agreed to stop importing shell in 1993, a significant illegal trade continues. It is believed individual hawksbill populations around the world will continue to disappear under the current regime of exploitation for eggs, meat, and tortoiseshell, loss of nesting and foraging habitat, incidental capture in fishing gear, ingestion of and entanglement in marine debris, oil pollution, and boat collisions. Hawksbills are closely associated with coral reefs, one of the most endangered of all marine ecosystem types.

Kemp's Ridley Sea Turtle

The decline of this species was primarily due to human activities, including the direct harvest of adults and eggs and incidental capture in commercial fishing operations. Today, under strict protection, the population appears to be in the early stages of recovery. The recent nesting increase can be attributed to full protection of nesting females and their nests in Mexico resulting from a bi-national effort between Mexico and the U.S. to prevent the extinction of the Kemp's ridley, and the requirement to use turtle excluder devices in shrimp trawls both in the United States and Mexico.

The Mexico government also prohibits harvesting and is working to increase the population through more intensive law enforcement, by fencing nest areas to diminish natural predation, and by relocating all nests into corrals to prevent poaching and predation. While relocation of nests

into corrals is currently a necessary management measure, this relocation and concentration of eggs into a “safe” area is of concern since it makes the eggs more susceptible to reduced viability due to movement-induced mortality, disease vectors, catastrophic events like hurricanes, and marine predators once the predators learn where to concentrate their efforts.

Analysis of the species/critical habitat likely to be affected

The proposed action has the potential to adversely affect nesting females, nests, and hatchlings within the proposed project area. The effects of the proposed action on sea turtles will be considered further in the remaining sections of this Biological Opinion. Potential effects include destruction of nests deposited within the boundaries of the proposed project, harassment in the form of disturbing or interfering with female turtles attempting to nest within the construction area or on adjacent beaches as a result of construction activities, and behavior modification of nesting females due to escarpment formation within the project area during the nesting season resulting in false crawls or situations where they choose marginal or unsuitable nesting areas to deposit eggs. The quality of the placed sand could affect the ability of female turtles to nest, the suitability of the nest incubation environment, and the ability of hatchling to emerge from the nest.

Critical habitat has not been designated in the continental U.S.; therefore, the proposed action would not result in an adverse modification.

ENVIRONMENTAL BASELINE

Palm Beach County is located within the most densely populated nesting range for three species of sea turtles: loggerhead, green, and leatherback sea turtles that regularly nest along the beaches of southeast Florida. Between 1998 and 2003, Palm Beach County beaches supported approximately 18 percent of the overall sea turtle nesting in Florida. Only one other Florida county, Brevard, supports a greater percentage of sea turtle nesting. In total, 11,893 and 13,327 sea turtle nests were recorded in 2004 and 2005, respectively, along the 42 miles of Palm Beach County beach included in the FWC’s Florida Statewide Nesting Beach Survey (Table 1). The distribution of nests among species in 2004 included 10,759 loggerhead sea turtles, 968 green sea turtles, and 166 leatherback sea turtles; and in 2005 the distribution included 10,791 loggerhead sea turtles, 2,252 green sea turtles, and 284 leatherback sea turtles (Table 1). Only false crawls have been recorded for Kemp’s Ridley sea turtles in Palm Beach County (Meylan et al. 1995).

Status of the species/critical habitat within the action area

In Palm Beach County, 283 and 317 sea turtle nests were laid per mile in 2004 and 2005, respectively (Table 1). The nesting density within the action area exceeded the County in 2004 and 2005, with a nesting density of 459 and 622 nests per mile in 2004 and 2005, respectively (Table 2).

Loggerhead Sea Turtle

The loggerhead sea turtle nesting and hatching season for Palm Beach County extends from March 15 through November 30. Incubation ranges from about 45 to 95 days.

Along the Florida east coast between 1995 and 2005, Palm Beach County had the second highest average nesting of loggerhead sea turtles, with approximately 13,000 nests laid (Palm Beach County 2006a). In 2004, a total of 10,759 loggerhead sea turtle nests, or 256 nests per mile, were laid in Palm Beach County (Table 1). In 2005, a total of 10,791 nests, or 257 nests per mile, were laid. Loggerhead sea turtles laid 637 nests in 2004 and 864 nests in 2005, along the beach between DEP monument R-13 and R-21 (Table 2).

Loggerhead sea turtles made 15,822 false crawls in 2004 and 14,345 in 2005 in Palm Beach County (Table 1). Within the action area, loggerhead turtles made 1,502 false crawls in 2004 and 1,429 in 2005 (Table 2). In both years, there were more false crawls than nests laid.

Green Sea Turtle

The green sea turtle nesting and hatching season for Palm Beach County extends from May 1 through November 30. Incubation ranges from about 45 to 75 days.

Between 1995 and 2005, Palm Beach County had the second highest average green sea turtle nesting along the Florida east coast; with a little more than 1,000 nests laid per year (Palm Beach County 2006b). In Palm Beach County, a total of 968 green sea turtle nests were laid in 2004, which is equivalent to 23 nests per mile (Table 1). A total of 2,252 green sea turtle nests were laid in 2005, for a nesting density of 54 nests per mile (Table 1). Between DEP monument R-13 and R-21, 41 green sea turtle nests were laid in 2004, and 57 in 2005 (Table 2). Only the nesting density in 2004 (27 nests per mile) exceeded that of the County.

In Palm Beach County, 1,283 green sea turtle false crawls were made in 2004 and 3,142 in 2005 (Table 1). Between DEP monument R-13 and R-21, there were 97 and 247 false crawls in 2004 and 2005, respectively, which was greater than the number of nests laid each year (Table 2).

Leatherback Sea Turtle

The leatherback sea turtle nesting and hatching season for Palm Beach County extends from February 15 through November 30. Incubation ranges from about 55 to 75 days.

Palm Beach County had the highest average leatherback sea turtle nesting along the Florida east coast from 1995 to 2005 (Palm Beach County 2006c). The total number of leatherback sea turtle nests laid in Palm Beach County was 166 in 2004, and 284 in 2005 (Table 1). The nesting density was four nests per mile in 2004, and seven nests per mile in 2005. Between DEP monument R-13 and R-21, 20 leatherback sea turtle nests were laid in 2004, and 25 in 2005 (Table 2). Leatherback sea turtles made 25 false crawls 2004 and 52 in 2005, in Palm Beach County (Table 1). Three and eight false crawls were recorded in 2004 and 2005, respectively, between DEP monument R-13 and R-21 (Table 2). In both years, there were fewer false crawls than nests laid.

Hawksbill Sea Turtle

The hawksbill sea turtle nesting and hatching season for Palm Beach County extends from June 1 through December 31. Incubation lasts about 60 days.

Nesting by this species has been documented five times in Palm Beach County since 1985, with the most recent report in 2004 (Meylan et al. 1995, FWC 2006). No nesting has been reported from the proposed action area.

Kemp's Ridley Sea Turtle

Although no nesting has been reported in Palm Beach County for Kemp's ridley turtles, four false crawls at Phipps Ocean Park were reported in 1989 (Meylan et al. 1995).

The majority of nesting surveys conducted in Florida occur during the morning hours and are based on interpretation of the tracks left by the turtles as they ascend and descend the beach; the turtles themselves are rarely observed. Because both hawksbill and Kemp's ridley turtle tracks are difficult to discern from loggerhead tracks, it is likely that nesting by both species is underreported (Meylan et al. 1995).

Factors affecting the species habitat within the action area

The project area includes 1.04 miles of shoreline which has an average beach width of 200 feet. Three government entities (Environmental Resources Management, Florida Inland Navigation District, and Jupiter Inlet District) routinely place sand along the project area south of Jupiter Inlet. Between 1995 and 2005, over 2,000,000 cy of sand has been dredged from the Intracoastal Waterway and the sand trap and placed within the project area.

A primary threat to sea turtles along nesting shorelines includes sea turtle hatchling disorientation as a result of artificial lighting along the beach. Typically, sea turtle hatchlings will emerge from the nest and orient themselves towards the brighter, open horizon of the ocean (Salmon et al. 1992). If artificial lights are visible from the beach, sea turtle hatchlings tend to travel toward the artificial lights instead of the ocean. Disorientation events often result in hatchling mortality as a result of dehydration, predation, and motor vehicle strikes. In addition, regular beach maintenance in the form of tractor tilling may disrupt or impact deposited nests and nesting females. Plastics, styrofoam, and fishing line are pollutants that may negatively impact nesting success and nearshore foraging.

As restored beaches equilibrate to a more natural profile, steep vertical escarpments often form along the seaward edge of the constructed beach berm and this presents a physical barrier to nesting turtles. Additionally, as beach profiles equilibrate, losses of nests laid in the seaward portions of the renourished beach due to erosion may be high. Steinitz et al. (1998) following long-term studies at Jupiter Inlet indicated that at 2 years post-renourishment, nesting success was considerably higher than pre-renourishment levels and similar to densities found on nearby non-eroded beaches. However, the nesting success declined as the renourished beach eroded and narrowed until the next renourishment event.

Lighting

Except for shoreline under public ownership, much of the remaining coast in Palm Beach County is developed. The majority of development consists of multi-family residences, with smaller sections containing commercial and single-family residences. Because many beach-front

residences are not fully occupied during much of the nesting season, impacts to nesting sea turtles and hatchlings may be reduced.

Palm Beach County has a Sea Turtle Protection Ordinance that includes measures to reduce impacts of coastal lighting on sea turtles. The proposed project area is subject to this ordinance.

EFFECTS OF THE ACTION

The analysis of the direct and indirect effects of the proposed action on sea turtles and the interrelated and interdependent activities of those effects was based on beneficial and detrimental factors.

Factors to be considered

The proposed action has the potential to adversely affect nesting females, nest, and hatchlings within the proposed project area through the placement of dredged material on the beach within Palm Beach County, Florida.

Analyses for effects of the action

Beneficial effects

The placement of sand on a beach with reduced dry fore-dune habitat may increase sea turtle nesting habitat if the placed sand is highly compatible (*e.g.*, grain size, shape, color, etc.) with naturally occurring beach sediments in the area, and compaction and escarpment remediation measures are incorporated into the project. In addition, a nourished beach that is designed and constructed to mimic a natural beach system may be more stable than the eroding one it replaces, thereby benefiting sea turtles.

Direct effects

Placement of up to 130,000 cy of sand along 1.04 miles of beach in and of itself may not provide suitable nesting habitat for sea turtles. Although placement of beach compatible material may increase the potential nesting area, significant negative impacts to sea turtles may result if protective measures are not incorporated during project construction. Placement of material during the nesting season, particularly on or near high density nesting beaches, can cause increased loss of eggs and hatchlings and along with other mortality sources, may impact the long-term survival of the species. For example, projects conducted during the nesting and hatching season could result in the loss of sea turtles through disruption of adult nesting activity and by burial or crushing of nests or hatchlings.

1. Nest relocation

Besides the potential for missing nests during a nest relocation program, there is a potential for eggs to be damaged by their movement, particularly if eggs are not relocated within 12 hours of

deposition (Limpus et al. 1979). Nest relocation can have adverse impacts on incubation temperature (and hence sex ratios), gas exchange parameters, hydric environment of nests, hatching success, and hatchling emergence (Limpus et al. 1979, Ackerman 1980, Parmenter 1980, Spotila et al. 1983, McGehee 1990). Relocating nests into sands deficient in oxygen or moisture can result in mortality, morbidity, and reduced behavioral competence of hatchlings. Water availability is known to influence the incubation environment of the embryos and hatchlings of turtles with flexible-shelled eggs, which has been shown to affect nitrogen excretion (Packard et al. 1984), mobilization of calcium (Packard and Packard 1986), mobilization of yolk nutrients (Packard et al. 1985), hatchling size (Packard et al. 1981, McGehee 1990), energy reserves in the yolk at hatching (Packard et al. 1988), and locomotory ability of hatchlings (Miller et al. 1987). In a 1994 Florida study comparing loggerhead hatching and emergence success of relocated nests with *in situ* nests, Moody (1998) found hatching success was lower in relocated nests at 9 of 12 beaches evaluated and emergence success was lower in relocated nests at 10 of 12 beaches surveyed in 1993 and 1994.

2. Missed nests

Although a nesting survey and nest marking program would reduce the potential for nests to be impacted by construction activities, nests may be inadvertently missed (when crawls are obscured by rainfall, wind, and/or tides) or misidentified as false crawls during daily patrols. Even under the best of conditions, about 7 percent of the nests can be misidentified as false crawls by experienced sea turtle nest surveyors (Schroeder 1994).

3. Equipment

The placement of pipelines and the use of heavy machinery on the beach during a construction project may also have adverse effects on sea turtles. They can create barriers to nesting females emerging from the surf and crawling up the beach, causing a higher incidence of false crawls and unnecessary energy expenditure.

4. Artificial lighting

Visual cues are the primary sea-finding mechanism for hatchling sea turtles (Mrosovsky and Carr 1967, Mrosovsky and Shettleworth 1968, Dickerson and Nelson 1989, Witherington and Bjorndal 1991). When artificial lighting is present on or near the beach, it can misdirect hatchlings once they emerge from their nests and prevent them from reaching the ocean (Philibosian 1976, Mann 1977). In addition, a significant reduction in sea turtle nesting activity has been documented on beaches illuminated with artificial lights (Witherington 1992). Therefore, construction lights along a project beach and on the dredging vessel may deter females from coming ashore to nest, misdirect females trying to return to the surf after a nesting event, and misdirect emergent hatchlings from adjacent non-project beaches. Any source of bright lighting can profoundly affect the orientation of hatchlings, both during the crawl from the beach to the ocean and once they begin swimming offshore. Hatchlings attracted to light sources on dredging barges may not only suffer from interference in migration, but may also experience higher probabilities of predation to predatory fishes that are also attracted to the barge lights. This impact could be reduced by using the minimum amount of light necessary (may require shielding) or low pressure sodium lighting during project construction.

Indirect effects

Many of the direct effects of beach nourishment may persist over time and become indirect impacts. These indirect effects include increased susceptibility of relocated nests to catastrophic events, the consequences of potential increased beachfront development, changes in the physical characteristics of the beach, the formation of escarpments, and future sand migration.

1. Increased susceptibility to catastrophic events

Nest relocation may concentrate eggs in an area making them more susceptible to catastrophic events. Hatchlings released from concentrated areas may also be subject to greater predation rates from both land and marine predators, because the predators learn where to concentrate their efforts (Glenn 1998, Wyneken et al. 1998).

2. Increased beachfront development

Pilkey and Dixon (1996) state that beach replenishment frequently leads to more development in greater density within shorefront communities that are then left with a future of further replenishment or more drastic stabilization measures. Dean (1999) also notes that the very existence of a beach nourishment project can encourage more development in coastal areas. Following completion of a beach nourishment project in Miami during 1982, investment in new and updated facilities substantially increased tourism in the area (National Research Council 1995). Increased building density immediately adjacent to the beach often resulted as older buildings were replaced by much larger ones that accommodated more beach users. Overall, shoreline management creates an upward spiral of initial protective measures resulting in more expensive development which leads to the need for more and larger protective measures. Increased shoreline development may adversely affect sea turtle nesting success. Greater development may support larger populations of mammalian predators, such as foxes and raccoons, than undeveloped areas (National Research Council 1990), and can also result in greater adverse effects due to artificial lighting, as discussed above.

3. Changes in the physical environment

Beach placement activities may result in changes in sand density (compaction), beach shear resistance (hardness), beach moisture content, beach slope, sand color, sand grain size, sand grain shape, and sand grain mineral content if the placed sand is dissimilar from the original beach sand (Nelson and Dickerson 1988a). These changes could result in adverse impacts on nest site selection, digging behavior, clutch viability, and emergence by hatchlings (Nelson and Dickerson 1987, Nelson 1988).

Beach compaction and unnatural beach profiles that may result from beach placement activities could negatively impact sea turtles regardless of the timing of projects. Very fine sand and/or the use of heavy machinery can cause sand compaction on nourished beaches (Nelson et al. 1987; Nelson and Dickerson 1988a). Significant reductions in nesting success (*e.g.*, increase in false crawls) have been documented on severely compacted nourished beaches (Fletemeyer

1980, Raymond 1984, Nelson and Dickerson 1987, Nelson et al. 1987), and increased false crawls may result in increased physiological stress to nesting females. Sand compaction may increase the length of time required for female sea turtles to excavate nests and also cause increased physiological stress to the animals (Nelson and Dickerson 1988b). Nelson and Dickerson (1988c) concluded that, in general, beaches nourished from offshore borrow sites are harder than natural beaches, and while some may soften over time through erosion and accretion of sand, others may remain hard for 10 years or more.

These impacts can be minimized by using suitable sand and by tilling compacted sand after project completion. The level of compaction of a beach can be assessed by measuring sand compaction using a cone penetrometer (Nelson 1987). Tilling of a nourished beach with a root rake may reduce the sand compaction to levels comparable to unnourished beaches. However, a pilot study by Nelson and Dickerson (1988b) showed that a tilled nourished beach will remain uncompacted for up to 1 year. Therefore, the Service requires multi-year beach compaction monitoring and, if necessary, tilling to ensure project impacts on sea turtles are minimized.

A change in sediment color on a beach could change the natural incubation temperatures of nests in an area, which, in turn, could alter natural sex ratios. To provide the most suitable sediment for nesting sea turtles, the color of the nourished sediments must resemble the natural beach sand in the area. Tilling, natural reworking of sediments, and bleaching from exposure to the sun would help to lighten dark nourishment sediments; however, the timeframe for sediment mixing and bleaching to occur could be critical to a successful sea turtle nesting season.

4. Escarpment formation

On nourished beaches, steep escarpments may develop along their water line interface as they adjust from an unnatural construction profile to a more natural beach profile (Coastal Engineering Research Center 1984, Nelson et al. 1987). These escarpments can hamper or prevent access to nesting sites (Nelson and Blihovde 1998). Researchers have shown that female turtles coming ashore to nest can be discouraged by the formation of an escarpment, leading to situations where they choose marginal or unsuitable nesting areas to deposit eggs (*e.g.*, in front of the escarpments, which often results in failure of nests due to prolonged tidal inundation). This impact can be minimized by leveling any escarpments prior to the nesting season.

Species' response to a proposed action

Ernest and Martin (1999) conducted a comprehensive study to assess the effects of beach nourishment on loggerhead sea turtle nesting and reproductive success. The following findings illustrate sea turtle responses to and recovery from a nourishment project. A significantly larger proportion of turtles emerging on nourished beaches abandoned their nesting attempts than turtles emerging on control or pre-nourished beaches. This reduction in nesting success was most pronounced during the first year following project construction and is most likely the result of changes in physical beach characteristics associated with the nourishment project (*e.g.*, beach profile, sediment grain size, beach compaction, frequency and extent of escarpments). During the first post-construction year, the time required for turtles to excavate an egg chamber on the

untilled, hard-packed sands of one treatment area increased significantly relative to Control and background conditions. However, in another treatment area, tilling was effective in reducing sediment compaction to levels that did not significantly prolong digging times. As natural processes reduced compaction levels on nourished beaches during the second post-construction year, digging times returned to background levels.

During the first post-construction year, nests on the nourished beaches were deposited significantly farther from both the toe of the dune and the tide line than nests on control beaches. Furthermore, nests were distributed throughout all available habitat and were not clustered near the dune as they were in the control area. As the width of nourished beaches decreased during the second year, among-treatment differences in nest placement diminished. More nests were washed out on the wide, flat beaches of the nourished treatments than on the narrower steeply sloped beaches of the control beach. This phenomenon persisted through the second post-construction year monitoring and resulted from the placement of nests near the seaward edge of the beach berm where dramatic profile changes, caused by erosion and scarping, occurred as the beach equilibrated to a more natural contour.

As with other beach nourishment projects, Ernest and Martin (1999) found the principal effect of nourishment on sea turtle reproduction was a reduction in nesting success during the first year following project construction. Although most studies have attributed this phenomenon to an increase in beach compaction and escarpment formation, Ernest and Martin (1999) indicate changes in beach profile may be more important. Regardless, as a nourished beach is reworked by natural processes in subsequent years and adjusts from an unnatural construction profile to a more natural beach profile, beach compaction and the frequency of escarpment formation decline, and nesting and nesting success return to levels found on natural beaches.

CUMULATIVE EFFECTS

Cumulative effects include the effects of future State, tribal, local, or private actions that are reasonably certain to occur in the action area considered in this Biological Opinion. Future Federal actions that are unrelated to the proposed action are not considered in this section because they require separate consultation pursuant to section 7 of the Act. The Service has considered potential cumulative effects of this project on sea turtles and, in this instance, there are no cumulative effects.

CONCLUSION

After reviewing the current status of the loggerhead, green, leatherback, hawksbill and Kemp's ridley sea turtles, the environmental baseline for the action area, the effects of the proposed dredging and beach nourishment, and the cumulative effects, it is the Service's biological opinion that the dredging and beach nourishment project, as proposed, is not likely to jeopardize the continued existence of these sea turtles. No critical habitat has been designated for the loggerhead, green, leatherback, Kemp's Ridley, and hawksbill sea turtles in the continental United States; therefore, none will be affected.

INCIDENTAL TAKE STATEMENT

Section 9 of the Act and Federal regulation pursuant to section 4(d) of the Act prohibit the take of endangered or threatened species, respectively, without special exemption. Take is defined as to harass, harm, pursue, hunt, shoot, wound, kill, trap, capture or collect, or to attempt to engage in any such conduct. Harm is further defined by the Service to include significant habitat modification or degradation that results in death or injury to listed species by significantly impairing essential behavioral patterns, including breeding, feeding, or sheltering. Harass is defined by the Service as intentional or negligent actions that create the likelihood of injury to listed species to such an extent as to significantly disrupt normal behavior patterns which include, but are not limited to, breeding, feeding, or sheltering. Incidental take is defined as take that is incidental to, and not the purpose of, carrying out an otherwise lawful activity. Under the terms of section 7(b)(4) and section 7(o)(2), taking that is incidental to and not intended as part of the agency action is not considered to be prohibited under the Act provided that such taking is in compliance with the terms and conditions of this incidental take statement.

The measures described below are non-discretionary, and must be implemented by the Corps so they become binding conditions of any grant or permit issued to the Applicant, as appropriate, for the exemption in section 7(o)(2) to apply. The Corps has a continuing duty to regulate the activity covered by this incidental take statement. If the Corps (1) fails to assume and implement the terms and conditions or, (2) fails to adhere to the terms and conditions of the incidental take statement through enforceable terms that are added to the permit or grant document, the protective coverage of section 7(o)(2) may lapse. In order to monitor the impact of incidental take, the Corps must report the progress of the action and its impacts on the species to the Service as specified in the incidental take statement [50 CFR §402.14(i)(3)].

AMOUNT OR EXTENT OF TAKE

The Service anticipates 1.04 miles of nesting beach habitat could be taken as a result of this proposed action. The take is expected to be in the form of: (1) destruction of all nests that may be constructed and eggs that may be deposited from March 1 through April 30 and from September 1 through September 30 and missed by a nest survey and egg relocation program within the boundaries of the proposed project; (2) destruction of all nests deposited from October 1 through February 28 (or 29 as applicable) when a nest survey and egg relocation program is not required to be in place within the boundaries of the proposed project; (3) reduced hatching success due to egg mortality during relocation and adverse conditions at the relocation site; (4) harassment in the form of disturbing or interfering with female turtles attempting to nest within the construction area or on adjacent beaches as a result of construction activities; (5) misdirection of hatchling turtles on beaches adjacent to the construction area as they emerge from the nest and crawl to the water as a result of project lighting; (6) behavior modification of nesting females due to escarpment formation within the project area during a nesting season, resulting in false crawls or situations where they choose marginal or unsuitable nesting areas to deposit eggs; and (7) destruction of nests from escarpment leveling within a nesting season when such leveling has been approved by the Service.

Incidental take is anticipated for only the 1.04 miles of beach that has been identified for sand placement. The Service anticipates incidental take of sea turtles will be difficult to detect for the following reasons: (1) the turtles nest primarily at night and all nests are not found because [a] natural factors, such as rainfall, wind, and tides may obscure crawls and [b] human-caused factors, such as pedestrian and vehicular traffic, may obscure crawls, and result in nests being destroyed because they were missed during a nesting survey and egg relocation program; (2) the total number of hatchlings per undiscovered nest is unknown; (3) the reduction in percent hatching and emerging success per relocated nest over the natural nest site is unknown; (4) an unknown number of females may avoid the project beach and be forced to nest in a less than optimal area; (5) lights may misdirect an unknown number of hatchlings and cause death; and (6) escarpments may form and cause an unknown number of females from accessing a suitable nesting site. However, the level of take of these species can be anticipated by the disturbance and nourishment of suitable turtle nesting beach habitat because: (1) turtles nest within the project site; (2) beach nourishment will likely occur during a portion of the nesting season; (3) the nourishment project will modify the incubation substrate, beach slope, and sand compaction; and (4) artificial lighting will deter and/or misdirect nesting females and hatchlings.

EFFECT OF THE TAKE

In the accompanying Biological Opinion, the Service determined this level of anticipated take is not likely to result in jeopardy to the species. Critical habitat has not been designated in the project area; therefore, the project will not result in destruction or adverse modification of critical habitat.

REASONABLE AND PRUDENT MEASURES

1. Beach quality sand suitable for sea turtle nesting, successful incubation, and hatchling emergence must be used on the project site;
2. Beach placement activities must not occur from May 1 through October 31, the period of peak sea turtle egg laying and egg hatching, to reduce the possibility of sea turtle nest burial, crushing of eggs, or nest excavation;
3. If the beach placement activities will be conducted during the period from March 1 through April 30, surveys for early nesting sea turtles must be conducted. If nests are constructed in the area of beach placement and dune restoration, the eggs must be relocated;
4. If the project will be conducted during the period from November 1 through November 30, surveys for late nesting turtles must be conducted. If nests are constructed in the area of beach placement, the eggs must be relocated;
5. Immediately after completion of the project and prior to the next 3 nesting seasons, beach compaction must be monitored and tilling must be conducted as required by March 1 to reduce the likelihood of impacting sea turtle nesting and hatching activities. The March 1 deadline is required to reduce impacts to leatherbacks that nest in greater frequency along the South Atlantic coast of Florida than elsewhere in the continental United States;

6. Immediately after completion of the project and prior to the next 3 nesting seasons, monitoring must be conducted to determine if escarpments are present and escarpments must be leveled as required to reduce the likelihood of impacting sea turtle nesting and hatching activities;
7. The Applicant must ensure contractors doing the beach placement and/or dune enhancement work fully understand the sea turtle protection measures detailed in this incidental take statement;
8. During the early (March 1 through April 30) and late (November 1 through November 30) portions of the nesting season, construction equipment and pipes must be stored in a manner that will minimize impacts to sea turtles to the maximum extent practicable; and
9. During the sea turtle nesting season (March 1 through April 30 and November 1 through November 30) lighting associated with the project must be minimized to reduce the possibility of disrupting and disorienting nesting and/or hatchling sea turtles.

TERMS AND CONDITIONS

In order to be exempt from the prohibitions of section 9 of the ESA, the Corps must comply with the following terms and conditions, which implement the reasonable and prudent measures described above and outline required reporting/monitoring requirements. These terms and conditions are non-discretionary.

1. All fill material placed must be sand similar to a native beach in the vicinity of the site that has not been affected by prior nourishment activities. The fill material must be similar in both coloration and grain size distribution (sand grain frequency, mean and median grain size, and sorting coefficient) to the native beach and must not contain:
 - 1a. Greater than 10 percent, by weight, silt, clay, or colloids passing the #230 sieve;
 - 1b. Greater than 10 percent, by weight, fine gravel retained on the #4 sieve;
 - 1c. Coarse gravel, cobbles, or other material retained on the ¾-in sieve in a percentage or size greater than found on the native beach; and
 - 1d. Construction debris, toxic material, or other foreign matter; and not result in contamination or cementation of the beach.
2. Project construction must be started after October 31 and be completed before May 1. During the May 1 through October 31 period, no construction equipment or pipes will be stored on the beach;
3. If the project will be conducted during the period from March 1 through April 30, daily early morning surveys for sea turtle nests must be conducted. In the event a sea turtle nest is excavated during construction activities, all work shall cease in that area immediately and the permitted person responsible for egg relocation for the project should be notified so the eggs can be relocated per the following requirements:

- 3a. Nesting surveys and egg relocations will only be conducted by personnel with prior experience and training in nesting survey and egg relocation procedures. Surveyors must perform under the supervision of a qualified professional with a valid FWC Marine Turtle Permit. Nesting surveys must be conducted daily between sunrise and 9 a.m. Surveys must be performed in such a manner so as to ensure that construction activity does not occur in any location prior to completion of the necessary sea turtle protection measures; and
- 3b. Only those nests that may be affected by construction activities will be relocated. Nests requiring relocation must be moved no later than 9 a.m. the morning following deposition to a nearby self-release beach site in a secure setting where artificial lighting will not interfere with hatchling orientation. Nest relocations in association with construction activities must cease when construction activities no longer threaten nests. Nests deposited within areas where construction activities have ceased or will not occur for 65 days must be marked and left in place unless other factors threaten the success of the nest. Any nests left in the active construction zone must be clearly marked, and all mechanical equipment must avoid nests by at least 10 feet.
4. If the project will be conducted during the period from November 1 through November 30, daily early morning sea turtle nesting surveys must be conducted 65 days prior to project initiation and continue through September 30. In the event a sea turtle nest is excavated during construction activities, all work shall cease in that area immediately and the permitted person responsible for egg relocation for the project should be notified so the eggs can be relocated per the preceding requirements;
5. Immediately after completion of the project (April 30) and prior to March 1 for 3 subsequent years, sand compaction must be monitored in the area of restoration in accordance with a protocol agreed to by the Service, the FWC, and the Applicant. At a minimum, the protocol provided under 5a and 5b below must be followed. If required, the area must be tilled to a depth of 36 inches. All tilling activity must be completed prior to March 1. An annual summary of compaction surveys and the actions taken must be submitted to the Service. (NOTE: The requirement for compaction monitoring can be eliminated if the decision is made to till regardless of post-construction compaction levels. Also, out-year compaction monitoring and remediation are not required if placed material no longer remains on the beach.)
- 5a. Compaction sampling stations must be located at 500-foot intervals along the project area. One station must be at the seaward edge of the dune/bulkhead line (when material is placed in this area), and one station must be midway between the dune line and the high water line (normal wrack line).

At each station, the cone penetrometer will be pushed to a depth of 6, 12, and 18 inches three times (three replicates). Material may be removed from the hole if necessary to ensure accurate readings of successive levels of sediment. The penetrometer may need to be reset between pushes, especially if sediment layering exists. Layers of

highly compact material may lie over less compact layers. Replicates will be located as close to each other as possible, without interacting with the previous hole and/or disturbed sediments. The three replicate compaction values for each depth will be averaged to produce final values for each depth at each station. Reports will include all 18 values for each transect line, and the final 6 averaged compaction values.

- 5b. If the average value for any depth exceeds 500 pounds per square inch (psi) for any two or more adjacent stations, then that area must be tilled prior to March 1. If values exceeding 500 psi are distributed throughout the project area, but in no case do those values exist at two adjacent stations at the same depth, then consultation with the Service will be required to determine if tilling is required. If a few values exceeding 500 psi are present randomly within the project area, tilling will not be required.
6. Visual surveys for escarpments along the project area must be made immediately after completion of the project and prior to March 1 for 3 subsequent years if placed sand still remains on the beach. All escarpments shall be leveled, or the beach profile shall be reconfigured, to minimize escarpment formation. In addition, weekly surveys of the project area shall be conducted during the 3 nesting seasons following completion of fill placement as follows:
 - 6a. The number of escarpments and their location relative to DEP reference monuments shall be recorded during each weekly survey and reported relative to the length of the beach surveyed (*e.g.*, 50 percent escarpments). Notations on the height of these escarpments shall be included (0 to 2 feet, 2 to 4 feet, and 4 feet or higher) as well as the maximum height of all escarpments; and
 - 6b. Escarpments that interfere with sea turtle nesting or that exceed 18 inches in height for a distance of 100 feet must be leveled to the natural beach contour by April 30. Any escarpment removal shall be reported relative to R-monument locations. The Service must be contacted immediately if subsequent reformation of escarpments that interfere with sea turtle nesting or that exceed 18 inches in height for a distance of 100 feet occurs during the nesting and hatching season (May 1 to October 31) to determine the appropriate action to be taken. If it is determined escarpment leveling is required during the nesting or hatching season, the Service will provide a brief written authorization that describes methods to be used to reduce the likelihood of impacting existing nests. An annual summary of escarpment surveys and actions taken must be submitted to the Service. (NOTE: Out-year escarpment monitoring and remediation are not required if placed material no longer remains on the dry beach.)
7. The Applicant must arrange a meeting between representatives of the contractor, the Service, the FWC, and the permitted person responsible for egg relocation at least 30 days prior to the commencement of work on this project. At least 10 days advance notice must be provided prior to conducting this meeting. This will provide an opportunity for explanation and/or clarification of the sea turtle protection measures.

8. From March 1 through April 30 and November 1 through November 30, staging areas for construction equipment must be located off the beach to the maximum extent practicable. Night-time storage of construction equipment not in use must be off the beach to minimize disturbance to sea turtle nesting and hatching activities during this period. In addition, all construction pipes placed on the beach must be located as far landward as possible without compromising the integrity of the existing or reconstructed dune system. Temporary storage of pipes must be off the beach to the maximum extent possible. Temporary storage of pipes on the beach must be in such a manner so as to impact the least amount of nesting habitat and must likewise not compromise the integrity of the dune systems (placement of pipes perpendicular to the shoreline is recommended as the method of storage).
9. From March 1 through April 30 and November 1 through November 30, all on-beach lighting associated with the project must be limited to the immediate area of active construction only and must be the minimal lighting necessary to comply with all safety requirements. Lighting on offshore or onshore equipment must be minimized through reduction, shielding, lowering, and appropriate placement of lights to avoid excessive illumination of the water, while meeting all U.S. Coast Guard and Occupational Safety and Health Administration (OSHA) requirements. Shielded low pressure sodium vapor lights are recommended for lights on offshore equipment that cannot be eliminated, and for illumination of the nesting beach and nearshore waters. Light intensity of lighting plants must be reduced to the minimum standard required by OSHA for General Construction areas, in order not to misdirect sea turtles. Shields must be affixed to the light housing and be large enough to block light from all lamps from being transmitted outside the construction area (Figure 2).
10. A lighting survey shall be conducted prior to April 30 of the first nesting season following beach placement activities and action taken to ensure no lights or light sources are visible from the newly elevated beach. A report summarizing all lights visible, using standard survey techniques for such surveys, shall be submitted to the Service by May 15 and documenting all compliance and enforcement action. Additional lighting surveys shall be conducted monthly through August and results reported by the 15th of each month of the first nesting season after project completion;
11. A report describing the actions taken to implement the terms and conditions of this incidental take statement must be submitted to the FWC, Imperiled Species Management Section, Tequesta office and the Tallahassee office as well as the South Florida Ecological Services Office, Vero Beach, Florida within 60 days of completion of the proposed work for each year when the activity has occurred. This report will include the dates of actual construction activities, names and qualifications of personnel involved in nest surveys and relocation activities, descriptions and locations of self-release beach sites, nest survey and relocation results, and hatching success of nests.
12. Upon locating a dead, injured, or sick endangered or threatened sea turtle specimen, initial notification must be made to the FWC at 1-888-404-3922, and the South Florida Ecological Services Office biologist listed at the end of this Biological Opinion. Care

should be taken in handling sick or injured specimens to ensure effective treatment and care and in handling dead specimens to preserve biological materials in the best possible state for later analysis of cause of death. In conjunction with the care of sick or injured endangered or threatened species or preservation of biological materials from a dead animal, the finder has the responsibility to ensure evidence intrinsic to the specimen is not unnecessarily disturbed.

The Service believes incidental take will be limited to 1.04 miles of beach that has been identified for sand placement. The reasonable and prudent measures, with their implementing terms and conditions, are designed to minimize the impact of incidental take that might otherwise result from the proposed action. The Service believes no more than the following types of incidental take will result from the proposed action: (1) destruction of all nests that may be constructed and eggs that may be deposited and missed by a nest survey and egg relocation program within the boundaries of the proposed project; (2) destruction of all nests deposited during the period when a nest survey and egg relocation program is not required to be in place within the boundaries of the proposed project; (3) reduced hatching success due to egg mortality during relocation and adverse conditions at the relocation site; (4) harassment in the form of disturbing or interfering with female turtles attempting to nest within the construction area or on adjacent beaches as a result of construction activities; (5) disorientation of hatchling turtles on beaches adjacent to the construction area as they emerge from the nest and crawl to the water as a result of project lighting; (6) behavior modification of nesting females due to escarpment formation within the project area during a nesting season, resulting in false crawls or situations where they choose marginal or unsuitable nesting areas to deposit eggs; and (7) destruction of nests from escarpment leveling within a nesting season when such leveling has been approved by the Service.

The amount or extent of incidental take for sea turtles will be considered exceeded if the project results in more than one placement of sand annually on the 1.04 miles of beach that has been identified for sand placement. If, during the course of the action, this level of incidental take is exceeded, such incidental take represents new information requiring reinitiation of consultation and review of the reasonable and prudent measures provided. The Corps must immediately provide an explanation of the causes of the taking and review with the Service the need for possible modification of the reasonable and prudent measures.

CONSERVATION RECOMMENDATIONS

Section 7(a)(1) of the Act directs Federal agencies to utilize their authorities to further the purposes of the Act by carrying out conservation programs for the benefit of endangered and threatened species. Conservation recommendations are discretionary agency activities to minimize or avoid adverse effects of a proposed action on listed species or critical habitat, to help implement recovery plans, or to develop information.

1. Appropriate native salt-resistant dune vegetation should be established on the restored dunes. DEP, Bureau of Beaches and Wetland Resources, can provide technical assistance on the specifications for design and implementation.

2. Surveys for nesting success of sea turtles should be continued for a minimum of 3 years following beach placement activities to determine whether sea turtle nesting success has been adversely impacted.
3. Educational signs should be placed where appropriate at beach access points explaining the importance of the area to sea turtles and/or the life history of sea turtle species that nest in the area.


In order for the Service to be kept informed of actions minimizing or avoiding adverse effects or benefiting listed species or their habitats, the Service requests notification of the implementation of any conservation recommendations.

REINITIATION NOTICE

This concludes formal consultation on the action outlined in the request. As provided in 50 CFR §402.16, reinitiation of formal consultation is required where discretionary Federal agency involvement or control over the action has been retained (or is authorized by law) and if: (1) the amount or extent of incidental take is exceeded; (2) new information reveals effects of the agency action that may affect listed species or critical habitat in a manner or to an extent not considered in this opinion; (3) the agency action is subsequently modified in a manner that causes an effect to the listed species or critical habitat not considered in this opinion; or (4) a new species is listed or critical habitat designated that may be affected by the action. In instances where the amount or extent of incidental take is exceeded, any operations causing such take must cease pending reinitiation.

Should you have additional questions or require clarification, please contact Jeff Howe at 772-562-3909, extension 283.

Sincerely yours,



Paul Souza
Field Supervisor
South Florida Ecological Services Office

cc:

Corps, Jacksonville, Florida (Catherine Brooks)
DEP, Tallahassee, Florida (Stephanie Gudeman)
EPA, West Palm Beach, Florida
FWC, Tallahassee, Florida (Robbin Trindell)
NOAA Fisheries, West Palm Beach, Florida (Jocelyn Karazsia)
Service, Jacksonville, Florida (Sandy MacPherson)
Service, Atlanta, Georgia (Joe Johnson) electronic copy
USGS, Florida Integrated Science Center, Gainesville, Florida (Susan Walls)

LITERATURE CITED

- Ackerman, R.A. 1980. Physiological and ecological aspects of gas exchange by sea turtle eggs. *American Zoologist* 20:575-583.
- Alexander, J., S. Deishley, K. Garrett, W. Coles, and D. Dutton. 2002. Tagging and nesting research on leatherback sea turtles (*Dermochelys coriacea*) on Sandy Point, St. Croix, U.S. Virgin Islands, 2002. Annual Report to the Fish and Wildlife Service. 41 pages.
- Bowen, B.W. 1994. Letter dated November 17, 1994, to Sandy MacPherson, National Sea Turtle Coordinator, U.S. Fish and Wildlife Service, Jacksonville, Florida. University of Florida. Gainesville, Florida.
- Bowen, B.W. 1995. Letter dated October 26, 1995, to Sandy MacPherson, National Sea Turtle Coordinator, U.S. Fish and Wildlife Service, Jacksonville, Florida. University of Florida. Gainesville, Florida.
- Bowen, B., J.C. Avise, J.I. Richardson, A.B. Meylan, D. Margaritoulis, and S.R. Hopkins-Murphy. 1993. Population structure of loggerhead turtles (*Caretta caretta*) in the northwestern Atlantic Ocean and Mediterranean Sea. *Conservation Biology* 7(4):834-844.
- Carr, A.F. 1963. Panspecific reproductive convergence in *Lepidochelys kempii*. *Ergebn. Biol.*, 26:298-303.
- Coastal Engineering Research Center. 1984. Shore protection manual, volumes I and II. U.S. Army Corps of Engineers Waterways Experiment Station, Vicksburg, Mississippi.
- Corliss, L.A., J.I. Richardson, C. Ryder, and R. Bell. 1989. The hawksbills of Jumby Bay, Antigua, West Indies. Pages 33-35 in Eckert, S.A., K.L. Eckert, and T.H. Richardson (compilers). Proceedings of the Ninth Annual Workshop on Sea Turtle Conservation and Biology. NOAA Technical Memorandum NMFS-SEFC-232.
- Dean, C. 1999. Against the tide: the battle for America's beaches. Columbia University Press; New York, New York.
- Dickerson, D.D. and D.A. Nelson. 1989. Recent results on hatchling orientation responses to light wavelengths and intensities. Pages 41-43 in Eckert, S.A., K.L. Eckert, and T.H. Richardson (compilers). Proceedings of the 9th Annual Workshop on Sea Turtle Conservation and Biology. NOAA Technical Memorandum NMFS-SEFC-232.
- Dodd, C.K., Jr. 1988. Synopsis of the biological data on the loggerhead sea turtle *Caretta caretta* (Linnaeus 1758). U.S. Fish and Wildlife Service, Biological Report 88(14).
- Ehrhart, L.M. 1989. Status report of the loggerhead turtle. Pages 122-139 in Ogren, L., F. Berry, K. Bjorndal, H. Kumpf, R. Mast, G. Medina, H. Reichart, and R. Witham (editors). Proceedings of the 2nd Western Atlantic Turtle Symposium. NOAA Technical Memorandum NMFS-SEFC-226.

- Encalada, S.E., K.A. Bjorndal, A.B. Bolten, J.C. Zurita, B. Schroeder, E. Possardt, C.J. Sears, and B.W. Bowen. 1998. Population structure of loggerhead turtle (*Caretta caretta*) nesting colonies in the Atlantic and Mediterranean as inferred from mitochondrial DNA control region sequences. *Marine Biology* 130:567-575.
- Ernest, R.G. and R.E. Martin. 1999. Martin County beach nourishment project: sea turtle monitoring and studies. 1997 annual report and final assessment. Unpublished report prepared for the Florida Department of Environmental Protection.
- Fletemeyer, J. 1980. Sea turtle monitoring project. Unpublished report prepared for the Broward County Environmental Quality Control Board, Florida.
- Florida Fish and Wildlife Conservation Commission (FWC). 2005. Standard Manatee Conditions for In-water Work. Tallahassee, Florida.
<http://myfwc.com/manatee/permits/StandardCondIn-waterWork.pdf>.
- Florida Fish and Wildlife Conservation Commission (FWC). 2006. FWC Marine Turtle Program. http://www.floridamarine.org/features/category_main.asp?id=1289.
- Glenn, L. 1998. The consequences of human manipulation of the coastal environment on hatchling loggerhead sea turtles (*Caretta caretta*, L.). Pages 58-59 in Byles, R., and Y. Fernandez (compilers). *Proceedings of the Sixteenth Annual Symposium on Sea Turtle Biology and Conservation*. NOAA Technical Memorandum NMFS-SEFSC-412.
- Hildebrand, H. 1963. Hallazgo del area de anidacion de la tortuga "lora" *Lepidochelys kempii* (Garman), en la costa occidental del Golfo de Mexico (Rept., Chel.). *Ciencia Mexicana* 22(4):105112.
- Hirth, H.F. 1997. Synopsis of the biological data on the green turtle *Chelonia mydas* (Linnaeus 1758). U.S. Fish and Wildlife Service, Biological Report 97(1).
- Hopkins, S.R. and J.I. Richardson (editors). 1984. Recovery plan for marine turtles. National Marine Fisheries Service, St. Petersburg, Florida.
- LeBuff, C.R., Jr. 1990. The loggerhead turtle in the eastern Gulf of Mexico. *Caretta Research, Inc.*; Sanibel Island, Florida.
- Lenarz, M.S., N.B. Frazer, M.S. Ralston, and R.B. Mast. 1981. Seven nests recorded for loggerhead turtle (*Caretta caretta*) in one season. *Herpetological Review* 12(1):9.
- Limpus, C.J., V. Baker, and J.D. Miller. 1979. Movement induced mortality of loggerhead eggs. *Herpetologica* 35(4):335-338.
- Limpus, C., J.D. Miller, and C.J. Parmenter. 1993. The northern Great Barrier Reef green turtle *Chelonia mydas* breeding population. Pages 47-50 in Smith, A.K. (compiler), K.H. Zevering and C.E. Zevering (editors). *Raine Island and Environs Great Barrier Reef: Quest to Preserve a Fragile Outpost of Nature*. Raine Island Corporation and Great Barrier Reef Marine Park Authority, Townsville, Queensland, Australia.

- Mann, T.M. 1977. Impact of developed coastline on nesting and hatchling sea turtles in southeastern Florida. M.S. thesis. Florida Atlantic University, Boca Raton, Florida.
- McDonald, D.L. and P.H. Dutton. 1996. Use of PIT tags and photoidentification to revise remigration estimates of leatherback turtles (*Dermochelys coriacea*) nesting in St. Croix, U.S. Virgin Islands, 1979-1995. *Chelonian Conservation and Biology* 2(2):148-152.
- McGehee, M.A. 1990. Effects of moisture on eggs and hatchlings of loggerhead sea turtles (*Caretta caretta*). *Herpetologica* 46(3):251-258.
- Meylan, A.B. 1992. Hawksbill turtle *Eretmochelys imbricata*. Pages 95-99 in Moler, P.E. (editor). Rare and endangered biota of Florida, volume III. University Press of Florida, Gainesville, Florida.
- Meylan, A.B. 1999. Status of the hawksbill turtle (*Eretmochelys imbricata*) in the Caribbean region. *Chelonian Conservation and Biology* 3(2):177-184.
- Meylan, A.B. and M. Donnelly. 1999. Status justification for listing the hawksbill turtle (*Eretmochelys imbricata*) as critically endangered on the 1996 IUCN *Red List of Threatened Animals*. *Chelonian Conservation and Biology* 3(2):200-224.
- Meylan, A., B. Schroeder, and A. Mosier. 1995. Sea turtle nesting activity in the State of Florida 1979-1992. Florida Marine Research Publications Number 52, St. Petersburg, Florida.
- Miller, K., G.C. Packard, and M.J. Packard. 1987. Hydric conditions during incubation influence locomotor performance of hatchling snapping turtles. *Journal of Experimental Biology* 127:401-412.
- Moody, K. 1998. The effects of nest relocation on hatching success and emergence success of the loggerhead turtle (*Caretta caretta*) in Florida. Pages 107-108 in Byles, R. and Y. Fernandez (compilers). Proceedings of the Sixteenth Annual Symposium on Sea Turtle Biology and Conservation. NOAA Technical Memorandum NMFS-SEFSC-412.
- Mrosovsky, N. and A. Carr. 1967. Preference for light of short wavelengths in hatchling green sea turtles (*Chelonia mydas*), tested on their natural nesting beaches. *Behavior* 28:217-231.
- Mrosovsky, N. and S.J. Shettleworth. 1968. Wavelength preferences and brightness cues in water finding behavior of sea turtles. *Behavior* 32:211-257.
- Murphy, T.M. and S.R. Hopkins. 1984. Aerial and ground surveys of marine turtle nesting beaches in the southeast region. Unpublished report prepared for the National Marine Fisheries Service.
- National Marine Fisheries Service (NOAA Fisheries) and U.S. Fish and Wildlife Service (Service). 1991a. Recovery plan for U.S. population of Atlantic green turtle (*Chelonia mydas*). National Marine Fisheries Service, Washington, D.C.

- National Marine Fisheries Service (NOAA Fisheries) and U.S. Fish and Wildlife Service (Service). 1991b. Recovery plan for U.S. population of loggerhead turtle (*Caretta caretta*). National Marine Fisheries Service, Washington, D.C.
- National Marine Fisheries Service (NOAA Fisheries) and U.S. Fish and Wildlife Service (Service). 1992. Recovery plan for leatherback turtles (*Dermochelys coriacea*) in the U.S. Caribbean, Atlantic, and Gulf of Mexico. National Marine Fisheries Service, Washington, D.C.
- National Marine Fisheries Service (NOAA Fisheries) and U.S. Fish and Wildlife Service (Service). 1993. Recovery plan for hawksbill turtle (*Eretmochelys imbricata*) in the U.S. Caribbean, Atlantic, and Gulf of Mexico. National Marine Fisheries Service, St. Petersburg, Florida.
- National Marine Fisheries Service (NOAA Fisheries) and U.S. Fish and Wildlife Service (Service). 1998a. Recovery plan for U.S. Pacific populations of the green turtle (*Chelonia mydas*). National Marine Fisheries Service, Silver Spring, MD. 84 pages.
- National Marine Fisheries Service (NOAA Fisheries) and U.S. Fish and Wildlife Service (Service). 1998b. Recovery plan for U.S. Pacific populations of the hawksbill turtle (*Eretmochelys imbricata*). National Marine Fisheries Service, Silver Spring, MD. 82 pages.
- National Research Council. 1990. Decline of the sea turtles: causes and prevention. National Academy Press; Washington, D.C.
- National Research Council. 1995. Beach nourishment and protection. National Academy Press; Washington, D.C.
- Nelson, D.A. 1987. The use of tilling to soften nourished beach sand consistency for nesting sea turtles. Unpublished report of the U.S. Army Corps of Engineers Waterways Experiment Station, Vicksburg, Mississippi.
- Nelson, D.A. 1988. Life history and environmental requirements of loggerhead turtles. U.S. Fish and Wildlife Service Biological Report 88(23). U.S. Army Corps of Engineers TR EL-86-2 (Rev.).
- Nelson, D.A. and B. Blihovde. 1998. Nesting sea turtle response to beach scarps. Page 113 in Byles, R., and Y. Fernandez (compilers). Proceedings of the Sixteenth Annual Symposium on Sea Turtle Biology and Conservation. NOAA Technical Memorandum NMFS-SEFSC-412.
- Nelson, D.A. and D.D. Dickerson. 1987. Correlation of loggerhead turtle nest digging times with beach sand consistency. Abstract of the 7th Annual Workshop on Sea Turtle Conservation and Biology.

- Nelson, D.A. and D.D. Dickerson. 1988a. Hardness of nourished and natural sea turtle nesting beaches on the east coast of Florida. Unpublished report of the U.S. Army Corps of Engineers Waterways Experiment Station, Vicksburg, Mississippi.
- Nelson, D.A. and D.D. Dickerson. 1988b. Response of nesting sea turtles to tilling of compacted beaches, Jupiter Island, Florida. Unpublished report of the U.S. Army Corps of Engineers Waterways Experiment Station, Vicksburg, Mississippi.
- Nelson, D.A. and D.D. Dickerson. 1988c. Hardness of nourished and natural sea turtle nesting beaches on the east coast of Florida. Unpublished report of the U.S. Army Corps of Engineers Waterways Experiment Station, Vicksburg, Mississippi.
- Nelson, D.A., K. Mauck, and J. Fletemeyer. 1987. Physical effects of beach nourishment on sea turtle nesting, Delray Beach, Florida. Technical Report EL-87-15. U.S. Army Corps of Engineers Waterways Experiment Station, Vicksburg, Mississippi.
- Ogren, L.H. 1989. Distribution of juvenile and subadult Kemp's ridley turtles: preliminary results from the 1984-1987 surveys. Pages 116-123 in Caillouet, C.W., Jr., and A.M. Landry, Jr. (eds.). Proceedings of the First International Symposium on Kemp's Ridley Sea Turtle Biology, Conservation and Management. Texas A&M University Sea Grant College Program TAMU-SG-89-105.
- Packard, M.J. and G.C. Packard. 1986. Effect of water balance on growth and calcium mobilization of embryonic painted turtles (*Chrysemys picta*). Physiological Zoology 59(4):398-405.
- Packard, G.C., M.J. Packard, and T.J. Boardman. 1984. Influence of hydration of the environment on the pattern of nitrogen excretion by embryonic snapping turtles (*Chelydra serpentina*). Journal of Experimental Biology 108:195-204.
- Packard, G.C., M.J. Packard, and W.H.N. Gutzke. 1985. Influence of hydration of the environment on eggs and embryos of the terrestrial turtle *Terrapene ornata*. Physiological Zoology 58(5):564-575.
- Packard, G.C., M.J. Packard, T.J. Boardman, and M.D. Ashen. 1981. Possible adaptive value of water exchange in flexible-shelled eggs of turtles. Science 213:471-473.
- Packard, G.C., M.J. Packard, K. Miller, and T.J. Boardman. 1988. Effects of temperature and moisture during incubation on carcass composition of hatchling snapping turtles (*Chelydra serpentina*). Journal of Comparative Physiology B 158:117-125.
- Palm Beach County. 2006a. Average loggerhead nesting along Florida's East Coast, 1995-2005. http://www.pbcgov.com/erm/protection/Images/PDF_Documents/cc_statewide.pdf Last accessed October 12, 2006.

- Palm Beach County. 2006b. Average green nesting along Florida's East Coast, 1995-2005. http://www.pbcgov.com/erm/protection/Images/PDF_Documents/cm_statewide.pdf
Last accessed October 12, 2006.
- Palm Beach County. 2006c. Average leatherback nesting along Florida's East Coast, 1995-2005. http://www.pbcgov.com/erm/protection/Images/PDF_Documents/dc_statewide.pdf
Last accessed October 12, 2006.
- Parmenter, C.J. 1980. Incubation of the eggs of the green sea turtle, *Chelonia mydas*, in Torres Strait, Australia: the effect of movement on hatchability. *Australian Wildlife Research* 7:487-491.
- Pearce, A.F. 2001. Contrasting population structure of the loggerhead turtle (*Caretta caretta*) using mitochondrial and nuclear DNA markers. M.S. thesis. University of Florida, Gainesville, Florida.
- Philibosian, R. 1976. Disorientation of hawksbill turtle hatchlings (*Eretmochelys imbricata*) by stadium lights. *Copeia* 1976:824.
- Pilkey, O.H. and K.L. Dixon. 1996. *The Corps and the shore*. Island Press; Washington, D.C.
- Pritchard, P.C.H. 1992. Leatherback turtle *Dermochelys coriacea*. Pages 214-218 in Moler, P.E. (editor). *Rare and Endangered Biota of Florida, Volume III*. University Press of Florida; Gainesville, Florida.
- Raymond, P.W. 1984. The effects of beach restoration on marine turtles nesting in south Brevard County, Florida. M.S. thesis. University of Central Florida, Orlando, Florida.
- Richardson, J.I. and T.H. Richardson. 1982. An experimental population model for the loggerhead sea turtle (*Caretta caretta*). Pages 165-176 in Bjorndal, K.A. (editor). *Biology and Conservation of Sea Turtles*. Smithsonian Institution Press; Washington, D.C.
- Ross, J.P. 1979. Sea turtles in the Sultanate of Oman. World Wildlife Fund Project 1320. May 1979 report. 53 pp.
- Ross, J.P. 1982. Historical decline of loggerhead, ridley, and leatherback sea turtles. Pages 189-195 in Bjorndal, K.A. (editor). *Biology and Conservation of Sea Turtles*. Smithsonian Institution Press; Washington, D.C.
- Ross, J.P. and M.A. Barwani. 1995. Review of sea turtles in the Arabian area. Pages 373-383 in Bjorndal, K.A. (editor). *Biology and Conservation of Sea Turtles, Revised Edition*. Smithsonian Institution Press, Washington, D.C. 615 pages.
- Salmon, M., J. Wyneken, E.U. Fritz, and M. Lucas. 1992. Ocean finding by hatchling sea turtles interplay of silhouette, slope, brightness as guideposts in orientation. Proceedings of the eleventh annual workshop in sea turtle biology and conservation. NOAA Technical Memorandum. NMFS-SEFSC-302. National Marine Fisheries Service, Southeast Fisheries Center. Miami, Florida.

- Schroeder, B.A. 1994. Florida index nesting beach surveys: are we on the right track? Pages 132-133 in Bjorndal, K.A., A.B. Bolten, D.A. Johnson, and P.J. Eliazar (compilers). Proceedings of the 14th Annual Symposium on Sea Turtle Biology and Conservation. NOAA Technical Memorandum NMFS-SEFSC-351.
- Spotila, J.R., E.A. Standora, S.J. Morreale, G.J. Ruiz, and C. Puccia. 1983. Methodology for the study of temperature related phenomena affecting sea turtle eggs. U.S. Fish and Wildlife Service Endangered Species Report 11.
- Spotila, J.R., A.E. Dunham, A.J. Leslie, A.C. Steyermark, P.T. Plotkin, and F.V. Paladino. 1996. Worldwide population decline of *Dermochelys coriacea*: are leatherback turtles going extinct? Chelonian Conservation and Biology 2(2):290-222.
- Steinitz, M.J., M. Salmon, and J. Wyneken. 1998. Beach renourishment and loggerhead turtle reproduction: A seven year study at Jupiter Island, Florida. Journal of Coastal Research. 14 (3) 1000-1013.
- Talbert, O.R., Jr., S.E. Stancyk, J.M. Dean, and J.M. Will. 1980. Nesting activity of the loggerhead turtle (*Caretta caretta*) in South Carolina I: a rookery in transition. Copeia 1980(4):709-718.
- Turtle Expert Working Group. 1998. An assessment of the Kemp's ridley (*Lepidochelys kempii*) and loggerhead (*Caretta caretta*) sea turtle populations in the western North Atlantic. NOAA Technical Memorandum NMFS-SEFSC-409.
- Turtle Expert Working Group. 2000. Assessment update for the Kemp's ridley and loggerhead sea turtle populations in the western North Atlantic. NOAA Technical Memorandum NMFS-SEFSC-444.
- U.S. Fish and Wildlife Service (Service). 2004. Draft Species Conservation Guidelines – South Florida – Wood Stork. U.S. Fish and Wildlife, South Florida Ecological Services, Office, Vero Beach, FL. 10pp.
- U.S. Fish and Wildlife Service (Service). 2005. Report on the Mexico/United States of America population restoration project for the Kemp's ridley sea turtle, *Lepidochelys kempii*, on the coasts of Tamaulipas and Veracruz, Mexico 2005. U.S. Fish and Wildlife Service Technical Report. 24pp.
- U.S. Fish and Wildlife Service (Service) and National Marine Fisheries Service (NOAA Fisheries). 1992. Recovery plan for the Kemp's ridley sea turtle (*Lepidochelys kempii*). National Marine Fisheries Service, St. Petersburg, FL. 40pp.
- Witherington, B.E. 1992. Behavioral responses of nesting sea turtles to artificial lighting. Herpetologica 48:31-39.
- Witherington, B.E. and K.A. Bjorndal. 1991. Influences of artificial lighting on the seaward orientation of hatchling loggerhead turtles (*Caretta caretta*). Biological Conservation 55:139-149.

- Witherington, B.E. and L.M. Ehrhart. 1989. Status and reproductive characteristics of green turtles (*Chelonia mydas*) nesting in Florida. Pages 351-352 in Ogren, L., F. Berry, K. Bjorndal, H. Kumpf, R. Mast, G. Medina, H. Reichart, and R. Witham (editors). Proceedings of the Second Western Atlantic Turtle Symposium. NOAA Technical Memorandum NMFS-SEFC-226.
- Wyneken, J., L. DeCarlo, L. Glenn, M. Salmon, D. Davidson, S. Weege., and L. Fisher. 1998. On the consequences of timing, location and fish for hatchlings leaving open beach hatcheries. Pages 155-156 in Byles, R. and Y. Fernandez (compilers). Proceedings of the Sixteenth Annual Symposium on Sea Turtle Biology and Conservation. NOAA Technical Memorandum NMFS-SEFSC-412.
- Zug, G.R. and J.F. Parham. 1996. Age and growth in leatherback turtles, *Dermochelys coriacea* (Testidines: Dermochelyidae): a skeletochronological analysis. Chelonian Conservation and Biology 2(2):244-249.

Table 1. Summary of sea turtle nesting data for Palm Beach County, Florida from 2000 to 2005.
Data accessed: http://research.myfwc.com/features/view_article.asp?id=7630 on December 12, 2006.

Year	Loggerhead Nests	Loggerhead False Crawls	Green Nests	Green False Crawls	Leatherback Nests	Leatherback False Crawls
2000	14,187	16,124	1,942	1,931	160	33
2001	13,757	12,957	175	103	334	36
2002	13,032	12,841	2,339	2,824	250	47
2003	12,963	15,050	767	846	306	50
2004	10,759	15,822	968	1,283	166	25
2005	10,791	14,345	2,252	3,142	284	52
Mean	12,581	14,523	1,407	1,688	250	41

Table 2. Summary of sea turtle nesting data (DEP monument R-13 to R-21), Palm Beach County, Florida from 2000 to 2005 (Corps).

Year	Loggerhead Nests	Loggerhead False Crawls	Green Nests	Green False Crawls	Leatherback Nests	Leatherback False Crawls
2000	1,107	1,563	61	153	12	4
2001	971	1,480	0	0	26	5
2002	675	1,221	89	186	27	13
2003	762	1,515	13	26	26	3
2004	637	1,502	41	97	20	3
2005	864	1,429	57	247	25	8
Mean	836	1,452	43	118	23	6

Aerial Location Map




Figure 1. Location of the proposed channel dredging and beach placement project along Jupiter Beach, Palm Beach County, Florida (Corps).




Figure 2. Beach lighting schematic.