




www.fws.gov/charleston www.fws.gov/southeast/candidateconservation

Conserving South Carolina's At-Risk Species:


Species facing threats to their survival

Raven's seedbox

(Ludwigia ravenii)







Illustrations from Peng, C.-I. 1984. Ludwigia Habitat top-bottom: plant, flower, and mature fruit)

Description

Raven's seedbox is a relatively newly de- occur in non-wetlands. scribed species. Superficially the plant resembles hairy primrose-willow **Status** covered with short, somewhat coarse ed. hairs. Numerous inconspicuous flowers are borne in the leaf axils. The flowers Threats have green sepals and no petals. It flowers The ditches and powerline rights-of-way Where it occurs, it does not appear to widening and paving. grow aggressively. Consequently, the populations consist of scattered individuals Management/Protection Needs pilosa and other species.

Range

eastern Virginia, eastern North Carolina, sites. southeastern South Carolina and northeastern Florida, the species has a some- References what spotty distribution. At least seven- NatureServe. 2017. NatureServe Explorteen of the nearly 30 occurrences known er: An online encyclopedia of life [web are historic; knowledge of current status application]. Version 7.1. NatureServe, of these populations (as well as additional Arlington, Virginia. Available http:// inventory of suitable habitat within the explorer.natureserve.org. range) is needed. At present, 5 to 6 occounties.

ravenii (Onagraceae), a new species from the Raven's seedbox is restricted to open, U.S. Fish & Wildlife Service Coastal Plain of the southeastern United States. wet, peaty places, such as ditches and the South Carolina Field Office Systematic Botany. 9(2): 129-132. (Pictured margins of swamps, ponds, and bogs. It 843/727-4707 almost always occurs in wetlands under april_punsalan@fws.gov

natural conditions, but may also rarely

(Ludwigia pilosa) and globefruit primrose- NatureServe's Rounded Global Status is willow (Ludwigia sphaerocarpa) but careful G1 - Critically Imperiled. Its Global Stamorphological study and field observa- tus was last reviewed on July 21, 2016. tions revealed that they differ significantly The U.S. Fish and Wildlife Service was in floral details and breeding system, petitioned to list the species in April 2010 Raven's seedbox is a branched, leafy per- and in September 2011 issued a 90-day ennial herb, usually 3.5-9 dm tall, densely finding indicating listing may be warrant-

from July-September and bears fruit from occupied by the species are subject to August-October. Seeds are light brown, many threats including herbicide use, exelliptic-oblong with slightly curved ends. cavation and deepening of ditches, road

instead of large colonies, as is typical in L. Identify and monitor existing populations of Raven's seedbox and protect from threats such as herbicide use and habitat alteration. Encourage increased survey Known from the Coastal Plain of south- efforts for the species. Protect known

currences are known to be in existence. Peng, C.-I. 1984. Ludwigia ravenii In South Carolina, the species is known (Onagraceae), a new species from the from Beaufort, Berkeley, and Hampton Coastal Plain of the southeastern United States. Systematic Botany. 9(2): 129-132.

Contact