Computation and analysis of rescued lethal gene deletions in a host-microbe model Note: This tutorial is a draft and needs completion. Contributions welcome!

Author: Almut Heinken, Molecular Systems Physiology Group, University of Luxembourg.

% Initialize the COMMA Toolbox.

r Checking if the repository is tracked using git ... Done.

r Define CB map output... set to sup. r Translate(DPG, in installed and work)

- [---] IDEAL PAIN -- set this path manually after installing the salver (see instruction - [---] ROBK PAIN -- set this path nameally after installing the salver (see instruction - [---] ROBK PAIN -- set this path nameally after installing the salver (see instruction - [----]

LP PELP OF H20F

r Nou can sobre IP problems usings "gurabi" - "Negaptes" - "glyb" - "mattab" - "deptimes" - "palas" - "quadtimes" - "fg_sobre" r Nou can sobre MCP problems usings "horobi" - "Ne cober" - "blab"

% tood the mouse reconstruction land("1552.002.not"); INSTRUMENTANCES (extree(INSTRE, 'Sineaux_es_1_os_glegio'))

% NOTE: Since dietary nutrients can also rescue many lethal gene defects, a diet

reduced better tracted for convenient of a convenient process and white-a conservate passe in the acceptance passe in the acce

made Lc={}; name TapoPtode Lc={};

land(streat(mode)Fath.epcropemode(.".ext")); model = convertEldStyleModel(model); mode Lt(2,1)-mode to boodD(1.1)-model.rxms(find(strncmp(model.rxms, 'bloomss',7))); nameTageMode to (1,1)-streat (a scrubeMode 1, '_');

b his the distinct with the same. Instability of confidence in the property of the property

All particularly addressing "9" in the small - address in the small - discourage destinating "Subpilly and its maint - address in the small - 2000000 Jupidly on "Discourage Substitution (Editorically) and its maint - address in the small - 24,200000000 Jupidlessing (2010) — 20100001010); International Computation (20100001010) and its maint - address in the small - 2010000101010 (20100001010) and its maint - address in the small - 2010000101010 (2010001010) and its maint - address in the small - 20100001010 (2010001010) and its maint - address in the small - 201000101010 (2010001010) and its maint - address in the small - 2010001010 (2010001010) and its maint - address in the small - 201000101010 (2010001010) and its maint - address in the small -

PHITTER PROVIDED - PROVIDED |
PHITTER PROVIDED | - PROVIDED |
PHITTER PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROVIDED | - PROVIDED |
EX PROVIDED | - PROV

Moving Setabolite h(k) and is model - added to the model 20022 2hk(k) = h(k) are h(k) = 2hk(k) EX. 2hk(k) = 2hk(k) are the facility function with the case same already models in the model,

Moreovery functions with the same same affirming matrix in the smint, and all M and M are size of the smint M and M and M and M are size of the smint M and M and M are size of the smint M and M and M are smint M and M and M are smint M and M and M are smint M and M and M and M are smint M and M and M and M are smint M and M and M and M are smint M and M and M and M are smint M and M and M and M are smint M and M and M and M are smint M and M and M and M are smint M and M and M and M are smint M and M and M and M are smint M and M and M and M are smint M and M and M and M are smint M and M and M are smint M and M and M and M are smint M and M and M are smint M and M and M and M are smint M and M are smint M and M are smint M and M are smint M and M and M are smint M and M and M are smint M

Morning Notificities serims([c]) and in model — admir in the and ACDMRICA Sulph — assumed [c] — and (c) — assumed (c) (c) — assum

Security Security and in some $-\sin \theta$ is the most temperature potential and $|\phi|$ and in some $-\sin \theta$ is the some $-\sin \theta$ in the some security Security $|\phi| = 1$, $|\phi| = 1$,

Moreing Medinicity $a_{11}(x_1)$ and is smooth - mine for the mass $M_{11}(x_1)$ and $A_{12}(x_1)$ an

TRAIN h(s) + h(s(s) - s(s(s) + h(s))universe Pershaville hat (s) and (s) mains t - added to the model<math>B(X(s), h(s)) + h(s(s) - s(s), h(s) + h(s(s))universe Pershaville scally not in model - added to the model B(X(s), h(s)) + adp(s) + ada(s) - ada(s) + ada(s) + h(s(s + h(s) + s)) DESize $h(k) = h2u(k) = dist_12(k) \implies sh4(k) = dist(k)$ GLENGE h(x) + h(b) = 3 $\operatorname{rad}(b) = gh_{a_{i}}h(b) \implies h(x) = 3$ $\operatorname{rad}(x) = gh_{a_{i}}h(x) = h(b)$ Marriago Relabello glogo(yb) and in model - added to the model

GPECHLER h(h) = gampshel(h) = cor h(s) = gampshel(s)GIVES h(k) = gan(k) over $h(\epsilon) = gan(\epsilon)$

MDFSSmin $12\pi(h) + 14\pi(h) \rightarrow 1(h) + \mu 1(h) + 14\mu(h)$

PRODUCTOR $h(k) = progly(k) \rightarrow h(k) = progly(k)$ SPERMIN 2 h(b) + 2 x2x(b) -+ h2x2(b) + x2(b) $\mathtt{ESLDMBE2b} \ \operatorname{hosb}[a] = \mathtt{hittenb}[b] \ \ \operatorname{vor} \ \mathtt{hittenb}[a] = \mathtt{hosb}[b]$ NEWCOLPELED h(x) = a1(x) + a2(b) + band(b) + cor band(c) + sa1(c) + h(b) + c1(b) ALANDWalled and [iv] + als [iv] + ral[iv] -- ral[iv] + als [iv] + ann [iv] $\frac{\text{ALEYMBERS } \exp_{i,k}(s) + \text{als}_{i,k}(s) + \text{all}(s) + \text{als}_{i,k}(s) + \text{als}_{i,k}(s) + \exp_{i,k}(s) }{\text{ALEXMBERS } \exp_{i,k}(s) + \text{als}_{i,k}(s) +$ ALBERTA $h(s) + ata_k L(b) + 2 \cdot aat(b) + cor 2 \cdot aat(s) + ata_k L(s) + h(b)$

Crisimation and Link = malk) = cyc_L(b) == malk) = cyc_L(c) = and_L(b) = cyc_L(c) = and_L(b) = cyc_L(c) = and_L(b) = cyc_L(c) = and_L(c) = and_

| Description | Proceedings |

quicking the remains $(SL(M)) \approx 2(1) + \ln (2) + \ln (2) + \ln (2) + \ln (2)$ $(SL(M)) \approx 2(1) + \ln (2) + \ln ($

In smally $-(aabb)_1$ we will $-(aabb)_2$ [12]. $(aabb)_1$ [12], $(aabb)_2$ [13], $(aabb)_3$ [13], $(aabb)_4$ [13], $(aabb)_4$ [14], $(aabb)_4$ [15], $(aabb)_4$ [16], $(aabb)_4$ [16], $(aabb)_4$ [17], $(aabb)_4$ [18], $(aabb)_$

PNTDSSh $h2a(x) + adp(x) + 2 \cdot ad(b) + poin_{a}B(b) \rightarrow adp(x) + b(x) + pi(x) + 2 \cdot ad(x) + poin_{a}B(x)$

Miller Studies 1 and 1 and 2 and 3 a

 $\begin{aligned} & \text{MASSIGNED}(x) = \{ \mathbf{x}_1(x), \mathbf{x}_2(x), \mathbf{x}_3(x) \} \cdot \| \mathbf{x}_{1,1}(x), \mathbf{x}_{1,2}(x) \| \mathbf{x}_{1,1}(x) \|$

splitting the resultion $BBRER h(s) + 2 \sin[s] + \cos r_s(s) = \cos 2 \sin[s] + \cos r_s(s) = \cos(s) = \sin[s] + \sin[s] + \cos s$ $\cos(s, 2s) \sin[s] + \sin[s] + \cos 2 \sin[s] + \sin[s] + \sin[s] + \sin[s]$ $\cos(s, 2s) \sin[s] + \cos[s] + \cos 2 \sin[s] + \sin[s] + \sin[s] + \sin[s]$ $\cos(s, 2s) \sin[s] + \sin[$ $\mathsf{GOMO}_{\mathsf{L}\mathsf{T}(k)} \; h(k) \; + \; \mathsf{Gigen}_{\mathsf{L}\mathsf{T}(k)} \; \; \mathsf{vor} \; h(k) \; + \; \mathsf{Gigen}_{\mathsf{L}\mathsf{T}(k)}$ different depts ((a) vor depts ((c)

SACTITITIONS is ladicisterane(b) our ladicisterane(c)

THEFTIGE abb(a) + themp(b) was themp(a) + abb(b)

```
The control of the co
```

ACMENDATE 2 h2e(s) + 2 alp(s) + 3 ach(s) - - 2 alp(s) + 2 h(s) + 2 pc(s) + 3 ach(b)

Diffield $h(k) + h2u(k) + dud_{\nu}2(k) \rightarrow sh4(k) + din(k)$

NEPhoto $h2a(k) = malp(k) \rightarrow h(k) = man(k) = adjorbp(k)$ MDFTends $h2a(k) + udp(k) \rightarrow h(k) + p1(k) + ump(k)$ NTDSeb h2u(b) = cap(b) -- pi(b) + cytd(b) NTDSeb h2u(b) = asp(b) -- adn(b) = pi(b) NTORek h2u(k) = $gap(k) \rightarrow pi(k) + gan(k)$

GAMING SZe(k) = g(ppoli(k) -- Tyr_gpo(k) = mali(k)

Recommendation of the control of the

UNISSIS h2a(k) + urea(k) + vor h2a(x) + urea(x)

sings function with the class came already models in the matel, atting the resulting the resulting the results of high | red high | red high | the high |

SHOM analysis and the "of such a smalled place of the such and a such as a s

MRMS is aat[u] = abt[h] = aat[h], were ship[u] = aat[u] = aat[u] = aat[u] whereing Fennium with the case are already exists in the mode, updating the resistion MRMTAP abt[u] = abt[u] was b[u] = abt[h]Mresings Fennium with the case case already exists in the model, updating the resistion MRMTAP abt[u] = abt[u] = abt[u] was $abt[u] + aat[u] + 2 \in b[u]$

odel - added to

```
APPERED admitted to admitted
NESSes 12a(k) + nad(k) \rightarrow h(k) + adjrib(k) + ncan(k)
ADMILPHENNELS 2 N2v(x) + 2 adp(x) + 3 advet(x) \rightarrow 2 adp(x) + 2 b(x) + 2 pa(x) + 3 advet(b)
ASSESS advalled your advalual
ALMERSKADOR gla_L[x] = ata_L[x] = nal[x] \rightarrow nal[x] = ata_L[x] = gla_L[x]
ALBERTANDO ser Lisi + ata Libi + nalibi -- nalisi + ata Lisi + ser Libi
ALETONNALS the L(x) = ata L(x) = rad(x) -= rad(x) + ata L(x) = the L(x)
```

 $ALBBER h(s) + ala_L(b) + 2 sal(b) + cor 2 sal(s) + ala_L(s) + h(b)$

AMMERIATION IN $f_{i}(s)$ + and $f_{i}(s)$ = and $f_{i}(s)$ = if $f_{i}(s)$ and an american substitution with the case case already solving in the substitution with the case case already solving in the substitution with the case case already solving in the substitution with the case case already solving in the substitution with the case case already solving in the substitution with the case case already solving in the substitution of the sub $ASSERTA \ h(x) + 2 \ mal(h) + ans_{\lambda}(h) + nor 2 \ mal(x) + ans_{\lambda}(x) + h(h)$ THEOREM and $L[n] = \min\{n\} = \lim_{n \to \infty} L[n] = \min\{n\} = \lim_{n \to \infty} L[n] = \max_{n \to \infty} L[n]$ Marriage Restine with the case case already exists in the made. CPULARATION at $a_i(x) = \max\{x\} = \exp(x/k)$ $\rightarrow \min\{x\} = \exp(x/k) = \max\{x\}$ CHROMATION gla_i(s) = mal(s) = $cys_i(s)s$ $\rightarrow mal(s)$ = $cys_i(s)$ = $gla_i(s)s$ CHINNMAD $\operatorname{thr}_{\lambda}[x] = \operatorname{rad}[k] = \operatorname{cyc}_{\lambda}[k] \longrightarrow \operatorname{rad}[x] + \operatorname{cyc}_{\lambda}[x] = \operatorname{thr}_{\lambda}[k]$

Criticalish gis $L[n] = \min\{n\} = \exp[L[n]] \Rightarrow \min\{n\} = \exp[L[n]] = gis [L[n]]$ Marriage Resction with the case case already exists in the made, CLASSWARDS $acc_{\lambda}(x) = cal(x) = gla_{\lambda}(x) \implies cal(x) + gla_{\lambda}(x) = acc_{\lambda}(x)$ $SERECLE with gia_L(x) = ser_L(x) = ser_L(x) = ser_L(x) + gia_L(x)$ CLIMBERGE Alo(s) = atp(s) = 3 g(y(s) -> adp(s) + b(s) = pc(s) = 3 g(y(b)samesomes name() = aip(s) = 3 qiy(s) == aip(s) + b(s) = pi(s) = Marriage Resition with the case case already exists in the model. GLN17_SII_rk 3 sal(k) = g(y(k) = c1(k) our 3 sal(k) = c1(k) = g(y(k))SERGLINARS sloid = ser S(k) our ser S(c) = a(c(k))

ALHERMATION $cor_{\perp}(x) = a(a_{\perp}(b) = nal(b) \rightarrow nal(c) = a(a_{\perp}(c) = cor_{\perp}(b)$ Crissmann ser List + notice + over Libt -- notice + over List + ser Libt

DESCRIPTION 2 madded + citied + the Libit -- 2 madded + citied + the Libit

SERBIB $h(s) + 2 \operatorname{mal}(b) + \operatorname{sev}_{L}(b) + \operatorname{vor}_{L}(a) + \operatorname{sev}_{L}(a) + h(b)$ ALETONNADA Uhr_L(x) + ala_L(b) + nal(b) \rightarrow nal(x) + ala_L(x) + thr_L(b) CFEDWHALLO Sir_L(c) = $\operatorname{rad}(k)$ = $\operatorname{cyc}_L(k)$ $\rightarrow \operatorname{rad}(c)$ + $\operatorname{cyc}_L(c)$ = $\operatorname{Sir}_L(k)$

and $(x_i, x_j) = (x_i, x_j)$

quality (in contains the conta

GRACHING arg.[10] = $80[1 + 90[1 + 90] + \gamma h_1] = 90[1,10] = 90[1,10]$ charing fraction with the teas and strategy scale in the small, quinting the restion spiritup for resting the strategy of the small, and the strategy continged from the strategy $h_1 = h_2 = h_3 = h_3$ SPECIA E NEUTR) + strate(h) --- E gtc_E(h) + strate(h)

COMLieb $h\lambda u(k) = g(ygnk(k)) \rightarrow g(u_{i}k(k) = g(ygnk(k))$

```
Apparent registration and arranged in a contract of the contra
```

STREE SCALE) + SHOUTE | -- SCALE) + 1pt_L(S)

PERSon account(b) = 2 fell(b) \rightarrow 2 h(b) = distant(b) = 2 fell(b)

 $ATP92 \pi h \ h2 \mu(h) \ + \ ad \mu(h) \ \rightarrow \ h(h) \ + \ am \mu(h) \ + \ \mu \lambda(h)$ NTETER $12\pi(k) = \exp(k) - - \sin(k) = \sin(k)$ A7992eb 2 h2u(b) = atp(b) -> 2 h(b) = amp(b) + 2 pi(b) NOPSked $12\pi(h) + 14\pi(h) \rightarrow 1(h) + \mu 1(h) + 14\mu(h)$ MDFTends $h2u(k) + udp(k) \rightarrow h(k) + p1(k) + ump(k)$ NTE2ek h2u(k) + ump(k) $\rightarrow p1(k) + ur1(k)$ NTDSet $12a(k) = cop(k) \rightarrow p1(k) + cyt4(k)$ NTFSek $12\pi(k)$ = q(p/k) $\rightarrow h(k)$ = p1(k) = q4p(k)ATPlet h2u(k) = g(p)(k) == h(k) = p(k) = g(p)(k)Murrians Resition with the case case already exists in the model $A7992 wh h2 u(h) + ad u(h) \rightarrow h(h) + au u(h) + \mu i(h)$

SIZECCURION $12a(x) + atp(x) + bildgless(x) \rightarrow adp(x) + b(x) + pi(x) + bildgless(b)$ ETMDet 12x(k) + biscyt(k) -- bis(k) + 1ps_1(k) $ETN(Sih h2a(x) + atp(x) + 2 \cdot nal(h) + btn(h) \rightarrow adp(x) + b(x) + pi(x) + 2 \cdot nal(x) + btn(x)$

CAPPIB $h2u(x) = adp(x) + capp(x) \rightarrow adp(x) + h(x) + px(x) + capp(b)$

COPPLA h2a(x) = adp(x) + 2hxpmp(x) -c adp(x) + b(x) + px(x) + 2hxpmp(k)DESTRUCTE $h2u(x) = atp(x) = absterol(x) \rightarrow adp(x) + b(x) = pi(x) = absterol(b)$

CLEREnch 2 for[s] = s1[b] -r s1[s] = 2 for[b] CLEREnch 2 for[s] = 2 s1[b] -r 2 s1[s] = for[b] NEWCOLUMN h(s) + st(s) + sal(b) + hosl(b) + or hosl(s) + sal(s) + h(b) + st(b) NEXTER $\operatorname{nal}(k) = h(k) = 2 \times l(k) + \operatorname{nal}(x) = 2 \times l(x) = h(x)$ CLPDRinois 2 for $(a) = a1(b) \rightarrow a1(a) = 2$ for (b) $000000 \cosh 2 \cosh(\epsilon) + \cos(k) \rightarrow \cos(\epsilon) + 2 \cosh(k)$

Officials 2 h2x(b) + shin(b) -r 3 aspec(b)

COMUNA coret(g) -r coret(k)

COMUNA careful) -r careful) COMMERCIAL COMMERCIA (b) vor commercia (c) COPTek cray_ba(c) vor cray_ba(b) OUTSTAND orisins(k) was orisins(s)

CONSTRUCT A $12\pi(k) + 5451(k) \rightarrow 4 + 54\pi(k) + 451(k)$

NTDieb $12a(b) = \exp(b) - \exp(b) + \exp(d/b)$ Nowing: Reaction with the case case already exists in the ma DESSES $h(b) = h2a(b) = dad_2(b) \longrightarrow sh4(b) = din(b)$ DESSES $h(b) = h2a(b) = dad_2(b) \longrightarrow sh4(b) = din(b)$

DOMESTICS 2 h2a(x) = 2 alp(a) = 3 dops(a) = 2 adp(a) = 2 b(a) = 2 pi(a) = 3 dops(b)

 $ESWEELELCtob \ hDu(x) = adp(x) = extriatg(x(x) - r \ adp(x) + h(x) + pi(x) = extriatg(x(b) - r \ adp(x) + h(x) + pi(x) = extriatg(x(b) - r \ adp(x) + h(x) + pi(x) = extriatg(x(b) - r \ adp(x) + h(x) = extriatg(x(b) + h(x) = extriatg(x(b$

EUTROWERZh 2 nal(b) = eutroneu(b) \rightarrow 2 nal(c) = eutroneu(c)

NETSCHO N2x(b) + 12_sis_retfa(b) -> b(b) + Eletat2(b) + retiret sis 11(b)

GAMING hOw(k) = $q(pqnh(k) \rightarrow Tyr_pqn(k) = mall(k)$ EX_adjorkp(n)k adjorkp(k) \rightarrow

NTDSet $12a(k) = cop(k) \rightarrow p1(k) + cyt4(k)$

```
Section 1. The content of the conten
```

hering; Sentian will the case name afrondy minit in the model, policing the resultion GCHEPob hCu[h] + bai[h] --> fur_i[h] + bai_mbgl[h] hering; Sentian with the case name afrondy minit in the model, policing the resultion

menings Mediakrilis Principplikografingsfogsfogsforides, belgi met in media diskel in Una media fogsforides for formatische principlikogsforides for formatische forfil) - v ROMENTE FORMATION (EUROPPLIKE UNITED BEST benegativngsforigensforides for forfil) - vor henregativngsforigensforides forfil) - vor henregativngsforigensforides forfil) - vor henregativngsforigensforides for

One mode!

Jeanneyalingsingsingstande hatelin framenyalangtoprightande hat[h] ==
CHEMIALENCEMENTERIES framenyalangtoprightande hat[h] ==
CHEMIALENCEMENTERIES framenyalangtoprightande hately ==
chemicalencement ==
chemicalencement

Annual Mediatrille framenyalangtoprightande hately ==
chemicalencement

Annual ==
chemical ==
c

_inchestpalidempingsipsication_in(e) to the incligations of the inclination _in(e) --VOCTIONALIZATION_INCLINATIONS of the inclination of the inc

Bernard, remaining trappication properties (trappication (trappication) trappication (trappication) (trappicati

__outputsenguagequates_rerer trapetatesquagequestes_rerer REGALIMENCEGERIZZENE fragetlesqtisqs(plantes_tab); wer fragetlesqtisqs(state_ta(s))

rerieq. Refabilite fragetlesqtisqs(spatjonide_tab), sat in madel
ded to the madel -

EL foregothingstangsjongstjoniske, belein brogethingstangstopstylaniske belj) — r PREDERFEREDLEGOLGEREERENDE hoppfrompt prodpostate by belje) — no foregothingstangstopstylaniske jusje skenings Heldhellie foregotjekiske jusje) ont in maket – added in the maket. Pri forestandstylaniske forestandstylaniske beljefer

updating the resition SLE(2,2k | 2 h| k) + gal(k) + one 2 h(s) + gal(s)Marsings Praction with the case case already exists in the mode updating the resition

GELSE mal(b) = gat(b) = or mal(c) = gat(c) Morning: Resition with the case mane already exists in the updating the resition

```
CARDA SADA() = SADA() = sap() = spa() = spa()
```

Mewings Relabelite galfullyaldangtagatgloside,hujb) mat in madet added in the madet Migaffullyaldangtagatgloside,hujeja galfullyaldangtagatgloside,hujb) ---

adale is tik madi.

R. grapsfilm hepsplangingstang (apstilangingstjanske, jadark, polyet for hepsplangingstangingstilangingstglanke, ja [k] ~~

ORASINETERSKERE CORRECTED SACCEST SACCEST SERVICE SACTOR before pringit for hepsplangingstangingstglanke, ja [k] ~~

DERSKERE PROFESSION CORRECTED SACCEST SACCEST SERVICE SACCEST SAC

ophiling the resultant CORLAND 2 salph) = phola(a) -> 2 sal(s) = pabela(s) CORLAND 52s(s) = sip(s) + pubula(s) -> asp(s) + b(s) = pub(s) = pubula(b)

COMELINE generally our general) Newwings Petalorite generally held not in model - added to the ma EX.generally generally or generally our generally helds COMELINE generally our generally helds

using function with the case mass already maints in the madel, similar the reaction $Pleads Na(k) = gle(k) \sim h(k) + pi(k) = gap(k)$ using function with the case mass already maints in the madel,

Within the |h| = q(p|h) - r h|h| = pi|h| = q(p|h)Newing Melabelle gloor |h| and in model — added in the model Explose r(h) gloor |h| = rGLOSSION gloor |h| = r

 $K, glastale(b) gloss[b] \sim star_{b}(s) + basyce(s)$ $MALLER gloss[b] \sim star_{b}(s) + basyce(s)$ deving Resition with the case case already exists in the signified the resition $MALLER alg_{a}(s) \sim gloss[c] \sim sg(s/b) + gloss[a]$

ophiling the resilian
(CHRLE_Z, 2) and[0] = glyb(b) = vor 2 and[o] = glyb(o)
EL_glyo(a)b glyo[b] = vor
GLYCE glyb(o) = glyb(b) = vor
GLYCE glyb(o) = vor glyb(b) = vor
GLYCE glyb(o) = vor glyb(b) = vor anne already exists in t

updating the resident UFGeb b2u(b) = $m_0 \ln |b|$ = h|b| = Hintal2|b| = glyc|b| Minimize Residential $glyc_0(b)$ and in model - added to the Hintal2[b] = $glyc_0(b)$ = $glyc_0(b)$ = $glyc_0(b$

Exploint of the give lib | -- |

Exploint field give lib | -- |

Exploint field give lib | -- |

Exploint field give lib | -- |

Noving fraction with the case case already exists in

GLECTER h(h) + gigglp(h) — nor h(s) + gigglp(s) EX_gigable(h giggb(h) — or Morning: Besilian with the case mass already mixts in the a updating the reaction

Define how(s) = glope(s) -- Tyr_ggn(s) = mail(s) develop: Section with the case case already exists in the s spining the reaction DESieb hou(s) = strength) -- six Dish = strengthing

Eighpheuleis glyleu(s) -Eighpheuleis glyphe(s) -Erghpheuleis glyphe(s) -Erwinn franklinn with the same name already minis in the s

Committee of the commit

MIFReb $12a(k) = q4p(k) \rightarrow h(k) = p1(k) = q4p(k)$

GPECHLES h(b) = gampshel(b) = cor h(s) = gampshel(s)

SPERMS 2 h(b) = 2 x2x(b) -> h2x2(b) = x2(b)

MDFSesh Nac(k) = $gip(k) \rightarrow b(k) + pi(k) + gsp(k)$ Manuago Festion with the case name already main updating the resistan NDSesh Nac(k) = $gsp(k) \rightarrow pi(k) + gan(k)$

Marriage Related the Marrial Data on model - added to the model

NOPSked $12\pi(h) + 14\pi(h) \rightarrow 1(h) + \mu 1(h) + 14\mu(h)$

LIMITATE Linewild vor Linewick $LIPStih\ h2u(s) + atp(s) + 2\ nal(h) + Lipsate(h) -- adp(s) + b(s) + pi(s) + 2\ nal(s) + Lipsate(s)$

Section 1. The control of the contro

LESTING challened[b] + point_ha[b] -- tpoint_ha[b] + satesi2_i LPOSCUS tasked halb! -- tasked halb! MADDE and $|\rangle$ = cas(c) [$|\langle L_i | ab| c \rangle$ is only = the case one strong matrix is the main', upstrong the reaction MADDE above $|\rangle$ = case $|\langle L_i | a \rangle$ = $|\langle L_i | a \rangle$ = case $|\langle L_i | a \rangle$

Expansion smalls $\gamma = 0$ through Smallmann with the case area attempt matrix in the main quantum part of the resultant quantum part of the resultant part of the part of the

any accompanies of a companies of the STOCOTOMIC and STOCOTOMIC an

densing Netherlike spatish) and in maint — mixed in the maint (Equation's spatish) = m = States spatish) = m = States spatish) = m = States spatish) = m

spating the results.

| Second | Second

And the second of the second o

noting function with the case one of roundy mainta in the mainty, since the form $|x-y| = -2 \log |y| = 2 \log$

ORDING underly (we underly)
[Explorately instances]: "
Non-inp function with the same case strong units in the underly
updating (no require (see) and see a see at easy update (see a see at easy update (see a see at easy))

ORDING underly) we underly)

ORDING underly) we underly)

Explorately underly underly and in underly added to the underly

Explorately unterly) or control or the underly underly underly underly underly underly underly underly underly).

El_magnizate(s) megosiste(s) ~ ~ OMPHANDITAN magnizate(s) ~ memprative(s) Norsing Statistite empti(s) est in endri - abbel to the s El_magnizate(s) magnit(s) ~ OMPHANDITAN magnit(s) ~ Norsing Statistite are(s) est in endri - abbel to the endr El_magnize(s) con(s) ~ OMPHANDITAN magnit(s) ~

 $(\mu_0 l, j_0 l)(h)$ $\mu_0 l, j_0 l)$ \sim which $\mu_0 l, j_0 l$ \sim which was the variant matrix in the an exact point $\mu_0 l$ \sim $\mu_0 l, j_0 l$ \sim $\mu_0 l$ \sim μ

Amendment of the property of t

ophicing the reaction PRODER high; perintribudipl; wo high; pertinductic; Norsian Returbitic pupics[b] ont in maint: albed to the maint: Expanylate(in peripola); ~ PROVING pupics[b] one peripola; International peripola; Expanylate(in peritola); on in smint: abide to the maint Expanylate(in peritola); ~ PRODERIM peritol(in); we PRODERIM peritol(in); we PRODERIM peritol(in); we

Rigardischio meritichio)

Rigardischio meritichio)

Rigardischio meritichio)

Rigardischio meritichio meritich

Within purple), \sim purple) and in maint \sim mind in the maint \sim form of the purple of the purple

EX.parieth panish. "

Within papish." "

Within papish." "

Martinia properties, and in model - added to the model

Experterior properties). "

MARTINIA properties). "

EXPLICATION properties."

EXPLICATION properties.

Whereing Papish model.

Whereing Papish."

Kino, Link) are L(h) are descript function with the case case already solits in the solet, splitting the resultion SMEATH h(h) = $\rho_{AL}(h)$ are h(a) = $\rho_{AL}(h)$ decrings function with the case case already solits in the solet, splitting the resultine SMEAH $\rho_{AL}(h)$ = $\rho_{AL}(h)$ = $\rho_{AL}(h)$ decrings function with the case case already solits in the solet, solit solet, where $\rho_{AL}(h)$ = $\rho_{AL}(h)$ = $\rho_{AL}(h)$

Mornings function with the same name already moints in the moint, updating the remains m(k) or p(x,k) and p(x) = p(x,k) are 2 and p(x) = p(x,k). When the function with the same name already moints in the moint, updating the remaint p(x) = p(x) and p(x)

Nowskap function with the case case already moints in the opinion in the resultion PRENTILLE house, a proving $\Omega(k)$ was proving $\Omega(k)$ = house, Expressing $\Omega(k)$ = proving $\Omega(k)$ = $\Omega(k)$ = house, thereing in function with the case case already moints in the opinion; the resultion PRENTILLE house $\Omega(k)$ = proving $\Omega(k)$ = house, PRENTILLE house $\Omega(k)$ = proving $\Omega(k)$ = house,

Morning Frantism with the case name already moists in the updating the resultion MORNINGTON provided (b) nor provided(s) = heal(h MORNINGTON provided(s) = nor provided(s) = heal(h MORNINGTON provided(s) = nor provided(s)

radp(x) = h(x) = pi) maining in the madel, e2(x) = heal(b) PRETERING h2u(x) = atp(x) = prestge2(x) = cadp(x) = h(x) = p2(x) = prestge2(x) Marsing: Relabelity prestgf2(x) set in model - added to the model

RESPUBLISH $12a(x) + a(p(x) + ribfle(b) \rightarrow adp(x) + b(x) + pi(x) + ribfle(b)$ EXECUTES $12a(x) = a(p(x) + r)a(r(x)) \rightarrow aap(x) + b(x) + pa(x) + r)a(r(x))$ EX_s2125c2a2aaco(a(x) s2125c2a2aaco(a(x) \rightarrow

THE Range of the state of the THEFTIGE shL[x] + themp[k] was themp[x] + shL[k] $TMD(4b \text{ nal}[b] = \text{thymd}[b] \rightarrow \text{nal}[c] = \text{thymd}[c]$ TWESTA 2 nal(k) = thyrei(k) \rightarrow 2 nal(c) = thyrei(c)

TETETHROWIGLER No. [x] = alp[x] + ininimum [x] [x] -x adp[x] = h[x] + h[x] + ininimum [x] [x]

THYPOELETIAN $pho_{i,i}(x) + i eg_{i,i}(x) \rightarrow i eg_{i,i}(x) + pho_{i,i}(x)$ Marriage Resilian with the case case already exists in the updating the resultant.

BYOLD and $[h] + i eg_{i,i}(x) \rightarrow end(x) + i eg_{i,i}(x)$

 $TREPORTATION plot ([a] + tyr_{a}([b] \rightarrow tyr_{a}([a] + plot ([b])$ MDFSeeds $h2u(k) = uip(k) \implies h(k) = pi(k) = uip(k)$ MDFTends $h2a(k) + udp(k) \rightarrow h(k) + p1(k) + ump(k)$ MDPBeak $h2a(k) + uip(k) \rightarrow h(k) + pi(k) + uip(k)$ NOPHeak $h2a(k) + uip(k) \rightarrow v(k) + pi(k) + uip(k)$ Norminus Retabolite vaca(b) not in model - added to the model

ELystell(e)s vitel(b) --

EX_text(e)& text(b) ---

```
A continue of the continue of
```

The control of the co

C+188) leaded by intl-coupletten; int Ptus (ended by int. model board, resulat reacts) named and else(1, 1), ended by int. runs(), other transfer and the first of the ended by int. runs(), other transfer and the ended by int. [modelJuint]=completentistIten(modelJuint, modelJuint, rans(strumtch("mone", modelJuint, rans)), "mone jiimus juin styn jijylo", s, w);

model2acint-changetormundcisadel2acint, 'Mouse EX a2(e)0',-288, 'L');

and things - change to the ode that the control of the control of

Implement the reduced diet.
 modelJointours@iet(modelJaint.reducedDietConstraints);

Section 1. A section of the control of the control

When the productions of rescond power, this will take that their lights body, pathod, plants Maria Jesus and their pathod in the pathod becoming framition fundamentals, and only J.T. values [MISSE], [JVM] and in many pathod in the pathod 726133.17 *19890.1*

7388347.37

A occasial for rescuing the defect is nowse DP cyclings.

[model_Apolifict.com(Company_delatedones] = delatedones(benesisadelmust.'22207.1');

modelSocial-changeExamineds(endelSulat, streat(sameTageModels(1), 'M_aty_c'), 0, 'l');

if colution.obj-m.memmemmem

fyrintf('%s \s',model3sint.rxms(microbetxchanges(1)),' is essential for recoving protic action(a.')