
emocha Health®
Support every patient
at every dose
www.emocha.com / @emochaHealth / info@emocha.com

Snapshot

Standard of Care for

450 Health Departments

Validated by over

15 peer-reviewed studies


Baltimore-based

Johns Hopkins spin out

Nurses, pharmacists, public health technologists

50-person team

emocha is the first comprehensive **Digital Medication Adherence Program**



emocha's model leverages a CDC-endorsed, public health practice to confirm dose-by-dose adherence

Defining Adherence

Possession does not equal ingestion



Medication Possession Ratio (MPR) is a standard but inadequate way to measure adherence.

VS.



emocha measures adherence by observing of every dose of medication ingested.

A significant portion of transplant patients are non-adherent to their anti-rejection medications

Non-adherence ranges from 20% to 70% across kidney, heart, and liver



40%[1]

Of rejections are linked to medication nonadherence

2/3[2]

⅔ of nonadherent patients will experience graft failure

$4-11K[3]


Hospitalization costs from rejection episodes

7x[4]

Non-adherent patients have 7x greater risk for late rejection



Post-transplant, patients face myriad challenges.



Digital Medication Adherence Program



Pharmacist-led Medication Review



3 months Daily Directly Observed Therapy

Low Possession

Concerning diagnostic measure


Hi there - I see that you didn't pick up your medication this month. Everything ok?

Ongoing Support & Medication Management

Platform & Service Overview

Patients

emocha Mobile App



Patients complete daily video check-ins to report side effects, take their medication, track progress, and engage with their adherence coach

Adherence Service

Engagement, Support, & Escalation



emocha's Clinical Adherence Team confirms daily ingestion, motivates and supports patient at every dose, and addresses any adherence challenges that arise by following our transplant-specific playbook.

Transplant Team

Analytics & Reporting



Transplant teams receive weekly and monthly reports on patient progress and adherence barriers, and – if desired – ability to view video check-ins using emocha's engagement platform.

Patient-facing Mobile App

emocha's patient app is designed to makes daily virtual check-ins feel easy and supportive. Patients open their emocha app daily to report side effects, take a video selfie of medication ingestion and engage with their adherence coach via in-app messaging and video engagement.

FEATURES

- Medication reminders
- Symptoms Capture
- Asynchronous & Live Video
- In-app messaging
- Adherence Metrics
- Financial Incentives
- Transplant resources
- Touch ID / Face ID / PIN Login
- 22 Languages



emocha's Transitional Care Team: Designed to Amplify Transplant Care Teams



Pharmacists

- Provide medication reconciliation and education at program outset
- Deliver medication summary and action plan to patient



Adherence Coaches

- Review all submitted video check-ins for medication adherence
- Compassionate, empowering, observant; support patients in overcoming every adherence barrier
- Provide linkage to existing services and resources, and maintain daily communication with patients



Registered Nurses

- Provide clinical review of all symptoms / side effects
- Liaise directly with the patient's care team regarding side effects or other concerning issues as needed

Clinical evidence generated through independent studies shows adherence rates, acceptability, and impact

Tuberculosis

92-95% Adherence

$8K Saved per patient

NIH

Partners:
Johns Hopkins, State of Maryland

Tuberculosis

87-92% Adherence

100% Treatment completion

CDC

Partners:
Puerto Rico Department of Health

Hepatitis C

98% Adherence among drug using population

CROI

Partners:
University of Colorado Anschutz Medical Campus, Gilead-supported

Asthma*

92% Patient Adoption

73% Reduction in ED visits

* Pending Publication:
Medicaid Population,
Baltimore, MD

Type 2 Diabetes*

86% Adherence

50% Reduction in ED visits

27% Diabetes-related hospital admissions post-program

* Pending Publication:
Medicaid Population,
Baltimore, MD

Holtzman SB, Zarolinski A, Shah M. Advancing Patient-Centered Care in Tuberculosis Management: A Mixed-Methods Appraisal of Video Directly Observed Therapy. JMIR 2018; Volume 5: Issue 4: e9540.

Olano Soler M, Thomas D, Joglar O, et al. Notes from the Field: Use of Asynchronous Video Directly Observed Therapy for Treatment of Tuberculosis and Latent Tuberculosis Infection in a Long-Term–Care Facility — Puerto Rico, 2016–2017. MMWR Morb Mortal Wkly Rep 2017;66:1284–1287.

Kiser, Jennifer J., Brooks, Kristina M., Castillo-Mancilla, Jose R., et al. (2020, March). Adherence in Active Drug Users with HCV: The INCLUD Trial. Poster session presented at the Conference on Retroviruses and Opportunistic Infections (CROI).

References

1 Burra et. al. Adherence in liver transplant recipients.
https://pubmed.ncbi.nlm.nih.gov/21384527/

2 Nevins et al. Understanding Medication Nonadherence after Kidney Transplant.
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5533244/

3 Pinsky et al. Transplant outcomes and economic costs associated with patient noncompliance to immunosuppression.
https://pubmed.ncbi.nlm.nih.gov/19843035/

4 Butler et al. Frequency and impact of nonadherence to immunosuppressants after renal transplantation: a systematic review.
https://pubmed.ncbi.nlm.nih.gov/15021846/


emocha Health®
Support every patient
at every dose
www.emocha.com / @emochaHealth / info@emocha.com

Transplant Opportunities & Program Value

- Minimize avoidable costs and poor outcomes due to non-adherence
- Decrease likelihood of rejection episodes, graft failure, mortality, or retransplantation
- Protect a second chance at life for transplant recipients
- Amplify and streamline care coordination post-discharge

Vision

Make technology-enabled directly observed therapy the standard of care for transplant recipients who need support


Transplant Partners


+ Johns Hopkins University
+ Florida State University
+ University of Florida
+ University of Virginia
+ University of Miami
+ National Institute of Diabetes and Digestive Kidney Diseases

Supported by the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health under Award Number R44DK133978. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Detailed patient and population-level analytics on adherence and medication challenges

75	92%	6.8%
Total Enrollment	Average adherence rate for currently active patients	Patients required escalation







Scalable messaging protocols with transplant and medication-specific context

emocha Adherence Coach



It's important to take your medication at the same time each day. Try doing this at the same time as something you do everyday (like brushing your teeth).

Sent 8:42 AM EST



John Doe

Thanks for the reminder. I think I'm getting back on track. Trying not to take meds more than 12 hours apart.

Sent 9:02 AM EST

Adherence Barrier | Motivational | Clinical Issue

Technical Issues | Protocol Issues | Adherence Streak

SUGGESTED

Hi! Keep up the great work taking your medication every day. Remember, it's also important to keep a healthy diet after your liver transplant.

Send

emocha Health®