EXERCISE 4

Exact Differential Equations

Original statement · ProblemsCh2.pdf, p. 2

Exactness test • Potential function

PROBLEM STATEMENT

Check that the following equations are exact and solve them.

(i) $(2xy - \sec^2 x) + (x^2 + 2y)\dfrac{dy}{dx} = 0,$

(ii) $(1 + e^x y + x e^x y) + (x e^x + 2)\dfrac{dy}{dx} = 0,$

(iii) $(x\cos y + \cos x)\dfrac{dy}{dx} + \sin y - y\sin x = 0,$

(iv) $e^x \sin y + y + (e^x \cos y + x + e^y)\dfrac{dy}{dx} = 0.$

Source: ProblemsCh2.pdf, Exercise 4 (page 2).

PHASE 1: CLASSIFICATION, HYPOTHESES AND DOMAIN

Each equation is written in the form

$$M(x,y) + N(x,y)\,\frac{dy}{dx} = 0, \qquad \text{equivalently} \qquad M\,dx + N\,dy = 0.$$

The equation is exact if there is a function $F(x,y)$ of class C^2 with $F_x = M$ and $F_y = N$. Along any solution $y(x)$ one then has $\frac{d}{dx}F(x,y(x)) = F_x + F_y y' = M + Ny' = 0$, so $F(x,y(x)) = C$.

Hypotheses: M, N and their first partial derivatives continuous on a rectangle (or any simply connected open set) Ω. Then (Robinson, Section 10.1, condition (10.5)) the equation is exact on Ω if and only if

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}.$$

Part	M	N	Domain used
(i)	$2xy - \sec^2 x$	$x^2 + 2y$	$x \in (-\pi/2, \pi/2)$ (or any strip between zeros of $\cos x$), $y \in \mathbb{R}$

Part	M	N	Domain used
(ii)	$1 + e^x y + x e^x y$	$x e^x + 2$	$\mathbb{R}^2$
(iii)	$\sin y - y \sin x$	$x \cos y + \cos x$	$\mathbb{R}^2$
(iv)	$e^x \sin y + y$	$e^x \cos y + x + e^y$	$\mathbb{R}^2$

PHASE 2: CHOICE OF METHOD AND CHANGE OF VARIABLES

Why this method: if the exactness criterion $M_y = N_x$ holds, a potential F exists and the solution curves are the level sets $F = C$, so no integrating factor or substitution is needed.

Construction of F, applied identically in each part:

- (a) Test: compute M_y and N_x and compare.
- (b) Integrate $F_x = M$ in x holding y fixed: $F = \int M\,dx + h(y)$, where the arbitrary function $h(y)$ plays the role of the constant of integration.
- (c) Differentiate in y and impose $F_y = N$: $h'(y) = N - \partial_y \int M\,dx$. The right side depends on y only because $M_y = N_x$.
- (d) Integrate h' and write the implicit solution $F(x, y) = C$.

PHASE 3: STEP-BY-STEP DERIVATION

Part (i): $M = 2xy - \sec^2 x,\ N = x^2 + 2y$

(a) $M_y = 2x$ and $N_x = 2x$. They coincide, so the equation is exact.

(b) $F = \int (2xy - \sec^2 x)\,dx + h(y) = x^2 y - \tan x + h(y)$ (using $\frac{d}{dx}\tan x = \sec^2 x$).

(c) $F_y = x^2 + h'(y)$ must equal $N = x^2 + 2y$, so $h'(y) = 2y$.

(d) $h(y) = y^2$ (the constant is absorbed in C). Therefore

$$F(x, y) = x^2 y - \tan x + y^2 = C.$$

Solving the quadratic $y^2 + x^2 y - (\tan x + C) = 0$ for y gives, where the discriminant is nonnegative,

$$y(x) = \frac{-x^2 \pm \sqrt{x^4 + 4(\tan x + C)}}{2}.$$

Part (ii): $M = 1 + e^x y + x e^x y,\ N = x e^x + 2$

(a) $M_y = e^x + x e^x$ and $N_x = e^x + x e^x$ (product rule on $x e^x$). They coincide: exact.

(b) $F = \int [1 + y\,(e^x + xe^x)]\,dx + h(y)$. Since $\frac{d}{dx}(xe^x) = e^x + xe^x$, this integral is $x + y\,xe^x$. So $F = x + xe^x y + h(y)$.

(c) $F_y = xe^x + h'(y)$ must equal $xe^x + 2$, so $h'(y) = 2$.

(d) $h(y) = 2y$. Therefore

$$F(x, y) = x + xe^x y + 2y = C.$$

This is linear in y, so it can be solved explicitly:

$$y(x) = \frac{C - x}{xe^x + 2}.$$

The denominator is never zero: $g(x) = xe^x$ has $g'(x) = (1 + x)e^x = 0$ only at $x = -1$, where $g(-1) = -1/e \approx -0.368$, its global minimum, so $xe^x + 2 \geq 2 - 1/e > 0$. The solutions are therefore defined on all of $\mathbb{R}$.

Part (iii): $M = \sin y - y\sin x,\ N = x\cos y + \cos x$

(a) $M_y = \cos y - \sin x$ and $N_x = \cos y - \sin x$. Exact.

(b) $F = \int (\sin y - y\sin x)\,dx + h(y) = x\sin y + y\cos x + h(y)$ (since $\int \sin x\,dx = -\cos x$, so $-y\int \sin x\,dx = y\cos x$).

(c) $F_y = x\cos y + \cos x + h'(y)$ must equal $N = x\cos y + \cos x$, so $h'(y) = 0$ and h is constant.

(d) Absorbing the constant:

$$F(x, y) = x\sin y + y\cos x = C.$$

Part (iv): $M = e^x \sin y + y,\ N = e^x\cos y + x + e^y$

(a) $M_y = e^x\cos y + 1$ and $N_x = e^x\cos y + 1$. Exact.

(b) $F = \int (e^x\sin y + y)\,dx + h(y) = e^x\sin y + xy + h(y)$.

(c) $F_y = e^x\cos y + x + h'(y)$ must equal $e^x\cos y + x + e^y$, so $h'(y) = e^y$.

(d) $h(y) = e^y$. Therefore

$$F(x, y) = e^x\sin y + xy + e^y = C.$$

PHASE 4: VERIFICATION, LIMITS AND INTERPRETATION

Check $F_x = M$ and $F_y = N$ for each potential:

- (i) $F = x^2y - \tan x + y^2$: $F_x = 2xy - \sec^2 x = M$, $F_y = x^2 + 2y = N$.
- (ii) $F = x + xe^x y + 2y$: $F_x = 1 + (e^x + xe^x)y = M$, $F_y = xe^x + 2 = N$. Also, for $y = (C - x)/(xe^x + 2)$ with $C = 1$, $x = 0$: $y(0) = 1/2$; direct insertion gives $M(0, \frac{1}{2}) + N(0, \frac{1}{2})y'(0) = \frac{3}{2} + 2y'(0)$, and $y'(0) = -\frac{3}{4}$ from the quotient rule, so the sum is 0.
- (iii) $F = x \sin y + y \cos x$: $F_x = \sin y - y \sin x = M$, $F_y = x \cos y + \cos x = N$.
- (iv) $F = e^x \sin y + xy + e^y$: $F_x = e^x \sin y + y = M$, $F_y = e^x \cos y + x + e^y = N$.

Caveat on the level sets: $F(x, y) = C$ defines y as a function of x only where $F_y = N \neq 0$ (implicit function theorem). At points where $N = 0$ the level curve can have a vertical tangent and cannot be written as $y(x)$; for (ii) this never happens on the solution branch because $N = xe^x + 2 > 0$.

RESULTS

Part	Exactness	Solution
(i)	$M_y = N_x = 2x$	$x^2y - \tan x + y^2 = C$
(ii)	$M_y = N_x = e^x(1 + x)$	$x + xe^x y + 2y = C$, i.e. $y = \frac{C - x}{xe^x + 2}$
(iii)	$M_y = N_x = \cos y - \sin x$	$x \sin y + y \cos x = C$
(iv)	$M_y = N_x = e^x \cos y + 1$	$e^x \sin y + xy + e^y = C$