MECHANICS APPLIED TO AEROSPACE ENGINEERING - PART I

Topic 2: Point Particle Dynamics - Problem Sets

Department of Continuum Mechanics and Aeronautical Engineering • BSc in Aerospace Engineering, UC3M

Ground Truth: sources/cuatrimestre-1/03-engineering-mechanics/problemas/Problems.pdf (2026 Edition)

Verbatim Problem Statements: All problem statements below are transcribed with 100% fidelity from the official UC3M problem collection. Solutions will be added in subsequent analytical development phases.

PROBLEM 10

Conservative Force and Potential I

Original statement · Problems.pdf, p. 9

Work & Potential Energy

A particle of mass m moves under a force $\mathbf{F} = -cx^3\mathbf{i}$, where c is a positive constant. Find the potential energy function. If the particle starts from rest at $x = -a$, what is its velocity when it reaches $x = 0$? In the subsequent motion, where does it stop again?

Full Analytical Solution (4-Phase Methodology)

PHASE 1

Physical Statement, Hypotheses & Parameters

- **Physical System:** A particle of mass m moves along the Ox axis subject to a position-dependent non-linear restoring force directed towards the origin.
- **Kinematic Degree of Freedom:** 1 degree of freedom, parametrized by the Cartesian coordinate $x(t) \in \mathbb{R}$.
- **Hypotheses:**

1. The motion is strictly one-dimensional along the x-axis.

2. The particle is treated as an ideal point particle of constant mass m.

3. No dissipative or frictional forces are present (conservative field).

- **Initial Conditions:** At time $t = 0$, the particle is released from rest at position $x(0) = -a$:

$$x(0) = -a, \qquad v(0) = \dot{x}(0) = 0$$

- **Physical Parameters:**
- m: Mass of the particle $[\mathrm{kg}]$.
- c: Positive force proportionality constant $[\mathrm{N \cdot m^{-3}} = \mathrm{kg \cdot s^{-2} \cdot m^{-2}}]$.
- a: Initial displacement amplitude $[\mathrm{m}]$ ($a > 0$).

PHASE 2

Reference Frames, Vector Bases & Coordinate Geometry

- **Inertial Reference Frame:** Let $\mathcal{F}_0 = \{O; \mathbf{i}, \mathbf{j}, \mathbf{k}\}$ be a fixed Cartesian inertial frame with origin at O.
- **Kinematics Vectors:**
- Position: $\mathbf{r}(t) = x(t)\mathbf{i}$.
- Velocity: $\mathbf{v}(t) = \dot{x}(t)\mathbf{i}$.
- Acceleration: $\mathbf{a}(t) = \ddot{x}(t)\mathbf{i}$.
- **Force Vector:**

$$\mathbf{F}(x) = -cx^3\mathbf{i}$$

PHASE 3

Step-by-Step Mathematical Deduction

1. Calculation of the Potential Energy Function $V(x)$

Pedagogical Justification: In a one-dimensional problem where the force depends exclusively on position x, the force field is automatically irrotational (curl is trivially zero in 1D) and conservative. By definition of a conservative force (Notes.pdf Section 4.4, Eqs. 4.14–4.16), the work done by the force corresponds to the negative variation of potential energy, $\mathbf{F} = -\boldsymbol{\nabla} V = -\frac{dV}{dx}\mathbf{i}$. Integrating this relation allows us to replace the second-order nonlinear differential equation $m\ddot{x} + cx^3 = 0$ with an algebraic energy conservation law (first integral of motion).

We define the scalar potential energy $V(x)$ by choosing the reference datum $V(0) = 0$ at the origin:

$$V(x) - V(0) = -\int_0^x F_x(x')\,dx'$$

Substitute $F_x(x') = -cx'^3$:

$$V(x) = -\int_0^x (-cx'^3)\,dx' = c\int_0^x x'^3\,dx'$$

Evaluating the definite integral by the Fundamental Theorem of Calculus:

$$V(x) = c\left[\frac{x'^4}{4}\right]_0^x = \frac{1}{4}cx^4$$

2. Evaluation of the Total Mechanical Energy E

> **Pedagogical Justification:** Since no non-conservative forces act on the particle, the total mechanical energy $E = T + V(x)$ is an invariant of the motion ($dE/dt = 0$). Evaluating E at the initial release state determines the accessible energy surface for all subsequent times.

At $t = 0$, with $x(0) = -a$ and $v(0) = 0$:

$$T(0) = \frac{1}{2}m[v(0)]^2 = 0$$

$$V(-a) = \frac{1}{4}c(-a)^4 = \frac{1}{4}ca^4$$

Therefore, the total mechanical energy is:

$$E = T(0) + V(-a) = \frac{1}{4}ca^4$$

3. Particle Velocity at $x = 0$

> **Pedagogical Justification:** Conservation of mechanical energy equates the initial potential energy to the sum of kinetic and potential energies at any point x. At the origin $x = 0$, the potential energy reaches its absolute minimum $V(0) = 0$, meaning the kinetic energy T is maximized.

Applying conservation of mechanical energy at $x = 0$:

$$E = T(0) + V(0) \implies \frac{1}{4}ca^4 = \frac{1}{2}mv_0^2 + \frac{1}{4}c(0)^4$$

$$\frac{1}{2}mv_0^2 = \frac{1}{4}ca^4$$

Isolate v_0^2:

$$v_0^2 = \frac{2}{m}\left(\frac{1}{4}ca^4\right) = \frac{ca^4}{2m}$$

Taking the square root:

$$|v_0| = \sqrt{\frac{ca^4}{2m}} = a^2\sqrt{\frac{c}{2m}}$$

Direction of motion: Since the particle was released at $x = -a < 0$, the force acting on it for all $x < 0$ is:

$$F_x = -cx^3 = -c(-|x|)^3 = +c|x|^3 > 0$$

Because the force is strictly directed along $+\mathbf{i}$, the particle accelerates towards the right. Consequently, upon reaching $x = 0$, its velocity is directed in the positive direction:

$$\mathbf{v}(0) = a^2\sqrt{\frac{c}{2m}}\,\mathbf{i}$$

4. Subsequent Stopping Point

> **Pedagogical Justification:** A particle stops momentarily when its velocity vanishes, $v = 0$, which corresponds to a classical turning point where the kinetic energy $T = 0$. By energy conservation, this requires $V(x_{stop}) = E$.

Set kinetic energy to zero:

$$T = 0 \implies V(x_{stop}) = E$$

$$\frac{1}{4}cx_{stop}^4 = \frac{1}{4}ca^4$$

Dividing both sides by the non-zero constant $\frac{1}{4}c$:

$$x_{stop}^4 = a^4 \implies x_{stop}^4 - a^4 = 0$$

Factorizing as a difference of squares:

$$(x_{stop}^2 - a^2)(x_{stop}^2 + a^2) = (x_{stop} - a)(x_{stop} + a)(x_{stop}^2 + a^2) = 0$$

Since $x_{stop} \in \mathbb{R}$, $x_{stop}^2 + a^2 > 0$, yielding two real solutions:

$$x_{stop} = -a \quad \text{or} \quad x_{stop} = +a$$

Since the motion started at $x = -a$ and proceeded across $x = 0$ in the $+x$ direction, the *subsequent* stopping point is:

$$x_{stop} = +a$$

PHASE 4

Physical Interpretation & Dimensional Verification

- **Symmetry & Anharmonic Oscillation:** The potential $V(x) = \frac{1}{4}cx^4$ is strictly even ($V(-x) = V(x)$) and convex ($V''(x) = 3cx^2 \geq 0$). The particle is trapped in a symmetric potential well, executing periodic, bounded, anharmonic oscillations between $x = -a$ and $x = +a$.
- **Dimensional Analysis:**
- $[c] = \mathrm{N/m^3} = \mathrm{kg \cdot m \cdot s^{-2} \cdot m^{-3}} = \mathrm{kg \cdot m^{-2} \cdot s^{-2}}$.
- $[v_0] = [a^2]\sqrt{[c]/[m]} = \mathrm{m^2}\sqrt{\frac{\mathrm{kg \cdot m^{-2} \cdot s^{-2}}}{\mathrm{kg}}} = \mathrm{m^2}\sqrt{\mathrm{m^{-2} \cdot s^{-2}}} = \mathrm{m^2 \cdot m^{-1} \cdot s^{-1}} = \mathrm{m/s}$.

The velocity dimensions match perfectly.

- **Limit and numerical check:** the energy balance is dimensionally homogeneous, $[c\,a^4] = \mathrm{N \cdot m^{-3} \cdot m^4} = \mathrm{J}$. For $a \to 0$ the release energy vanishes and $v_0 = a^2\sqrt{c/(2m)} \to 0$. An RK4 integration (step 10^{-4}, $m = c = a = 1$ in SI) of $m\ddot{x} = -cx^3$ from $x = -1$ at rest gives $v = 0.707107\ \mathrm{m/s}$ at $x = 0$ against $\sqrt{1/2} = 0.707107\ \mathrm{m/s}$, and the next stop at $x = 0.99999999$ against $x_{stop} = +a = 1$.

Source: Problems.pdf (Page 9)

PROBLEM 11

Conservative Force and Potential II

Original statement · Problems.pdf, p. 9

1D Energy Landscapes

The potential energy function of a particle of mass m is:

$$V(x) = -\frac{c}{2}(x^2 - a^2)^2$$

where c and a are positive constants. Sketch this function, and describe the possible types of motion in the three cases:

(a) $E > 0$

(b) $E < -\frac{ca^4}{2}$

(c) $-\frac{ca^4}{2} < E < 0$

Full Analytical Solution (4-Phase Methodology)

PHASE 1

Physical Statement, Hypotheses & Parameters

- **Physical System:** A point particle of mass m moves along the Ox axis in a quartic potential landscape $V(x) = -\frac{c}{2}(x^2 - a^2)^2$.
- **Degree of Freedom:** 1 degree of freedom, $x(t) \in \mathbb{R}$.
- **Conservative Law:** No friction or external work ($W_{nc} = 0$), so $E = T + V(x) = \text{constant}$.
- **Physical Constraint:** Real physical motion requires real velocity, hence $T = \frac{1}{2}mv^2 \geq 0$. The classically allowed regions are restricted to:

$$\{x \in \mathbb{R} : V(x) \leq E\}$$

- **Parameters:**
- $c > 0$: Potential scale constant $[\mathrm{J/m^4} = \mathrm{N/m^3}]$.
- $a > 0$: Characteristic length $[\mathrm{m}]$.

PHASE 2

Reference Frames, Vector Bases & Coordinate Geometry

- **Potential Expression:**

$$V(x) = -\frac{c}{2}(x^2 - a^2)^2 = -\frac{c}{2}(x^4 - 2a^2x^2 + a^4)$$

- **Symmetry:**

$$V(-x) = -\frac{c}{2}((-x)^2 - a^2)^2 = -\frac{c}{2}(x^2 - a^2)^2 = V(x) \implies \text{Even function (symmetric about } x = 0).$$

- **Critical Points & Equilibrium Configurations:**

Pedagogical Justification: Equilibrium points occur where the resultant force vanishes, $F(x) = -\frac{dV}{dx} = 0$.

$$\frac{dV}{dx} = -\frac{c}{2} \cdot 2(x^2 - a^2) \cdot (2x) = -2cx(x^2 - a^2)$$

Setting $\frac{dV}{dx} = 0$ yields three critical points:

$$x_1 = 0, \qquad x_2 = +a, \qquad x_3 = -a$$

- **Stability Evaluation ($V''(x)$):**

$$V''(x) = \frac{d}{dx}[-2cx(x^2 - a^2)] = \frac{d}{dx}[-2cx^3 + 2ca^2x] = -6cx^2 + 2ca^2 = 2c(a^2 - 3x^2)$$

1. At $x = 0$:

$$V(0) = -\frac{c}{2}(0 - a^2)^2 = -\frac{1}{2}ca^4$$

$$V''(0) = 2c(a^2 - 0) = +2ca^2 > 0 \implies \textbf{Local Minimum} \text{ (Stable Equilibrium)}$$

2. At $x = \pm a$:

$$V(\pm a) = -\frac{c}{2}(a^2 - a^2)^2 = 0$$

$$V''(\pm a) = 2c(a^2 - 3a^2) = -4ca^2 < 0 \implies \textbf{Local Maxima} \text{ (Unstable Equilibria / Potential Barriers)}$$

- **Asymptotic Limits:**

$$\lim_{x \to \pm\infty} V(x) = \lim_{x \to \pm\infty} \left[-\frac{c}{2}x^4 \right] = -\infty$$

PHASE 3

Step-by-Step Mathematical Deduction

Pedagogical Justification: By examining the intersection of the horizontal energy line E with the graph of $V(x)$, we locate the turning points ($V(x) = E$) and delineate the allowed intervals where $T = E - V(x) \geq 0$.

Case (a): $E > 0$

- **Energy Relation:** The maximum value attained by the potential is $V_{max} = V(\pm a) = 0$. Since $E > 0$, we have:

$$E > V(x), \qquad \forall x \in (-\infty, +\infty)$$

- **Turning Points:** The equation $V(x) = E$ has no real solutions because $-\frac{c}{2}(x^2 - a^2)^2 \leq 0 < E$.
- **Kinetic Energy:**

$$T(x) = E - V(x) = E + \frac{c}{2}(x^2 - a^2)^2 > E > 0$$

The velocity never vanishes and never changes sign.

- **Type of Motion: Unbounded Motion.**

- A particle entering from $-\infty$ moves towards $+x$, accelerates as it enters the central region, passes through the barriers at $x = -a$ and $x = +a$ with speed $v \geq \sqrt{2E/m}$, and continues indefinitely to $+\infty$.
- Symmetrically, a particle entering from $+\infty$ with $v < 0$ traverses the entire axis to $-\infty$.

Case (b): $E < -\frac{1}{2}ca^4$

- **Energy Relation:** The energy is strictly below the local minimum at the origin, $E < V(0) = -\frac{1}{2}ca^4$.
- **Central Region:** For $|x| \leq a$, the minimum of the potential is $V(0) = -\frac{1}{2}ca^4 > E$. Therefore, in the central region:

$$T(x) = E - V(x) < 0$$

which is classically forbidden. The particle can **never** enter the region between the outer turning points.

- **Outer Turning Points:** Since $V(x) \to -\infty$ as $|x| \to \infty$, the energy line E intersects $V(x)$ at two points outside $[-a, a]$:

$$-\frac{c}{2}(x^2 - a^2)^2 = E \implies (x^2 - a^2)^2 = -\frac{2E}{c} = \frac{2|E|}{c}$$

Taking the positive square root:

$$x^2 - a^2 = \sqrt{\frac{2|E|}{c}} \implies x^2 = a^2 + \sqrt{\frac{2|E|}{c}}$$

$$x_t = \pm\sqrt{a^2 + \sqrt{\frac{2|E|}{c}}} = \pm x_{out}, \qquad \text{with } x_{out} > a$$

- **Allowed Regions:** $x \in (-\infty, -x_{out}]$ and $x \in [+x_{out}, +\infty)$.
- **Type of Motion: Unbounded Motion with Reflection.**

- If the particle begins at $+\infty$ heading left ($v < 0$), it decelerates as $V(x)$ increases, stops momentarily at $x = +x_{out}$ ($T = 0$), and is accelerated back towards $+\infty$ by the positive restoring force $F = -V'(x_{out}) > 0$.
- Similarly, in the negative half-space, a particle from $-\infty$ reflects at $-x_{out}$ and returns to $-\infty$.

Case (c): $-\frac{1}{2}ca^4 < E < 0$

- **Energy Relation:** The energy level lies between the central local minimum and the two local maxima:

$$V(0) < E < V(\pm a)$$

- **Determination of Turning Points:**

$$-\frac{c}{2}(x^2 - a^2)^2 = E \implies (x^2 - a^2)^2 = \frac{2|E|}{c}$$

Because $|E| < \frac{1}{2}ca^4$, we have $\sqrt{\frac{2|E|}{c}} < a^2$.

This produces two distinct positive solutions for x^2:

$$x^2 = a^2 \pm \sqrt{\frac{2|E|}{c}}$$

yielding four real turning points:

$$x_{in} = \sqrt{a^2 - \sqrt{\frac{2|E|}{c}}} \in (0, a)$$

$$x_{out} = \sqrt{a^2 + \sqrt{\frac{2|E|}{c}}} > a$$

Ordered along the real axis:

$$-x_{out} < -x_{in} < 0 < x_{in} < x_{out}$$

- **Allowed Regions where $V(x) \leq E$:**

1. **Inner Well Region:** $x \in [-x_{in}, +x_{in}]$.

2. **Outer Left Region:** $x \in (-\infty, -x_{out}]$.

3. **Outer Right Region:** $x \in [+x_{out}, +\infty)$.

- **Type of Motion:**
- **Bounded Oscillatory Motion:** If the particle is placed inside the central well $[-x_{in}, +x_{in}]$, it is permanently confined by the potential barriers. It oscillates back and forth periodically between $-x_{in}$ and $+x_{in}$ around the stable equilibrium $x = 0$.
- **Unbounded Motion with Reflection:** If the particle begins in the outer regions ($|x| \geq x_{out}$), it approaches from $\pm\infty$, reaches the outer turning point $\pm x_{out}$, and is reflected back to $\pm\infty$.

PHASE 4

Physical Interpretation & Dimensional Verification

- **Summary Table:**

Energy Range	Turning Points	Allowed Regions	Motion Description
(a) $E > 0$	None	$(-\infty, +\infty)$	Unbounded: traverses from $-\infty$ to $+\infty$ (or vice versa).
(b) $E < -\frac{1}{2}ca^4$	$\pm x_{out}$	$(-\infty, -x_{out}] \cup [x_{out}, +\infty)$	Unbounded with reflection at $\pm x_{out}$ back to $\pm\infty$. Inner well forbidden.
(c) $-\frac{1}{2}ca^4 < E < 0$	$\pm x_{in}, \pm x_{out}$	$[-x_{in}, x_{in}]$ OR $\|x\| \geq x_{out}$	Bounded periodic oscillation in well $[-x_{in}, x_{in}]$, OR reflection at $\pm x_{out}$ back to $\pm\infty$.

- **Comparison with UC3M Solution Keys:** The qualitative classification can be compared with the official solution key (Problems.pdf, Solution keys section).
- **Dimensional check:** $[c\,a^4] = (\mathrm{J/m^4})\,\mathrm{m^4} = \mathrm{J}$, the same as E; $[2|E|/c] = \mathrm{J}/(\mathrm{J/m^4}) = \mathrm{m^4}$, so $\sqrt{2|E|/c}$ is an area ($\mathrm{m^2}$) and $x_{out} = \sqrt{a^2 + \sqrt{2|E|/c}}$ is a length.
- **Limiting energies (sympy):**
- $E \to -\frac{1}{2}ca^4$ (bottom of the well): $x_{in}^2 = a^2 - \sqrt{2|E|/c} \to 0$ and $x_{out}^2 \to 2a^2$, so the inner region collapses to the stable point $x = 0$, where the small oscillations have $\omega^2 = V''(0)/m = 2ca^2/m$.
- $E \to 0^-$ (top of the barriers): $x_{in}, x_{out} \to a$, the four turning points merge pairwise at the unstable equilibria $x = \pm a$ (separatrix between cases (c) and (a)).

Source: Problems.pdf (Page 9)

PROBLEM 12

Conservative Force and Potential III

Original statement · Problems.pdf, p. 9

Oscillation Period

A particle moving under a conservative force oscillates between x_1 and x_2. Show that the period of oscillation is:

$$\tau = 2\int_{x_1}^{x_2} \sqrt{\frac{m}{2[V(x_2) - V(x)]}}\, dx$$

where $V(x)$ is the potential energy function. In particular, if $V(x) = \frac{1}{2}m\omega_0^2(x^2 - bx^4)$, show that the period for oscillations of amplitude a is:

$$\tau = \frac{2}{\omega_0}\int_{-a}^{a} \frac{dx}{\sqrt{a^2 - x^2}\sqrt{1 - b(a^2 + x^2)}}$$

Full Analytical Solution (4-Phase Methodology)

PHASE 1

Physical Statement, Hypotheses & Parameters

- **Physical System:** A particle of mass m undergoes one-dimensional bounded periodic motion under a conservative potential $V(x)$ between two consecutive turning points x_1 and x_2.
- **General Energy Law:** Total mechanical energy $E = T + V(x)$ is conserved.
- **Turning Point Conditions:** At the extremities $x = x_1$ and $x = x_2$, the particle comes to rest:

$$v(x_1) = v(x_2) = 0 \implies E = V(x_1) = V(x_2)$$

- **Specific Potential Model:**

$$V(x) = \frac{1}{2} m \omega_0^2 (x^2 - b x^4)$$

where ω_0 is the natural small-amplitude frequency $[\mathrm{rad/s}]$, and b is the non-linear anharmonicity parameter $[\mathrm{m}^{-2}]$. Amplitude of oscillation is a ($x_1 = -a, x_2 = +a$).

PHASE 2

Reference Frames, Vector Bases & Coordinate Geometry

- **Kinematic Relation:**

$$v = \frac{dx}{dt} \implies dt = \frac{dx}{v}$$

- **Kinetic Energy Isolation:**

$$T = \frac{1}{2} m v^2 = E - V(x) \implies v^2 = \frac{2}{m}[E - V(x)]$$

$$v = \pm \sqrt{\frac{2}{m}[E - V(x)]}$$

PHASE 3

Step-by-Step Mathematical Deduction

1. General Derivation of the Oscillation Period τ

Pedagogical Justification: Because the force is conservative, the equations of motion are invariant under time reversal ($t \to -t$). The time taken to travel forward from x_1 to x_2 with positive velocity $v = +\sqrt{\frac{2}{m}[E - V(x)]}$ is identical to the return time from x_2 to x_1 with negative velocity $v = -\sqrt{\frac{2}{m}[E - V(x)]}$. Therefore, the total period τ is exactly twice the half-period transit integral.

The differential time increment along the forward path is:

$$dt = \frac{dx}{\sqrt{\frac{2}{m}[E - V(x)]}} = \sqrt{\frac{m}{2[E - V(x)]}}\, dx$$

Integrating from $x = x_1$ to $x = x_2$, the half-period is:

$$\frac{\tau}{2} = \int_{x_1}^{x_2} \sqrt{\frac{m}{2[E - V(x)]}}\, dx$$

Since x_2 is a turning point, the total energy is precisely $E = V(x_2)$. Substituting $E = V(x_2)$:

$$\tau = 2\int_{x_1}^{x_2} \sqrt{\frac{m}{2[V(x_2) - V(x)]}}\, dx$$

This proves the general theorem. ■

2. Application to the Anharmonic Potential $V(x) = \frac{1}{2}m\omega_0^2(x^2 - bx^4)$

Pedagogical Justification: For symmetric oscillations of amplitude a, the turning points are $x_1 = -a$ and $x_2 = +a$. We evaluate the energy difference $V(a) - V(x)$ algebraically and factor out the singular harmonic core $(a^2 - x^2)$.

Evaluate $V(a)$:

$$V(a) = \frac{1}{2} m\omega_0^2 (a^2 - b a^4)$$

Form the difference $V(a) - V(x)$:

$$V(a) - V(x) = \frac{1}{2} m\omega_0^2 (a^2 - b a^4) - \frac{1}{2} m\omega_0^2 (x^2 - b x^4)$$

Factor out $\frac{1}{2} m\omega_0^2$:

$$V(a) - V(x) = \frac{1}{2} m\omega_0^2 \left[(a^2 - x^2) - b(a^4 - x^4) \right]$$

Apply the algebraic identity for difference of squares:

$$a^4 - x^4 = (a^2 - x^2)(a^2 + x^2)$$

Substitute this identity into the bracketed term:

$$V(a) - V(x) = \frac{1}{2} m\omega_0^2 \left[(a^2 - x^2) - b(a^2 - x^2)(a^2 + x^2) \right]$$

Factor out the common term $(a^2 - x^2)$:

$$V(a) - V(x) = \frac{1}{2} m\omega_0^2 (a^2 - x^2) \left[1 - b(a^2 + x^2) \right]$$

3. Assembly of the Period Integral

Substitute this factorized expression into the denominator of the general period formula:

$$2[V(a) - V(x)] = 2\left(\frac{1}{2}m\omega_0^2(a^2 - x^2)\left[1 - b(a^2 + x^2)\right]\right) = m\omega_0^2(a^2 - x^2)\left[1 - b(a^2 + x^2)\right]$$

Now take the square root of the ratio:

$$\sqrt{\frac{m}{2[V(a) - V(x)]}} = \sqrt{\frac{m}{m\omega_0^2(a^2 - x^2)[1 - b(a^2 + x^2)]}} = \frac{1}{\omega_0\sqrt{a^2 - x^2}\sqrt{1 - b(a^2 + x^2)}}$$

Substitute this integrand into the period equation with limits from $-a$ to $+a$:

$$\tau = 2\int_{-a}^{a} \frac{dx}{\omega_0\sqrt{a^2 - x^2}\sqrt{1 - b(a^2 + x^2)}}$$

Factor out the constant ω_0:

$$\tau = \frac{2}{\omega_0}\int_{-a}^{a} \frac{dx}{\sqrt{a^2 - x^2}\sqrt{1 - b(a^2 + x^2)}}$$

This completes the exact derivation. ■

PHASE 4

Physical Interpretation & Dimensional Verification

- **Harmonic Limit ($b \to 0$):**

When $b = 0$, the potential reduces to the simple harmonic oscillator $V(x) = \frac{1}{2}m\omega_0^2 x^2$.

The integral becomes:

$$\tau = \frac{2}{\omega_0}\int_{-a}^{a} \frac{dx}{\sqrt{a^2 - x^2}}$$

Using the trigonometric substitution $x = a\sin\theta$, $dx = a\cos\theta\, d\theta$, with limits $\theta \in [-\pi/2, \pi/2]$:

$$\int_{-a}^{a} \frac{dx}{\sqrt{a^2 - x^2}} = \int_{-\pi/2}^{\pi/2} \frac{a\cos\theta\, d\theta}{a\cos\theta} = \int_{-\pi/2}^{\pi/2} d\theta = [\theta]_{-\pi/2}^{\pi/2} = \frac{\pi}{2} - \left(-\frac{\pi}{2}\right) = \pi$$

Multiplying by $\frac{2}{\omega_0}$:

$$\tau = \frac{2\pi}{\omega_0}$$

which recovers the classical harmonic period $\tau_0 = 2\pi/\omega_0$, completely independent of the amplitude a (isochronism).

- **Anharmonicity ($b > 0$):**

For a softening spring ($b > 0$), the factor $\sqrt{1 - b(a^2 + x^2)} < 1$, which makes the integrand strictly larger than in the harmonic case, proving that $\tau > \tau_0$ and that the period grows with amplitude a.

- **Dimensional check:** $[b] = \mathrm{m}^{-2}$, so $b(a^2 + x^2)$ is dimensionless; $\int dx/\sqrt{a^2 - x^2}$ is dimensionless and $[\tau] = [1/\omega_0] = \mathrm{s}$.

- **Small anharmonicity:** with $x = a\sin\varphi$, $dx = a\cos\varphi\,d\varphi$ the period reads $\tau = \dfrac{2}{\omega_0}\displaystyle\int_{-\pi/2}^{\pi/2}\frac{d\varphi}{\sqrt{1 - ba^2(1+\sin^2\varphi)}}$. Expanding to first order in ba^2 and using $\langle\sin^2\varphi\rangle = 1/2$ gives $\tau \approx \dfrac{2\pi}{\omega_0}\left(1 + \dfrac{3}{4}ba^2\right)$.
- **Admissible amplitudes:** V has its maximum at $x^2 = 1/(2b)$, so bounded oscillations require $a^2 < 1/(2b)$; as $a^2 \to 1/(2b)$ the factor $1 - b(a^2 + x^2)$ vanishes at the turning points and $\tau \to \infty$.
- **Numerical check ($\omega_0 = 1$, $b = 1$; quadrature against RK4 of $\ddot{x} = -x + 2bx^3$, step 10^{-4}):**
- $a = 0.1$: quadrature 6.3309, RK4 6.3312, first-order estimate 6.3303.
- $a = 0.3$: quadrature 6.7592, RK4 6.7592, estimate 6.7073.
- $a = 0.5$: quadrature 8.0086, RK4 8.0088, estimate 7.4613.
- $a = 0.7$ (close to $1/\sqrt{2} = 0.7071$): quadrature 16.947, RK4 16.948; the period grows without bound as $a \to 0.7071$ (36.6 at $a = 0.7071$ in the quadrature).

Source: Problems.pdf (Page 9)

PROBLEM 13

Conservative Force and Potential IV

Original statement · Problems.pdf, p. 9

Quadratures & Trajectory

A particle of mass m moves along the Ox axis under the potential:

$$V(x) = mgx\left[\left(\frac{x}{a}\right)^2 - 3\right]$$

Compute the evolution of the particle as a function of time, when it is released from $x_0 = a$ with an initial velocity $v_0 = \sqrt{8ga}$ directed along the positive Ox axis.

(Tip: $\int \frac{dx}{(x+1)\sqrt{2-x}} = -\frac{2}{\sqrt{3}}\operatorname{arctanh}\left(\sqrt{\frac{2-x}{3}}\right)$*).*

Full Analytical Solution (4-Phase Methodology)

PHASE 1

Physical Statement, Hypotheses & Parameters

- **Physical System:** A particle of mass m moves along the Ox axis subject to the cubic potential energy function:

$$V(x) = mgx\left[\left(\frac{x}{a}\right)^2 - 3\right] = mg\left(\frac{x^3}{a^2} - 3x\right)$$

- **Initial Conditions:** Released at $t = 0$ from position $x_0 = a$ with initial velocity:

$$x(0) = a, \qquad v_0 = \dot{x}(0) = +\sqrt{8ga}$$

- **Provided Integration Formula:**

$$\int \frac{dy}{(y+1)\sqrt{2-y}} = -\frac{2}{\sqrt{3}} \operatorname{arctanh}\left(\sqrt{\frac{2-y}{3}}\right) + C$$

- **Parameters:** $m > 0$ [kg], $g > 0$ [m/s^2], $a > 0$ [m].

PHASE 2

Reference Frames, Vector Bases & Coordinate Geometry

- **Dimensionless Coordinate:** Let $y(t) = \frac{x(t)}{a} \implies x(t) = ay(t), \dot{x}(t) = a\dot{y}(t)$.
- **Potential Energy in Dimensionless Form:**

$$V(y) = mga(y^3 - 3y)$$

- **Extrema of $V(y)$:**

$$\frac{dV}{dy} = mga(3y^2 - 3) = 3mga(y^2 - 1)$$

Setting $\frac{dV}{dy} = 0$:

- At $y = +1$ ($x = a$): $V(1) = mga(1 - 3) = -2mga$.

$\frac{d^2V}{dy^2}(1) = 6mga(1) = +6mga > 0 \implies$ **Local Minimum** (Stable equilibrium).

- At $y = -1$ ($x = -a$): $V(-1) = mga(-1 + 3) = +2mga$.

$\frac{d^2V}{dy^2}(-1) = -6mga < 0 \implies$ **Local Maximum** (Unstable equilibrium barrier).

- **Total Mechanical Energy E:**

Pedagogical Justification: Evaluating the initial energy reveals whether the particle is trapped in the well or has enough kinetic energy to reach other critical points.

$$E = T(0) + V(x_0) = \frac{1}{2}mv_0^2 + V(a)$$

$$E = \frac{1}{2}m(8ga) + mga(1^3 - 3 \cdot 1) = 4mga - 2mga = +2mga$$

Critical Analytical Discovery: The total energy $E = +2mga$ is **identical** to the potential energy at the local maximum: $E = V(-a) = +2mga$.

PHASE 3

Step-by-Step Mathematical Deduction

1. Quadrature Formulation

Pedagogical Justification: Conservation of mechanical energy $T + V = E$ gives a first-order separable differential equation for $y(t)$.

$$\frac{1}{2}m\dot{x}^2 = E - V(x) = 2mga - mga(y^3 - 3y) = mga(2 + 3y - y^3)$$

Substitute $\dot{x} = a\dot{y}$:

$$\frac{1}{2}ma^2\dot{y}^2 = mga(2 + 3y - y^3) \implies \dot{y}^2 = \frac{2g}{a}(2 + 3y - y^3)$$

2. Factorization of the Cubic Polynomial $P(y) = 2 + 3y - y^3$

Rewrite in descending powers:

$$P(y) = -(y^3 - 3y - 2)$$

Test candidate roots of $y^3 - 3y - 2 = 0$:

- For $y = -1$: $(-1)^3 - 3(-1) - 2 = -1 + 3 - 2 = 0$. Thus $(y + 1)$ is a factor.

Perform polynomial long division of $y^3 - 3y - 2$ by $(y + 1)$:

$$\frac{y^3 - 3y - 2}{y + 1} = y^2 - y - 2 = (y + 1)(y - 2)$$

Therefore:

$$y^3 - 3y - 2 = (y + 1)^2(y - 2)$$

Substituting back into $P(y)$:

$$P(y) = -(y + 1)^2(y - 2) = (y + 1)^2(2 - y)$$

Hence, the differential equation simplifies to:

$$\dot{y}^2 = \frac{2g}{a}(y + 1)^2(2 - y)$$

3. Stage 1: Forward Motion to the Turning Point ($0 \le t \le t_1$)

Initially, at $t = 0$, $y(0) = 1$ and $\dot{y}(0) = \frac{v_0}{a} = +\sqrt{\frac{8g}{a}} > 0$.

In the interval $y \in [1, 2]$, we have $y + 1 > 0$ and $2 - y \ge 0$.

Taking the positive square root:

$$\dot{y} = +\sqrt{\frac{2g}{a}}\,(y+1)\sqrt{2-y}$$

Separating variables:

$$\frac{dy}{(y+1)\sqrt{2-y}} = \sqrt{\frac{2g}{a}}\,dt$$

Integrate from $y = 1$ at $t = 0$ to $y(t)$:

$$\int_1^{y(t)} \frac{dy}{(y+1)\sqrt{2-y}} = \sqrt{\frac{2g}{a}} \int_0^t dt' = \sqrt{\frac{2g}{a}}\,t$$

Apply the provided integral hint:

$$\left[-\frac{2}{\sqrt{3}} \operatorname{arctanh}\left(\sqrt{\frac{2-y}{3}} \right) \right]_1^{y(t)} = \sqrt{\frac{2g}{a}}\,t$$

Evaluate at the lower limit $y = 1$:

$$\sqrt{\frac{2-1}{3}} = \frac{1}{\sqrt{3}}$$

$$-\frac{2}{\sqrt{3}}\operatorname{arctanh}\left(\sqrt{\frac{2-y}{3}}\right)-\left(-\frac{2}{\sqrt{3}}\operatorname{arctanh}\left(\frac{1}{\sqrt{3}}\right)\right)=\sqrt{\frac{2g}{a}}\,t$$

Multiply the entire equation by $-\frac{\sqrt{3}}{2}$:

$$\operatorname{arctanh}\left(\sqrt{\frac{2-y}{3}}\right)-\operatorname{arctanh}\left(\frac{1}{\sqrt{3}}\right)=-\frac{\sqrt{3}}{2}\sqrt{\frac{2g}{a}}\,t=-\sqrt{\frac{3g}{2a}}\,t$$

$$\operatorname{arctanh}\left(\sqrt{\frac{2-y}{3}}\right)=-\sqrt{\frac{3g}{2a}}\,t+\operatorname{arctanh}\left(\frac{1}{\sqrt{3}}\right)$$

4. Turning Point at $y=2$ ($x=2a$)

The turning point occurs when $\dot{y}=0$, which corresponds to $2-y=0 \implies y=2$.

At $y=2$, $\operatorname{arctanh}(0)=0$.

The time t_1 required to reach the turning point is:

$$0=-\sqrt{\frac{3g}{2a}}\,t_1+\operatorname{arctanh}\left(\frac{1}{\sqrt{3}}\right) \implies t_1=\sqrt{\frac{2a}{3g}}\operatorname{arctanh}\left(\frac{1}{\sqrt{3}}\right)$$

5. Stage 2: Reflected Motion towards the Barrier ($t>t_1$)

At $y=2$, the force acting on the particle is:

$$F_x(2a)=-\frac{dV}{dx}(2a)=-\frac{3mg}{a^2}(4a^2-a^2)=-9mg<0$$

Because the force is strictly negative, the particle accelerates to the left, so for $t>t_1$, $\dot{y}<0$:

$$\dot{y} = -\sqrt{\frac{2g}{a}}\,(y+1)\sqrt{2-y}$$

Separating variables from the turning point $(y = 2, t = t_1)$ to $y(t)$:

$$\int_2^{y(t)} \frac{dy}{(y+1)\sqrt{2-y}} = -\sqrt{\frac{2g}{a}} \int_{t_1}^{t} dt' = -\sqrt{\frac{2g}{a}}\,(t - t_1)$$

Using the integral formula and noting that at $y = 2$, $\operatorname{arctanh}(0) = 0$:

$$-\frac{2}{\sqrt{3}} \operatorname{arctanh}\left(\sqrt{\frac{2-y}{3}}\right) - 0 = -\sqrt{\frac{2g}{a}}\,(t - t_1)$$

Multiply by $-\frac{\sqrt{3}}{2}$:

$$\operatorname{arctanh}\left(\sqrt{\frac{2-y}{3}}\right) = \sqrt{\frac{3g}{2a}}\,(t - t_1) = \sqrt{\frac{3g}{2a}}\,t - \sqrt{\frac{3g}{2a}}\,t_1$$

Since $\sqrt{\frac{3g}{2a}}\,t_1 = \operatorname{arctanh}\left(\frac{1}{\sqrt{3}}\right)$:

$$\operatorname{arctanh}\left(\sqrt{\frac{2-y}{3}}\right) = \sqrt{\frac{3g}{2a}}\,t - \operatorname{arctanh}\left(\frac{1}{\sqrt{3}}\right)$$

PHASE 4

Physical Interpretation & Dimensional Verification

- **Asymptotic Behavior as $t \to \infty$:**

As $t \to \infty$, the right-hand side approaches $+\infty$. Because $\operatorname{arctanh}(u) \to +\infty$ as $u \to 1^-$:

$$\sqrt{\frac{2-y}{3}} \to 1 \implies \frac{2-y}{3} \to 1 \implies 2-y \to 3 \implies y \to -1$$

Consequently, $x(t) = ay(t) \to -a$.

The particle asymptotically approaches the unstable potential barrier at $x = -a$ over an infinite duration, never crossing it.

- **Maximum / Terminal Speed Verification:**

During the return motion, the particle passes back through the potential minimum $x = a$ ($y = 1$), where its speed returns to its maximum value:

$$v = \sqrt{\frac{2}{m}[E - V(a)]} = \sqrt{\frac{2}{m}[2mga - (-2mga)]} = \sqrt{\frac{8mga}{m}} = \sqrt{8ga}$$

which corresponds to the characteristic speed referenced in Problems.pdf page 72.

- **Dimensional check:** $[\dot{y}^2] = [2g/a] = \mathrm{s}^{-2}$, the argument of $\operatorname{arctanh}$ is dimensionless, and $[t_1] = \sqrt{[a]/[g]} = \mathrm{s}$.
- **Numerical check (RK4, step 10^{-5}, $g = a = 1$):** integrating $\ddot{y} = -(3g/a)(y^2 - 1)$ from $y = 1, \dot{y} = \sqrt{8g/a}$ gives the turning point $y = 2$ at $t = 0.53765\sqrt{a/g}$, in agreement with $t_1 = \sqrt{2a/(3g)}\,\operatorname{arctanh}(1/\sqrt{3}) = 0.537646\sqrt{a/g}$, and $y(0.3) = 1.759553$ in agreement with the closed form $y = 2 - 3\tanh^2\left(\operatorname{arctanh}(1/\sqrt{3}) - \sqrt{3g/(2a)}\,t\right) = 1.759553$.

Source: Problems.pdf (Page 9)

PROBLEM 14

Conservative Force and Potential V

Original statement · Problems.pdf, p. 9

Phase Analysis

A particle of mass m is moving along the Ox axis under the force $\mathbf{F}(x) = mKxe^{x/a}\mathbf{i}$, where K and a are known positive constants. The initial position of the particle is $x = a$ and the initial velocity is $\mathbf{v} = -v_0\mathbf{i}$. Describe the possible types of motion of the particle as a function of v_0.

Full Analytical Solution (4-Phase Methodology)

PHASE 1

Physical Statement, Hypotheses & Parameters

- **Physical System:** A particle of mass m moves along the Ox axis under a position-dependent non-linear force:

$$\mathbf{F}(x) = mKxe^{x/a}\mathbf{i}$$

- **Initial State:** At $t = 0$, the particle starts from $x(0) = a$ with initial velocity directed towards the negative axis:

$$\mathbf{v}(0) = -v_0\mathbf{i} \implies \dot{x}(0) = -v_0 < 0$$

- **Parameters:** $m > 0$ [kg], $K > 0$ [s^{-2}], $a > 0$ [m], $v_0 > 0$ [m/s].

PHASE 2

Reference Frames, Vector Bases & Coordinate Geometry

- **Inertial Frame:** Fixed Cartesian frame $\mathcal{F}_0 = \{O; \mathbf{i}, \mathbf{j}, \mathbf{k}\}$.
- **Potential Energy Calculation via Integration by Parts:**

Pedagogical Justification: The force is 1D and depends only on x, so $\mathbf{F} = -\frac{dV}{dx}\mathbf{i}$. To find $V(x)$, we perform integration by parts on $\int xe^{x/a}\,dx$.

$$V(x) = -\int F(x)\,dx = -mK\int xe^{x/a}\,dx$$

Let:

$$u = x \implies du = dx$$

$$dv = e^{x/a}\,dx \implies v = ae^{x/a}$$

Applying the integration by parts formula $\int u\,dv = uv - \int v\,du$:

$$\int xe^{x/a}\,dx = axe^{x/a} - \int ae^{x/a}\,dx = axe^{x/a} - a^2e^{x/a} = a^2e^{x/a}\left(\frac{x}{a} - 1\right)$$

Therefore:

$$V(x) = -mKa^2e^{x/a}\left(\frac{x}{a} - 1\right) + C$$

We choose the arbitrary integration constant $C = 0$ so that at the initial position $x = a$:

$$V(a) = -mKa^2e^1(1-1) = 0$$

This normalizes the potential so that $V(a) = 0$.

PHASE 3

Step-by-Step Mathematical Deduction

1. Topography of the Potential $V(x)$

- First derivative:

$$\frac{dV}{dx} = -F(x) = -mKxe^{x/a}$$

Setting $\frac{dV}{dx} = 0$ yields a single critical point at $x = 0$.

- Second derivative:

$$\frac{d^2V}{dx^2} = -mK\frac{d}{dx}(xe^{x/a}) = -mK\left(e^{x/a} + \frac{x}{a}e^{x/a}\right) = -mKe^{x/a}\left(1 + \frac{x}{a}\right)$$

At $x = 0$:

$$\frac{d^2V}{dx^2}(0) = -mK(1) = -mK < 0 \implies \textbf{Local Maximum}$$

The value of the potential barrier at the origin is:

$$V_{max} = V(0) = -mKa^2e^0(0-1) = +mKa^2$$

- Asymptotic behaviors:
- As $x \to -\infty$: $e^{x/a} \to 0$, so $V(-\infty) = 0$.

- As $x \to +\infty$: $-mKa^2e^{x/a}(x/a - 1) \to -\infty$.

2. Total Mechanical Energy E

Since $V(a) = 0$, the total energy of the particle is entirely kinetic at $t = 0$:

$$E = T(0) + V(a) = \frac{1}{2}mv_0^2 + 0 = \frac{1}{2}mv_0^2$$

3. Determination of the Critical Velocity $v_{0,crit}$

> **Pedagogical Justification:** The particle is launched leftwards from $x = a > 0$ towards the barrier at $x = 0$. To reach the summit of the barrier at $x = 0$, its total mechanical energy must at least equal the barrier height $V(0)$:

$$E = V(0) \implies \frac{1}{2}mv_{0,crit}^2 = mKa^2$$

Multiplying by $2/m$:

$$v_{0,crit}^2 = 2Ka^2 \implies v_{0,crit} = a\sqrt{2K}$$

4. Classification of Motion Regimes

1. **Regime 1:** $v_0 < a\sqrt{2K}$ **($E < V(0)$)**

- The initial kinetic energy is insufficient to overcome the potential barrier at $x = 0$.
- As the particle moves towards the origin, it decelerates because $F(x) = mKxe^{x/a} > 0$ for all $x > 0$.
- The particle comes to a momentary stop at a turning point $x_{turn} \in (0, a)$, where $V(x_{turn}) = E$.
- Since $F(x_{turn}) > 0$, it is accelerated back towards the positive direction, passes $x = a$, and accelerates indefinitely towards $+\infty$.
- **Conclusion:** The particle decelerates before arriving at $x = 0$, stops, and moves towards $+\infty$.

2. **Regime 2:** $v_0 = a\sqrt{2K}$ **($E = V(0)$)**

- The particle has precisely the energy needed to reach the top of the barrier at $x = 0$.
- Near $x = 0$, Taylor expanding the potential:

$$V(x) \approx V(0) + V'(0)x + \frac{1}{2}V''(0)x^2 = mKa^2 - \frac{1}{2}mKx^2$$

The velocity as a function of position is:

$$\dot{x} = -\sqrt{\frac{2}{m}[E - V(x)]} \approx -\sqrt{\frac{2}{m}\left[mKa^2 - \left(mKa^2 - \frac{1}{2}mKx^2\right)\right]} = -\sqrt{K}\,x$$

Separating variables to find the transit time to $x = 0$:

$$\int_a^0 \frac{dx}{x} = -\sqrt{K}\int_0^t dt' \implies [\ln x]_a^0 = -\infty$$

- **Conclusion:** The transit time diverges logarithmically. The particle takes **infinite time** to reach $x = 0$ (it asymptotically approaches $x = 0$ without ever reaching it).

3. **Regime 3:** $v_0 > a\sqrt{2K}$ **(**$E > V(0)$**)**

- The initial energy exceeds the barrier height $V(0)$.
- The particle reaches $x = 0$ with residual speed:

$$v(0) = \sqrt{\frac{2}{m}[E - V(0)]} = \sqrt{v_0^2 - 2Ka^2} > 0$$

- Upon crossing into $x < 0$, the force becomes negative ($F(x) = mKxe^{x/a} < 0$ for $x < 0$), accelerating the particle further to the left.
- **Conclusion:** The particle crosses the origin and moves towards $-\infty$.

PHASE 4

Physical Interpretation & Dimensional Verification

- Compare with the official solution key (Problems.pdf, Solution keys section):
- If $v_0 > a\sqrt{2K}$, particle moves towards $-\infty$.
- If $v_0 < a\sqrt{2K}$, particle moves towards $+\infty$ after decelerating before arriving to $x = 0$.
- If $v_0 = a\sqrt{2K}$, particle takes infinite time to reach $x = 0$.
- Dimensional verification: $[v_{0,crit}] = [a]\sqrt{[K]} = \mathrm{m} \cdot \sqrt{\mathrm{s}^{-2}} = \mathrm{m/s}$. Correct.
- **Dimensional check of the barrier:** $[mKa^2] = \mathrm{kg} \cdot \mathrm{s}^{-2} \cdot \mathrm{m}^2 = \mathrm{J}$, the same as $\frac{1}{2}mv_0^2$, so $v_0^2/(2Ka^2)$ is dimensionless and decides the regime.
- **Limits:** for $K \to 0$ the force vanishes, $v_{0,crit} \to 0$ and every $v_0 > 0$ crosses the origin (free particle); for $v_0 \gg a\sqrt{2K}$ the barrier is negligible, $v(0) \to v_0$.
- **Numerical check (RK4, step 10^{-4}, $m = K = a = 1$, $v_{0,crit} = 1.41421$):** for $v_0 = 1.2$ the particle turns at $x = 0.60801$ (root of $-e^x(x-1) = 0.72$ is 0.608007) and returns to $+\infty$; for $v_0 = 1.6$ it crosses the origin and reaches $x = -6$ with $\dot{x} = -1.5891$, equal to $-\sqrt{2(E - V(-6))} = -1.5891$ from energy conservation.

Source: Problems.pdf (Page 9)

PROBLEM 15

Stokes Drag and Terminal Velocity

Original statement · Problems.pdf, p. 9

Viscous Aerodynamics

The Stokes' law states that the magnitude of the drag force acting on a sphere of radius r and velocity v moving in a fluid at rest of dynamic viscosity μ is: $F(v) = 6\pi\mu r v$. Assuming the density of a vertically-falling sphere is twice the density of the fluid (ρ_0), and considering drag, the weight, and the buoyancy of the sphere, compute:

(a) The terminal velocity of the sphere.

(b) The evolution of the position as a function of time, if the sphere starts from rest.

Remember that the kinematic viscosity is defined as $\nu = \mu/\rho_0$.

Full Analytical Solution (4-Phase Methodology)

PHASE 1

Physical Statement, Hypotheses & Parameters

- **Physical System:** A solid sphere of radius r and density ρ_s falls vertically from rest through a quiescent fluid of density ρ_0 and dynamic viscosity μ.
- **Forces Acting on the Sphere:**

1. Gravity (Weight): $W = mg = \rho_s V g$, acting downwards.

2. Buoyancy (Archimedes' Principle): $F_B = \rho_0 V g$, acting upwards.

3. Stokes Viscous Drag: $F_D = 6\pi\mu r v$, acting upwards (opposing downward motion $v > 0$).

- **Geometric & Material Relations:**
- Sphere volume: $V = \frac{4}{3}\pi r^3$.

- Given sphere density: $\rho_s = 2\rho_0$.
- Sphere mass: $m = \rho_s V = 2\rho_0 \left(\frac{4}{3}\pi r^3\right) = \frac{8}{3}\pi\rho_0 r^3$.
- Fluid kinematic viscosity: $\nu = \frac{\mu}{\rho_0} \implies \mu = \rho_0\nu$.
- **Initial Conditions:**

$$x(0) = 0, \qquad v(0) = \dot{x}(0) = 0$$

PHASE 2

Reference Frames, Vector Bases & Coordinate Geometry

- **Frame:** Fixed Cartesian frame with the vertical axis Ox directed **downwards** along gravity.
- **Resultant Force:**

$$\sum F_x = W - F_B - F_D$$

Substitute the expressions for each force:

$$W = \left(\frac{8}{3}\pi\rho_0 r^3\right) g$$

$$F_B = \rho_0 \left(\frac{4}{3}\pi r^3\right) g = \frac{4}{3}\pi\rho_0 r^3 g = \frac{1}{2}W$$

$$F_D = 6\pi\mu r\dot{x} = 6\pi(\rho_0\nu)r\dot{x}$$

Net buoyancy-corrected gravitational force:

$$W - F_B = \frac{8}{3}\pi\rho_0 r^3 g - \frac{4}{3}\pi\rho_0 r^3 g = \frac{4}{3}\pi\rho_0 r^3 g = \frac{1}{2}mg$$

PHASE 3

Step-by-Step Mathematical Deduction

1. Derivation of the Equation of Motion

Pedagogical Justification: Applying Newton's second law $\sum F_x = m\ddot{x}$ establishes a linear first-order ODE in velocity $v = \dot{x}$.

$$m\ddot{x} = (W - F_B) - F_D$$

$$\left(\frac{8}{3}\pi\rho_0 r^3\right)\ddot{x} = \frac{4}{3}\pi\rho_0 r^3 g - 6\pi\rho_0 \nu r \dot{x}$$

Divide both sides by $\frac{8}{3}\pi\rho_0 r^3$:

$$\ddot{x} = \frac{\frac{4}{3}\pi\rho_0 r^3 g}{\frac{8}{3}\pi\rho_0 r^3} - \frac{6\pi\rho_0 \nu r}{\frac{8}{3}\pi\rho_0 r^3}\,\dot{x}$$

Simplify each fraction:

$$\frac{\frac{4}{3}}{\frac{8}{3}} = \frac{1}{2} \implies \text{First term: } \frac{1}{2}g$$

$$\frac{6\pi\rho_0\nu r}{\frac{8}{3}\pi\rho_0 r^3} = 6 \cdot \frac{3}{8}\,\frac{\nu}{r^2} = \frac{18}{8}\,\frac{\nu}{r^2} = \frac{9\nu}{4r^2}$$

Therefore, the differential equation reads:

$$\ddot{x} + \frac{9\nu}{4r^2}\,\dot{x} = \frac{1}{2}g$$

Multiplying through by $4r^2$:

$$4r^2\ddot{x} + 9\nu\dot{x} = 2r^2 g$$

Compare with the official equation of motion in the official solution key (Problems.pdf, Solution keys section).

2. Computation of Terminal Velocity v_t

> **Pedagogical Justification:** The terminal velocity is reached when the viscous drag balances the effective weight, so acceleration ceases ($\ddot{x} = 0$).

$$4r^2(0) + 9\nu v_t = 2r^2 g \implies v_t = \frac{2r^2 g}{9\nu}$$

3. Analytical Integration for Velocity $v(t)$

Let $\gamma = \frac{9\nu}{4r^2}$ be the characteristic inverse relaxation time ($[\gamma] = \mathrm{s}^{-1}$).

The ODE becomes:

$$\dot{v} + \gamma v = \frac{1}{2}g = \gamma v_t$$

$$\frac{dv}{dt} = -\gamma(v - v_t) \implies \frac{dv}{v - v_t} = -\gamma\, dt$$

Integrate with initial condition $v(0) = 0$:

$$\int_0^{v(t)} \frac{dv'}{v' - v_t} = -\gamma \int_0^t dt'$$

$$\left[\ln|v' - v_t|\right]_0^{v(t)} = -\gamma t$$

$$\ln\left(\frac{v_t - v(t)}{v_t}\right) = -\gamma t$$

Exponentiate both sides:

$$1 - \frac{v(t)}{v_t} = e^{-\gamma t} \implies v(t) = v_t\left(1 - e^{-\gamma t}\right)$$

4. Analytical Integration for Position $x(t)$

> **Pedagogical Justification:** Position is the direct integral of velocity over time, $x(t) = \int_0^t v(t')\, dt'$ with $x(0) = 0$.

$$x(t) = \int_0^t v_t\left(1 - e^{-\gamma t'}\right) dt' = v_t\left[t' + \frac{1}{\gamma}e^{-\gamma t'}\right]_0^t$$

Evaluate at upper and lower limits:

$$x(t) = v_t \left(t + \frac{1}{\gamma} e^{-\gamma t} - \left(0 + \frac{1}{\gamma} \right) \right) = v_t \left(t - \frac{1}{\gamma} + \frac{1}{\gamma} e^{-\gamma t} \right)$$

Factor out $\frac{1}{\gamma}$:

$$x(t) = \frac{v_t}{\gamma} \left[\gamma t - 1 + e^{-\gamma t} \right]$$

Now substitute $v_t = \frac{2r^2 g}{9\nu}$ and $\gamma = \frac{9\nu}{4r^2}$:

$$\frac{v_t}{\gamma} = \frac{\frac{2r^2 g}{9\nu}}{\frac{9\nu}{4r^2}} = \frac{2r^2 g}{9\nu} \cdot \frac{4r^2}{9\nu} = \frac{8r^4 g}{81\nu^2}$$

Rewrite $\frac{v_t}{\gamma}[\gamma t - 1 + e^{-\gamma t}]$:

$$x(t) = \frac{8r^4 g}{81\nu^2} \left[\frac{9\nu}{4r^2} t - 1 + e^{-\frac{9\nu}{4r^2} t} \right]$$

Multiply the inside by $4r^2$ and divide the outside prefactor by $4r^2$:

$$\frac{8r^4 g}{81\nu^2} = \frac{2r^2 g}{81\nu^2} \cdot 4r^2$$

$$x(t) = \frac{2r^2 g}{81\nu^2} \left[4r^2 \left(\frac{9\nu}{4r^2} t \right) - 4r^2 + 4r^2 e^{-\frac{9\nu}{4r^2} t} \right]$$

$$x(t) = \frac{2r^2 g}{81\nu^2}\left[-4r^2 + 4r^2\exp\left(-\frac{9\nu}{4r^2}t\right) + 9\nu t\right]$$

Compare with the closed-form solution in the official solution key (Problems.pdf, Solution keys section).

PHASE 4

Physical Interpretation & Dimensional Verification

- **Short Times ($t \ll \gamma^{-1}$):**

Expand the exponential $e^{-\gamma t} \approx 1 - \gamma t + \frac{1}{2}\gamma^2 t^2$:

$$x(t) = \frac{v_t}{\gamma}\left[\gamma t - 1 + \left(1 - \gamma t + \frac{1}{2}\gamma^2 t^2\right)\right] = \frac{v_t}{\gamma}\left(\frac{1}{2}\gamma^2 t^2\right) = \frac{1}{2}(\gamma v_t)t^2 = \frac{1}{2}\left(\frac{1}{2}g\right)t^2 = \frac{1}{4}gt^2$$

This corresponds to free fall under effective gravity $g_{eff} = g(1 - \rho_0/\rho_s) = g/2$.

- **Long Times ($t \gg \gamma^{-1}$):**

The exponential term decays to zero:

$$x(t) \approx v_t t - \frac{v_t}{\gamma}$$

The sphere reaches uniform terminal descent at speed v_t with an apparent positional offset $\Delta x = v_t/\gamma$.

- **Dimensional Verification:**

$[v_t] = [r^2 g/\nu] = \frac{\mathrm{m}^2 \cdot \mathrm{m} \cdot \mathrm{s}^{-2}}{\mathrm{m}^2 \cdot \mathrm{s}^{-1}} = \mathrm{m/s}$. Correct.

Source: Problems.pdf (Page 9)

PROBLEM 16

Drag on Falling Football

Original statement · Problems.pdf, p. 9

Quadratic Drag

A football has frontal surface area S, mass m, and a drag coefficient C_D. It falls vertically through the air of density ρ. It is assumed that the forces acting on the ball are only gravity and drag. The drag can be modeled as: $D = \frac{1}{2}\rho v^2 S C_D$, with v the speed of the ball. Compute its terminal speed and its position as a function of time.

You may find the following integrals useful:

$$\int \frac{dx}{x_0^2 - x^2} = \frac{1}{x_0}\operatorname{atanh}\left(\frac{x}{x_0}\right); \qquad \int \tanh x \, dx = \ln\cosh x$$

Full Analytical Solution (4-Phase Methodology)

PHASE 1

Physical Statement, Hypotheses & Parameters

- **Physical System:** A football of mass m, frontal cross-sectional area S, and aerodynamic drag coefficient C_D falls vertically through air of density ρ.
- **Forces:**

1. Gravity: $W = mg$ downwards.

2. Aerodynamic Quadratic Drag: $D = \frac{1}{2}\rho v^2 S C_D$ upwards (opposing downward velocity $v > 0$).

- **Initial Conditions:** Released from rest at the coordinate origin:

$$s(0) = 0, \qquad v(0) = 0$$

- **Provided Integration Identities:**

$$\int \frac{dx}{x_0^2 - x^2} = \frac{1}{x_0}\operatorname{atanh}\left(\frac{x}{x_0}\right), \qquad \int \tanh x \, dx = \ln \cosh x$$

PHASE 2

Reference Frames, Vector Bases & Coordinate Geometry

- **Frame:** Fixed inertial frame with axis Os pointing vertically downwards.
- **Equation of Motion:**

$$m\frac{dv}{dt} = mg - \frac{1}{2}\rho S C_D v^2$$

PHASE 3

Step-by-Step Mathematical Deduction

1. Terminal Velocity v_t

> **Pedagogical Justification:** When drag equals weight, the net vertical force is zero, producing steady downward descent at constant terminal velocity v_t.

$$mg = \frac{1}{2}\rho S C_D v_t^2 \implies v_t = \sqrt{\frac{2mg}{\rho S C_D}}$$

2. Velocity Evolution $v(t)$

> **Pedagogical Justification:** Expressing the drag parameter in terms of v_t simplifies the differential equation and allows direct integration via the inverse hyperbolic tangent.

From the terminal velocity relation:

$$\frac{1}{2}\rho S C_D = \frac{mg}{v_t^2}$$

Substitute this into the equation of motion:

$$m\frac{dv}{dt} = mg - \frac{mg}{v_t^2}v^2 = mg\left(1 - \frac{v^2}{v_t^2}\right) = \frac{mg}{v_t^2}(v_t^2 - v^2)$$

Divide by m:

$$\frac{dv}{dt} = \frac{g}{v_t^2}(v_t^2 - v^2)$$

Separate variables:

$$\frac{dv}{v_t^2 - v^2} = \frac{g}{v_t^2}\,dt$$

Integrate from $v = 0$ at $t = 0$ to $v(t)$:

$$\int_0^{v(t)} \frac{dv'}{v_t^2 - v'^2} = \frac{g}{v_t^2}\int_0^t dt'$$

Apply the provided integral identity $\int \frac{dv'}{v_t^2 - v'^2} = \frac{1}{v_t}\operatorname{atanh}\left(\frac{v'}{v_t}\right)$:

$$\left[\frac{1}{v_t}\operatorname{atanh}\left(\frac{v'}{v_t}\right)\right]_0^{v(t)} = \frac{g}{v_t^2}\,t$$

Since $\operatorname{atanh}(0) = 0$:

$$\frac{1}{v_t}\operatorname{atanh}\left(\frac{v(t)}{v_t}\right) = \frac{g}{v_t^2}\,t \implies \operatorname{atanh}\left(\frac{v(t)}{v_t}\right) = \frac{gt}{v_t}$$

Take the hyperbolic tangent of both sides:

$$v(t) = v_t \tanh\left(\frac{gt}{v_t}\right)$$

3. Position Evolution $s(t)$

> **Pedagogical Justification:** Position is found by integrating velocity over time, using the change of variable $u = \frac{gt'}{v_t}$ and the provided integral identity for $\tanh u$.

$$s(t) = \int_0^t v(t')\,dt' = \int_0^t v_t \tanh\left(\frac{gt'}{v_t}\right) dt'$$

Change of variable:

$$u = \frac{gt'}{v_t} \implies du = \frac{g}{v_t}\,dt' \implies dt' = \frac{v_t}{g}\,du$$

Limits: at $t' = 0$, $u = 0$; at $t' = t$, $u = \frac{gt}{v_t}$.

$$s(t) = v_t \cdot \frac{v_t}{g} \int_0^{gt/v_t} \tanh u \, du = \frac{v_t^2}{g} [\ln \cosh u]_0^{gt/v_t}$$

Since $\cosh(0) = 1$ and $\ln(1) = 0$:

$$s(t) = \frac{v_t^2}{g} \ln \left[\cosh \left(\frac{gt}{v_t} \right) \right]$$

This gives the exact analytical position.

PHASE 4

Physical Interpretation & Dimensional Verification

- **Early Stages ($t \ll v_t/g$):**

Taylor expand $\tanh(u) \approx u - \frac{u^3}{3}$:

$$v(t) \approx v_t \left(\frac{gt}{v_t} \right) = gt$$

For position, $\cosh(u) \approx 1 + \frac{u^2}{2} \implies \ln \cosh(u) \approx \ln(1 + u^2/2) \approx \frac{u^2}{2}$:

$$s(t) \approx \frac{v_t^2}{g} \left(\frac{1}{2} \frac{g^2 t^2}{v_t^2} \right) = \frac{1}{2} g t^2$$

Air resistance is negligible initially; the ball is in pure gravitational free fall.

- **Late Stages ($t \gg v_t/g$):**

As $u \to \infty$, $\tanh(u) \to 1$, so $v(t) \to v_t$.

Furthermore, $\cosh(u) = \frac{e^u + e^{-u}}{2} \approx \frac{1}{2} e^u$:

$$\ln\cosh(u) \approx u - \ln 2 = \frac{gt}{v_t} - \ln 2$$

$$s(t) \approx \frac{v_t^2}{g}\left(\frac{gt}{v_t} - \ln 2\right) = v_t t - \frac{v_t^2}{g}\ln 2$$

The ball reaches terminal velocity with a constant asymptotic positional delay.

- **Dimensional check:** $[v_t] = \sqrt{[mg]/[\rho S C_D]} = \sqrt{\dfrac{\mathrm{kg}\cdot\mathrm{m}\cdot\mathrm{s}^{-2}}{\mathrm{kg}\cdot\mathrm{m}^{-3}\cdot\mathrm{m}^2}} = \mathrm{m/s}$; the argument gt/v_t of tanh is dimensionless; $[v_t^2/g] = \mathrm{m}$, so $s(t)$ is a length.
- **Numerical check (RK4, step 10^{-4}, units with $g = v_t = 1$):** integrating $\dot v = 1 - v^2$, $\dot s = v$ up to $t = 1$ gives $v = 0.761594$ and $s = 0.433781$, equal to $\tanh 1 = 0.761594$ and $\ln\cosh 1 = 0.433781$.

Source: Problems.pdf (Page 9)

PROBLEM 17

Linear Central Force

Original statement · Problems.pdf, p. 9

Newtonian Restoring Force

A force attracts a heavy particle M of mass m to a point O. The attractive force is proportional to the mass of the particle and to the distance OM. The proportionality coefficient is k^2. The coordinate system $Oxyz$ has the origin at the centre of attraction O and the axis Oz is vertically oriented. Initially, the coordinates of M are:

$$(x_0, y_0, z_0) = \left(\frac{\sqrt{3}g}{k^2}, 0, 0 \right)$$

and its initial velocity is:

$$(\dot{x}_0, \dot{y}_0, \dot{z}_0) = \left(0, \frac{2g}{k}, 0 \right)$$

(a) Describe the motion of M, specifying clearly the shape of the path.

(b) What is the velocity of the particle as a function of t?

Full Analytical Solution (4-Phase Methodology)

PHASE 1

Physical Statement, Hypotheses & Parameters

- **Physical System:** A particle M of mass m is subject to:

1. An attractive 3D linear central force directed towards origin O, proportional to mass m and distance $r = OM$, with coefficient k^2:

$$\mathbf{F}_c = -mk^2\mathbf{r}$$

2. Uniform vertical gravity along $-Oz$:

$$\mathbf{F}_g = -mg\mathbf{k}$$

- **Initial State ($t = 0$):**
- Position: $(x_0, y_0, z_0) = \left(\frac{\sqrt{3}g}{k^2}, 0, 0\right)$.
- Velocity: $(\dot{x}_0, \dot{y}_0, \dot{z}_0) = \left(0, \frac{2g}{k}, 0\right)$.
- **Parameters:** $m > 0$ [kg], $g > 0$ [m/s^2], $k > 0$ [s^{-1}].

PHASE 2

Reference Frames, Vector Bases & Coordinate Geometry

- **Inertial Reference Frame:** Fixed Cartesian frame $\mathcal{F}_0 = \{O; \mathbf{i}, \mathbf{j}, \mathbf{k}\}$ with origin at O and Oz pointing vertically upwards.
- **Position Vector:** $\mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k}$.
- **Newton's Second Law:**

$$m\ddot{\mathbf{r}} = \mathbf{F}_c + \mathbf{F}_g = -mk^2\mathbf{r} - mg\mathbf{k}$$

Divide by m:

$$\ddot{\mathbf{r}} + k^2\mathbf{r} = -g\mathbf{k}$$

PHASE 3

Step-by-Step Mathematical Deduction

1. Decoupled Component Differential Equations

Pedagogical Justification: Because the system is linear and isotropic in 3D Cartesian coordinates, the vector ODE uncouples into three independent scalar second-order ODEs with constant coefficients.

$$\ddot{x} + k^2 x = 0$$

$$\ddot{y} + k^2 y = 0$$

$$\ddot{z} + k^2 z = -g$$

2. Exact Integration of Each Component

1. x**-Component:**

General solution:

$$x(t) = A_x \cos(kt) + B_x \sin(kt)$$

Velocity: $\dot{x}(t) = -kA_x \sin(kt) + kB_x \cos(kt)$.

Applying initial conditions:

$$x(0) = A_x = \frac{\sqrt{3}g}{k^2}$$

$$\dot{x}(0) = kB_x = 0 \implies B_x = 0$$

$$x(t) = \frac{\sqrt{3}g}{k^2}\cos(kt)$$

2. y**-Component:**

General solution:

$$y(t) = A_y\cos(kt) + B_y\sin(kt)$$

Velocity: $\dot{y}(t) = -kA_y\sin(kt) + kB_y\cos(kt)$.

Applying initial conditions:

$$y(0) = A_y = 0$$

$$\dot{y}(0) = kB_y = \frac{2g}{k} \implies B_y = \frac{2g}{k^2}$$

$$y(t) = \frac{2g}{k^2}\sin(kt)$$

3. z**-Component:**

Non-homogeneous equation with constant forcing $-g$.

Particular solution: $z_p = -\frac{g}{k^2}$.

General solution:

$$z(t) = -\frac{g}{k^2} + A_z \cos(kt) + B_z \sin(kt)$$

Velocity: $\dot{z}(t) = -kA_z \sin(kt) + kB_z \cos(kt)$.

Applying initial conditions:

$$z(0) = -\frac{g}{k^2} + A_z = 0 \implies A_z = \frac{g}{k^2}$$

$$\dot{z}(0) = kB_z = 0 \implies B_z = 0$$

$$z(t) = \frac{g}{k^2}(\cos(kt) - 1)$$

3. Particle Trajectory Vector

$$\mathbf{r}_M(t) = \frac{\sqrt{3}g}{k^2} \cos(kt)\,\mathbf{i} + \frac{2g}{k^2} \sin(kt)\,\mathbf{j} + \frac{g}{k^2}(\cos(kt) - 1)\,\mathbf{k}$$

4. Velocity Vector and Modulus

Differentiating with respect to time:

$$\dot{x}(t) = -\frac{\sqrt{3}g}{k} \sin(kt)$$

$$\dot{y}(t) = \frac{2g}{k}\cos(kt)$$

$$\dot{z}(t) = -\frac{g}{k}\sin(kt)$$

Velocity vector:

$$\mathbf{v}_M(t) = -\frac{\sqrt{3}g}{k}\sin(kt)\,\mathbf{i} + \frac{2g}{k}\cos(kt)\,\mathbf{j} - \frac{g}{k}\sin(kt)\,\mathbf{k}$$

Modulus of velocity:

$$|\mathbf{v}_M(t)|^2 = \dot{x}^2 + \dot{y}^2 + \dot{z}^2 = \left(-\frac{\sqrt{3}g}{k}\sin(kt)\right)^2 + \left(\frac{2g}{k}\cos(kt)\right)^2 + \left(-\frac{g}{k}\sin(kt)\right)^2$$

$$|\mathbf{v}_M(t)|^2 = \frac{3g^2}{k^2}\sin^2(kt) + \frac{4g^2}{k^2}\cos^2(kt) + \frac{g^2}{k^2}\sin^2(kt)$$

Factor out $\frac{g^2}{k^2}$:

$$|\mathbf{v}_M(t)|^2 = \frac{g^2}{k^2}\left[(3+1)\sin^2(kt) + 4\cos^2(kt)\right] = \frac{g^2}{k^2}\left[4\sin^2(kt) + 4\cos^2(kt)\right] = \frac{4g^2}{k^2}$$

Taking the square root:

$$|\mathbf{v}_M(t)| = \frac{2g}{k} = \text{constant}$$

The particle travels with strictly **constant speed**!

5. Geometric Nature and Shape of the Trajectory

- **Planarity Test:**

Notice the direct linear relationship between $x(t)$ and $z(t)$:

$$x(t) = \frac{\sqrt{3}g}{k^2}\cos(kt) \implies \cos(kt) = \frac{k^2}{\sqrt{3}g}x(t)$$

Substitute this into $z(t)$:

$$z(t) = \frac{g}{k^2}\cos(kt) - \frac{g}{k^2} = \frac{g}{k^2}\left(\frac{k^2}{\sqrt{3}g}x(t)\right) - \frac{g}{k^2} = \frac{x(t)}{\sqrt{3}} - \frac{g}{k^2}$$

Multiply through by $\sqrt{3}$:

$$\sqrt{3}z = x - \frac{\sqrt{3}g}{k^2} \implies x - \sqrt{3}z = \frac{\sqrt{3}g}{k^2}$$

This is the equation of a **plane** in $\mathbb{R}^3$. The trajectory is strictly planar!

- **Spherical Surface Test:**

Consider the distance of M from the point $C\left(0, 0, -\frac{g}{k^2}\right)$:

$$x^2 + y^2 + \left(z + \frac{g}{k^2}\right)^2 = \left(\frac{\sqrt{3}g}{k^2}\cos(kt)\right)^2 + \left(\frac{2g}{k^2}\sin(kt)\right)^2 + \left(\frac{g}{k^2}\cos(kt)\right)^2$$

$$= \frac{3g^2}{k^4}\cos^2(kt) + \frac{4g^2}{k^4}\sin^2(kt) + \frac{g^2}{k^4}\cos^2(kt) = \frac{4g^2}{k^4}(\cos^2(kt) + \sin^2(kt)) = \frac{4g^2}{k^4}$$

This defines a **sphere** centered at $C\left(0, 0, -\frac{g}{k^2}\right)$ with radius:

$$R = \frac{2g}{k^2}$$

- **Intersection & Circle Characterization:**

Check if the center of the sphere $C\left(0, 0, -\frac{g}{k^2}\right)$ lies on the plane $x - \sqrt{3}z = \frac{\sqrt{3}g}{k^2}$:

$$0 - \sqrt{3}\left(-\frac{g}{k^2}\right) = +\frac{\sqrt{3}g}{k^2}$$

The plane passes directly through the center of the sphere! Therefore, the intersection of the sphere and the plane is a **great circle** of radius $R = \frac{2g}{k^2}$.

PHASE 4

Physical Interpretation & Dimensional Verification

- The motion of the particle is a **uniform circular motion** along a great circle of radius $R = \frac{2g}{k^2}$ with constant orbital speed $v = \frac{2g}{k}$ and angular frequency $\omega = \frac{v}{R} = k$.
- The effective center of attraction is shifted from the origin O to $C(0, 0, -g/k^2)$ because the linear restoring force combines with uniform gravity into an equivalent spring centered at the equilibrium point where $k^2 z_{eq} = -g$.

- **Dimensional check:** $[x] = [g/k^2] = \dfrac{\mathrm{m/s^2}}{\mathrm{s^{-2}}} = \mathrm{m}$; $[v] = [g/k] = \mathrm{m/s}$; $[R] = [2g/k^2] = \mathrm{m}$; and $\omega = v/R = k$ has units $\mathrm{s^{-1}}$.
- **Limit $k \to \infty$ (stiff attraction):** $R = 2g/k^2 \to 0$ and the equilibrium offset $z_{eq} = -g/k^2 \to 0$, so gravity becomes negligible against the central force. The centripetal consistency check holds for any k: $v^2/R = (4g^2/k^2)/(2g/k^2) = 2g$ equals the effective spring force per unit mass $k^2 R = 2g$.
- **Symbolic check (sympy):** the trajectory of Phase 3 satisfies $\ddot{\mathbf{r}} + k^2\mathbf{r} = -g\mathbf{k}$ with zero residual and the initial data $\mathbf{r}(0)$, $\mathbf{v}(0)$; $\|\mathbf{v}\|^2 = 4g^2/k^2$, the squared distance to C is $4g^2/k^4$, and $x - \sqrt{3}z = \sqrt{3}g/k^2$ for all t (the centre C lies on this plane).

Source: Problems.pdf (Page 10)

PROBLEM 21

Rotating Planet & Apparent Weight

Original statement · Problems.pdf, p. 10

Gravitation & Centrifugal Balance

On a spinning spherical planet of radius R and gravitational acceleration g_p, the bodies weight twice as much on the poles as on the equator. Determine the duration of the day on that planet.

Full Analytical Solution (4-Phase Methodology)

PHASE 1

Physical Statement, Hypotheses & Parameters

- **Physical System:** A spherical planet of radius R spins with constant angular velocity $\mathbf{\Omega} = \Omega\mathbf{k}$ around its polar axis.
- **Gravitational Field:** Uniform surface gravitational acceleration g_p directed toward the center of the planet.
- **Apparent Weight:** The weight measured by a spring balance or scale is the normal reaction force $\mathbf{N}$ exerted by the surface to support a static mass m on the surface.
- **Problem Condition:** A body on the pole weighs twice as much as on the equator:

$$W_{pole} = 2W_{equator}$$

- **Objective:** Determine the duration of the planet's sidereal day $T_{day} = \frac{2\pi}{\Omega}$.

PHASE 2

Reference Frames, Vector Bases & Coordinate Geometry

- **Inertial Frame $\mathcal{F}_0$:** Frame centered at the planet center O.
- **Rotating Frame $\mathcal{F}_1$:** Planet-fixed reference frame rotating with angular velocity $\boldsymbol{\omega}_{1/0} = \Omega\mathbf{k}$.
- **Acceleration of a Stationary Point on the Surface:**

For a body resting on the surface, relative velocity and acceleration vanish: $\mathbf{v}' = \mathbf{0}, \mathbf{a}' = \mathbf{0}$.

The inertial acceleration is purely centripetal:

$$\mathbf{a}_0 = \boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \mathbf{r}) = -\Omega^2 \boldsymbol{\rho}$$

where $\boldsymbol{\rho}$ is the projection of the position vector onto the equatorial plane.

- **Change of basis (planet-fixed rotating basis).** Let $\mathcal{B}_0 = \{\mathbf{i}, \mathbf{j}, \mathbf{k}\}$ be the inertial basis and $\mathcal{B}_1 = \{\mathbf{i}_1, \mathbf{j}_1, \mathbf{k}\}$ the basis fixed to the planet, rotating at Ω about the polar axis $\mathbf{k}$, with $\mathbf{i}_1 = \cos\Omega t\,\mathbf{i} + \sin\Omega t\,\mathbf{j}$. The columns are the components of $\mathbf{i}_1, \mathbf{j}_1, \mathbf{k}$ in $\mathcal{B}_0$:

$$[{}_0R_1] = \begin{pmatrix} \cos\Omega t & -\sin\Omega t & 0 \\ \sin\Omega t & \cos\Omega t & 0 \\ 0 & 0 & 1 \end{pmatrix}, \qquad \det[{}_0R_1] = \cos^2\Omega t + \sin^2\Omega t = 1, \qquad [{}_0R_1][{}_0R_1]^T = I$$

- A body fixed at the equator at longitude φ_0 has $\mathbf{r} = R\,[{}_0R_1](\cos\varphi_0, \sin\varphi_0, 0)^T = R\,\mathbf{e}_r$, with $\mathbf{e}_r = \cos(\Omega t + \varphi_0)\,\mathbf{i} + \sin(\Omega t + \varphi_0)\,\mathbf{j}$. Poisson's formula $\dot{\mathbf{e}}_r = \Omega\,\mathbf{e}_\varphi$, $\dot{\mathbf{e}}_\varphi = -\Omega\,\mathbf{e}_r$ gives $\mathbf{a}_0 = -\Omega^2 R\,\mathbf{e}_r$, the centripetal acceleration used in Phase 3. The determinant, RR^T and this acceleration were verified symbolically with sympy.

PHASE 3

Step-by-Step Mathematical Deduction

1. Dynamic Equilibrium in the Rotating Frame

> **Pedagogical Justification:** In the non-inertial reference frame of the planet, static equilibrium requires the sum of real gravitational force $\mathbf{F}_g$, reaction force $\mathbf{N}$, and fictitious centrifugal force $\mathbf{F}_{cf} = -m\boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \mathbf{r}) = m\Omega^2\boldsymbol{\rho}$ to vanish.

$$\mathbf{F}_g + \mathbf{N} + \mathbf{F}_{cf} = \mathbf{0} \implies \mathbf{N} = -\mathbf{F}_g - \mathbf{F}_{cf}$$

The apparent weight is the magnitude of the normal force, $W = |\mathbf{N}|$.

2. Apparent Weight at the Pole

At the geographic poles, the position vector lies along the rotation axis $\mathbf{k}$:

$$\mathbf{r}_{pole} = \pm R\mathbf{k} \implies \boldsymbol{\rho}_{pole} = \mathbf{0}$$

The centrifugal force vanishes at the poles:

$$\mathbf{F}_{cf,pole} = \mathbf{0}$$

Therefore:

$$\mathbf{N}_{pole} - mg_p\mathbf{k} = \mathbf{0} \implies W_{pole} = mg_p$$

3. Apparent Weight at the Equator

At the equator, the position vector lies entirely in the equatorial plane:

$$|\boldsymbol{\rho}_{eq}| = R$$

Gravity acts radially inwards: $\mathbf{F}_{g,eq} = -mg_p\mathbf{e}_r$.

The centrifugal force acts radially outwards: $\mathbf{F}_{cf,eq} = +m\Omega^2 R\mathbf{e}_r$.

Normal reaction from the surface: $\mathbf{N}_{eq} = N_{eq}\mathbf{e}_r$.

Equilibrium equation:

$$-mg_p\mathbf{e}_r + N_{eq}\mathbf{e}_r + m\Omega^2 R\mathbf{e}_r = \mathbf{0}$$

$$N_{eq} = m(g_p - \Omega^2 R)$$

Thus, the apparent weight at the equator is:

$$W_{equator} = m(g_p - \Omega^2 R)$$

4. Computation of the Rotation Rate Ω

Imposing the given condition $W_{pole} = 2W_{equator}$:

$$mg_p = 2m(g_p - \Omega^2 R)$$

Divide both sides by m:

$$g_p = 2g_p - 2\Omega^2 R$$

Rearrange to isolate Ω^2:

$$2\Omega^2 R = 2g_p - g_p = g_p$$

$$\Omega^2 = \frac{g_p}{2R} \implies \Omega = \sqrt{\frac{g_p}{2R}}$$

5. Duration of the Day

The duration of one full planetary rotation (the day) is:

$$T_{day} = \frac{2\pi}{\Omega} = \frac{2\pi}{\sqrt{\frac{g_p}{2R}}} = 2\pi\sqrt{\frac{2R}{g_p}} = 2\sqrt{2}\pi\sqrt{\frac{R}{g_p}}$$

PHASE 4

Physical Interpretation & Dimensional Verification

- **Dimensional Analysis:**

$[T_{day}] = \sqrt{\frac{[R]}{[g_p]}} = \sqrt{\frac{\mathrm{m}}{\mathrm{m/s^2}}} = \sqrt{\mathrm{s^2}} = \mathrm{s}.$

The result is dimensionally sound.

- **Centrifugal Breakup Limit:**

A planet becomes gravitationally unbounded at the equator (weightlessness) when $\Omega_{crit} = \sqrt{g_p/R}$. Here $\Omega = \sqrt{g_p/(2R)} = \frac{1}{\sqrt{2}}\Omega_{crit} \approx 0.707\,\Omega_{crit}$, ensuring that objects remain resting on the equatorial surface.

- **Numerical illustration (Python):** for $g_p = 9.81\ \mathrm{m/s^2}$ and $R = 6.371 \times 10^6\ \mathrm{m}$, $\Omega = \sqrt{g_p/(2R)} = 8.774 \times 10^{-4}\ \mathrm{rad/s}$ and $T_{day} = 2\pi\sqrt{2R/g_p} = 7.161 \times 10^3\ \mathrm{s} = 1.99\ \mathrm{h}$, which satisfies $\Omega^2 R/g_p = 0.5$ (so $W_{equator} = \frac{1}{2}W_{pole}$ as imposed).

Source: Problems.pdf (Page 10)

PROBLEM 23

Simple Pendulum (Unilateral Constraint)

Original statement · Problems.pdf, p. 11

Pendulum & Liftoff

A simple pendulum consists of a mass m hanging from a string of length L and fixed at a pivot point O. The simple pendulum is equivalent to the problem of a heavy particle constrained to move along a smooth circumference. A unilateral constraint is considered, such that the particle moves along the interior of the circumference and it might fall under certain conditions.

Analyze the motion of the particle when it is pushed from the lowest point of the circumference with velocity v_0. At what point does the particle leave the circumference? Is this point ever reached?

If the circumference rotates around the vertical axis with a constant angular velocity equal to ω, write down the new equations of motion. Compute the locations along the circumference in which the particle is in equilibrium.

Full Analytical Solution (4-Phase Methodology)

PHASE 1

Physical Statement, Hypotheses & Parameters

- **Physical System:** A particle of mass m is suspended by a string of length L from pivot O, or equivalently moves along the smooth interior of a vertical circular track of radius L.
- **Unilateral Constraint:** The particle cannot cross the circular boundary outward, but may detach if the inward contact force drops to zero ($N = 0$, or string slack $T = 0$).
- **Initial Motion:** Pushed from the lowest point ($\theta = 0$) with velocity v_0.
- **Part (b) Modification:** The circular track rotates about its vertical diameter with constant angular velocity ω.
- **Parameters:** $m > 0$ [kg], $L > 0$ [m], $g > 0$ [m/s^2], $v_0 > 0$ [m/s], $\omega \geq 0$ [rad/s].

PHASE 2

Reference Frames, Vector Bases & Coordinate Geometry

- **Inertial Frame $\mathcal{F}_0 = \{O; \mathbf{i}, \mathbf{j}, \mathbf{k}\}$:** Oz points vertically upward ($-\mathbf{k}$ is downward).
- **Polar Coordinate Basis $\mathcal{B}_1 = \{\mathbf{e}_r, \mathbf{e}_\theta, \mathbf{j}\}$:**

Angle θ measured from the downward vertical ($-\mathbf{k}$):

$$\mathbf{e}_r = \sin\theta\,\mathbf{i} - \cos\theta\,\mathbf{k} \quad \text{(pointing radially outward)}$$

$$\mathbf{e}_\theta = \cos\theta\,\mathbf{i} + \sin\theta\,\mathbf{k} \quad \text{(tangential, along increasing } \theta\text{)}$$

- **Kinematics:**
- Position: $\mathbf{r} = L\mathbf{e}_r$.
- Velocity: $\mathbf{v} = L\dot{\theta}\mathbf{e}_\theta$.
- Acceleration: $\mathbf{a} = -L\dot{\theta}^2\mathbf{e}_r + L\ddot{\theta}\mathbf{e}_\theta$.
- **Change of basis.** The triple $\{\mathbf{e}_r, \mathbf{e}_\theta, \mathbf{j}\}$ listed above has $\mathbf{e}_r \times \mathbf{e}_\theta = -\mathbf{j}$, so it is left-handed (determinant -1); the right-handed polar triple is $\{\mathbf{e}_r, \mathbf{e}_\theta, \mathbf{e}_3\}$ with $\mathbf{e}_3 = -\mathbf{j}$ (the third vector plays no role in the planar kinematics). The columns are the components in $\mathcal{B}_0 = \{\mathbf{i}, \mathbf{j}, \mathbf{k}\}$:

$$[{}_0R_1] = \begin{pmatrix} \sin\theta & \cos\theta & 0 \\ 0 & 0 & -1 \\ -\cos\theta & \sin\theta & 0 \end{pmatrix}, \qquad \det[{}_0R_1] = 1, \qquad [{}_0R_1][{}_0R_1]^T = I$$

- The angular velocity of $\mathcal{B}_1$ is $\boldsymbol{\omega}_{10} = \dot{\theta}\,\mathbf{e}_3 = -\dot{\theta}\,\mathbf{j}$, and Poisson's formula gives $\dot{\mathbf{e}}_r = \dot{\theta}\,\mathbf{e}_\theta$, $\dot{\mathbf{e}}_\theta = -\dot{\theta}\,\mathbf{e}_r$, the identities behind $\mathbf{v} = L\dot{\theta}\,\mathbf{e}_\theta$ and $\mathbf{a} = -L\dot{\theta}^2\mathbf{e}_r + L\ddot{\theta}\,\mathbf{e}_\theta$.
- For part (b) the track rotates about $\mathbf{k}$ at rate ω: the rotating basis $\{\mathbf{e}_\rho, \mathbf{e}_\phi, \mathbf{k}\}$ is related to $\mathcal{B}_0$ by $\mathrm{R}_z(\omega t)$ (a rotation matrix with the same properties, $\det = 1$, $RR^T = I$), and inside it the pair $\{\mathbf{e}_r, \mathbf{e}_\theta\}$ is written with $\mathbf{e}_\rho$ in place of $\mathbf{i}$. The determinant and the product RR^T were verified symbolically with sympy.

PHASE 3

Step-by-Step Mathematical Deduction

Part (a): Planar Motion & Detachment Analysis

1. Mechanical Energy Conservation

> **Pedagogical Justification:** The normal constraint force $\mathbf{N} = -N\mathbf{e}_r$ is always perpendicular to velocity $\mathbf{v} = L\dot{\theta}\mathbf{e}_\theta$ ($\mathbf{N} \cdot \mathbf{v} = 0$), so it does no work. Mechanical energy is strictly conserved.

Taking the lowest point $\theta = 0$ as the gravitational potential reference ($V(0) = 0$):

$$V(\theta) = mg(L - L\cos\theta) = mgL(1 - \cos\theta)$$

$$E = \frac{1}{2}mv_0^2 = \frac{1}{2}mv^2 + mgL(1 - \cos\theta)$$

Isolate $v^2(\theta)$:

$$v^2(\theta) = v_0^2 - 2gL(1 - \cos\theta)$$

2. Normal Reaction Force $N(\theta)$

> **Pedagogical Justification:** Newton's second law along the inward radial normal $\mathbf{e}_n = -\mathbf{e}_r$ provides an explicit expression for the contact reaction force $N(\theta)$ as a function of angle θ.

The forces acting on the particle are gravity $\mathbf{F}_g = -mg\mathbf{k}$ and track reaction $\mathbf{N} = -N\mathbf{e}_r$ ($N > 0$ pushes inward):

$$\mathbf{F}_g = -mg\mathbf{k} = -mg(-\cos\theta\,\mathbf{e}_r + \sin\theta\,\mathbf{e}_\theta) = mg\cos\theta\,\mathbf{e}_r - mg\sin\theta\,\mathbf{e}_\theta$$

Projecting Newton's second law along the inward normal $-\mathbf{e}_r$:

$$\sum F_n = N - mg\cos\theta = ma_n = m\frac{v^2}{L}$$

$$N(\theta) = mg\cos\theta + \frac{m}{L}v^2(\theta)$$

Substitute the energy relation for $v^2(\theta)$:

$$N(\theta) = mg\cos\theta + \frac{m}{L}\left[v_0^2 - 2gL(1-\cos\theta)\right]$$

Expand and combine terms:

$$N(\theta) = mg\cos\theta + \frac{mv_0^2}{L} - 2mg + 2mg\cos\theta = \frac{mv_0^2}{L} - 2mg + 3mg\cos\theta$$

$$N(\theta) = \frac{mv_0^2}{L} - mg(2 - 3\cos\theta)$$

3. Detachment Condition ($N = 0$)

The particle leaves the circumference if N drops to zero while $v > 0$:

$$N(\theta_d) = 0 \implies \frac{mv_0^2}{L} - 2mg + 3mg\cos\theta_d = 0$$

$$3mg\cos\theta_d = 2mg - \frac{mv_0^2}{L} \implies \cos\theta_d = \frac{2gL - v_0^2}{3gL}$$

4. Motion Regimes and Question Resolution: "Is this point ever reached?"

- **Regime 1:** $v_0 \leq \sqrt{2gL}$ **(Low energy, oscillation in lower semicircle)**

Velocity reaches zero when $v^2 = 0$:

$$v_0^2 - 2gL(1 - \cos\theta_{max}) = 0 \implies \cos\theta_{max} = 1 - \frac{v_0^2}{2gL} \geq 0 \implies \theta_{max} \leq \frac{\pi}{2}$$

At this maximum angle, the reaction force is:

$$N(\theta_{max}) = mg\cos\theta_{max} \geq 0$$

The particle comes to rest and reverses direction before N can vanish.

Conclusion: The detachment point is **never reached**. The particle oscillates periodically in the lower semicircle.

- **Regime 2:** $\sqrt{2gL} < v_0 < \sqrt{5gL}$ **(Intermediate energy, detachment occurs)**

Since $v_0^2 > 2gL$, the particle crosses the horizontal position $\theta = \pi/2$ with positive velocity.

In the upper semicircle ($\pi/2 < \theta < \pi$), $\cos\theta < 0$.

Evaluate the detachment angle $\cos\theta_d$:

$$\cos\theta_d = \frac{2gL - v_0^2}{3gL}$$

Because $2gL < v_0^2 < 5gL$:

$$-1 = \frac{2gL - 5gL}{3gL} < \cos\theta_d < \frac{2gL - 2gL}{3gL} = 0$$

So $\cos\theta_d \in (-1, 0)$, meaning $\theta_d \in (\pi/2, \pi)$ is physically accessible!

At this point, the speed is:

$$v^2(\theta_d) = v_0^2 - 2gL\left(1 - \frac{2gL - v_0^2}{3gL}\right) = v_0^2 - 2gL + \frac{2}{3}(2gL - v_0^2) = \frac{v_0^2 - 2gL}{3} > 0$$

Conclusion: The detachment point **is reached**. The particle leaves the circular track at angle $\theta_d = \arccos\left(\frac{2gL - v_0^2}{3gL}\right)$ and transitions into free parabolic flight under gravity alone.

- **Regime 3:** $v_0 \geq \sqrt{5gL}$ **(High energy, complete vertical loops)**

At the very top of the circle ($\theta = \pi$, $\cos\pi = -1$):

$$N(\pi) = \frac{mv_0^2}{L} - mg(2 - 3(-1)) = \frac{mv_0^2}{L} - 5mg \geq 0$$

The reaction force remains positive throughout the entire loop.

Conclusion: The particle **never leaves** the circumference and performs complete revolutions.

Part (b): Rotating Circumference with Angular Velocity ω

1. Equations of Motion in the Rotating Frame

> **Pedagogical Justification:** When the circular track rotates at constant angular speed ω about the vertical diameter Oz, we analyze the motion in the rotating frame $\mathcal{F}_1$. The centrifugal force $\mathbf{F}_{cf} = m\omega^2\boldsymbol{\rho}$ adds a horizontal outward component.

Let cylindrical coordinates be (ρ, ϕ, z), with $\rho = R\sin\theta$, $z = -R\cos\theta$.

In the rotating frame, the unit tangent along increasing θ is:

$$\mathbf{e}_\theta = \cos\theta\,\mathbf{e}_\rho + \sin\theta\,\mathbf{k}$$

Forces acting on the particle:

1. Gravity: $\mathbf{F}_g = -mg\mathbf{k}$.

2. Centrifugal force: $\mathbf{F}_{cf} = m\omega^2\rho\,\mathbf{e}_\rho = m\omega^2 R\sin\theta\,\mathbf{e}_\rho$.

3. Reaction force $\mathbf{N}$ (normal to the wire, along $\mathbf{e}_r$ and $\mathbf{e}_\phi$).

Projecting along the tangent $\mathbf{e}_\theta$:

$$mR\ddot{\theta} = (\mathbf{F}_g + \mathbf{F}_{cf}) \cdot \mathbf{e}_\theta$$

$$\mathbf{F}_g \cdot \mathbf{e}_\theta = (-mg\mathbf{k}) \cdot (\cos\theta\,\mathbf{e}_\rho + \sin\theta\,\mathbf{k}) = -mg\sin\theta$$

$$\mathbf{F}_{cf} \cdot \mathbf{e}_\theta = (m\omega^2 R\sin\theta\,\mathbf{e}_\rho) \cdot (\cos\theta\,\mathbf{e}_\rho + \sin\theta\,\mathbf{k}) = m\omega^2 R\sin\theta\cos\theta$$

Therefore:

$$mR\ddot{\theta} = -mg\sin\theta + m\omega^2 R\sin\theta\cos\theta$$

Divide by mR:

$$\ddot{\theta} + \frac{g}{R}\sin\theta - \omega^2\sin\theta\cos\theta = 0$$

$$\ddot{\theta} + \left(\frac{g}{R} - \omega^2\cos\theta\right)\sin\theta = 0$$

2. Computation of Equilibrium Locations

> **Pedagogical Justification:** Equilibrium configurations in the rotating frame correspond to static solutions where $\ddot{\theta} = 0$ and $\dot{\theta} = 0$.

Setting the angular acceleration to zero:

$$\left(\frac{g}{R} - \omega^2 \cos\theta \right) \sin\theta = 0$$

This equation is satisfied if either factor vanishes:

1. **Vertical Alignments** ($\sin\theta = 0$)**:**

$$\theta_1 = 0 \quad \text{(lowest point / bottom)}$$

$$\theta_2 = \pi \quad \text{(highest point / top)}$$

2. **Off-Axis Equilibrium** ($\cos\theta = \frac{g}{\omega^2 R}$)**:**

This solution exists physically if and only if:

$$\left| \frac{g}{\omega^2 R} \right| \leq 1 \iff \omega^2 \geq \frac{g}{R}$$

When $\omega > \sqrt{\frac{g}{R}}$, two symmetric equilibrium positions emerge:

$$\theta_{3,4} = \pm \arccos\left(\frac{g}{\omega^2 R} \right)$$

PHASE 4

Physical Interpretation & Dimensional Verification

- **Supercritical Pitchfork Bifurcation:**
- For slow rotation $\omega < \omega_c = \sqrt{g/R}$, the bottom $\theta = 0$ is the only stable equilibrium.
- When the rotation rate crosses the critical threshold ω_c, the bottom becomes unstable, and the particle naturally climbs up the wire to settle at the tilted angle $\theta_0 = \arccos\left(\frac{g}{\omega^2 R}\right)$, where the upward component of the centrifugal force balances gravity.
- As $\omega \to \infty$, $\cos\theta_0 \to 0 \implies \theta_0 \to \pi/2$ (the particle moves toward the equator).
- **Dimensional check:** $[N] = [mv_0^2/L] = \text{kg}\cdot\text{m}\cdot\text{s}^{-2} = \text{N}$; $\cos\theta_d = (2gL - v_0^2)/(3gL)$ is dimensionless; $\omega_c = \sqrt{g/R}$ has units s^{-1}.
- **Limits of the regimes:** at $v_0^2 = 2gL$, $\cos\theta_d = 0$ ($\theta_d = \pi/2$), the boundary between oscillation and detachment; at $v_0^2 = 5gL$, $\cos\theta_d = -1$ ($\theta_d = \pi$) and $N(\pi) = 0$, the boundary between detachment and full loops. For $\omega \to 0$ in part (b) the equation reduces to the simple pendulum $\ddot{\theta} + (g/R)\sin\theta = 0$.
- **Numerical example ($g = L = 1$, $v_0^2 = 3.5$, regime 2):** $\cos\theta_d = (2 - 3.5)/3 = -0.5$, $\theta_d = 120^\circ$; $v^2(\theta_d) = (v_0^2 - 2gL)/3 = 0.5$ and $N(\theta_d) = v_0^2/L - g(2 - 3\cos\theta_d) = 3.5 - 3.5 = 0$, as required.

Source: Problems.pdf (Page 11)