PROBLEM 4 | Problems T2_CrystStruct.pdf | Crystalline Structures & Miller Indices

Planar and Linear Densities in the BCC Lattice

Original statement · Problems T2_CrystStruct.pdf

4. For a BCC structure, determine the surface (planar) density for the planes (100)**,** (110) ***and*** (111)***; the linear density in the direction*** $[100]$***, and the volume density.*** *(Solution:* $\rho_{(100)} = 1/a^2$*,* $\rho_{(110)} = \quad 2/a^2$*,* $\rho_{(111)} = 1/(a^2 \quad 3)$*,* $\rho_{[100]} = 1/a$*,* $\rho_v = 2/a^3$*)*

Solution: $\rho_{(100)} = 1/a^2$, $\rho_{(110)} = \quad 2/a^2$, $\rho_{(111)} = 1/(a^2 \quad 3)$, $\rho_{[100]} = 1/a$, $\rho_v = 2/a^3$

PHASE 1: HYPOTHESIS & PARAMETERS

Physical Context:

A Body-Centred Cubic (BCC) lattice with lattice parameter a is studied.

BCC Lattice Data:

- Atoms per unit cell: $n = 2$ (8 at vertices $\times 1/8$ + 1 central).
- Volume of the cubic cell: $V_C = a^3$.
- Atomic radius: $R = \frac{a \quad 3}{4}$ (contact along the cube diagonal $\langle 111 \rangle$).

PHASE 2: FUNDAMENTAL PHYSICS & FORMULA SELECTION

Applying the formal definitions of the course [Session 4 Slides 7-9]:

- **Planar Density ($\rho_{(hkl)}$):**

$$\rho_{(hkl)} = \frac{N_{\text{atoms contained in the face or section}}}{A_{(hkl)}}$$

- **Linear Density ($\rho_{[uvw]}$):**

$$\rho_{[uvw]} = \frac{N_{\text{atoms cut diametrically}}}{L_{[uvw]}}$$

- **Volumetric Density (ρ_v):**

$$\rho_v = \frac{n}{V_C} = \frac{2}{a^3}$$

PHASE 3: ANALYTICAL DERIVATION & CALCULATION STEP-BY-STEP

1. Planar Density on the (100) Plane:

- **Plane geometry:** Outer face of the cube, of dimensions $a \times a$.
- **Area:** $A_{(100)} = a^2$.
- **Atoms contained:** Only the 4 atoms located at the vertices of the square (the central atom of the BCC cell is at $z = a/2$, so it does not belong to the plane $z = 0$ or $x = 0$).

$$N_{\text{atoms}} = 4 \times \frac{1}{4} = 1 \text{ atom}$$

- **Planar density:**

$$\rho_{(100)} = \frac{1}{a^2} = \mathbf{\frac{1}{a^2}}$$

2. Planar Density on the (110) Plane:

- **Plane geometry:** Diagonal rectangle crossing the cube through two opposite edges.
- **Dimensions:** Base $= a \quad 2$ (face diagonal), Height $= a$.
- **Area:**

$$A_{(110)} = a \quad 2 \times a = a^2 \quad 2$$

- **Atoms contained:**
- 4 vertices of the rectangle, each contributing $\frac{1}{4}$: $4 \times \frac{1}{4} = 1$ atom.
- 1 atom at the geometric centre of the cell (the (110) plane passes exactly through the centre of the cube): it contributes a whole 1 atom.
- Total atoms: $N_{\text{atoms}} = 1 + 1 = 2$ atoms.
- **Planar density:**

$$\rho_{(110)} = \frac{2}{a^2 \quad 2} = \frac{2 \quad 2}{2a^2} = \boxed{\frac{\mathbf{2}}{\mathbf{a^2}}}$$

3. Planar Density on the (111) Plane:

- **Geometry and spacing:**

In the BCC lattice, the parallel (111) planes have an interplanar spacing $d_{111} = \frac{a}{1^2+1^2+1^2} = \frac{a}{3}$.

However, because the central atom sits exactly at the midpoint of the cube diagonal along $[111]$, the real (111) planes alternate at an effective distance $d_{\text{eff}} = \frac{a}{2 \quad 3}$ (corresponding to the reflected family $\{222\}$).

- **Direct calculation by cell or density relation:**

Using the fundamental conservation relation between densities [Session 4 Slide 9]:

$$\rho_v = \frac{\rho_{(hkl)}}{d_{\text{eff}}} \implies \rho_{(111)} = \rho_v \cdot d_{\text{eff}} = \left(\frac{2}{a^3}\right) \cdot \left(\frac{a}{2 \quad 3}\right) = \boxed{\frac{\mathbf{1}}{\mathbf{a^2} \quad \mathbf{3}}}$$

- **Direct check by geometric area:**

The plane that cuts the vertices $(a,0,0), (0,a,0), (0,0,a)$ forms an equilateral triangle of side $L = a \quad 2$ and area $A = \frac{3}{4}(a \quad 2)^2 = \frac{3}{2}a^2$.

The plane does not contain the central atom (it lies at $(a/2, a/2, a/2)$, whose distance to the origin is $\frac{a \quad 3}{2}$, while the plane is at $\frac{a}{3}$).

Therefore, the plane contains only the 3 vertices: $N_{\text{atoms}} = 3 \times \frac{1}{6} = \frac{1}{2}$ atom.

$$\rho_{(111)} = \frac{1/2}{\frac{3}{2}a^2} = \boxed{\frac{\mathbf{1}}{\mathbf{a^2} \quad \mathbf{3}}}$$

4. Linear Density in the $[100]$ Direction:

- **Edge segment length:** $L_{[100]} = a$.
- **Atoms cut:** It cuts the two vertex atoms at their ends:

$$N_{\text{atoms}} = 2 \times \frac{1}{2} = 1 \text{ atom}$$

- **Linear density:**

$$\rho_{[100]} = \frac{1}{a} = \boxed{\frac{\mathbf{1}}{\mathbf{a}}}$$

5. Volumetric Density (ρ_v):

- It contains $n = 2$ atoms in a cell volume $V_C = a^3$:

$$\rho_v = \frac{\mathbf{2}}{\mathbf{a^3}}$$

PHASE 4: PHYSICAL INTERPRETATION & VERIFICATION

Study notes written from the official course material (Problems T2_CrystStruct.pdf; Problems T2 defects.pdf).
Online version: subjects/aerospace-materials-1/problemas/topic-2-problems.html