PROBLEM 44 | Angular Momentum + Vis-Viva

The Little Prince Throwing Seeds

Original statement · Problems.pdf, p. 31

The little Prince is on his spherical asteroid, which has radius R and mass M, and does not rotate. While he sits at the equator, he throws a baobab seed with an initial velocity of magnitude v_0 and angle $\beta = 60^\circ$ with respect to the local horizontal. Knowing that it reaches an apocenter of $r_a = 3R$ before falling back, determine v_0 as a function of the parameters of the problem and the universal gravitational constant G.

Worked Solution (4-Phase Methodology)

PHASE 1

Physical Statement, Hypotheses & Parameters

- **System:** a seed of mass $m \ll M$ launched from the surface ($r_0 = R$) with speed v_0 at 60° above the local horizontal; it reaches the apocenter $r_a = 3R$.
- **Hypotheses:** spherically symmetric, non-rotating asteroid, so the gravitational force $\mathbf{F} = -\dfrac{GMm}{r^2}\mathbf{e}_r$ is central and conservative with $\mu = GM$; vacuum (no drag); inertial frame S_0 centred on the asteroid.
- **Degrees of freedom:** free particle, planar motion by central-force conservation of $\mathbf{H}_O$; coordinates (r, θ).
- **Initial state:** $r = R$, radial component $v_{r0} = v_0 \sin 60^\circ = \frac{3}{2} v_0$ and transverse component $v_{\theta 0} = v_0 \cos 60^\circ = \frac{1}{2} v_0$.

Symbol	Meaning	SI unit
M, R	mass and radius of the asteroid	kg, m
G	universal gravitational constant, 6.6743×10^{-11}	$\mathrm{m^3/(kg\ s^2)}$
$\mu = GM$	gravitational parameter	$\mathrm{m^3/s^2}$
β	launch angle above the horizontal	60°
r_a	apocenter radius	$3R$

PHASE 2

Reference Frames, Vector Bases & Coordinate Geometry

- **Change of basis (polar basis in the orbital plane).** $\mathcal{B}_1 = \{\mathbf{e}_r, \mathbf{e}_\theta, \mathbf{k}\}$ is the inertial basis rotated by the true anomaly θ about $\mathbf{k}$ (along $\mathbf{H}_O$):

$$[{}_0R_1] = \begin{pmatrix} \cos\theta & -\sin\theta & 0 \\ \sin\theta & \cos\theta & 0 \\ 0 & 0 & 1 \end{pmatrix}, \qquad \det[{}_0R_1] = 1, \qquad [{}_0R_1]^T[{}_0R_1] = I$$

- **Velocity in $\mathcal{B}_1$:** $\mathbf{v} = \dot{r}\,\mathbf{e}_r + r\dot{\theta}\,\mathbf{e}_\theta$; at launch $\dot{r} = v_0 \sin 60^\circ$ and $r\dot{\theta} = v_0 \cos 60^\circ$. At the apocenter $\dot{r} = 0$, so the velocity is purely transverse.

PHASE 3

Step-by-Step Mathematical Deduction

1. Conservation of specific angular momentum

Pedagogical Justification: The gravitational force passes through O, so $\mathbf{M}_O = \mathbf{0}$ and $h = \|\mathbf{r} \times \mathbf{v}\| = r^2\dot{\theta}$ is constant. Evaluate it at launch and at the apocenter, where $v_a = r_a\dot{\theta}_a$.

$$h = R\,(v_0 \cos 60^\circ) = \frac{1}{2}Rv_0, \qquad h = r_a v_a = 3Rv_a \implies v_a = \frac{v_0}{6}$$

2. Conservation of specific energy

Pedagogical Justification: Gravity is conservative with $V_0 = -\mu/r$ (derived with Barrow's rule, reference at infinity), so $\xi = \frac{1}{2}v^2 - GM/r$ takes the same value at launch and at the apocenter.

$$\frac{1}{2}v_0^2 - \frac{GM}{R} = \frac{1}{2}\left(\frac{v_0}{6}\right)^2 - \frac{GM}{3R} = \frac{v_0^2}{72} - \frac{GM}{3R}$$

3. Solving for v_0

$$\left(\frac{1}{2}-\frac{1}{72}\right)v_0^2=\left(1-\frac{1}{3}\right)\frac{GM}{R}\implies\frac{35}{72}v_0^2=\frac{2}{3}\frac{GM}{R}\implies v_0^2=\frac{2}{3}\cdot\frac{72}{35}\frac{GM}{R}=\frac{48}{35}\frac{GM}{R}$$

$$\boxed{v_0=\sqrt{\frac{48}{35}\frac{GM}{R}}}$$

PHASE 4

Physical Interpretation & Dimensional Verification

- **Bound orbit:** the escape speed at the surface is $v_e=\sqrt{2GM/R}$, i.e. $v_e^2=\frac{70}{35}GM/R$. Since $\frac{48}{35}<2$, $v_0<v_e$ and the orbit is an ellipse: the seed falls back.
- **Orbital parameters:** $h^2=\frac{1}{4}R^2v_0^2=\frac{12}{35}GMR$, $p=h^2/\mu=\frac{12}{35}R$. From $r_a=p/(1-e)=3R$: $1-e=\frac{4}{35}$ and $e=\frac{31}{35}\approx0.8857$. The periapsis of the full conic is $r_p=p/(1+e)=\frac{2}{11}R<R$, inside the asteroid; only the arc from the launch point to the apocenter is physical, and the seed lands before reaching the periapsis.
- **Independent check with the Binet solution.** With $u=\frac{\mu}{h^2}(1+e\cos\theta)$ and the launch values $u(\theta_0)=1/R$, $u'(\theta_0)=-\dot r/h$ (here $\dot r=\frac{3}{2}v_0$), the eccentricity formula of the theory page gives

$$e=\frac{h^2}{\mu}\sqrt{\left(\frac{1}{R}-\frac{\mu}{h^2}\right)^2+\left(\frac{\dot r}{h}\right)^2}=\frac{12R}{35}\sqrt{\frac{529}{144R^2}+\frac{3}{R^2}}=\frac{12R}{35}\cdot\frac{31}{12R}=\frac{31}{35}$$

and the launch angle follows from $\tan\beta=\dot r/(r\dot\theta)=\frac{R}{p}\,e\sin\theta_0$ with $\cos\theta_0=\frac{p/R-1}{e}=-\frac{23}{31}$, $\sin\theta_0=\frac{432}{31}$:

$$\tan\beta=\frac{35}{12}\cdot\frac{31}{35}\cdot\frac{432}{31}=\frac{432}{12}=\sqrt{3}\implies\beta=60^\circ$$

- **Numerical check** (Python, velocity-Verlet integration of $\ddot{\mathbf{r}} = -\mu\mathbf{r}/r^3$ with $\mu = R = 1$ and $v_0 = \sqrt{48/35}$): the maximum radius reached is $2.99999999981R$, in agreement with $r_a = 3R$.
- **Dimensions:** $[v_0] = \sqrt{[G][M]/[R]} = \sqrt{\dfrac{\mathrm{m}^3}{\mathrm{kg\,s}^2}\cdot\dfrac{\mathrm{kg}}{\mathrm{m}}} = \mathrm{m/s}$.
- **Key:** the official key (Problems.pdf) has no entry for Problem 44.

Worked Solution

Source: Problems.pdf (PDF page 31); no key available

Study notes written from the official course material (Problems.pdf · p. 9). Online version: subjects/engineering-mechanics/problemas/topic-4.html