




## A Python/Zig optimized and customizable implementation for the $\rho_{DCCA}$ and $DMC_x^2$ methods

Fernando Ferraz Ribeiro   
Universidade Federal da Bahia

Gilney Figueira Zebende   
Universidade Estadual  
de Feira de Santana

---

### Abstract

This paper presents the **Zebende**, a Python package written in Python and Zig, that calculates the *DFA*, *DCCA*  $\rho_{DCCA}$  and the  $DMC_x^2$ . The package presents an optimized algorithm that significantly improves the calculations speed. A comparison with other packages that calculates the .The package is also the first to implement the  $DMC_x^2$  coefficient for any number of series.

*Keywords:*  $\rho_{DCCA}$ ,  $DMC_x^2$ , optimization, Python, Zig.

---

## 1. introduction

The  $\rho_{DCCA}$  (Zebende 2011) is a widely used coefficient that measures the cross-correlation between two non-stationary time series. It's an extension of the Detrended Fluctuation Analysis (*DFA*) (Peng, Buldyrev, Havlin, Simons, Stanley, and Goldberger 1994) and the Detrended Cross-correlation Analysis (*DCCA*) (Podobnik and Stanley 2008): while the *DFA* calculates the self-affinity and long-memory properties of a time series data, and the *DCCA* analyses power-law cross correlations between two different non-stationarity time series, the  $\rho_{DCCA}$  coefficient quantifies this cross-correlation in simple values ranging from  $-1$  to  $1$ , where  $-1$  indicates a perfect anti-correlation between the series,  $1$  a perfect correlation and zero ( $0$ ) no correlation at all.

The Detrended Multiple Cross-Correlation Coefficient (Zebende and Silva 2018) ( $DMC_x^2$ ) is a generalization of the  $\rho_{DCCA}$  coefficient that correlates one time series (dependent variable) a number of time series (independent variables). The  $DMC_x^2$  values ranges from zero ( $0$ ), indicating no correlation to  $1$ , meaning perfect correlation or anti-correlation between the dependent and the independent variables.

This paper presents the **Zebende** Python package, an implementation of the *DFA*, *DCCA*,  $\rho_{DCCA}$ ,  $DMC_x^2$  and utility functions related to the methods. In section 2 the steps for calculating the  $\rho_{DCCA}$  and  $DMC_x^2$  are presented and discussed. Section 3 shows how this library was implemented, the optimization technics and the recommended steps to use the library. In Section 4 the **Zebende** package is compared with other packages for Python and R that calculates the  $\rho_{DCCA}$  in terms of performance and usability, leading to the conclusions in 5.

## 2. Algorithms of the coefficients

$$DMC_x^2 \equiv \rho_{Y,X_i}(n)^T \times \rho^{-1}(n) \times \rho_{Y,X_i}(n) \quad (1)$$

$$\rho^{-1}(n) = \begin{pmatrix} 1 & \rho_{X_1,X_2}(n) & \rho_{X_1,X_3}(n) & \dots & \rho_{X_1,X_j}(n) \\ \rho_{X_2,X_1}(n) & 1 & \rho_{X_2,X_3}(n) & \dots & \rho_{X_2,X_j}(n) \\ \vdots & \vdots & \vdots & \dots & \vdots \\ \rho_{X_j,X_1}(n) & \rho_{X_j,X_2}(n) & \rho_{X_j,X_3}(n) & \dots & 1 \end{pmatrix}^{-1} \quad (2)$$

$$\rho_{Y,X_k}(n)^T = [\rho_{Y,X_1}(n), \rho_{Y,X_2}(n), \dots, \rho_{Y,X_j}(n)] \quad (3)$$

$$\rho_{DCCA}(n) = \frac{F_{DCCA}^2(x\alpha, x\beta)(n)}{F_{DFA}(x\alpha)(n) \times F_{DFA}(x\beta)(n)} \quad (4)$$

Taking a time series  $\{x_i\}$  with  $i$  ranging from 1 to  $N$ , the integrated series  $X_k$  is calculated by  $X_k = \sum_{i=1}^k [x_i - \langle x \rangle]$  with  $k$  also ranging from 1 to  $N$ ;

the  $X_k$  series is divided in  $N - n$  boxes of size  $n$ (time scale), each box containing  $n + 1$  values, starting in  $i$  up to  $i + n$ ;

for each box, a polynomial (usually of degree 1) best fit is calculated, getting  $\tilde{X}_{k,i}$  with  $i \leq k \leq (i + n)$  (detrended values);

in each box is calculated:  $f_{DFA}^2(n, i) = \frac{1}{1+n} \sum_{k=i}^{i+n} (X_k - \tilde{X}_{k,i})^2$

for all the boxes of a time scale, the DFA is calculated as:

$$F_{DFA}(n) = \sqrt{\frac{1}{N-n} \sum_{i=1}^{N-n} f_{DFA}^2(n, i)};$$

for a number of different timescales ( $n$ ), with possible values  $4 \leq n \leq \frac{N}{4}$  the  $F_{DFA}$  function is calculated to find a relation among  $F_{DFA} \times n$

Taking two time series with the same length  $\{x\alpha_i\}$  and  $\{x\beta_i\}$  with  $i$  ranging from 1 to  $N$ , the integrated series  $X\alpha_k$  and  $X\beta_k$  are calculated by  $X_k = \sum_{i=1}^k [x_i - \langle x \rangle]$  for each series, with  $k$  also ranging from 1 to  $N$ ;

$X\alpha_k$  and  $X\beta_k$  series are divided in  $N - n$  boxes of size  $n$ (time scale), each box containing  $n + 1$  values, starting in  $i$  up to  $i + n$ ;

for each box, a polynomial (usually of degree 1) best fit is calculated, getting  $\widetilde{X\alpha_{k,i}}$  and  $\widetilde{X\beta_{k,i}}$ , for series  $\{x\alpha_i\}$  and  $\{x\beta_i\}$  respectively, with  $i \leq k \leq (i + n)$  (detrended values);

in each box is calculated:  $f_{DCCA}^2(n, i) = \frac{1}{1+n} \sum_{k=i}^{i+n} (X\alpha_k - \widetilde{X\alpha_{k,i}}) \times (X\beta_k - \widetilde{X\beta_{k,i}})$

for all the boxes of a time scale, the DFA is calculated as:

$$F_{DCCA}^2(n) = \frac{1}{N-n} \sum_{i=1}^{N-n} f_{DCCA}^2(n, i);$$

for a number of different timescales ( $n$ ), with possible values  $4 \leq n \leq \frac{N}{4}$  the  $F_{DCCA}^2$  function is calculated to find a relation among  $F_{DCCA}^2 \times n$

### 3. Zebende package: implementation and optimization

## 4. Results

## 5. Summary and discussion

■ As usual ...

## References

- Peng CK, Buldyrev SV, Havlin S, Simons M, Stanley HE, Goldberger AL (1994). “Mosaic Organization of DNA Nucleotides.” **49**(2), 1685–1689.
- Podobnik B, Stanley HE (2008). “Detrended cross-correlation analysis: A new method for analyzing two nonstationary time series.” *Physical Review Letters*, **100**(8). ISSN 00319007. doi:10.1103/PhysRevLett.100.084102. 0709.0281.
- Zebende GF (2011). “DCCA cross-correlation coefficient: Quantifying level of cross-correlation.” *Physica A: Statistical Mechanics and its Applications*, **390**(4), 614–618. ISSN 03784371. doi:10.1016/j.physa.2010.10.022. URL <http://dx.doi.org/10.1016/j.physa.2010.10.022>.
- Zebende GF, Silva AM (2018). “Detrended Multiple Cross-Correlation Coefficient.” *Physica A*, **510**, 91–97. ISSN 0378-4371. doi:10.1016/j.physa.2018.06.119.

**Affiliation:**

Fernando Ferraz Ribeiro  
Universidade Federal da Bahia  
Faculty of Architecture  
Universität Innsbruck  
Universitätsstr. 15  
6020 Innsbruck, Austria  
E-mail: [fernando.ribeiro@ufba.br](mailto:fernando.ribeiro@ufba.br)  
*and*  
Centro Universitário Senai-Cimatec