

Measure Theoretic Probability Theory

Ryan Schumm

Durrett ed. 4, Problem 1.1.3

Let $\mathcal{S}_d = \{(a_1, b_1] \times \cdots \times (a_d, b_d] : a_i, b_i \in \mathbb{R}, 1 \leq i \leq d\}$. Show that $\sigma(\mathcal{S}_d) = \mathcal{R}^d$ where \mathcal{R}^d is the Borel subsets of \mathbb{R}^d .

Solution:

Let \mathcal{U} be the set of all open sets of \mathbb{R}^d and $S \in \mathcal{S}_d$. By definition, we can write

$$S = (a_1, b_1] \times \cdots \times (a_d, b_d], \quad a_i, b_i \in (-\infty, \infty).$$

Take $\eta_i > 0$ such that $a_i + \eta_i < b_i$ for all $i = 1, \dots, d$ and set

$$U = (a_1, b_1) \times \cdots \times (a_d, b_d), \tag{1}$$

$$V = [a_1 + \eta_1, b_1] \times \cdots \times [a_d + \eta_d, b_d] \tag{2}$$

It follows that $S = U \cup V \in \sigma(\mathcal{U})$ since the complement of any open set is a closed set and sigma algebras are closed under unions. By definition of a generated sigma algebra and since S was arbitrary, we have $\mathcal{S}_d \subset \sigma(\mathcal{U}) \implies \sigma(\mathcal{S}_d) \subset \sigma(\mathcal{U})$.

Now let $U \in \mathcal{U}$ and define

$$\mathcal{S} = \{S = (a_1, b_1] \times \cdots \times (a_d, b_d] : a_i, b_i \in \mathbb{Q}, S \subset U\}. \tag{3}$$

Note that \mathcal{S} is a countable subset of \mathcal{S}_d which implies that $\cup_{S \in \mathcal{S}} S \in \sigma(\mathcal{S}_d)$. For any $x \in U$, we have that

$$x \in (c_1, e_1) \times \cdots \times (c_d, e_d) \subset U \tag{4}$$

since the open sets \mathcal{U} are generated by open boxes. Since \mathbb{Q} is dense in \mathbb{R} , there exists $a_i, b_i \in \mathbb{Q}$ such that $x_i \in (a_i, b_i] \subset (c_i, e_i)$ for all $i = 1, \dots, d$. It follows that

$$x \in \cup_{S \in \mathcal{S}} S \implies U \subset \cup_{S \in \mathcal{S}} S \tag{5}$$

since x was arbitrary. We also have that $\cup_{S \in \mathcal{S}} S \subset U \implies U = \cup_{S \in \mathcal{S}} S$. Therefore, we have shown that $\mathcal{U} \subset \sigma(\mathcal{S}_d) \implies \sigma(\mathcal{U}) \subset \sigma(\mathcal{S}_d)$. Hence, $\sigma(\mathcal{S}_d) = \sigma(\mathcal{U}) = \mathcal{R}^d$.

Durrett ed. 4, Problem 1.1.4

A sigma field \mathcal{F} is said to be countably generated if there is a countable collection of subsets $\mathcal{C} \subset \mathcal{F}$ so that $\sigma(\mathcal{C}) = \mathcal{F}$. Show that \mathcal{R}^d is a countably generated.

Solution:

Let the countable set \mathcal{C} be given by

$$\mathcal{C} = \{(a_1, b_1] \times \cdots \times (a_d, b_d] : a_i, b_i \in \mathbb{Q}\} \quad (6)$$

and observe that $\mathcal{C} \subset \mathcal{S}_d \implies \sigma(\mathcal{C}) \subset \sigma(\mathcal{S}_d) = \mathcal{R}^d$. In the previous problem, we showed that $U = \cup_{S \in \mathcal{S}} S \in \mathcal{C}$ for all $U \in \mathcal{U}$ where $\mathcal{S} \subset \mathcal{C}$ is defined in (3). It follows that $\mathcal{U} \subset \mathcal{C} \implies \sigma(\mathcal{U}) = \mathcal{R}^d \subset \sigma(\mathcal{C})$. Thus, we have shown that \mathcal{R}^d is generated by the countable set \mathcal{C} .