

Kolmogorov Complexity and Diophantine Approximation

Jan Reimann

Institut für Informatik
Universität Heidelberg

Algorithmic Information Theory

- Algorithmic Information Theory classifies objects by their **descriptive complexity**.

Algorithmic Information Theory

- Algorithmic Information Theory classifies objects by their **descriptive complexity**.
- **Examples:**
 - A sequence of 10000 zeroes has low complexity, as there is a much shorter description.
 - Any (infinite) sequence computable by an algorithm such as π has a short description (the algorithm).
 - An outcome of an (unbiased) coin toss has high complexity. (There is no shorter description than the sequence itself.)

Algorithmic Information Theory

- Algorithmic Information Theory classifies objects by their **descriptive complexity**.
- **Examples:**
 - A sequence of 10000 zeroes has low complexity, as there is a much shorter description.
 - Any (infinite) sequence computable by an algorithm such as π has a short description (the algorithm).
 - An outcome of an (unbiased) coin toss has high complexity. (There is no shorter description than the sequence itself.)
- **Rigorous Formulation:** **Kolmogorov complexity**.

Kolmogorov Complexity

- **Kolmogorov complexity:** U a universal Turing-machine. Def. for a binary string $\sigma \in \{0, 1\}^*$,

$$C(\sigma) = C_U(\sigma) = \min\{|p| : p \in \{0, 1\}^*, U(p) = \sigma\},$$

i.e. $C(\sigma)$ is the length of the shortest program (for U) that outputs σ .

Kolmogorov Complexity

- **Kolmogorov complexity:** U a universal Turing-machine. Def. for a binary string $\sigma \in \{0, 1\}^*$,

$$C(\sigma) = C_U(\sigma) = \min\{|p| : p \in \{0, 1\}^*, U(p) = \sigma\},$$

i.e. $C(\sigma)$ is the length of the shortest program (for U) that outputs σ .

- **Kolmogorov's invariance theorem:** C is independent of U (up to a constant).

Kolmogorov Complexity

- **Kolmogorov complexity:** U a universal Turing-machine. Def. for a binary string $\sigma \in \{0, 1\}^*$,

$$C(\sigma) = C_U(\sigma) = \min\{|p| : p \in \{0, 1\}^*, U(p) = \sigma\},$$

i.e. $C(\sigma)$ is the length of the shortest program (for U) that outputs σ .

- **Kolmogorov's invariance theorem:** C is independent of U (up to a constant).
- The **pigeonhole principle** yields that for any length there are **incompressible** strings, $C(\sigma) \geq |\sigma|$ (in fact, most of them are).

Digression: Euclid Revisited

- Suppose there are only finitely many primes p_1, \dots, p_n .

Digression: Euclid Revisited

- Suppose there are only finitely many primes p_1, \dots, p_n .
- Then each number N can be factored

$$N = p_1^{k_1} \cdots p_n^{k_n}, \text{ each } k_i \leq \log N.$$

Digression: Euclid Revisited

- Suppose there are only finitely many primes p_1, \dots, p_n .
- Then each number N can be factored

$$N = p_1^{k_1} \cdots p_n^{k_n}, \text{ each } k_i \leq \log N.$$

- Therefore, N can be described by a program of length $O(n \log \log N)$. (Identify natural numbers with their binary representation).

Digression: Euclid Revisited

- Suppose there are only finitely many primes p_1, \dots, p_n .
- Then each number N can be factored

$$N = p_1^{k_1} \cdots p_n^{k_n}, \text{ each } k_i \leq \log N.$$

- Therefore, N can be described by a program of length $O(n \log \log N)$. (Identify natural numbers with their binary representation).
- Yields a contradiction if N is incompressible ($C(N) \geq \log N$).

Kolmogorov complexity and Coding

- A **prefix-free Turing machine** is a TM with **prefix-free domain**. The prefix-free version of C (use only prefix free TMs) is denoted by K .

Kolmogorov complexity and Coding

- A **prefix-free Turing machine** is a TM with **prefix-free domain**. The prefix-free version of C (use only prefix free TMs) is denoted by K .
- **Kraft-Chaitin Theorem:** $\{\sigma_i\}_{i \in \mathbb{N}}$ set of strings, $\{l_i, l_2, \dots\}$ sequence of natural numbers ('lengths') such that

$$\sum_{i \in \mathbb{N}} 2^{-l_i} \leq 1,$$

then one can construct (primitive recursively) a prefix-free TM M and strings $\{\tau_i\}_{i \in \mathbb{N}}$, such that

$$|\tau_i| = l_i \quad \text{and} \quad M(\tau_i) = \sigma_i.$$

Kolmogorov complexity and Coding

- **Semimeasures:** $m : \{0, 1\}^* \rightarrow [0, \infty)$ with

$$\sum_{\sigma \in \{0, 1\}^*} m(\sigma) \leq 1.$$

Kolmogorov complexity and Coding

- **Semimeasures:** $m : \{0, 1\}^* \rightarrow [0, \infty)$ with

$$\sum_{\sigma \in \{0, 1\}^*} m(\sigma) \leq 1.$$

- There exists a **maximal enumerable semimeasure** \tilde{m} , i.e. \tilde{m} is enumerable from below, and for any enumerable semimeasure m it holds that $m \leq c_m \tilde{m}$ for some constant c_m .

Kolmogorov complexity and Coding

- **Semimeasures:** $m : \{0, 1\}^* \rightarrow [0, \infty)$ with

$$\sum_{\sigma \in \{0, 1\}^*} m(\sigma) \leq 1.$$

- There exists a **maximal enumerable semimeasure** \tilde{m} , i.e. \tilde{m} is enumerable from below, and for any enumerable semimeasure m it holds that $m \leq c_m \tilde{m}$ for some constant c_m .
- **Coding Theorem:** (Zvonkin-Levin)

$$K(\sigma) = -\log \tilde{m}(\sigma) + c.$$

Kolmogorov complexity and Coding

- **Semimeasures:** $m : \{0, 1\}^* \rightarrow [0, \infty)$ with

$$\sum_{\sigma \in \{0,1\}^*} m(\sigma) \leq 1.$$

- There exists a **maximal enumerable semimeasure** \tilde{m} , i.e. \tilde{m} is enumerable from below, and for any enumerable semimeasure m it holds that $m \leq c_m \tilde{m}$ for some constant c_m .
- **Coding Theorem: (Zvonkin-Levin)**

$$K(\sigma) = -\log \tilde{m}(\sigma) + c.$$

- Identify **randomness** with **incompressibility**: Say an infinite binary sequence ξ is **random** if for all n , $K(\xi \upharpoonright_n) \geq n - c$, for some constant c .

Diophantine Approximation

- Diophantine Approximation classifies real numbers by how well they may be approximated by rational numbers.

Diophantine Approximation

- Diophantine Approximation classifies real numbers by how well they may be approximated by rational numbers.
- Fundamental **Theorem** by **Dirichlet**: For any irrational α there exist infinitely many p/q (rel. prime) such that

$$|\alpha - p/q| \leq 1/q^2. \quad (3)$$

Diophantine Approximation

- Diophantine Approximation classifies real numbers by how well they may be approximated by rational numbers.
- Fundamental **Theorem** by **Dirichlet**: For any irrational α there exist infinitely many p/q (rel. prime) such that

$$|\alpha - p/q| \leq 1/q^2. \quad (5)$$

- Such a series of rationals is obtained by the **continued fraction expansion**:

$$\alpha = [a_0, a_1, \dots] = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}} \quad a_i \in \mathbb{N} \quad (6)$$

Diophantine Approximation

- Diophantine Approximation classifies real numbers by how well they may be approximated by rational numbers.
- Fundamental **Theorem** by **Dirichlet**: For any irrational α there exist infinitely many p/q (rel. prime) such that

$$|\alpha - p/q| \leq 1/q^2. \quad (7)$$

- Such a series of rationals is obtained by the **continued fraction expansion**:

$$\alpha = [a_0, a_1, \dots] = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}} \quad a_i \in \mathbb{N} \quad (8)$$

- The expansion is **finite** if and only if α is **rational**.

Diophantine Approximation

- In general, one cannot improve the factor 2 in Dirichlet's theorem.

Diophantine Approximation

- In general, one cannot improve the factor 2 in Dirichlet's theorem.
- A number β is **badly approximable** if there exists a K such that

$$\forall p/q \in \mathbb{Q} \quad |\beta - p/q| \geq K/q^2.$$

Diophantine Approximation

- In general, one cannot improve the factor 2 in Dirichlet's theorem.
- A number β is **badly approximable** if there exists a K such that

$$\forall p/q \in \mathbb{Q} \quad |\beta - p/q| \geq K/q^2.$$

- **Examples of badly approximable numbers:**
 - Golden mean $(1 + \sqrt{5})/2 = [1, 1, 1, 1, \dots]$. $0 < K < \sqrt{5}$.
 - $\sqrt{2}/2 = [1, 2, 2, 2, \dots]$, in fact all irrational square roots.
 - $e \bmod 1 = [1, 2, 1, 1, 4, 1, 1, 6, \dots]$ not badly approximable.
 - $\pi \bmod 1 = [7, 15, 1, 292, 1, 1, \dots]$???

Diophantine Approximation

- Algebraic numbers are close to badly approximable:
Roth's Theorem: For any algebraic α , for any $\varepsilon > 0$,

$$\left| \alpha - \frac{p}{q} \right| \leq \frac{1}{q^{2+\varepsilon}} \quad (9)$$

has only finitely many solutions.

Diophantine Approximation

- Algebraic numbers are close to badly approximable:
Roth's Theorem: For any algebraic α , for any $\varepsilon > 0$,

$$\left| \alpha - \frac{p}{q} \right| \leq \frac{1}{q^{2+\varepsilon}} \quad (10)$$

has only finitely many solutions.

- However, there are numbers which are **very well approximable**.

A **Liouville number** is an irrational α for which

$$\forall n \exists \frac{p}{q} \left| \alpha - \frac{p}{q} \right| \leq \frac{1}{q^n}.$$

Example: $\sum 10^{-n!}$.

Metric Diophantine Approximation

- What's the situation **typically**? Are most numbers rather well or rather badly approximable?

Metric Diophantine Approximation

- What's the situation **typically**? Are most numbers rather well or rather badly approximable?
- For almost all numbers, the exponent 2 cannot be improved much:

Khintchine's Theorem: Let $\psi : \mathbb{N} \rightarrow \mathbb{R}^+$ be such that $\lim_n \psi(n) = 0$. Define

$$W_\psi = \{\alpha : \exists^\infty (p/q) \mid |\alpha - (p/q)| < \psi(q)\}.$$

Then it holds, for Lebesgue measure λ ,

$$\lambda W_\psi = \begin{cases} 0, & \text{if } \sum k\psi(k) < \infty, \\ 1, & \text{if } \sum k\psi(k) = \infty. \end{cases}$$

Metric Diophantine Approximation

- What's the situation **typically**? Are most numbers rather well or rather badly approximable?
- For almost all numbers, the exponent 2 cannot be improved much:

Khintchine's Theorem: Let $\psi : \mathbb{N} \rightarrow \mathbb{R}^+$ be such that $\lim_n \psi(n) = 0$. Define

$$W_\psi = \{\alpha : \exists^\infty (p/q) \mid |\alpha - (p/q)| < \psi(q)\}.$$

Then it holds, for Lebesgue measure λ ,

$$\lambda W_\psi = \begin{cases} 0, & \text{if } \sum k\psi(k) < \infty, \\ 1, & \text{if } \sum k\psi(k) = \infty. \end{cases}$$

- So, well-approximable numbers are rather rare. Can we tell **how rare**?

Hausdorff Measures

- **Caratheodory-Hausdorff construction** on metric spaces: let $A \subseteq X$, X some separable metric space, $h : \mathbb{R} \rightarrow \mathbb{R}$ a monotone, increasing, continuous on the right function with $h(0) = 0$, and let $\delta > 0$.

Hausdorff Measures

- **Caratheodory-Hausdorff construction** on metric spaces: let $A \subseteq X$, X some separable metric space, $h : \mathbb{R} \rightarrow \mathbb{R}$ a monotone, increasing, continuous on the right function with $h(0) = 0$, and let $\delta > 0$.
- Define a set function

$$\mathcal{H}_\delta^h(A) = \inf \left\{ \sum_i h(\text{diam}(U_i)) : A \subseteq \bigcup_i U_i, \text{diam}(U_i) \leq \delta \right\}.$$

Hausdorff Measures

- **Caratheodory-Hausdorff construction** on metric spaces: let $A \subseteq X$, X some separable metric space, $h : \mathbb{R} \rightarrow \mathbb{R}$ a monotone, increasing, continuous on the right function with $h(0) = 0$, and let $\delta > 0$.
- Define a set function

$$\mathcal{H}_\delta^h(A) = \inf \left\{ \sum_i h(\text{diam}(U_i)) : A \subseteq \bigcup_i U_i, \text{diam}(U_i) \leq \delta \right\}.$$

- Letting $\delta \rightarrow 0$ yields an (outer) measure.

Hausdorff Measures

- **Caratheodory-Hausdorff construction** on metric spaces: let $A \subseteq X$, X some separable metric space, $h : \mathbb{R} \rightarrow \mathbb{R}$ a monotone, increasing, continuous on the right function with $h(0) = 0$, and let $\delta > 0$.
- Define a set function

$$\mathcal{H}_\delta^h(A) = \inf \left\{ \sum_i h(\text{diam}(U_i)) : A \subseteq \bigcup_i U_i, \text{diam}(U_i) \leq \delta \right\}.$$

- Letting $\delta \rightarrow 0$ yields an (outer) measure.
- The h -**dimensional Hausdorff measure** \mathcal{H}^h is defined as

$$\mathcal{H}^h(A) = \lim_{\delta \rightarrow 0} \mathcal{H}_\delta^h(A)$$

Properties of Hausdorff Measures

- \mathcal{H}^h is **Borel regular**:
all Borel sets are measurable and for every $Y \subseteq X$ there is a Borel set $B \subseteq Y$ such that $\mathcal{H}^h(B) = \mathcal{H}^h(Y)$.

Properties of Hausdorff Measures

- \mathcal{H}^h is **Borel regular**:
all Borel sets are measurable and for every $Y \subseteq X$ there is a Borel set $B \subseteq Y$ such that $\mathcal{H}^h(B) = \mathcal{H}^h(Y)$.
- An obvious choice for h is $h(x) = x^s$ for some $s \geq 0$. For such h , denote the corresponding Hausdorff measure by \mathcal{H}^s .

Properties of Hausdorff Measures

- \mathcal{H}^h is **Borel regular**:
all Borel sets are measurable and for every $Y \subseteq X$ there is a Borel set $B \subseteq Y$ such that $\mathcal{H}^h(B) = \mathcal{H}^h(Y)$.
- An obvious choice for h is $h(x) = x^s$ for some $s \geq 0$. For such h , denote the corresponding Hausdorff measure by \mathcal{H}^s .
- For $s = 1$, \mathcal{H}^1 is the usual Lebesgue measure λ on $2^{\mathbb{N}}$.

Properties of Hausdorff Measures


- \mathcal{H}^h is **Borel regular**:
all Borel sets are measurable and for every $Y \subseteq X$ there is a Borel set $B \subseteq Y$ such that $\mathcal{H}^h(B) = \mathcal{H}^h(Y)$.
- An obvious choice for h is $h(x) = x^s$ for some $s \geq 0$. For such h , denote the corresponding Hausdorff measure by \mathcal{H}^s .
- For $s = 1$, \mathcal{H}^1 is the usual Lebesgue measure λ on $2^{\mathbb{N}}$.
- For $0 \leq s < t < \infty$ and $Y \subseteq X$,

$$\mathcal{H}^s(Y) < \infty \text{ implies } \mathcal{H}^t(Y) = 0,$$

$$\mathcal{H}^t(Y) > 0 \text{ implies } \mathcal{H}^s(Y) = \infty.$$


Hausdorff dimension

- The situation is depicted in the following graph:



Hausdorff dimension

- The situation is depicted in the following graph:



- The Hausdorff dimension of A is

$$\begin{aligned}\dim_H(A) &= \inf\{s \geq 0 : \mathcal{H}^s(A) = 0\} \\ &= \sup\{t \geq 0 : \mathcal{H}^t(A) = \infty\}\end{aligned}$$

Hausdorff Dimension and Diophantine Approximation

- Let

$$W_\delta = W_\psi \text{ for } \psi(q) = \frac{1}{q^\delta}.$$

Hausdorff Dimension and Diophantine Approximation

- Let

$$W_\delta = W_\psi \text{ for } \psi(q) = \frac{1}{q^\delta}.$$

- So, the Liouville numbers are just the ones contained in every W_n .

Hausdorff Dimension and Diophantine Approximation

- Let

$$W_\delta = W_\psi \text{ for } \psi(q) = \frac{1}{q^\delta}.$$

- So, the Liouville numbers are just the ones contained in every W_n .
- **Jarnik-Besicovitch Theorem:** For $\delta \geq 2$,

$$\dim_H W_\delta = 2/\delta.$$

Hausdorff Dimension and Diophantine Approximation

- Let

$$W_\delta = W_\psi \text{ for } \psi(q) = \frac{1}{q^\delta}.$$

- So, the Liouville numbers are just the ones contained in every W_n .
- **Jarnik-Besicovitch Theorem:** For $\delta \geq 2$,

$$\dim_H W_\delta = 2/\delta.$$

- In particular, the set of **Liouville numbers** has dimension zero.

Random Continued Fractions

- Other questions: What does a **typical** continued fraction look like? Do the components reflect some approximation properties?

Random Continued Fractions

- Other questions: What does a **typical** continued fraction look like? Do the components reflect some approximation properties?
- Interesting from a different point of view: What is the **complexity** of a **function from natural numbers to natural numbers**?

Random Continued Fractions

- Other questions: What does a **typical** continued fraction look like? Do the components reflect some approximation properties?
- Interesting from a different point of view: What is the **complexity** of a **function from natural numbers to natural numbers**?
- In the following, identify an initial segment $[a_1, \dots, a_n]$ of a continued fraction with the **n-convergent**

$$\frac{p_n}{q_n} = \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots \frac{1}{a_n}}}}.$$

Random Continued Fractions

- Using the **ergodic theorem**, one can show, for instance, that

$$\frac{1}{n} \log q_n \longrightarrow \frac{\pi^2}{12 \log 2}$$

for almost every α .

Random Continued Fractions

- Using the **ergodic theorem**, one can show, for instance, that

$$\frac{1}{n} \log q_n \longrightarrow \frac{\pi^2}{12 \log 2}$$

for almost every α .

- Entropy of the **Gauss map** $x \mapsto \frac{1}{x} \bmod 1$.

Measure on Baire Space

- **Basic open cylinders:** convergents of continued fractions, $[a_0, \dots a_n]$.

Measure on Baire Space

- **Basic open cylinders**: convergents of continued fractions, $[a_0, \dots, a_n]$.
- The **diameter** of these (as a subset of $[0, 1]$) can be determined:

$$\text{diam}[a_0, \dots, a_n] = \frac{1}{q_n(q_n + q_{n-1})}$$

Measure on Baire Space

- **Basic open cylinders**: convergents of continued fractions, $[a_0, \dots a_n]$.
- The **diameter** of these (as a subset of $[0, 1]$) can be determined:

$$\text{diam}[a_0, \dots a_n] = \frac{1}{q_n(q_n + q_{n-1})}$$

- Gives rise to a Borel measure in the usual way (extension from algebra to σ -algebra). Equivalently, Hausdorff measures.

Effective (Hausdorff) measure

- Introduce effective coverings and define the notion of effective \mathcal{H}^s -measure 0.

Effective (Hausdorff) measure

- Introduce effective coverings and define the notion of **effective \mathcal{H}^s -measure** 0.
- Let $X = 2^{\mathbb{N}}$ or $X = \mathbb{N}^{\mathbb{N}}$, let $s \geq 0$ be rational. $A \subseteq X$ is Σ_1 - \mathcal{H}^s **null**, Σ_1^0 - $\mathcal{H}^s(A) = 0$, if there is a recursive sequence (C_n) of enumerable sets such that for each n ,

$$A \subseteq \bigcup_{w \in C_n} [w] \quad \text{and} \quad \sum_{w \in C_n} \text{diam}[w] < 2^{-n}.$$

Effective (Hausdorff) measure

- Introduce effective coverings and define the notion of **effective \mathcal{H}^s -measure** 0.
- Let $X = 2^{\mathbb{N}}$ or $X = \mathbb{N}^{\mathbb{N}}$, let $s \geq 0$ be rational. $A \subseteq X$ is Σ_1 - \mathcal{H}^s **null**, Σ_1^0 - $\mathcal{H}^s(A) = 0$, if there is a recursive sequence (C_n) of enumerable sets such that for each n ,

$$A \subseteq \bigcup_{w \in C_n} [w] \quad \text{and} \quad \sum_{w \in C_n} \text{diam}[w] < 2^{-n}.$$

- For $s = 1$, one obtains an **effective version of Lebesgue measure** λ .

Effective (Hausdorff) measure

- Introduce effective coverings and define the notion of **effective \mathcal{H}^s -measure** 0.
- Let $X = 2^{\mathbb{N}}$ or $X = \mathbb{N}^{\mathbb{N}}$, let $s \geq 0$ be rational. $A \subseteq X$ is Σ_1 - \mathcal{H}^s **null**, Σ_1^0 - $\mathcal{H}^s(A) = 0$, if there is a recursive sequence (C_n) of enumerable sets such that for each n ,

$$A \subseteq \bigcup_{w \in C_n} [w] \quad \text{and} \quad \sum_{w \in C_n} \text{diam}[w] < 2^{-n}.$$

- For $s = 1$, one obtains an **effective version of Lebesgue measure** λ .
- **Theorem:** (**Schnorr**) A sequence $\xi \in 2^{\mathbb{N}}$ is random if and only if $\{\xi\}$ is not Σ_1 - λ **null**.

Random Continued Fractions

- Does a similar characterization hold for continued fractions?

Random Continued Fractions

- Does a similar characterization hold for continued fractions?
- **Theorem:** (Reimann, Gacs) A continued fraction α is not Σ_1 - λ -null if and only if

$$\sup_n \{-K(\langle a_1, \dots, a_n \rangle) - \log \lambda[a_1, \dots, a_n]\} < \infty$$

Random Continued Fractions

- Is the real represented by a random binary sequence (via its dyadic expansion) also a random continued fraction (in terms of Σ_1 -measure)?

Random Continued Fractions

- Is the real represented by a **random binary sequence** (via its dyadic expansion) also a **random continued fraction** (in terms of Σ_1 -measure)?
- Problem: The continued fraction expansion might code things more efficiently.

Random Continued Fractions

- Surprisingly, it does not.

Theorem: (Lochs) For $\xi \in 2^{\mathbb{N}}$, denote by $\pi_n(\xi)$ the number of partial convergents a_i of the continued fraction expansion of ξ obtained from the first n digits of ξ . Then it holds for almost every ξ ,

$$\lim_{n \rightarrow \infty} \frac{|\pi_n(\xi)|}{n} = \frac{6 \log^2 2}{\pi^2}.$$

Random Continued Fractions

- Surprisingly, it does not.

Theorem: (Lochs) For $\xi \in 2^{\mathbb{N}}$, denote by $\pi_n(\xi)$ the number of partial convergents a_i of the continued fraction expansion of ξ obtained from the first n digits of ξ . Then it holds for almost every ξ ,

$$\lim_{n \rightarrow \infty} \frac{|\pi_n(\xi)|}{n} = \frac{6 \log^2 2}{\pi^2}.$$

- Lochs' Theorem **holds effectively**, that is, for any random ξ .

Random Continued Fractions

- Surprisingly, it does not.

Theorem: (Lochs) For $\xi \in 2^{\mathbb{N}}$, denote by $\pi_n(\xi)$ the number of partial convergents a_i of the continued fraction expansion of ξ obtained from the first n digits of ξ . Then it holds for almost every ξ ,

$$\lim_{n \rightarrow \infty} \frac{|\pi_n(\xi)|}{n} = \frac{6 \log^2 2}{\pi^2}.$$

- Lochs' Theorem **holds effectively**, that is, for any random ξ .
- Proof requires some effort to avoid the use of the ergodic theorem (which is not an effective law of probability).

Random Continued Fractions

- Every random continued fraction satisfies

$$\frac{1}{n} \log q_n \longrightarrow \frac{\pi^2}{12 \log 2}.$$

Random Continued Fractions

- Every random continued fraction satisfies

$$\frac{1}{n} \log q_n \longrightarrow \frac{\pi^2}{12 \log 2}.$$

- The proof also yields that a random c.f. $\alpha = [a_1, a_2, \dots]$ must have arbitrary large partial convergents a_i .

Random Continued Fractions

- Every random continued fraction satisfies

$$\frac{1}{n} \log q_n \longrightarrow \frac{\pi^2}{12 \log 2}.$$

- The proof also yields that a random c.f. $\alpha = [a_1, a_2, \dots]$ must have arbitrary large partial convergents a_i .
- **Theorem:** An irrational α is badly approximable if and only if its continued fraction expansion is bounded.

Random Continued Fractions

- Every random continued fraction satisfies

$$\frac{1}{n} \log q_n \longrightarrow \frac{\pi^2}{12 \log 2}.$$

- The proof also yields that a random c.f. $\alpha = [a_1, a_2, \dots]$ must have arbitrary large partial convergents a_i .
- **Theorem:** An irrational α is **badly approximable** if and only if its **continued fraction expansion** is **bounded**.
- Hence, from our point of view, badly approximable numbers must be **compressible**.

Effective Hausdorff Dimension

- Based on effective measure, we can also define **effective Hausdorff dimension**:

$$\dim_1 A = \inf\{s \geq 0 : \Sigma_1^0\text{-}\mathcal{H}^s A = 0\}.$$

Effective Hausdorff Dimension

- Based on effective measure, we can also define **effective Hausdorff dimension**:

$$\dim_1 A = \inf\{s \geq 0 : \Sigma_1^0\text{-}\mathcal{H}^s A = 0\}.$$

- Yields a notion of Hausdorff dimension for **individual sequences**.

Effective Hausdorff Dimension

- Based on effective measure, we can also define **effective Hausdorff dimension**:

$$\dim_1 A = \inf\{s \geq 0 : \Sigma_1^0\text{-}\mathcal{H}^s A = 0\}.$$

- Yields a notion of Hausdorff dimension for **individual sequences**.
- It turns out that the effective dimension of a set is determined by the **dimension of its most complex members**:

$$\dim_1 A = \sup_{\xi \in A} \dim_1 \xi.$$

Effective Hausdorff Dimension

- Based on effective measure, we can also define **effective Hausdorff dimension**:

$$\dim_1 A = \inf\{s \geq 0 : \Sigma_1^0\text{-}\mathcal{H}^s A = 0\}.$$

- Yields a notion of Hausdorff dimension for **individual sequences**.
- It turns out that the effective dimension of a set is determined by the **dimension of its most complex members**:

$$\dim_1 A = \sup_{\xi \in A} \dim_1 \xi.$$

- Theorem:** For any sequence $\xi \in 2^{\mathbb{N}}$ it holds that

$$\dim_1 \xi = \liminf_{n \rightarrow \infty} \frac{K(\xi \upharpoonright_n)}{n}.$$

Badly approximable numbers

- Badly approximable numbers **are** compressible.

Badly approximable numbers

- Badly approximable numbers **are** compressible.
- **Jarnik's Theorem:** Let

$$E_n = \{\alpha = [a_0, a_1, \dots] \in [0, 1] : \forall i \ a_i \leq n\}.$$

Then, for $n \geq 8$, $1 - \frac{4}{n \log 2} \leq \dim_H E_n \leq 1 - \frac{1}{8n \log n}$.

Badly approximable numbers

- Badly approximable numbers **are** compressible.
- **Jarnik's Theorem:** Let

$$E_n = \{\alpha = [a_0, a_1, \dots] \in [0, 1] : \forall i \ a_i \leq n\}.$$

Then, for $n \geq 8$, $1 - \frac{4}{n \log 2} \leq \dim_H E_n \leq 1 - \frac{1}{8n \log n}$.

- Later, **Bumby** and **Hensley** gave good approximations of $\dim_H E_n$ for smaller values of n , e.g. $\dim_H E_2 = 0.5312050 \dots$.

Badly approximable numbers

- Badly approximable numbers **are** compressible.
- **Jarnik's Theorem:** Let

$$E_n = \{\alpha = [a_0, a_1, \dots] \in [0, 1] : \forall i \ a_i \leq n\}.$$

Then, for $n \geq 8$, $1 - \frac{4}{n \log 2} \leq \dim_H E_n \leq 1 - \frac{1}{8n \log n}$.

- Later, **Bumby** and **Hensley** gave good approximations of $\dim_H E_n$ for smaller values of n , e.g. $\dim_H E_2 = 0.5312050 \dots$.
- Using complexity theoretic characterization of dimension, we can give a simpler, essentially combinatorial proof.