


What is the reproductive system?


The reproductive systems in both the male and female consist of primary and secondary sex organs and sex glands.




The primary function of the reproductive systems is to perpetuate the species through sexual or germ cell fertilization and reproduction.

Female reproductive system: the uterus (womb).

The uterus is a hollow, muscular, pear-shaped organ about the size of a woman's clenched fist. The top is tipped forward in a normal 'anteflexion' position. It can be divided into the body or corpus, and the bottom cervix. The rounded top portion, above the fallopian tubes, is called the fundus.




Female reproductive system: the uterus (womb)






The uterus is supported and held in position by a number of ligaments. Trauma, disease, or multiple pregnancies can weaken these ligaments and result in abnormal positioning.



the uterus...

The uterus wall has 3 layers: the outside layer called the perimetrium, the muscular inside layer called the myometrium, and the mucous membrane lining the uterus called the endometrium.



The cervical opening is about the diameter of a pencil.

the uterus...

The uterus has 3 functions:

- 1. The endometrium sheds the lining of the uterus every 21 to 40 days by menstruation
- 2. It provides a place for the protection and nourishment of the fetus during pregnancy



3. It contracts during labor to expel the fetus

Female reproductive system: the fallopian tubes...

Tube widens to form the ampulla (am pyu lah) <


The isthmus is the portion that connects to uterus


Fimbria (fihm bree ah) are the finger-like projections around the opening that trap the egg as it leaves the ovary

End of tube is called the infundibulum (in fun DIB yū lum)

Opening is called the ostium (ah stē um)

fallopian tubes...






The fallopian tube is 4-6 inches long. The egg, released from the ovary, is captured by the fimbria and brought into the fallopian tube. The egg is moved along inside the tube by muscular contractions and the waving action of cilia. It takes an egg about 3-4 days to travel the length of the tube. If an egg is fertilized, it occurs here.

the ovaries... The two ovaries are attached to each side of the uterus by a ligament. They are oval-shaped, about the size of a large olive, and lie close to the fimbria at the end of the fallopian

tubes.




Each ovary is filled, already at birth, with egg-containing sacs called follicles. Each egg is called an ovum.

Produces mature eggs and female sex hormones (estrogen and progesterone)

ovaries and ova


Once every 21 days, one follicle in one ovary ripens. This mature follicle is a graafian (GRAW fee un) follicle. The follicle ruptures in response to hormones from the pituitary gland, releasing the second oocyte... a process called ovulation.



After the follicle ruptures, it becomes a mass of yellow cells called the corpus luteum. This is a temporary, progesterone-producing structure.

vagina...

The vagina extends from the uterus to the outside of the body. It is a 3 ½ inch long muscular tube that expands in length and width during sexual arousal.




The vagina is the female organ for copulation (sexual intercourse), receiving the seminal fluid from the male penis. It is also a passageway for menstruation or the birth of a fetus.

vulva...

The vulva is 5 organs making up the external genitalia of the female:

1. Mons pubis: triangular-shaped pad of fatty tissue over the pubis bone, covered with pubic hair



- 2. Labia majora: 2 large folds of adipose tissue on the sides of the vaginal opening
- 3. Labia minora: 2 smaller folds of adipose tissue on the inside of the labia majora
- 4. Vestibule: area between labia with openings for the vagina, urethra, and two excretory ducts for Bartholin's glands (provide lubricant)
- 5. Clitoris: sensitive fold of tissue partially covered by hood

vulva...


The perineum is the area located between the vaginal opening and the anus. It is a muscular sheet that can be torn during childbirth.





Some doctors avoid uncontrolled tearing of the perineum by making a surgical incision called an episiotomy.


breasts...



The breasts are the mammary glands, varying in size according to age, heredity, and the amount of fatty tissue present. Each breast has 15-20 glandular lobes separated by connective tissue. After childbirth, the pituitary gland stimulates these lobules with the hormone prolactin... and they produce milk.


breasts...

The dark-colored circle at the tip of the breast is called the areola (ah REE ah lah). It contains sebaceous glands to keep the skin conditioned. In the center is the nipple, where ducts from the lobules open. The first secretion from the breast is not a true milk, but a thin yellowish substance called colostrum.



Colostrum contains nutrients and the mother's immunities that can protect baby.


Female reproductive system: the menstrual cycle...



the process of sloughing off the inner lining of the uterus, once monthly

The onset of the menstrual cycle, or menarche, begins at puberty and ceases at menopause.

Female reproductive system: the menstrual cycle...




The cycle has 3 phases:

Follicular Phase
Ovulatory Phase
Luteal Phase


The follicular phase is characterized by menstruation, when the thickened lining of the endometrium is shed because no egg was fertilized or implanted.

Female reproductive system: the menstrual cycle...


Ovulatory Phase




Luteal Phase



The fertilized egg implants in the uterus. **Progesterone** production increases to signal a pregnancy; it and blood.




The resulting cell, a zygote, now has 46 chromosomes. The cell begins dividing and is also can be detected in urine called a blastocyst. From week 2-8, it is called an embryo.



The AMNIOTIC sac is fluid-filled bag that protects and supports the embryo during the 280 day gestation period. The umbilical cord attached at the navel connects it to the placenta, where it gets nutrients and oxygen. During the 1st trimester, all parts of the embryo are formed.

During the second trimester all parts start to function; during the last trimester the embryo is now called a fetus and the main task is growth.




The umbilical cord is cut and clamped, and placenta Caesarian is delivery through (afterbirth) is delivered following birth of the baby.

Labor, characterized by muscle contractions, dilation (to 10 cm) and effacement (thinning)of the cervix, and expulsion of the mucous plug that formed in the cervix, signals the onset of parturition... the childbirth process. The cephalic, or head-first delivery, is the most common. Breech is a backward presentation;

an incision in the abdomen.



The newborn may be covered with traces of vernix caseosa (cheesy coating) or lanugo (downy hair) that protected the skin before birth. The health of the baby is immediately evaluated on the APGAR scale. Color, heartbeat, reflexes, muscle tone, and breathing are scored on a scale of 0-10.




Pregnancy can be complicated by certain conditions:

- Placenta previa is the development of the placenta over the opening of the cervix
- 2. Preeclampsia is a pregnancy- induced hypertension (high blood pressure)
- 3. Spontaneous abortion or miscarriage is the loss of a fetus during the first 20 weeks, often due to abnormalities, trauma, or lifestyle choices.


Male reproductive system: scrotum

The scrotum is a pouch of skin that The testicle protects and holds the testes outside the body to maintain a lower temperature for sperm development.



The scrotal tissue contracts in the absence of sufficient heat, pulling the testes up closer to the body where the temperature maintains the viability of sperm.

Male reproductive system: penis




The urethra runs the length of the penis from the bladder to the meatus (me A tus). The flaccid penis fills with blood during arousal, causing an erection and allowing for the function of copulation/intercourse.

Penis delivers sperm into the vagina of the female during ejaculation. This foreskin may be circumcised/ removed for hygiene or religious reasons. It protect against foreign bodies from entering the urethra.

Male reproductive system: testes


and epididymis...

Testes produces sperm and male hormones (testosterone)




Male reproductive system: testes and epididymis...

Epididymis
where sperm are
stored
temporarily,
found on the
back of testes



Male reproductive system: sperm...




Sperm carry either an X/female OR Y/male chromosome. Since all ova carry the X/female chromosome, the male sperm does influence the baby's sex.

The acrosome (AK roh zome) COVEring the head of the sperm contains enzymes that help it penetrate the ova. The head carries the genetic material. The midpiece supplies energy. The tail or flagellum (flah JELL um) provides motility.


Male reproductive system: vas deferens and seminal vesicles...

The vas deferens are muscular tubes that conducts sperm from the testes to the urethra.



Male reproductive system: vas deferens and seminal vesicles...


The seminal vesicles produce a slightly alkaline fluid that mixes with the sperm and becomes part of the semen (a mixture of fluid produced by the male sex glands and sperm) ejaculated during the male orgasm. It protects the sperm from the acidity of the urethra.



Male reproductive system: Prostate and Cowper's glands

The prostate gland also secretes an alkaline fluid that mixes with the sperm as part of the semen. Enlargement of the prostate, benign prostatic hyperplasia, obstructs the flow of urine through the

urethra.



The Cowper's or bulbourethral glands (bull boh yur REE thral) produce another mucous-alkaline secretion in the urethra, excreted from the glands just before ejaculation.