

1 Gaussian elimination and matrices

1.1 Solving linear systems using elimination method

Recall: Common method to solve linear systems using Gaussian/Gauss-Jordan elimination:

- 1) Construct a matrix (called augmented matrix) corresponding to the system
- 2) Using *elementary row operations* on matrices to reduced the matrix to the (reduced) row-echelon form.
- 3) Using back-substitution elimination to solve.

Exercises:

1. Which of the following are linear equations in x_1, x_2 and x_3 ?

(a) $x_1 + 5x_2 - \sqrt{x_3} = 1$

(c) $\pi x_1 - \sqrt{2}x_2 + \frac{1}{3}x_3 = 7^{1/3}$

(b) $x_1^{-2} + x_2 + 8x_3 = 5$

(d) $x_1 - 3x_2 + \frac{\sqrt{4-\sqrt{32}}}{\sqrt{5}}x_3 = 2$

2. Determine coefficient matrix and augmented matrix for the following linear equation systems. Circle their pivot positions and determine their free variables. Using back-substitution to solve them.

(a)

$$\begin{aligned}x_1 - 7x_2 + 2x_3 - 5x_4 + 8x_5 &= 10 \\x_2 - 3x_3 + 3x_4 + x_5 &= -5 \\x_4 - x_5 &= 4\end{aligned}$$

(b)

$$\begin{aligned}x_1 - 2x_2 - x_3 + 3x_4 &= 0 \\3x_3 + x_4 &= 3 \\x_4 &= 5\end{aligned}$$

3. Given augmented matrices of linear equation systems. Determine which matrices are in row-echelon form. If it is not, please transform it to row-echelon form and use back-substitution method for solving the corresponding linear equation systems.

(a) $\begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$

(c) $\begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$

(b) $\begin{bmatrix} 1 & -2 & -1 & 3 & 0 \\ 0 & 0 & 3 & 1 & 3 \\ 0 & 0 & 0 & 0 & 5 \end{bmatrix}$

$$(d) \begin{bmatrix} 1 & 1 & 0 & 1 & 1 \\ 0 & 2 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 0 & 4 \end{bmatrix}$$

4. Rewriting the matrix-forms into systems of equations. Then solving the systems

(a)

$$\begin{bmatrix} 1 & 0 & 0 & 5 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 4 \end{bmatrix}$$

(c)

$$\begin{bmatrix} 1 & 6 & 0 & 0 & 4 & -2 \\ 0 & 0 & 1 & 0 & 3 & 1 \\ 0 & 0 & 0 & 1 & 5 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

(b)

$$\begin{bmatrix} 1 & 0 & 0 & 4 & -1 \\ 0 & 1 & 0 & 2 & 6 \\ 0 & 0 & 1 & 3 & 2 \end{bmatrix}$$

(d)

$$\begin{bmatrix} 1 & 2 & 0 & 3 & 0 & 7 \\ 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 2 \end{bmatrix}$$

5. Solving the following linear systems using Gaussian or Gauss-Jordan (back-substitution) elimination. Write clearly all elementary row operations you used.

(a)

$$\begin{aligned} x + y + 2z &= 0 \\ 2x + 4y - 3z &= 1 \\ 3x + 6y - 5z &= 0. \end{aligned}$$

(d)

$$\begin{aligned} x - 2y + z - 4t &= 1 \\ x + 3y + 7z + 2t &= 2 \\ x - 12y - 11z - 16t &= 5. \end{aligned}$$

(b)

$$\begin{aligned} 2x + 2y + 2z &= 0 \\ -2x + 5y + 2z &= 1 \\ 8x + y + 4z &= -1. \end{aligned}$$

(e)

$$\begin{aligned} 2I_1 - I_2 + 3I_3 + 4I_4 &= 9 \\ I_1 - 2I_3 + 7I_4 &= 11 \\ 3I_1 - 3I_2 + I_3 + 5I_4 &= 8 \\ 2I_1 + I_2 + 4I_3 + 4I_4 &= 10. \end{aligned}$$

(c)

$$\begin{aligned} x + 2y - t + w &= 1 \\ 3y + z - w &= 2 \\ z + 7t &= 1. \end{aligned}$$

(f)

$$\begin{aligned} 2u - v &= 0 \\ -u + 2v - w &= 0 \\ -v + 2w - z &= 0 \\ -w + 2z &= 5. \end{aligned}$$

6. For which values of a will the following system have no solutions? Exactly one solution? Infinitely

many solutions?

$$x + 2y - 3z = 4$$

$$3x - y + 5z = 2$$

$$4x + y + (a^2 - 14)z = a + 2.$$

1.2 Operations on matrices

Recall:

- 1) Essential matrices: Identity, zero matrix, symmetric matrix...
- 2) Operations: addition, scalar multiply, transposition, multiplication, inverse.
- 3) General properties: Associative, commutative, distributive

Exercises:

7. Suppose that A, B, C, D, E are matrices of the following sizes: $A : 4 \times 5$; $B : (4 \times 5)$; $C : 5 \times 2$; $D : 4 \times 2$; $E : 5 \times 5$. Determine (if there exists) the size of the following matrices:

- | | | |
|----------------|-------------------|--------------------|
| (a) BA ; | (d) $AB + B$; | (g) $E^t A$; |
| (b) $AC + D$; | (e) $2E(A + B)$; | |
| (c) $AE + B$; | (f) $E(AC)$; | (h) $(A^t + E)D$. |

8. Let

$$A = \begin{bmatrix} 3 & 0 \\ -1 & 2 \\ 1 & 1 \end{bmatrix}, B = \begin{bmatrix} 4 & -1 \\ 0 & 2 \end{bmatrix}, C = \begin{bmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{bmatrix}, D = \begin{bmatrix} 1 & 5 & 2 \\ -1 & 0 & 1 \\ 3 & 2 & 4 \end{bmatrix}, E = \begin{bmatrix} 6 & 1 & 3 \\ -1 & 1 & 2 \\ 4 & 1 & 3 \end{bmatrix}.$$

Compute the following (where possible)

- | | |
|---------------------------------------|-----------------------------------------|
| (a) $2B - C$; | (e) AC and CA ; |
| (b) $3D - 2E^t$; | (f) $(C^t B)A^t$ and $Tr((C^t B)A^t)$; |
| (c) $3D^t - 2E$ and $Tr(3D - 2E^t)$; | (g) $tr(DD^t)$. |
| (d) AB and BA ; | (h) $D^t E^t - (ED)^t$. |
9. Write down the 2 by 2 matrices A and B that have entries $a_{ij} = i + j$ and $b_{ij} = (-1)^{i+j}$. Multiply them to find AB and BA . Is the product of A and B commutative?
10. True or false? Give a specific counterexample when false.
- (a) If columns 1 and 3 of B are the same, so are columns 1 and 3 of AB .
 - (b) If rows 1 and 3 of B are the same, so are rows 1 and 3 of AB .
 - (c) If rows 1 and 3 of A are the same, so are rows 1 and 3 of AB .
 - (d) $(AB)^2 = A^2 B^2$.
11. Which of the following matrices are guaranteed to equal $(A + B)^2$

- (a) $A^2 + 2AB + B^2$ (c) $(A + B)(B + A)$
 (b) $A(A + B) + B(A + B)$ (d) $A^2 + AB + BA + B^2$

12. By trial and error find examples of 2 by 2 matrices such that

- (a) $A^2 = -I$, A having only real entries.
 (b) $B^2 = 0$, although $B \neq 0$;
 (c) $CD = -DC$, not allowing the case $CD = 0$.
 (d) $EF = 0$, although no entries of E or F are zero.

13. Suppose A commutes with every 2 by 2 matrix (that is $AB = BA$), and in particular

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, B_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, B_2 = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix},$$

Show that $a = d$ and $b = c = 0$. Consequently, prove that if $AB = BA$ for all matrices B , then A is a multiple of the identity.

14. In each part find matrices A, X, B which express the given system of linear equations as a single matrix equation $AX = B$. Solve those equations.

(a)

$$\begin{aligned} x_1 - 3x_2 + 5x_3 &= 7 \\ 9x_1 - x_2 + x_3 &= -1 \\ x_1 + 5x_2 + 4x_3 &= 0 \end{aligned}$$

(b)

$$\begin{aligned} x_1 - 3x_3 + x_4 &= 7 \\ 5x_1 + x_2 - 8x_4 &= 3 \\ 2x_1 - 5x_2 + 9x_3 - x_4 &= 0 \\ 3x_2 - x_3 + 7x_4 &= 2 \end{aligned}$$

15. Find the powers $A^2, A^3, B^2, B^3, C^2, C^3$. What are A^k, B^k and C^k for a given k ?

$$A = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{bmatrix}, B = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \text{ and } C = AB = \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{bmatrix}.$$

1.3 Inverse matrix

Recall: For finding the inverse of a matrix, we can either

- use elementary row operations to bring $[A|I]$ into $[I|A^{-1}]$, or
- use determinants and calculate adjoint matrices.

Exercise:

16. Show that if A and B are invertible matrices then

- A^t is invertible and $(A^t)^{-1} = (A^{-1})^t$
- AB are invertible and $(AB)^{-1} = B^{-1}A^{-1}$.

17. Use the Gauss-Jordan method to invert the following matrices then solve the equations $Ax = b$ for $b = (-1, 2, 7)$.

(a)

$$A_1 = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix},$$

(c)

$$A_3 = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 2 \\ 1 & 2 & 3 \end{bmatrix},$$

(e)

$$A_5 = \begin{bmatrix} \sqrt{2} & 3\sqrt{2} & 0 \\ -4\sqrt{2} & \sqrt{2} & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

(b)

$$A_2 = \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix},$$

(d)

$$A_4 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix},$$

(f)

$$A_6 = \begin{bmatrix} \frac{1}{5} & \frac{1}{5} & \frac{-2}{5} \\ \frac{1}{5} & \frac{1}{5} & \frac{1}{10} \\ \frac{-4}{5} & \frac{1}{5} & \frac{1}{10} \end{bmatrix}$$

18. True or false (with a counterexample if false and a reason if true):

- A 4 by 4 matrix with a row of zeros is not invertible.
- If A is invertible then A^{-1} is invertible
- If A^t is invertible, then A is invertible.

19. If a matrix A has row 1 + row 2 = row 3, show that A is not invertible:

- Explain why $Ax = (1, 0, 0)$ cannot have a solution.
- Which right-hand sides (b_1, b_2, b_3) might allow a solution to $Ax = b$?
- What happens to row 3 in elimination?

20. Find the inverse (in any legal way) of

(a)

$$A_1 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 2 & 0 \\ 0 & 3 & 0 & 0 \\ 4 & 0 & 0 & 0 \end{bmatrix},$$

(b)

$$A_2 = \begin{bmatrix} a & b & 0 & 0 \\ c & d & 0 & 0 \\ 0 & 0 & a & b \\ 0 & 0 & c & d \end{bmatrix}$$

(c)

$$A_3 = \begin{bmatrix} k & 0 & 0 & 0 \\ 1 & k & 0 & 0 \\ 0 & 1 & k & 0 \\ 0 & 0 & 1 & k \end{bmatrix}$$

21. For which three numbers c is this matrix not invertible, and why not?

$$A = \begin{bmatrix} 2 & c & c \\ c & c & c \\ 8 & 7 & c \end{bmatrix}$$

22. Give examples of matrices A and B such that

- (a) $A + B$ is not invertible although A and B are invertible.
- (b) $A + B$ is invertible although A and B are not invertible.
- (c) all of A , B , and $A + B$ are invertible.
- (d) In the last case use $A^{-1}(A + B)B^{-1} = B^{-1} + A^{-1}$ to show that $C = B^{-1} + A^{-1}$ is also invertible and find a formula for C

23. Show that $A^2 = 0$ is possible but $A^t A = 0$ is not possible (unless $A =$ zero matrix).

24. If the inverse of A^2 is B , show that the inverse of A is AB . Thus, A is invertible whenever A^2 is invertible.

25. If $A = A^t$ and $B = B^t$, which of these matrices are certainly symmetric?

- (a) $A^2 - B^2$;
- (b) $(A + B)(A - B)$;
- (c) ABA ;
- (d) $ABAB$.

1.4 Determinants

26. Let $A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$. Assume that $\det(A) = 7$, find

- (a) $\det(3A)$; (d) $\det A^t$;
 (b) $\det(2A^{-1})$;
 (c) $\det((2A)^{-1})$; (e) $\det A^2$;

(f) $\begin{vmatrix} a & g & d \\ b & h & e \\ c & i & f \end{vmatrix}$.

27. Prove the identity without evaluating the determinants

- (a) $\det \begin{bmatrix} b+c & c+a & b+a \\ a & b & c \\ 1 & 1 & 1 \end{bmatrix} = 0$;
 (b) $\begin{bmatrix} a_1 & b_1 & a_1+b_1+c_1 \\ a_2 & b_2 & a_2+b_2+c_2 \\ a_3 & b_3 & a_3+b_3+c_3 \end{bmatrix} = \begin{bmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{bmatrix}$;
 (c) $\begin{bmatrix} a_1+b_1 & a_1-b_1 & c_1 \\ a_2+b_2 & a_2-b_2 & c_2 \\ a_3+b_3 & a_3-b_3 & c_3 \end{bmatrix} = -2 \begin{bmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{bmatrix}$;

28. Evaluate determinants by cofactor expansion along a row or column of your choice:

- (a) $\begin{bmatrix} -3 & 0 & 7 \\ 2 & 5 & 1 \\ -1 & 0 & 5 \end{bmatrix}$ (d) $\begin{bmatrix} k+1 & k-1 & 7 \\ 2 & k-3 & 4 \\ 5 & k+1 & k \end{bmatrix}$ (f) $\begin{bmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{bmatrix}$
 (b) $\begin{bmatrix} 3 & 3 & 1 \\ 1 & 0 & -4 \\ 1 & -3 & 5 \end{bmatrix}$ (g) $\begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 \end{bmatrix}$
 (c) $\begin{bmatrix} 1 & k & k^2 \\ 1 & k & k^2 \\ 1 & k & k^2 \end{bmatrix}$ (e) $\begin{bmatrix} 3 & 3 & 0 & 5 \\ 2 & 2 & 0 & -2 \\ 4 & 1 & -3 & 0 \\ 2 & 10 & 3 & 2 \end{bmatrix}$

29. Find the inverse of the following matrices by calculating its cofactors

- (a) $\begin{bmatrix} 2 & 5 & 5 \\ -1 & -1 & 0 \\ 2 & 4 & 3 \end{bmatrix}$ (b) $\begin{bmatrix} 2 & 0 & 3 \\ 0 & 3 & 2 \\ -2 & 0 & -4 \end{bmatrix}$

30. Let $A = \begin{bmatrix} 1 & 3 & 1 & 1 \\ 2 & 5 & 2 & 2 \\ 1 & 3 & 8 & 9 \\ 1 & 3 & 2 & 2 \end{bmatrix}$.

- (a) Evaluate A^{-1} using its cofactors;

- (b) Evaluate A^{-1} using the elimination method.
31. Solving equations using Cramer's Rule

$$(a) \begin{cases} 7x_1 - 2x_2 &= 3 \\ 3x_1 + x_2 &= 5 \end{cases}$$

$$(c) \begin{cases} 2x_1 + x_2 &= 3 \\ x_1 + 2x_2 + x_3 &= 70 \\ x_2 + 2x_3 &= 0 \end{cases}$$

$$(b) \begin{cases} 4x + 5y &= 3 \\ 11x + y + 2z &= 3 \\ x + 5y + 2z &= 1 \end{cases}$$

$$(d) \begin{cases} x_1 - 3x_2 + x_3 &= 4 \\ 2x_1 - x_2 &= -2 \\ 4x_1 - 3x_3 &= 0 \end{cases}$$

32. Let $v = (3, 2)$ and $w = (1, 4)$.

- (a) Find the area of the parallelogram with edges v and w ;
- (b) Find the area of the triangle with sides v, w and $v + w$. Draw it.
- (c) Find the area of the triangle with sides v, w and $w - v$. Draw it.
- (d) The corners of a triangle are $(2, 1)$, $(3, 4)$ and $(0, 5)$. What is its area?

2 Vector spaces

2.1 Vector spaces and subspaces

33. By checking the vector space axioms, determine which sets together with the two operations (scalar and addition) defined respectively are vector spaces:
- (a) The set of all triples of real numbers (x, y, z) with the following operations $(x, y, z) + (x', y', z') = (x + x', y + y', z + z')$ and $k \cdot (x, y, z) = (kx, y, z)$;
 - (b) The set of all triples of real numbers (x, y, z) with the following operations $(x, y, z) + (x', y', z') = (x + x', y + y', z + z')$ and $k \cdot (x, y, z) = (0, 0, 0)$;
 - (c) The set of all triples of real numbers (x, y, z) with the following operations $(x_1, y_1, z_1) + (x_2, y_2, z_2) = (x_1 + x_2 + 1, y_1 + y_2 + 1, z_1 + z_2 + 1)$ and $k \cdot (x, y, z) = (kx, ky, kz)$;
 - (d) The set of all pairs of real numbers (x, y) with the following operations $(x, y) + (x', y') = (x + x', y + y')$ and $k \cdot (x, y) = (2kx, 2ky)$;
 - (e) The set of all pairs of real numbers of the form (x, y) where $x \geq 0$, with the standard operations on \mathbb{R}^2 ;
 - (f) All 2 by 2 matrices with the matrix addition and scalar multiplication;
 - (g) The set of singular 2 by 2 matrices with the matrix addition and scalar multiplication;
 - (h) The set of non-singular 2 by 2 matrices with the matrix addition and scalar multiplication;
 - (i) All non-singular matrices 2 by 2 with the matrix scalar multiplication and the addition is the matrix multiplication.
 - (j) The set of all 2 by 2 matrices of the form $\begin{bmatrix} a & 1 \\ 1 & b \end{bmatrix}$ with the matrix addition and scalar multiplications;
 - (k) The set of all 2 by 2 matrices of the form $\begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}$ with the matrix addition and scalar multiplications;
 - (l) All one variable polynomials of degree 2 with the scalar multiplication and addition are the scalar multiplication and additions in polynomials.
34. Which of the following subsets of \mathbb{R}^3 are actually subspaces?
- (a) The plane of vectors (b_1, b_2, b_3) with first component $b_1 = 0$.
 - (b) The plane of vectors b with $b_1 = 1$.
 - (c) The vectors b with $b_2 b_3 = 0$ (this is the union of two subspaces, the plane $b_2 = 0$ and the plane $b_3 = 0$).
 - (d) All combinations of two given vectors $(1, 1, 0)$ and $(2, 0, 1)$.
 - (e) The plane of vectors (b_1, b_2, b_3) that satisfy $b_3 - b_2 + 3b_1 = 0$.
35. Which of the followings are subspaces of \mathbb{R}^∞ ?
- (a) All sequences like $(1, 0, 1, 0, \dots)$ that include infinitely many zeros?

- (b) All sequences x_1, x_2, \dots with $x_j = 0$ from some point onward?
- (c) All decreasing sequences $x_{j+1} \leq x_j$ for each j ?
- (d) All convergent sequences?
- (e) All arithmetic progression $x_{j+1} - x_j$ is the same for all j ?
- (f) All geometric progression $(x_1, kx_1, k^2x_1, \dots)$ allowing all k and x_1 ?

36. Given

$$W = \{(6a - b, a + b, -7a) \in \mathbb{R}^3 : a, b \in \mathbb{R}\}.$$

- (a) Prove that W is a subspace of \mathbb{R}^3 .
- (b) Find a spanning set for W .

37. Given vectors in \mathbb{R}^3 :

$$v_1 = (1, -1, -2), \quad v_2 = (5, -4, -7), \quad v_3 = (-3, 1, 0), \quad v = (-4, 3, h).$$

For which value of h will $v \in \text{span}\{v_1, v_2, v_3\}$?

38. Which of the following descriptions are correct? The solutions x of

$$Ax = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

form

- (a) a plane;
- (b) a line;
- (c) a point;
- (d) a subspace;
- (e) the nullspace of A ;
- (f) the column space of A .

39. Describe the smallest subspace of the 2 by 2 matrix space M that contains

- (a) $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ and $\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$;
- (b) $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ and $\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$;
- (c) $\begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix}$;
- (d) $\begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix}$, $\begin{bmatrix} 0 & -1 \\ 0 & 2 \end{bmatrix}$, $\begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$.

40. Describe the null space of the matrices

$$A = \begin{bmatrix} 1 & -1 \\ 0 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 0 & 3 \\ 1 & 2 & 3 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix},$$

$$D = \begin{bmatrix} 1 & 1 & -2 \\ 1 & 1 & -2 \\ 1 & 1 & -2 \\ 1 & 1 & -2 \end{bmatrix}, \quad E = \begin{bmatrix} -4 & 6 & 1 \\ -1 & 4 & 1 \\ 5 & 6 & 7 \\ 4 & 7 & 1 \end{bmatrix}.$$

41. For which vectors (b_1, b_2, b_3) is each system below consistent?

$$(a) \begin{bmatrix} 1 & 2 & 4 \\ 0 & 3 & 6 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

$$(b) \begin{bmatrix} 1 & 4 & 2 \\ 2 & 8 & 4 \\ -1 & -4 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}.$$

$$(c) \begin{bmatrix} 1 & 4 \\ 2 & 9 \\ -1 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

2.2 Solving $Ax = 0$ and $Ax = b$

42. Using Gauss or Gauss-Jordan elimination to find the rank of each given matrix. In each case, specify which variables are free? Describe the nullspaces. Solve the complete solution of the system

$$Ax = \begin{bmatrix} 1 \\ 0 \\ 3 \end{bmatrix}.$$

(a) $\begin{bmatrix} 1 & 2 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 1 & 2 & 0 & 0 \end{bmatrix};$

(c) $\begin{bmatrix} 1 & 3 & -2 & 2 \\ -1 & -2 & -1 & -1 \\ -1 & -5 & 8 & -3 \end{bmatrix};$

(b) $\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}.$

(d) $\begin{bmatrix} 1 & 2 & 1 & 2 \\ 1 & 1 & 4 & 0 \\ -1 & -1 & -4 & 1 \end{bmatrix}.$

43. What conditions on b_1, b_2, b_3, b_4 make each system solvable? Solve for x .

$$\begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 2 & 5 \\ 3 & 9 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{bmatrix}, \quad \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 2 & 5 & 7 \\ 3 & 9 & 12 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{bmatrix}.$$

44. If $Ax = b$ has two distinct solutions x_1 and x_2 , find two solutions to $Ax = 0$. Then find another solution to $Ax = b$.
45. Suppose that you know x_p (free variables of equation $Ax = 0$) and all special solutions for $Ax = b$. Find x_p and all special solutions for these systems:

$$Ax = 2b, \quad [A \quad A] \begin{bmatrix} x \\ x \end{bmatrix} = b, \quad \begin{bmatrix} A \\ A \end{bmatrix} [x] = \begin{bmatrix} b \\ b \end{bmatrix}.$$

46. Write all known relations among r, m and n if $Ax = b$ has

- (a) no solution for some b ;
- (b) infinite many solutions for every b ;
- (c) exactly one solution for some b , no solution for other b .
- (d) exactly one solution for every b .

2.3 Linear independence, dependence, spanning, bases and dimension

47. By using the definitions, show that v_1, v_2, v_3 are linearly independent but v_1, v_2, v_3, v_4 are dependent:

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \quad v_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \quad v_4 = \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix}.$$

48. Which of the followings are bases for \mathbb{R}^3 ?

- (a) $(1, 2, 0)$ and $(0, 1, -1)$?
- (b) $(1, 1, -1), (2, 3, 4), (4, 1, -1), (0, 1, -1)$?
- (c) $(1, 2, 2), (-1, 2, 1), (0, 8, 0)$?
- (d) $(1, 2, 2), (-1, 2, 1), (0, 8, 6)$?

49. True or False

- (a) The vectors $(1, 3, 2), (2, 1, 3)$ and $(3, 2, 1)$ are independent;
- (b) The vectors $(1, -3, 2), (2, 1, -3), (-3, 2, 1)$ are independent;
- (c) The two vectors $(1, 1, -1)$ and $(-1, -1, 1)$ span \mathbb{R}^3 ; Describe the subspace spanned by two these vectors. Find its dimension.
- (d) The three vectors $(1, 1, 0), (1, 0, 1), (0, 1, 1)$ span \mathbb{R}^3 and it forms a basis of \mathbb{R}^3 ;
- (e) \mathbb{R}^3 is spanned by the columns of a 3 by 5 echelon matrix with 2 pivots?
- (f) All vectors with positive components forms a spanning system for \mathbb{R}^3 .

50. Given vectors $v_1 = (1, 0, 0)$ and $v_2 = (0, 1, 0)$ in \mathbb{R}^3 . Let $H = \{(s, s, 0) : s \in \mathbb{R}\}$.

- (a) Prove that H is a subspace of $\text{span}\{v_1, v_2\}$.
- (b) Is $\{v_1, v_2\}$ a basis for H ?
- (c) Find a basis for H .

51. Given $v_1 = (0, 2, -1), v_2 = (2, 2, 0), v_3 = (6, 16, -5)$.

- (a) Is $\{v_1, v_2, v_3\}$ linearly independent?
- (b) Find a basis for $\text{span}\{v_1, v_2, v_3\}$.

52. Find a basis for

- (a) the set of points on the line $y = -3x$
- (b) the space of solutions of $x - 3y + 2z = 0$

53. Given vectors $S = \{1 + t, 1 - t, 2\}$ in $P_1(t)$.

- (a) Prove that S is linear dependent.
- (b) Find a basis for $\text{span}(S)$.

54. Let w_1, w_2, w_3 be linearly independent vectors. Are the following vectors independent or dependent? If they are dependent, find a their non-zero linear combination that gives zero. Does the claims hold if the given vectors w_1, w_2, w_3 are dependent.

- (a) $v_1 = w_1 - w_2, v_2 = w_2 - w_3, v_3 = w_3 - w_1$;
 (b) $v_1 = w_1 + w_2, v_2 = w_2 + w_3, v_3 = w_3 + w_1$.

55. Decide whether or not the following vectors are linearly independent, by solving the equation $c_1v_1 + c_2v_2 + c_3v_3 + c_4v_4 = 0$, where

$$v_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \quad v_2 = \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \quad v_4 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix}.$$

Then decide also if they span \mathbb{R}^4 . If it not, find the largest number of independent vectors among them and find the dimension of the subspace spanned by these vectors.

56. To decide whether b is in the subspace spanned by w_1, w_2, \dots, w_m , let A be matrix having column vectors w and try to solve the equation $Ax = b$. What is the result for

- (a) $w_1 = (1, 1, 0), w_2 = (2, 2, 1), w_3 = (0, 0, 2), b = (3, 4, 5)$;
 (b) $w_1 = (1, 2, 0), w_2 = (2, 5, 0), w_3 = (0, 0, 2), w_4 = (0, 0, 0)$ and any b ?

57. For which numbers c and d do these matrices have rank 2?

$$A = \begin{bmatrix} 1 & 2 & 5 & 0 & 5 \\ 0 & 0 & c & 2 & 2 \\ 0 & 0 & 0 & d & 2 \end{bmatrix}, \quad U = \begin{bmatrix} c & d \\ d & c \end{bmatrix}.$$

58. Find a basis for each of these subspaces of \mathbb{R}^4 :

- (a) All vectors whose components are equal;
 (b) All vectors whose components add to zero.
 (c) All vectors that are perpendicular to $(1, 1, 0, 1)$ and $(1, 0, 1, 1)$.
 (d) The column space (in \mathbb{R}^2) and null space (in \mathbb{R}^5) of $U = \begin{bmatrix} 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 \end{bmatrix}$. Find their dimensions.

2.4 Linear Transformations

Recall: Let T be a map from \mathbb{R}^n to \mathbb{R}^n . T is linear transformation if T satisfies three properties:

- i) $T(0) = 0$;
- ii) $T(x + y) = T(x) + T(y)$ for any $x, y \in \mathbb{R}^n$;
- iii) $T(cx) = cT(x)$.

Exercises:

59. Which of these transformations is not linear? The input is $v = (v_1, v_2)$. Find its matrix with respect to the standard basis $(1, 0)$ and $(0, 1)$?

- (a) $T(v) = (v_2, v_1)$,
- (b) $T(v) = (v_1, v_1)$,
- (c) $T(v) = (0, v_1)$,
- (d) $T(v) = (0, 1)$,
- (e) $T(v) = (2v_1 + v_2, v_1^2)$.

60. Suppose a linear T transforms $(1, 1)$ to $(2, 2)$ and $(2, 0)$ to $(0, 0)$. Find $T(v)$ when

- (a) $v = (2, 2)$
- (b) $v = (3, 1)$
- (c) $v = (-1, 1)$
- (d) $v = (a, b)$.

61. For these transformations of $V = \mathbb{R}^2$ to $W = \mathbb{R}^2$. Find $T(T(v))$

- (a) $T(v) = -v$;
- (b) $T(v) = v + (1, 1)$;
- (c) $T(v) = 90^\circ \text{ rotation} = (-v_2, v_1)$;

62. Find the range and kernel of T :

- (a) $T(v_1, v_2) = (v_2, v_1)$;
- (b) $T(v_1, v_2, v_3) = (v_1, v_2)$;
- (c) $T(v_1, v_2) = (0, 0)$;
- (d) $T(v_1, v_2) = (v_1, v_1)$.

63. From the cubics \mathbb{P}_3 to the fourth-degree polynomials \mathbb{P}_4 , what matrix represents multiplication by $2 + 3t$? The columns of the 5 by 4 matrix A come from applying the transformation to $1, t, t^2, t^3$.

64. The space of all 2 by 2 matrices has the four basis "vectors":

$$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}.$$

For the linear transformation of "*transposing*", find its matrix A with respect to this matrix. Why is $A^2 = I$?

2.5 Reviews

Goal: Understand the equation $Ax = b$: the existence of solutions, the space of solutions (or nullspace) in case it has infinite solutions (find the dimension, the basis of the solution space); For which b the equation has a solution, the space of the vectors b such that $Ax = b$ (or the column space) has at least one solution. Find solutions of a linear equation, the inverse matrix, the dimension, the basis...

65. Find the basis and dimension for the following subspaces of \mathbb{R}^4 :

- (a) The vectors for which $x_1 = 2x_4$;
- (b) The vectors for which $x_1 + x_2 + x_3 = 0$ and $x_3 + x_4 = 0$;
- (c) The subspace spanned by $(1, 1, 1, 1), (1, 2, 3, 4), (2, 3, 4, 5)$;

66. What is the echelon form U of A ? Find the dimensions of its four fundamental subspaces?

$$\begin{bmatrix} 1 & 2 & 0 & 2 & 1 \\ -1 & -2 & 1 & 1 & 0 \\ 1 & 2 & -3 & -7 & -2 \end{bmatrix}.$$

67. (a) $Ax = b$ has a solution under what conditions on b , for the following A and b :

$$A = \begin{bmatrix} 1 & 2 & 0 & 3 \\ 0 & 0 & 0 & 0 \\ 2 & 4 & 0 & 1 \end{bmatrix} \quad b = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}.$$

- (b) Find a basis for the null-space of A ;
- (c) Find a basis for the column space of A ;
- (d) Find the rank of A^T ;
- (e) Find the general solution to $Ax = b$, when a solution exists;

68. Find the inverse of the matrix $A = \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix}$. Then solve the equation $Ax = (0, 1, 2)$.

69. Write down the matrix representation of the following linear maps relative to the natural bases of \mathbb{R}^n :

- (a) $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$ given by $T(x_1, x_2, x_3, x_4) = (x_2, x_3)$;
- (b) $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ given by $T(x_1, x_2, x_3, x_4) = (x_1 + x_2 - x_3, 0, 3x_4)$;
- (c) $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ given by $T(x) = 3x$;