

Chapter 1

Problem 9.1. The objects of **Rel** are sets, and an arrow $A \rightarrow B$ is a relation from A to B , that is, a subset $R \subseteq A \times B$. The equality relation $\{\langle a, a \rangle \in A \times A \mid a \in A\}$ is the identity arrow on a set A . Composition in **Rel** is to be given by

$$S \circ R = \{\langle a, c \rangle \in A \times C \mid \exists b (\langle a, b \rangle \in R \& \langle b, c \rangle \in S)\}$$

for $R \subseteq A \times B$ and $S \subseteq B \times C$.

Solution. (a) Show that **Rel** is a category.

Composability:

Shown above.

Identity:

Shown above.

Associativity:

Let $D \in \mathbf{Rel}$ such that $A \xrightarrow{R} B \xrightarrow{S} C \xrightarrow{T} D$. Consider $\langle a, d \rangle \in T \circ (S \circ R)$. I need to show that $\langle a, d \rangle \in (T \circ S) \circ R$. By composition, there exists a c such that $\langle a, c \rangle \in S \circ R$ and $\langle c, d \rangle \in T$. Again, by composition, there exists a b such that $\langle a, b \rangle \in R$ and $\langle b, c \rangle \in S$. Since $\langle c, d \rangle \in T$, I have that $\langle b, d \rangle \in T \circ S$. Finally, since $\langle a, b \rangle \in R$, then $\langle a, d \rangle \in (T \circ S) \circ R$.

Showing $(T \circ S) \circ R \subset T \circ (S \circ R)$ is done symmetrically.

Unit:

Let $\langle a, b \rangle \in R$. By *identity*, $\langle a, a \rangle \in 1_A$ and $\langle b, b \rangle \in 1_B$.

Hence, $\langle a, b \rangle \in R \circ 1_A, 1_B \circ R$. If $\langle a, b \rangle \in R \circ 1_A, 1_B \circ R$; then definitionally $\langle a, b \rangle \in R$. This shows that $R \circ 1_A = R = 1_B \circ R$.

(b) Show also that there is a functor $G : \mathbf{Sets} \rightarrow \mathbf{Rel}$ taking objects to themselves and each function $f : A \rightarrow B$ to its graph,

$$G(f) = \{\langle a, f(a) \rangle \in A \times B \mid a \in A\}$$

(c) Finally, show that there is a functor $C : \mathbf{Rel}^{\text{Op}} \rightarrow \mathbf{Rel}$ taking each relation $R \subseteq A \times B$ to its converse $R^c \subseteq B \times A$, where,

$$\langle a, b \rangle \in R^c \Leftrightarrow \langle b, a \rangle \in R$$

□

Problem 9.2.

Solution.

