

HUB REMOVAL IN RANDOMLY ORIENTED DIVISOR GRAPHS: THE GIANT STRONG COMPONENT IS THE SMOOTH FIBRE

[AUTHOR]

ABSTRACT. Let $\mathcal{D}_\rho(N)$ be the randomly oriented divisor graph of Kim and Phillips: the divisor graph on $\{1, \dots, N\}$, with each edge oriented from multiple to divisor and then independently reversed with probability ρ . Let Φ_K be the size of the largest strongly connected component after the vertices $1, \dots, K$ are deleted. We prove that for every fixed $\rho \in (0, 1)$,

$$0 \leq \Psi\left(N, \frac{N}{K+1}\right) - \mathbb{E}[\Phi_K] \leq \frac{30N}{\log \log N} \quad (0 \leq K < N, N \geq N_0(\rho)),$$

where $\Psi(x, y)$ is the number of y -smooth integers up to x . Consequently, if $K = N^{\theta+o(1)}$ with $0 \leq \theta \leq 1$ (that is, $\log(K+1)/\log N \rightarrow \theta$), then $\mathbb{E}[\Phi_K]/N \rightarrow \rho_{\text{Dick}}(1/(1-\theta))$, where ρ_{Dick} is Dickman's function and $\rho_{\text{Dick}}(\infty) = 0$. For $0 \leq \theta < 1$ the same limit holds if the K vertices of largest degree are deleted instead, whether degrees are computed once or recomputed after each deletion. This answers Question 20 of Kim and Phillips. Moreover, the symmetric difference of the largest strong component and the set of $N/(K+1)$ -smooth vertices has expected size at most $2K + o(N)$, and the second largest strong component has expected size at most $K + o(N)$.

The upper bound comes from the decomposition of the divisor graph into fibres of the rough part. The lower bound rests on one observation. Every divisibility edge $\{x, x'\}$ with $x' \leq N^{1-2\delta}$ is shadowed by at least $N^{2\delta} - 2$ independent directed 2-paths through common multiples. So with high probability the strong components are at least as coarse as the connected components of a deterministic divisor graph on $(K, N^{1-2\delta}]$, and a doubling argument shows that the smooth part of that graph is connected. The proof is elementary and self-contained. It uses the estimates of Chebyshev and Mertens, a second-moment bound for the number of small prime factors, and, to identify the limit, Dickman's asymptotic for Ψ ; all three are proved here from first principles. Every numbered statement, including the remarks, has been formally verified in Lean 4 with Mathlib, and the proofs given here follow the formal ones.

CONTENTS

1. Introduction	2
2. Arithmetic preliminaries	5
3. The upper bound: fibres of the rough part	8
4. The lower bound	9
5. Proofs of the main results	13
6. Removing the vertices of highest degree	15
7. Further remarks	17
Appendix A. The sub-sum lemma	18
Appendix B. Status of all claims	18
Appendix C. Proofs of Lemmas 2.1 and 2.2	21
References	24

1. INTRODUCTION

1.1. The model and the question. For $N \geq 2$ the *divisor graph* G_N has vertex set $[N] = \{1, \dots, N\}$ and an edge $\{a, b\}$ whenever $a \neq b$ and $a \mid b$ or $b \mid a$. Kim and Phillips [9] introduced the *randomly oriented divisor graph* $\mathcal{D}_\rho(N)$. Each edge $\{d, m\}$ with $d \mid m$, $d < m$, is first oriented $m \rightarrow d$ (multiple to divisor) and then independently reversed with probability $\rho \in [0, 1]$. Thus every edge independently points $m \rightarrow d$ with probability $1 - \rho$ and $d \rightarrow m$ with probability ρ . Write $\#\Phi(D)$ for the number of vertices of a largest strongly connected component (SCC) of a directed graph D . The main theorem of [9] states that $\mathbb{E}[\#\Phi(\mathcal{D}_\rho(N))]/N \rightarrow 1$ for every fixed $\rho \in (0, 1)$. The authors ask:

Question 1.1 ([9, Question 20]). What can be said about the expected size of the largest strongly connected component as we remove the vertices of highest degrees, one by one. In other words, how will the expected size of the largest strongly connected component change as we remove the vertices labeled 1, 2, 3, 4, and so on?

For $0 \leq K < N$ let $V_{N,K} = \{K+1, \dots, N\}$, let $G_{N,K} = G_N[V_{N,K}]$ be the induced subgraph, and let

$$\mathcal{D}_\rho(N, K) := \mathcal{D}_\rho(N)[V_{N,K}]$$

be the random directed graph obtained from $\mathcal{D}_\rho(N)$ by deleting the vertices $1, \dots, K$. Its edges are exactly the edges of $G_{N,K}$, oriented independently as above. Put

$$\Phi_K := \#\Phi(\mathcal{D}_\rho(N, K)).$$

Throughout, $\rho \in (0, 1)$ is the reversal probability, and we write

$$q := \rho(1 - \rho) \in (0, \frac{1}{4}], \quad \lambda := \max(\rho, 1 - \rho) \in [\frac{1}{2}, 1).$$

1.2. Notation. p always denotes a prime. $P^+(n)$ is the largest prime factor of n , with $P^+(1) = 1$. For real $x \geq 0$ and $y \geq 1$,

$$\Psi(x, y) := \#\{1 \leq n \leq x : P^+(n) \leq y\}.$$

For $z \geq 2$, $\omega_z(n) := \#\{p \leq z : p \mid n\}$. We write ρ_{Dick} for Dickman's function (§2); the letter ρ alone always means the reversal probability. For $0 \leq K < N$ put

$$B = B(N, K) := \left\lfloor \frac{N}{K+1} \right\rfloor, \quad F_1 = F_1(N, K) := \{m \in V_{N,K} : P^+(m) \leq B\}.$$

Note that $P^+(m) \leq N/(K+1)$ if and only if $P^+(m) \leq B$, so $\Psi(N, N/(K+1)) = \Psi(N, B)$. We call F_1 the *smooth fibre*. Logarithms are natural. “With high probability” (w.h.p.) means with probability $1 - o(1)$ as $N \rightarrow \infty$.

1.3. Results. Each statement carries a status tag. [PROVEN] means a complete proof is given in this paper. A line [Lean: name] gives the names of the corresponding results in the Lean 4 formalisation (directory `lean/` of the repository; Lean and Mathlib v4.34.1). There the statement, as given in this paper, is proved with no `sorry`, and `#print axioms` shows only the standard axioms `propext`, `Classical.choice` and `Quot.sound`. Appendix B explains how the formal model matches the definitions here. The proofs in this paper follow the formal proofs. Where the formal statement has weaker hypotheses or explicit constants, we state the formal version. Every numbered statement of the paper, including the remarks and the appendices, is formalised in this sense.

Theorem 1.2 (Main theorem). [PROVEN]

Lean: `thm_main`

Fix $\rho \in (0, 1)$. For all $N \geq 2$ and $0 \leq K < N$ we have $\Phi_K \leq \Psi(N, N/(K+1))$ deterministically, and

$$\lim_{N \rightarrow \infty} \max_{0 \leq K < N} \frac{1}{N} \left(\Psi\left(N, \frac{N}{K+1}\right) - \mathbb{E}[\Phi_K] \right) = 0.$$

Theorem 1.3 (Rate). [PROVEN]Lean: *thm_rate*

For every $\rho \in (0, 1)$ there is $N_0(\rho)$ such that for all $N \geq N_0(\rho)$ and all $0 \leq K < N$,

$$0 \leq \Psi\left(N, \frac{N}{K+1}\right) - \mathbb{E}[\Phi_K] \leq \frac{30N}{\log \log N}.$$

The bound is non-trivial only when $\log \log N > 30$. Its content is the uniformity in K and the explicit rate, not the numerical constant, which we have not tried to optimise.

For $K \leq \sqrt{N} - 1$ this becomes fully explicit (Remark 5.1):

$$(1) \quad 1 - \frac{\mathbb{E}[\Phi_K]}{N} = \log \frac{\log N}{\log B} + O_\rho\left(\frac{1}{\log \log N}\right), \quad B = \left\lfloor \frac{N}{K+1} \right\rfloor.$$

So, up to an additive error $O_\rho(1/\log \log N)$ that is uniform in $0 \leq K \leq \sqrt{N} - 1$, the expected fraction of vertices lost after removing the hubs $1, \dots, K$ equals $\log \frac{\log N}{\log B}$. This main term dominates the error only when $\log K \gg \log N / \log \log N$. For smaller K , in particular for bounded K , we do not determine the loss to within a factor $1 + o(1)$.

Corollary 1.4 (Limit profile). [PROVEN]Lean: *cor_profile, dickLim_continuous, dickLim_strictAntiOn, dickLim_of_le_half*

Fix $\rho \in (0, 1)$, and let $K = K(N)$ satisfy $0 \leq K(N) < N$ for all large N and $\log(K+1)/\log N \rightarrow \theta$. (Necessarily $\theta \in [0, 1]$.) Then

$$\lim_{N \rightarrow \infty} \frac{\mathbb{E}[\Phi_K]}{N} = \rho_{\text{Dick}}\left(\frac{1}{1-\theta}\right),$$

with the convention $\rho_{\text{Dick}}(\infty) = 0$. The profile $\theta \mapsto \rho_{\text{Dick}}(1/(1-\theta))$, extended by 0 for $\theta \geq 1$, is continuous on $\mathbb{R}$ and strictly decreasing on $[0, 1]$. It equals $1 - \log \frac{1}{1-\theta}$ for $0 \leq \theta \leq \frac{1}{2}$.

Corollary 1.5 (Structure of the giant). [PROVEN]Lean: *cor_structure, cor_structure_prob, cor_structure_whp*

Fix $\rho \in (0, 1)$. Let Φ_K^* be a largest SCC of $\mathcal{D}_\rho(N, K)$ (chosen by any rule), and let $\Phi_K^{(2)}$ be the size of a largest SCC other than Φ_K^* (0 if there is none). For $N \geq N_0(\rho)$ and all $0 \leq K < N$:

- (a) $\mathbb{E}|\Phi_K - \Psi(N, \frac{N}{K+1})| \leq 30N/\log \log N$. In particular $\Phi_K/N - \Psi(N, \frac{N}{K+1})/N \rightarrow 0$ in probability, uniformly in K .
- (b) $\mathbb{E}|F_1 \triangle \Phi_K^*| \leq 2K + 30N/\log \log N$.
- (c) $\mathbb{E}[\Phi_K^{(2)}] \leq K + 30N/\log \log N$.

So when $K = o(N)$, the following holds for every fixed $\varepsilon > 0$ with probability $1 - o(1)$, by Markov's inequality applied to (b) and (c) (*cor_structure_whp*). The largest SCC differs from the set of $N/(K+1)$ -smooth integers in $(K, N]$ in at most εN vertices, and every other SCC has at most εN vertices. The term K in (c) cannot be dropped in general; see Remark 5.2.

Corollary 1.6 (Removing the vertices of highest degree). [PROVEN]Lean: *cor_degree_static, cor_degree_adaptive, static_eq_Icc*

Fix $\rho \in (0, 1)$ and let $\log(K+1)/\log N \rightarrow \theta \in [0, 1]$. Let $T = T(N) \subset [N]$ with $|T| = K$ be chosen from the undirected graph G_N alone (and so independently of the orientation), for all large N , in either of the following ways:

- (i) static attack: $\min_{t \in T} \deg_{G_N}(t) \geq \max_{v \notin T} \deg_{G_N}(v)$;
- (ii) adaptive attack: $T = \{u_1, \dots, u_K\}$, where for each j the vertex u_{j+1} has maximal degree in $G_N - \{u_1, \dots, u_j\}$.

Ties may be broken arbitrarily. Then $\mathbb{E}[\#\Phi(\mathcal{D}_\rho(N) - T)]/N \rightarrow \rho_{\text{Dick}}(1/(1-\theta))$. Moreover, for fixed $\kappa \in (0, \frac{1}{2})$ and $N \geq N_1(\kappa)$, if $1 \leq K \leq N^{1/2-\kappa}$ then the static set T is exactly $\{1, \dots, K\}$.

Since the total degree of a vertex in any orientation equals its degree in G_N , (i) and (ii) cover attacks by total degree in the directed graph. Attacks by in- or out-degree depend on the orientation and are not covered (see the end of §6).

θ	0.1	0.2	1/3	0.4	1/2	0.6
$u = \frac{1}{1-\theta}$	1.111	1.25	1.5	1.667	2	2.5
$\rho_{\text{Dick}}(u)$	0.8946	0.7769	0.5945	0.4892	0.3069	0.1303
θ	2/3	3/4	0.8	0.9	0.95	1
$u = \frac{1}{1-\theta}$	3	4	5	10	20	∞
$\rho_{\text{Dick}}(u)$	0.04861	0.004911	$3.547 \cdot 10^{-4}$	$2.770 \cdot 10^{-11}$	$2.46 \cdot 10^{-29}$	0

TABLE 1. The limit profile $\lim \mathbb{E}[\Phi_K]/N$ when $\log(K+1)/\log N \rightarrow \theta$. For $u \leq 2$, $\rho_{\text{Dick}}(u) = 1 - \log u$ (Lemma 2.2). The other values were computed numerically from $u\rho_{\text{Dick}}(u) = \int_{u-1}^u \rho_{\text{Dick}}(t) dt$ and agree with the standard tables. The decimal values are numerical approximations and are not part of the formal verification.

1.4. Discussion. *Answer to Question 1.1.* Deleting any $K = N^{o(1)}$ hubs costs nothing asymptotically; for instance $\mathbb{E}[\Phi_K]/N \rightarrow 1$ for every fixed K . Deleting $K = N^\theta$ hubs, a vanishing fraction of the vertices, leaves a giant of relative size $\rho_{\text{Dick}}(1/(1-\theta))$. For example, deleting the $\sqrt{N}$ hubs leaves $1 - \log 2 \approx 30.7\%$. Deleting $N^{1-o(1)}$ hubs destroys the giant. The vertices $1, \dots, K$ carry a θ -fraction of the $N \log N + O(N)$ edges (Remark 7.1), so for moderate θ hub removal is far more damaging than the loss of edges alone suggests: at $\theta = \frac{1}{2}$ about half of the edges disappear, but only 30.7% of the vertices remain in the giant. This is the fragility to targeted attacks familiar from scale-free networks [1, 2]. In the divisor graph, however, the surviving giant has an explicit arithmetic description: the smooth fibre. At finite N the error bound $30N/\log \log N$ of Theorem 1.3 tends to 0 only slowly relative to N .

What is new. To our knowledge Theorems 1.2 and 1.3 and Corollaries 1.4–1.6 are new. The fibre decomposition behind the upper bound (Lemma 3.1) is a simple and essentially known observation; a closely related statement appears in McNew’s work on the divisor graph [10]. The inputs from analytic number theory (Mertens, Chebyshev, Dickman, a Turán-type variance bound, and the divisor bound) are classical. The lower-bound mechanism (Lemmas 4.1 and 4.2) is, as far as we know, new. It differs from the triangle argument of [9], which uses paths through the vertex 1, and it does not need a strongly connected “hub clique”. Such a clique, for instance the interval $(K, N^{1/2-\delta}]$, whose elements pairwise have many common multiples, can capture only the integers with a divisor in the clique. For $1/3 < \theta < 1/2$ this misses a positive proportion of F_1 (Remark 7.2).

1.5. Idea of the proof. *Upper bound.* If $d \mid m$ with $K < d < m \leq N$, then $m/d \leq B$. Hence the B -rough part of a vertex is constant on connected components, and every SCC lies in a fibre $\{v : v = ns, P^+(s) \leq B\}$, which has at most $\Psi(N, B)$ elements.

Lower bound. Fix a small $\delta > 0$ and put $M = N^{1-2\delta}$.

- (1) *Robust edges.* If $K < x < x' \leq M$ and $x \mid x'$, then each of the $\geq N^{2\delta} - 2$ multiples y of x' in $(x', N]$ gives a directed path $x \rightarrow y \rightarrow x'$ with probability q , independently over y ; likewise for $x' \rightarrow y \rightarrow x$. A union bound shows that w.h.p. every edge of the *deterministic* graph $G_{M,K}$ joins two vertices of the same SCC. So every connected component of $G_{M,K}$ lies inside one SCC.
- (2) *Doubling.* All $x \in (K, M]$ with $P^+(x) \leq M/(2(K+1))$ lie in one component of $G_{M,K}$. Multiply by 2 until just below M , divide by an odd prime, and repeat; this ends at the power of 2 in $(M/2, M]$. Call this set S_M .

- (3) *Attachment.* A vertex $m \in (M, N]$ joins the SCC containing S_M as soon as, among the edges from m to its divisors in S_M , at least one points each way. This has probability $1 - \rho^s - (1 - \rho)^s$, where s is the number of such divisors. An elementary counting argument shows that all but $o(N)$ elements of F_1 have $s \geq s_0$ for any fixed s_0 . It uses a reserve of s_0 small prime factors of m (supplied by Turán’s variance bound), a deterministic case analysis, and Mertens’ theorem.

Organisation. §2 collects the arithmetic inputs. §3 proves the upper bound, §4 the general lower bound (Proposition 4.6), and §5 the theorems and corollaries. §6 treats degree-based removal, and §7 contains further remarks. Appendix A proves the sub-sum lemma from the preliminary notes, which is no longer needed. Appendix B lists every claim with its status and its Lean name. Appendix C proves Lemmas 2.1 and 2.2.

2. ARITHMETIC PRELIMINARIES

We use the following classical facts. All of them are proved in this paper, with the explicit constants given by the formal proofs. The longer proofs of Lemmas 2.1 and 2.2 are in Appendix C. For real arguments, $\Psi(x, y) = \Psi(\lfloor x \rfloor, \lfloor y \rfloor)$, and a sum over $p \leq y$ is a sum over $p \leq \lfloor y \rfloor$.

Lemma 2.1 (Mertens [11], Chebyshev; see [6, Ch. XXII], [12, Part I, Ch. 1]). [PROVEN]

Lean: `lemma_2_1`, `mertens_a`, `mertens_a_real`, `mertens_b`, `mertens_b_nat`, `mertens_c`, `mertens_d`

Put $c_0 := \log 4 + 2$ and $C_1 := 18$. Then:

- (a) $\sum_{p \leq y} \frac{\log p}{p-1} \leq \log y + c_0$ for every integer $y \geq 1$ and every real $y \geq 2$;
- (b) $\left| \sum_{y < p \leq w} \frac{1}{p} - \log \frac{\log w}{\log y} \right| \leq \frac{C_1}{\log y}$ for all real $2 \leq y \leq w$; for integers $2 \leq y \leq w$ the bound $8/\log y$ holds;
- (c) $\sum_{p \leq z} \frac{1}{p} \geq \log \log z - 13 \geq \log \log z - C_1$ for all real $z \geq 2$;
- (d) $\pi(x) \leq 5x/\log x \leq C_1 x/\log x$ for all real $x \geq 2$.

Proof. See Appendix C. The proof uses only Chebyshev’s bound $\prod_{p \leq n} p \leq 4^n$, Legendre’s formula for $v_p(n!)$, the inequalities $n^n/e^n \leq n! \leq n^n$, and Abel summation. $\square$

Lemma 2.2 (Dickman’s function [4, 3, 7]; see [12, Part III, Ch. 5], [5, 8]). [PROVEN]

Lean: `dickman`, `dickman_unique`, `dickman_integral`, `mul_dickman_eq_integral`, `dickman_hasDerivAt`, `dickman_pos`, `dickman_le_one`, `dickman_eq_one_sub_log`, `dickman_strictAntiOn`, `dickman_asymptotic`, `dickman_tendstoUniformlyOn`

There is a unique function $\rho_{\text{Dick}} : [0, \infty) \rightarrow \mathbb{R}$ that is continuous, equals 1 on $[0, 1]$, and is differentiable at every $u > 1$ with $u \rho'_{\text{Dick}}(u) = -\rho_{\text{Dick}}(u-1)$. For $u \geq 1$ it satisfies

$$\rho_{\text{Dick}}(u) = 1 - \int_1^u \frac{\rho_{\text{Dick}}(w-1)}{w} dw \quad \text{and} \quad u \rho_{\text{Dick}}(u) = \int_{u-1}^u \rho_{\text{Dick}}(v) dv.$$

Moreover $0 < \rho_{\text{Dick}} \leq 1$, $\rho_{\text{Dick}}(u) = 1 - \log u$ for $1 \leq u \leq 2$, and ρ_{Dick} is strictly decreasing on $[1, \infty)$. For every fixed $U \geq 1$,

$$\sup_{1 \leq u \leq U} \left| \frac{\Psi(x, x^{1/u})}{x} - \rho_{\text{Dick}}(u) \right| \rightarrow 0 \quad (x \rightarrow \infty).$$

Proof. See Appendix C. The function is constructed recursively on the intervals $[n, n+1]$. The asymptotic is proved by induction on $\lceil U \rceil$ from Buchstab’s identity (Lemma C.1), Lemma 2.1(b),(d) and a Riemann-sum estimate for sums over primes (Lemma C.2). No result from the literature on smooth numbers is used. The pointwise asymptotic is due to Dickman [4]; de Bruijn [3] and Hildebrand [7] give error terms. $\square$

Lemma 2.3 (Elementary smooth-number bound). [PROVEN, standard]

Lean: `psi_upper`, `psi_upper'`, `sum_log_smooth_le`, `dickman_le_two_div`, `dickman_tendsto_zero`

For integers $x \geq 2$ and $y \geq 1$, with $c_0 = \log 4 + 2$,

$$\Psi(x, y) \leq \sqrt{x} + \frac{2x(\log y + c_0)}{\log x}.$$

In particular $\rho_{\text{Dick}}(u) \leq 2/u$ for all $u \geq 1$, and $\rho_{\text{Dick}}(u) \rightarrow 0$ as $u \rightarrow \infty$.

Proof. For $n \geq 1$ we have $\log n = \sum_{p^k | n, k \geq 1} \log p$. Hence

$$\begin{aligned} \sum_{\substack{n \leq x \\ P^+(n) \leq y}} \log n &= \sum_{\substack{p \leq y \\ k \geq 1}} \log p \cdot \#\{n \leq x : P^+(n) \leq y, p^k | n\} \\ &\leq x \sum_{p \leq y} \log p \sum_{k \geq 1} p^{-k} = x \sum_{p \leq y} \frac{\log p}{p-1} \leq x(\log y + c_0) \end{aligned}$$

by Lemma 2.1(a) for the integer $y \geq 1$. On the other hand, at least $\Psi(x, y) - \sqrt{x}$ of the n counted satisfy $n > \sqrt{x}$, and each of them has $\log n > \frac{1}{2} \log x$. Therefore $(\Psi(x, y) - \sqrt{x}) \cdot \frac{1}{2} \log x \leq x(\log y + c_0)$.

For the last assertion, fix $u \geq 1$. For an integer $N \geq 2$ put $y := \lfloor N^{1/u} \rfloor \geq 1$, so $\Psi(N, N^{1/u}) = \Psi(N, y)$ and $\log y \leq \frac{1}{u} \log N$. The bound gives

$$\frac{\Psi(N, N^{1/u})}{N} \leq N^{-1/2} + \frac{2}{u} + \frac{2c_0}{\log N}.$$

As $N \rightarrow \infty$ the left side tends to $\rho_{\text{Dick}}(u)$ by Lemma 2.2, and the right side tends to $2/u$. So $\rho_{\text{Dick}}(u) \leq 2/u$, and $0 < \rho_{\text{Dick}}(u) \leq 2/u$ gives $\rho_{\text{Dick}}(u) \rightarrow 0$. $\square$

Uses: Lemma 2.1(a); for the consequence, Lemma 2.2.

Lemma 2.4 (Variance of ω_z ; the special case of Turán–Kubilius we need). [PROVEN, standard]

Lean: `turan_variance`, `turan_count`

Let $N \geq 1$ and $2 \leq z \leq N$ be integers, and $L := \sum_{p \leq z} \frac{1}{p}$. Then

$$\sum_{m \leq N} (\omega_z(m) - L)^2 \leq 3NL.$$

(The variance bound needs only $z \leq N$. For real z nothing changes, since $\omega_z = \omega_{\lfloor z \rfloor}$ and L depends only on $\lfloor z \rfloor$.) Consequently, for every real $s \leq L/2$,

$$\#\{m \leq N : \omega_z(m) < s\} \leq \frac{12N}{L}.$$

Proof. Write $f = \omega_z$. We have $S_1 := \sum_{m \leq N} f(m) = \sum_{p \leq z} \lfloor N/p \rfloor \geq NL - \pi(z)$. Also

$$S_2 := \sum_{m \leq N} f(m)^2 = \sum_{p \leq z} \lfloor N/p \rfloor + \sum_{\substack{p, p' \leq z \\ p \neq p'}} \lfloor N/(pp') \rfloor \leq NL + NL^2,$$

because m is divisible by the distinct primes p, p' if and only if $pp' | m$. Hence

$$\sum_{m \leq N} (f(m) - L)^2 = S_2 - 2LS_1 + NL^2 \leq NL + NL^2 - 2L(NL - \pi(z)) + NL^2 = NL + 2L\pi(z) \leq 3NL,$$

using $\pi(z) \leq z \leq N$. If $f(m) < s \leq L/2$, then $|f(m) - L| > L/2$. By Chebyshev's inequality the number of such m is at most $3NL/(L/2)^2 = 12N/L$. $\square$

Uses: nothing beyond counting.

Remark 2.5 (The Turán–Kubilius inequality). [PROVEN]

Lean: `remark_2_5`, `turan_kubilius`, `tk_core`, `tk_E`, `tk_AG`

Lemma 2.4 is the special case $f = \omega_z$ of the Turán–Kubilius inequality [13]; see [12, Part III, Ch. 3]. That inequality states the following. There is an absolute constant C (one may take $C = 130$) such that for every complex additive function f (that is, $f(mn) = f(m) + f(n)$ whenever $(m, n) = 1$) and every $x \geq 2$,

$$\sum_{n \leq x} |f(n) - A(x)|^2 \leq C x B(x)^2,$$

where $A(x) = \sum_{p^\nu \leq x} f(p^\nu) p^{-\nu} (1 - p^{-1})$ and $B(x)^2 = \sum_{p^\nu \leq x} |f(p^\nu)|^2 p^{-\nu}$. We use only the self-contained special case proved above.

Proof with $C = 130$. Put $N = \lfloor x \rfloor$. For a prime power $q = p^\nu \leq N$ ($\nu \geq 1$) put $w_q = q^{-1} (1 - p^{-1})$ and $X_q(n) = \mathbf{1}[v_p(n) = \nu]$, so that $f(n) = \sum_{q \leq N} f(q) X_q(n)$ for $1 \leq n \leq N$. Splitting f into real and imaginary parts, and each of these into positive and negative parts, costs a factor 2. It reduces the claim to $\sum_{n \leq N} (F(n) - A)^2 \leq 65 N B^2$, where $F = \sum_q g_q X_q$ with weights $g_q \geq 0$, $A = \sum_q g_q w_q$ and $B^2 = \sum_q g_q^2 / q$. Now $\#\{n \leq N : v_p(n) = \nu\} \geq N w_q - 1$. For $q = p^\nu$ and $q' = r^\mu$ with $p \neq r$, inclusion–exclusion gives $\#\{n \leq N : v_p(n) = \nu, v_r(n) = \mu\} \leq N w_q w_{q'} + 2 \cdot \mathbf{1}[qq' \leq N]$; for $p = r$ and $\nu \neq \mu$ this count is 0. Expanding the square therefore gives

$$\sum_{n \leq N} (F(n) - A)^2 \leq N B^2 + 2E + 2AG, \quad E = \sum_{qq' \leq N} g_q g_{q'}, \quad G = \sum_{q \leq N} g_q.$$

The prime powers $p^\nu \leq Y$ with $\nu \geq 2$ have $p \leq \sqrt{Y}$ and $\nu \leq \log_2 Y$. With Lemma 2.1(d) this shows that there are at most $58Y / \log 2Y$ prime powers $q \leq Y$. Also $\sum_{q \leq M} 1/q \leq 1 + \log M$. By Cauchy–Schwarz, $E^2 \leq B^4 \sum_{qq' \leq N} qq' \leq (2NB^2)^2$, where each pair is split according to which of q, q' is at most $\sqrt{N}$; and $(AG)^2 \leq B^2 (1 + \log N) \cdot B^2 \sum_{q \leq N} q \leq (10NB^2)^2$. Hence the sum is at most $(1 + 44 + 20)NB^2$. $\square$

Lemma 2.6 (Divisor bound [6, Thm. 315]). [PROVEN, standard]

Lean: `divisor_bound_explicit`, `divisor_bound`

For every $\varepsilon > 0$ we have $\tau(n) \leq C_\varepsilon n^\varepsilon$ for all $n \geq 1$, where $C_\varepsilon := \max(1, \frac{1}{\varepsilon \log 2})^{\lceil 2^{1/\varepsilon} \rceil}$.

Proof. Put $c := \max(1, 1/(\varepsilon \log 2))$ and $B := 2^{1/\varepsilon}$. We have $\tau(n) = \prod_{p^a \parallel n} (a+1)$ and $n^\varepsilon = \prod_{p^a \parallel n} p^{a\varepsilon}$. If $p \geq B$, then $p^{a\varepsilon} \geq 2^a \geq a+1$. If $p < B$, then $2^{a\varepsilon} = e^{a\varepsilon \log 2} \geq 1 + a\varepsilon \log 2$, so $c p^{a\varepsilon} \geq c 2^{a\varepsilon} \geq c + a c \varepsilon \log 2 \geq 1 + a$. Hence $\tau(n) \leq c^{\#\{p|n: p < B\}} n^\varepsilon \leq c^{\lceil B \rceil} n^\varepsilon$, since at most $\lceil B \rceil$ primes are $< B$ and $c \geq 1$. $\square$

Lemma 2.7 (Limit of the smooth count). [PROVEN, standard]

Lean: `psi_limit`, `psi_eq_PsiR_rpow`, `psi_div_le`

Let $K = K(N)$ satisfy $0 \leq K(N) < N$ for all large N and $\log(K+1)/\log N \rightarrow \theta$. (Necessarily $\theta \in [0, 1]$.) Then

$$\lim_{N \rightarrow \infty} \frac{\Psi(N, N/(K+1))}{N} = \rho_{\text{Dick}}\left(\frac{1}{1-\theta}\right), \quad \text{with } \rho_{\text{Dick}}(\infty) = 0.$$

Proof. Put $r_N := \log(K+1)/\log N$ and $y := N/(K+1)$. For large N we have $N \geq 2$ and $K < N$, so $0 \leq r_N \leq 1$ and $B = \lfloor y \rfloor \geq 1$.

If $\theta < 1$, then $r_N < 1$ for large N . Put $u_N := 1/(1 - r_N) \geq 1$. Then $N^{1/u_N} = N^{1-r_N} = \exp(\log N - \log(K+1)) = y$ exactly, so $\Psi(N, N/(K+1)) = \Psi(N, N^{1/u_N})$. Also $u_N \rightarrow u := 1/(1-\theta)$, so $u_N \in [1, u+1]$ for large N . Lemma 2.2 with $U = u+1$ gives $\Psi(N, N^{1/u_N})/N - \rho_{\text{Dick}}(u_N) \rightarrow 0$, and $\rho_{\text{Dick}}(u_N) \rightarrow \rho_{\text{Dick}}(u)$ by continuity of ρ_{Dick} .

If $\theta \geq 1$, then $\theta = 1$, since $r_N \leq 1$. As $B \geq 1$ and $\log B \leq \log N - \log(K + 1)$, Lemma 2.3 gives

$$\frac{\Psi(N, B)}{N} \leq N^{-1/2} + 2(1 - r_N) + \frac{2c_0}{\log N} \rightarrow 0.$$

□

Uses: Lemmas 2.2, 2.3.

3. THE UPPER BOUND: FIBRES OF THE ROUGH PART

For $n \geq 1$ let $R_B(n) := \prod_{p^v \parallel n, p > B} p^v$ be the B -rough part of n .

Lemma 3.1 (Fibres). [PROVEN; essentially known, cf. [10]]

Lean: `cofactor_le_B`, `roughPart_eq_of_edge`, `roughPart_eq_of_reachable`, `card_fibre_le`, `card_fibre_le_psi`, `fibre_one`, `card_smoothFibre`, `card_fibre_le_K`

Let $0 \leq K < N$ and $B = \lfloor N/(K + 1) \rfloor$.

- (a) If $\{d, m\}$ is an edge of $G_{N,K}$ with $d < m$, then $m/d \leq B$, and hence $R_B(d) = R_B(m)$.
- (b) R_B is constant on every connected component of $G_{N,K}$.
- (c) For $n \geq 1$ let $V_n := \{v \in V_{N,K} : R_B(v) = n\}$. Then $|V_n| \leq \Psi(N/n, B) \leq \Psi(N, B)$.
Moreover $V_1 = F_1$, $|F_1| = \Psi(N, B) - \Psi(K, B)$, and $|V_n| \leq K$ for $n > 1$.

Proof. (a) Since $d \geq K + 1$, we have $m/d \leq N/(K + 1)$. As m/d is an integer, $m/d \leq B$. So every prime factor of m/d is at most B , and $v_p(m) = v_p(d)$ for every $p > B$.

(b) This follows from (a) by induction along paths.

(c) Each $v \in V_n$ factors uniquely as $v = ns$ with $s = v/n$ being B -smooth. Since $s \leq N/n$, the map $v \mapsto s$ injects V_n into $\{s \leq N/n : P^+(s) \leq B\}$. Clearly $V_1 = \{v \in (K, N] : P^+(v) \leq B\} = F_1$, and $|F_1| = \Psi(N, B) - \Psi(K, B)$. If $n > 1$, then n has a prime factor $> B$, so $n \geq B + 1 > N/(K + 1)$. Hence $|V_n| \leq N/n < K + 1$. □

Uses: nothing.

Proposition 3.2 (Upper bound). [PROVEN]

Lean: `stronglyConnected_subset_fibre`, `stronglyConnected_card_le_psi`, `stronglyConnected_card_le_max`, `PhiK_le_psi`, `PhiK_le_max`

For every orientation of $G_{N,K}$, every SCC is contained in a single fibre V_n . Hence

$$\Phi_K \leq \Psi(N, B) \quad \text{and} \quad \Phi_K \leq \max(|F_1|, K).$$

Proof. An SCC is connected in the underlying undirected graph, so it lies in a connected component of $G_{N,K}$, and hence in a fibre by Lemma 3.1(b). Now apply Lemma 3.1(c). □

Uses: Lemma 3.1.

Corollary 3.3. [PROVEN]

Lean: `cor_upper_profile`, `cor_upper_profile_eventually`

Let $\rho \in [0, 1]$ be arbitrary. If $0 \leq K(N) < N$ for all large N and $\log(K + 1)/\log N \rightarrow \theta$, then $\limsup_N \mathbb{E}[\Phi_K]/N \leq \rho_{\text{Dick}}(1/(1 - \theta))$, with $\rho_{\text{Dick}}(\infty) = 0$.

Proof. By Proposition 3.2, $\mathbb{E}[\Phi_K]/N \leq \Psi(N, N/(K + 1))/N$, and the right side tends to $\rho_{\text{Dick}}(1/(1 - \theta))$ by Lemma 2.7. □

Remark 3.4 (Equivalent forms). [PROVEN]

Lean: `remark_3_4`, `remark_3_4_holds`

Let $\rho \in [0, 1]$ and suppose $K = o(N)$. The following are equivalent: (i) $\mathbb{E}[\Phi_K] = \Psi(N, B) + o(N)$; (ii) $\mathbb{E}[|F_1| - \Phi_K] = o(N)$; (iii) for every $\varepsilon > 0$, w.h.p. the largest SCC has at least $|F_1| - \varepsilon N$ vertices. Theorem 1.2 establishes (i), and hence all three forms, for $\rho \in (0, 1)$.

Proof. Pointwise, $|F_1| \leq \Psi(N, B) \leq |F_1| + K$ by Lemma 3.1(c), and $\Phi_K \leq \Psi(N, B)$ and $\Phi_K \leq \max(|F_1|, K)$ by Proposition 3.2. Hence $||F_1| - \Phi_K| \leq (\Psi(N, B) - \Phi_K) + K$, and taking expectations gives (i) $\Rightarrow$ (ii). Markov's inequality gives (ii) $\Rightarrow$ (iii). Finally, $\Psi(N, B) - \Phi_K \leq K + \varepsilon N + N \cdot \mathbf{1}[\Phi_K < |F_1| - \varepsilon N]$, which gives (iii) $\Rightarrow$ (i). $\square$

4. THE LOWER BOUND

Throughout this section $N \geq 2$, $0 \leq K < N$ and $\delta \in (0, \frac{1}{8}]$ are fixed, and

$$M := N^{1-2\delta} \quad (\text{a real number}), \quad Y := \frac{M}{2(K+1)}, \quad S_M := \{x \in \mathbb{Z} \cap (K, M] : P^+(x) \leq Y\}.$$

We write $G_{M,K}$ for the induced subgraph of G_N on $\mathbb{Z} \cap (K, M]$. This is a subgraph of $G_{N,K}$, and it is not random.

4.1. Robust edges.

Lemma 4.1 (Robust edges). [PROVEN; new]

Lean: `prob_not_robust`, `prob_not_robust_exp`, `robust_pair`, `robust_adj`, `robust_reachable`, `card_multiples_Ioc_ge`

Let $\mathcal{R}$ be the event that for all integers x, x' with $K < x < x' \leq M$ and $x \mid x'$ there are vertices $y, y' \in V_{N,K}$, both multiples of x' , such that $x \rightarrow y \rightarrow x'$ and $x' \rightarrow y' \rightarrow x$ in $\mathcal{D}_\rho(N, K)$. Then

$$\mathbb{P}(\mathcal{R}^c) \leq M^2(1-q)^{N^{2\delta}-2} \leq N^2 \exp(-q(N^{2\delta}-2)).$$

On $\mathcal{R}$, the two endpoints of every edge of $G_{M,K}$ lie in the same SCC of $\mathcal{D}_\rho(N, K)$. Consequently every connected component of $G_{M,K}$ lies in a single SCC. (The formal version allows any real $M \in (0, N]$, with N/M in place of $N^{2\delta}$.)

Proof. Fix x, x' as in the statement and let $\mathcal{Y} := \{y \in \mathbb{Z} : x' < y \leq N, x' \mid y\}$. Then

$$|\mathcal{Y}| = \lfloor N/x' \rfloor - 1 \geq N/M - 2 = N^{2\delta} - 2.$$

For $y \in \mathcal{Y}$ we have $K < x < x' < y \leq N$ and $x \mid x' \mid y$. So $\{x, y\}$ and $\{x', y\}$ are edges of $G_{N,K}$, and their default orientations are $y \rightarrow x$ and $y \rightarrow x'$. Let E_y be the event $\{x \rightarrow y \text{ and } y \rightarrow x'\}$, which says that $\{x, y\}$ is reversed and $\{x', y\}$ is not. Then $\mathbb{P}(E_y) = \rho(1-\rho) = q$. For distinct y the pairs of edges involved are disjoint, so the events $E_y, y \in \mathcal{Y}$, are independent. Hence

$$\mathbb{P}\left(\bigcap_{y \in \mathcal{Y}} E_y^c\right) = (1-q)^{|\mathcal{Y}|} \leq (1-q)^{N^{2\delta}-2}.$$

The same bound holds for the events $E'_y = \{x' \rightarrow y \text{ and } y \rightarrow x\}$, which say that $\{x', y\}$ is reversed and $\{x, y\}$ is not. The pairs (x, x') are the edges of $G_{M,K}$, and $2|E(G_{M,K})| \leq \lfloor M \rfloor^2$. Each pair has two directions, so the union bound gives $\mathbb{P}(\mathcal{R}^c) \leq M^2(1-q)^{N^{2\delta}-2}$. The second inequality uses $M \leq N$ and $(1-q)^t \leq e^{-qt}$ for $t = N^{2\delta} - 2 \geq 0$. If $N^{2\delta} < 2$, the right-hand side exceeds 1 and there is nothing to prove.

On $\mathcal{R}$, if $\{x, x'\}$ is an edge of $G_{M,K}$ with $x < x'$, then $K < x < x' \leq M$ and $x \mid x'$. So directed paths $x \rightarrow y \rightarrow x'$ and $x' \rightarrow y' \rightarrow x$ exist, and x, x' lie in the same SCC. Lying in the same SCC is an equivalence relation, so it propagates along paths of $G_{M,K}$. $\square$

Uses: independence of distinct edges; union bound.

Referee check. (1) Independence is used only for the events $E_y, y \in \mathcal{Y}$, for a fixed pair (x, x') ; these involve pairwise disjoint sets of edges. Across different pairs no independence is used, only the union bound. (2) The bound $|\mathcal{Y}| \geq N^{2\delta} - 2$ holds for every $x' \leq M$, so the union bound is uniform. (3) The vertices y may lie above M ; this is harmless, since paths may use any vertex of $V_{N,K}$. (4) If $K = 0$, the vertex 1 is included and the argument is unchanged.

4.2. Doubling.

Lemma 4.2 (Doubling). [PROVEN; new in this form, elementary]

Lean: `doubling, doubling_connected, reachable_two_pow_mul`

Assume $M \geq 4(K+1)$, so that $Y \geq 2$. Then every $x \in S_M$ is joined in $G_{M,K}$ to the largest power of 2 not exceeding M . In particular S_M lies in a single connected component of $G_{M,K}$.

Proof. Let 2^b be the power of 2 with $M/2 < 2^b \leq M$. Then $2^b > M/2 \geq 2(K+1) > K$ and $P^+(2^b) = 2 \leq Y$, so $2^b \in S_M$ (in particular $S_M \neq \emptyset$, which is not needed below). For $x \in S_M$ let $\Omega'(x)$ be the number of odd prime factors of x counted with multiplicity. We show by induction on $\Omega'(x)$ that x is joined to 2^b by a path in $G_{M,K}$ that stays inside S_M .

Let $a \geq 0$ be maximal with $2^a x \leq M$, and put $y := 2^a x$. Then $M/2 < y \leq M$. The vertices $x, 2x, \dots, 2^a x = y$ lie in $(K, M]$, each divides the next, and they all lie in S_M (their odd part is that of x , and $2 \leq Y$). So they form a path in $G_{M,K}$. If $\Omega'(x) = 0$, then y is a power of 2 in $(M/2, M]$, so $y = 2^b$. Otherwise let p be an odd prime dividing x , and put $x_1 := y/p$. Then $x_1 \mid y$, $x_1 < y \leq M$, and

$$x_1 = \frac{y}{p} > \frac{M/2}{Y} = K+1,$$

using $p \leq P^+(x) \leq Y$. So $\{x_1, y\}$ is an edge of $G_{M,K}$. Moreover $P^+(x_1) \leq \max(P^+(x), 2) \leq Y$, so $x_1 \in S_M$, and $\Omega'(x_1) = \Omega'(x) - 1$. By induction x_1 is joined to 2^b , hence so is x . $\square$

Uses: nothing.

Corollary 4.3. [PROVEN]

Lean: `sigma_mutual, goodSet_mem_smoothFibre, sigma_subset_smoothFibre, sigmaSet_mutual, sigmaSet_closed, sigmaSet_subset`

Assume $M \geq 4(K+1)$, and let Σ be the set of vertices $v \in V_{N,K}$ that are mutually reachable in $\mathcal{D}_\rho(N, K)$ with some element of S_M . On $\mathcal{R}$, any two elements of S_M lie in the same SCC. Hence, on $\mathcal{R}$, Σ is strongly connected and closed under mutual reachability, so it is the SCC containing S_M (recall $S_M \neq \emptyset$). Since $M \leq N$, for every outcome $\Sigma \subset F_1$.

Proof. The first claim follows from Lemmas 4.1 and 4.2. If $a, b \in \Sigma$ are mutually reachable with $d, d' \in S_M$, then $a \sim d \sim d' \sim b$ on $\mathcal{R}$. Closure under mutual reachability is immediate from the definition. For $x \in S_M$ we have $P^+(x) \leq Y < N/(K+1)$, so $P^+(x) \leq B$ and $S_M \subset F_1 = V_1$. If $v \in \Sigma$ is mutually reachable with $d \in S_M$, then $\{v, d\}$ is strongly connected, so it lies in a single fibre by Proposition 3.2, which must be V_1 . Hence $v \in F_1$. $\square$

4.3. Attachment. For $m \in V_{N,K}$ with $m > M$ let $D(m) := \{d \in S_M : d \mid m\}$ and $s(m) := |D(m)|$. Let A_m be the event that there are $d, d' \in D(m)$ with $m \rightarrow d$ and $d' \rightarrow m$.

Lemma 4.4 (Attachment). [PROVEN]

Lean: `prob_attachEv, prob_attachEv_eq, prob_attachEv_ge, prob_all_bit, attach_mutual, prob_attach`

Assume $M \geq 4(K+1)$ and let $m \in V_{N,K}$ with $m > M$. Then $\mathbb{P}(A_m) \geq 1 - \rho^{s(m)} - (1 - \rho)^{s(m)} \geq 1 - 2\lambda^{s(m)}$, with equality in the first step if $s(m) \geq 1$. On $A_m \cap \mathcal{R}$ we have $m \in \Sigma$.

Proof. Every $d \in D(m)$ satisfies $d \leq M < m$, so the edges $\{d, m\}$, $d \in D(m)$, are $s(m)$ distinct edges of $G_{N,K}$ with independent orientations. The complement A_m^c is contained in the union of the events “all point $m \rightarrow d$ ” and “all point $d \rightarrow m$ ”, which are cylinders with probabilities $(1 - \rho)^{s(m)}$ and $\rho^{s(m)}$; for $s(m) \geq 1$ the union is disjoint and equals A_m^c . On $A_m \cap \mathcal{R}$ we have $m \rightarrow d$, a directed path $d \rightsquigarrow d'$ in $\mathcal{D}_\rho(N, K)$ (Corollary 4.3, since $d, d' \in S_M$), and $d' \rightarrow m$. This is a closed walk through m and d , so m is mutually reachable with $d \in S_M$, that is, $m \in \Sigma$. $\square$

Uses: Corollary 4.3; independence of distinct edges.

Referee check. The events A_m and $\mathcal{R}$ are *not* independent: the edges $\{d, m\}$ may be among those used in the definition of $\mathcal{R}$. We never use independence between them. Below we only use $\mathbb{P}(A_m \cap \mathcal{R}) \geq \mathbb{P}(A_m) - \mathbb{P}(\mathcal{R}^c)$.

4.4. The arithmetic core.

Lemma 4.5 (Many divisors in the lower layer). [PROVEN; new]

Lean: `core_a, core_b, core_c, core_divisors, core_d, core_d'`

Let $\eta \in (0, 1)$, let $s_0 \geq 1$ be an integer and $\delta \in (0, \eta/8]$, and put $z := N^{\delta/s_0}$. Assume

$$(H1) \ K \leq N^{1-\eta}, \quad (H2) \ N^{\eta/4} \geq 4, \quad (H3) \ z \geq 2.$$

Let

$$\mathcal{U} := \{m \in \mathbb{Z} : N^{1-\delta} < m \leq N, \ P^+(m) \leq Y, \ \omega_z(m) \geq s_0\}.$$

Then:

- (a) $M \geq 4(K+1)$;
- (b) $\mathcal{U} \subset F_1$, and every $m \in \mathcal{U}$ satisfies $m > M$;
- (c) $s(m) \geq s_0 + 1$ for every $m \in \mathcal{U}$;
- (d) $|F_1 \setminus \mathcal{U}| \leq N^{1-\delta} + N \left(\frac{4\delta}{\eta} + \frac{2(1+C_1)}{\eta \log N} \right) + \#\{m \leq N : \omega_z(m) < s_0\}.$

Proof. Note first that $K+1 \leq 2N^{1-\eta}$ by (H1), and that $\eta - 2\delta \geq \frac{3}{4}\eta$ and $\eta - 6\delta \geq \frac{1}{4}\eta$ because $\delta \leq \eta/8$.

(a) $M/(K+1) \geq N^{1-2\delta}/(2N^{1-\eta}) = \frac{1}{2}N^{\eta-2\delta} \geq \frac{1}{2}(N^{\eta/4})^3 \geq 32$, by (H2).

(b) Let $m \in \mathcal{U}$. Then $m > N^{1-\delta} > N^{1-2\delta} = M$ and $m > N^{1-\delta} \geq N^{1-\eta} \geq K$, so $m \in V_{N,K}$. Also $P^+(m) \leq Y < N/(K+1)$, so $P^+(m) \leq B$ and $m \in F_1$.

(c) Fix $m \in \mathcal{U}$ and put $E_0 := N^{2\delta}$ and $E_1 := N^{1-\delta}/(K+1)$.

Step 1: divisors in $[E_0, E_1]$ give elements of $D(m)$. Let $e \mid m$ with $E_0 \leq e \leq E_1$ and put $d := m/e$. Then $d \leq N/E_0 = M$ and $d \geq m/E_1 > N^{1-\delta}(K+1)/N^{1-\delta} = K+1$. Also $P^+(d) \leq P^+(m) \leq Y$. Hence $d \in S_M$ and $d \mid m$, so $d \in D(m)$. Distinct e give distinct d .

Step 2: a reserve of small primes. Since $\omega_z(m) \geq s_0$, we may choose distinct primes $p_1, \dots, p_{s_0} \leq z$ dividing m . Put $r := p_1 \cdots p_{s_0}$, so $r \mid m$ and $r \leq z^{s_0} = N^\delta$. Then $m' := m/r$ is an integer with $m' \geq m/N^\delta > N^{1-2\delta}$.

Step 3: a base divisor $e_0 \mid m'$ with $E_0 \leq e_0 \leq E_1/z$. First note that

$$\frac{E_1}{z} = \frac{N^{1-\delta-\delta/s_0}}{K+1} \geq \frac{N^{1-2\delta}}{K+1} = 2Y.$$

Case (i): m' has a prime factor $q \geq E_0$. Then $E_0 \leq q \leq P^+(m) \leq Y \leq E_1/z$, and we take $e_0 := q$.

Case (ii): every prime factor of m' is $< E_0$. Write $m' = q_1 q_2 \cdots q_t$ with primes q_i (with multiplicity, in any order), and let $\pi_i := q_1 \cdots q_i$. Then $\pi_0 = 1 < E_0 \leq N^{1-2\delta} < m' = \pi_t$, where $E_0 \leq N^{1-2\delta}$ because $\delta \leq \frac{1}{4}$. Let j be the least index with $\pi_j \geq E_0$. Then $j \geq 1$ and $\pi_j = \pi_{j-1} q_j < E_0^2 = N^{4\delta}$. Moreover

$$\frac{E_1}{z} \geq \frac{N^{1-2\delta}}{K+1} \geq \frac{1}{2}N^{\eta-2\delta} = N^{4\delta} \cdot \frac{1}{2}N^{\eta-6\delta} \geq N^{4\delta} \cdot \frac{1}{2}N^{\eta/4} \geq N^{4\delta},$$

by (H2). So $e_0 := \pi_j$ satisfies $E_0 \leq e_0 \leq E_1/z$.

Step 4. Since $e_0 \mid m'$, we have $e_0 r \mid m$, so $e_0 p_i \mid m$ for each i . The s_0+1 numbers $e_0, e_0 p_1, \dots, e_0 p_{s_0}$ are distinct divisors of m , and they all lie in $[E_0, E_1]$ because $e_0 p_i \leq e_0 z \leq E_1$. By Step 1, $s(m) \geq s_0 + 1$.

(d) If $m \in F_1 \setminus \mathcal{U}$, then $m \leq N^{1-\delta}$, or $Y < P^+(m) \leq B$, or $\omega_z(m) < s_0$. The first set has at most $N^{1-\delta}$ elements. The second is empty if $Y \geq B$. Otherwise its size is at most $\sum_{Y < p \leq B} [N/p] \leq$

$N \sum_{Y < p \leq B} \frac{1}{p}$. We have $Y \geq 2$ by (a). Moreover

$$\log B - \log Y \leq \log \frac{N}{K+1} - \log \frac{N^{1-2\delta}}{2(K+1)} = 2\delta \log N + \log 2,$$

and

$$\log Y \geq \log \frac{N^{1-2\delta}}{4N^{1-\eta}} = (\eta - 2\delta) \log N - \log 4 \geq \frac{3}{4}\eta \log N - \log 4 \geq \frac{1}{2}\eta \log N,$$

by (H2). By Lemma 2.1(b) with $C_1 = 18$ and $\log(1+t) \leq t$,

$$\sum_{Y < p \leq B} \frac{1}{p} \leq \frac{\log B - \log Y}{\log Y} + \frac{C_1}{\log Y} \leq \frac{2\delta \log N + 1 + C_1}{\frac{1}{2}\eta \log N} = \frac{4\delta}{\eta} + \frac{2(1+C_1)}{\eta \log N}. \quad \square$$

Uses: Lemma 2.1(b).

Referee check. (1) No “almost all” statement is used inside the proof. Every $m \in \mathcal{U}$ has $s(m) \geq s_0 + 1$ deterministically, and the exceptional set is counted explicitly in (d). (2) The divisor e_0 is found by a deterministic case analysis. Case (i) covers a prime factor of any size in $[E_0, P^+(m)]$, including sizes close to Y ; this is where the reserve primes must be tiny ($\leq z$), because e_0 may be as large as Y . (3) The bounds are uniform in $K \leq N^{1-\eta}$; K enters only through (H1).

4.5. The lower bound.

Proposition 4.6 (Lower bound). [PROVEN]

Lean: `lower_bound`, `lower_bound_err'`, `fixedN_sandwich'`, `eq_4_2_core`, `eq_4_2`

Assume the hypotheses of Lemma 4.5, and put

$$\text{Err} := N^{1-\delta} + N \left(\frac{4\delta}{\eta} + \frac{2(1+C_1)}{\eta \log N} \right) + \#\{m \leq N : \omega_z(m) < s_0\} + 2\lambda^{s_0+1}N.$$

Then

$$(2) \quad \mathbb{E}[\Phi_K] \geq \Psi(N, B) - K - \text{Err} - N\mathbb{P}(\mathcal{R}^c),$$

$$(3) \quad \mathbb{E}[|F_1 \setminus \Sigma| \mathbf{1}_{\mathcal{R}}] \leq \text{Err}.$$

Here $C_1 = 18$, Σ is as in Corollary 4.3, and $\mathbb{P}(\mathcal{R}^c) \leq N^2 \exp(-q(N^{2\delta} - 2))$ by Lemma 4.1.

Proof. By Lemma 4.5(a), Corollary 4.3 and Lemma 4.4 apply. Let $X := \mathbf{1}_{\mathcal{R}} \cdot \#\{m \in \mathcal{U} : A_m \text{ occurs}\}$. On $\mathcal{R}$, every $m \in \mathcal{U}$ for which A_m occurs lies in Σ (Lemmas 4.5(b) and 4.4), so $X \leq |\Sigma| \leq \Phi_K$. On $\mathcal{R}^c$, $X = 0 \leq \Phi_K$. By Lemmas 4.5(c) and 4.4,

$$\mathbb{E}[\Phi_K] \geq \mathbb{E}[X] = \sum_{m \in \mathcal{U}} \mathbb{P}(A_m \cap \mathcal{R}) \geq \sum_{m \in \mathcal{U}} (\mathbb{P}(A_m) - \mathbb{P}(\mathcal{R}^c)) \geq |\mathcal{U}|(1 - 2\lambda^{s_0+1}) - N\mathbb{P}(\mathcal{R}^c).$$

Moreover $|\mathcal{U}| \geq |F_1| - |F_1 \setminus \mathcal{U}|$ and $|F_1| = \Psi(N, B) - \Psi(K, B) \geq \Psi(N, B) - K$. Also $2\lambda^{s_0+1}|\mathcal{U}| \leq 2\lambda^{s_0+1}N$. Combining these with Lemma 4.5(d) proves (2). For (3): on $\mathcal{R}$, every $m \in \mathcal{U}$ for which A_m occurs lies in Σ (Lemma 4.4), so $F_1 \setminus \Sigma \subset (F_1 \setminus \mathcal{U}) \cup \{m \in \mathcal{U} : A_m^c\}$. Taking cardinalities pointwise, $|F_1 \setminus \Sigma| \mathbf{1}_{\mathcal{R}} \leq |F_1 \setminus \mathcal{U}| + \sum_{m \in \mathcal{U}} \mathbf{1}_{A_m^c}$ for every outcome, and $\mathbb{P}(A_m^c) \leq 2\lambda^{s_0+1}$ by Lemmas 4.5(c) and 4.4. So

$$\mathbb{E}[|F_1 \setminus \Sigma| \mathbf{1}_{\mathcal{R}}] \leq |F_1 \setminus \mathcal{U}| + \sum_{m \in \mathcal{U}} \mathbb{P}(A_m^c) \leq |F_1 \setminus \mathcal{U}| + 2\lambda^{s_0+1}N \leq \text{Err}. \quad \square$$

Uses: Lemmas 4.1, 4.2, 4.4, 4.5; Corollary 4.3.

Referee check. (1) The only probabilistic inputs are exact computations for independent edges (Lemmas 4.1 and 4.4), the union bound, and $\mathbb{P}(A \cap \mathcal{R}) \geq \mathbb{P}(A) - \mathbb{P}(\mathcal{R}^c)$. (2) The inequality $\Phi_K \geq X$ holds pointwise on the whole probability space. (3) Proposition 4.6 is a statement for each fixed N , with no limits, so there is no interchange of limits. Uniformity in $K \leq N^{1-\eta}$ is explicit. Note that the term $\#\{m \leq N : \omega_z(m) < s_0\}$ does not depend on K .

5. PROOFS OF THE MAIN RESULTS

Proof of Theorem 1.3. The upper bound $\Phi_K \leq \Psi(N, B)$ is Proposition 3.2. For the lower bound, let N be large and put

$$\ell := \log \log N, \quad \eta := \frac{1}{\ell}, \quad \delta := \frac{1}{8\ell^2}, \quad s_0 := \max\left(1, \left\lceil \frac{\log \ell}{\log(1/\lambda)} \right\rceil\right), \quad z := N^{\delta/s_0}.$$

Then $\delta \leq \eta/8$ and $\lambda^{s_0} \leq 1/\ell$. Write $t := \log N$, so $\ell = \log t$. Since $s_0 = O_\rho(\log \ell)$, and $A(\log t)^k \leq t$ for large t for every fixed A and k , there is $N_0(\rho)$ such that for $N \geq N_0(\rho)$ all of the following hold:

- (R1) $\ell > 1$, $N^{\eta/4} = \exp(\frac{t}{4\ell}) \geq 4$ and $z = \exp(\frac{t}{8\ell^2 s_0}) \geq 2$;
- (R2) $L := \sum_{p \leq z} \frac{1}{p} \geq \log \log z - 13 = \ell - \log(8\ell^2 s_0) - 13 \geq \ell/2 + 2s_0$ (Lemma 2.1(c)), so $L \geq \max(2s_0, \ell/2)$;
- (R3) $N^{1-\eta} \leq N/\ell$ and $N^{1-\delta} \leq N/\ell$, that is, $\ell \log \ell \leq t$ and $8\ell^2 \log \ell \leq t$;
- (R4) $2(1 + C_1)/(\eta \log N) = 38\ell/t \leq 1/(2\ell)$, that is, $76\ell^2 \leq t$;
- (R5) $N \cdot N^2 \exp(-q(N^{2\delta} - 2)) \leq 1$, that is, $3t \leq q(\exp(t/(4\ell^2)) - 2)$, which holds because $N^{2\delta} = \exp(t/(4\ell^2))$ grows faster than any power of t ;
- (R6) $\sqrt{N} \leq N/(2\ell)$ and $2c_0 N/\log N \leq N/(2\ell)$.

These are the conditions `GoodT` of the formalisation (`eventually_goodN`). Note also $N/\ell \geq 1$. *Case* $K \leq N^{1-\eta}$. Hypotheses (H1)–(H3) of Lemma 4.5 hold by (R1). By Lemma 2.4 and (R2), $\#\{m \leq N : \omega_z(m) < s_0\} \leq 12N/L \leq 24N/\ell$. Also $4\delta/\eta = 1/(2\ell)$ and $2\lambda^{s_0+1} \leq 2/\ell$. Proposition 4.6 together with (R3)–(R5) gives

$$\Psi(N, B) - \mathbb{E}[\Phi_K] \leq \frac{N}{\ell} + \frac{N}{\ell} + N\left(\frac{1}{2\ell} + \frac{1}{2\ell}\right) + \frac{24N}{\ell} + \frac{2N}{\ell} + 1 \leq \frac{29N}{\ell} + 1 \leq \frac{30N}{\ell}.$$

In this case the terms of `Err` satisfy $\text{Err} \leq 28N/\ell$, and $N\mathbb{P}(\mathcal{R}^c) \leq 1$; we use this again in the proof of Corollary 1.5.

Case $K > N^{1-\eta}$. Then $B \leq N/(K+1) < N^\eta$, so $\log B < \eta \log N = t/\ell$. Since $K < N$, $B \geq 1$, and Lemma 2.3 and (R6) give

$$\Psi(N, B) \leq \sqrt{N} + \frac{2N \log B}{\log N} + \frac{2c_0 N}{\log N} \leq \frac{N}{2\ell} + \frac{2N}{\ell} + \frac{N}{2\ell} = \frac{3N}{\ell}.$$

Hence $0 \leq \Psi(N, B) - \mathbb{E}[\Phi_K] \leq \Psi(N, B) \leq 30N/\ell$. □

Uses: Propositions 3.2, 4.6; Lemmas 2.1, 2.3, 2.4, 4.5.

Referee check. (1) The parameters η, δ, s_0, z depend on N only, not on K . So the two cases cover all K with one common bound. (2) The threshold $N_0(\rho)$ depends on ρ through s_0 (via λ) and through q in (R5). (3) Proposition 4.6 is applied at each fixed N , so no limit is interchanged.

Proof of Theorem 1.2. The deterministic bound is Proposition 3.2. The limit follows from Theorem 1.3, since $30/\log \log N \rightarrow 0$. □

Remark 5.1 (Explicit form for $K \leq \sqrt{N} - 1$). [PROVEN]

Lean: `remark_5_1, smallK_psi, N_sub_psi_eq`

For $N \geq N'_0(\rho)$ and $0 \leq K \leq \sqrt{N} - 1$,

$$\left| 1 - \frac{\mathbb{E}[\Phi_K]}{N} - \log \frac{\log N}{\log B} \right| \leq \frac{31}{\log \log N}.$$

Proof. Since $N/(K+1) \geq \sqrt{N}$, we have $B \geq \lfloor \sqrt{N} \rfloor$, and every prime $p > B$ satisfies $p > \sqrt{N}$. So each $n \leq N$ has at most one prime factor $> B$, and that factor divides n exactly once.

Hence $N - \Psi(N, B) = \sum_{B < p \leq N} \lfloor N/p \rfloor$, which differs from $N \sum_{B < p \leq N} \frac{1}{p}$ by at most $\pi(N)$. By Lemma 2.1(b),(d) and $\log B \geq \frac{1}{4} \log N$ (for $N \geq 16$),

$$\left| 1 - \frac{\Psi(N, B)}{N} - \log \frac{\log N}{\log B} \right| \leq \frac{C_1}{\log B} + \frac{\pi(N)}{N} \leq \frac{4C_1 + 5}{\log N} = \frac{77}{\log N} \leq \frac{1}{\log \log N}$$

for large N . Combine this with $0 \leq (\Psi(N, B) - \mathbb{E}[\Phi_K])/N \leq 30/\log \log N$ (Theorem 1.3). $\square$

Proof of Corollary 1.4. Write $\mathbb{E}[\Phi_K]/N = \Psi(N, N/(K+1))/N - \text{gap}/N$ with $\text{gap} := \Psi(N, N/(K+1)) - \mathbb{E}[\Phi_K]$. By Theorem 1.3, $0 \leq \text{gap}/N \leq 30/\log \log N \rightarrow 0$, and $\Psi(N, N/(K+1))/N \rightarrow \rho_{\text{Dick}}(1/(1-\theta))$ by Lemma 2.7.

Write $P(\theta)$ for the profile. On $(-\infty, 1)$, P is continuous because ρ_{Dick} and $\theta \mapsto 1/(1-\theta)$ are; on $(1, \infty)$ it is 0. At $\theta = 1$ it is continuous because $0 \leq P(\theta) \leq 2(1-\theta)$ for $\theta \in [0, 1)$, by $\rho_{\text{Dick}}(u) \leq 2/u$ (Lemma 2.3). On $[0, 1)$, P is strictly decreasing because $\theta \mapsto 1/(1-\theta)$ is strictly increasing with values in $[1, \infty)$, where ρ_{Dick} is strictly decreasing (Lemma 2.2); and $P(1) = 0 < P(\theta)$ for $\theta < 1$ since $\rho_{\text{Dick}} > 0$. For $0 \leq \theta \leq \frac{1}{2}$ we have $u = 1/(1-\theta) \in [1, 2]$, where $\rho_{\text{Dick}}(u) = 1 - \log u$. $\square$

Uses: Theorem 1.2; Lemmas 2.2, 2.3, 2.7.

Proof of Corollary 1.5. Use the parameters and $N_0(\rho)$ from the proof of Theorem 1.3. The SCCs are the classes of mutual reachability; for $K < N$ a largest one exists, so rules Φ_K^* exist. Write $S := \Phi_K^*$, so $|S| = \Phi_K$.

(a) By Proposition 3.2, $|\Phi_K - \Psi(N, B)| = \Psi(N, B) - \Phi_K$ pointwise, so $\mathbb{E}|\Phi_K - \Psi(N, B)| = \Psi(N, B) - \mathbb{E}[\Phi_K] \leq 30N/\ell$ by Theorem 1.3. For $\varepsilon > 0$, Markov's inequality gives $\mathbb{P}(|\Phi_K/N - \Psi(N, B)/N| \geq \varepsilon) \leq 30/(\varepsilon\ell)$ for every $K < N$, and this tends to 0.

(b), (c): *Case $K \leq N^{1-\eta}$.* Then the hypotheses of Lemma 4.5 hold, so $M \geq 4(K+1)$ and $M \leq N$. Work on $\mathcal{R}$. By Corollary 4.3, $\Sigma \subset F_1$ is strongly connected and closed under mutual reachability. Hence every SCC C either equals Σ or is disjoint from it: if $x \in C \cap \Sigma$, then $C = \Sigma$. Every SCC lies in a fibre V_n (Proposition 3.2); if $n = 1$ then $C \subset F_1$, and if $n > 1$ then $|C| \leq K$ (Lemma 3.1(c)). So every SCC $C \neq \Sigma$ has $|C| \leq W + K$, where $W := |F_1 \setminus \Sigma|$. Also $|\Sigma| \leq \Phi_K = |S|$, and $|F_1| = W + |\Sigma|$.

- If $S = \Sigma$, then $|F_1 \triangle S| = |F_1 \setminus \Sigma| = W$.
- If $S \neq \Sigma$ and $S \subset F_1$, then $|F_1 \triangle S| = |F_1| - |S| \leq |F_1| - |\Sigma| = W$.
- If $S \neq \Sigma$ and $|S| \leq K$, then $|F_1 \triangle S| \leq |F_1| + |S| = W + |\Sigma| + |S| \leq W + 2K$.

For $\Phi_K^{(2)}$, let $C \neq S$ be an SCC. If $C \neq \Sigma$, then $|C| \leq W + K$. If $C = \Sigma$, then $S \neq \Sigma$, so $|C| = |\Sigma| \leq |S| \leq W + K$.

Thus, on $\mathcal{R}$, $|F_1 \triangle S| \leq W + 2K$ and $\Phi_K^{(2)} \leq W + K$. For every outcome both quantities are at most N , since all sets involved lie in $V_{N,K}$. Hence

$$\mathbb{E}|F_1 \triangle S| \leq 2K + \mathbb{E}[W \mathbf{1}_{\mathcal{R}}] + N\mathbb{P}(\mathcal{R}^c), \quad \mathbb{E}[\Phi_K^{(2)}] \leq K + \mathbb{E}[W \mathbf{1}_{\mathcal{R}}] + N\mathbb{P}(\mathcal{R}^c).$$

By (3) and the estimates in the proof of Theorem 1.3, $\mathbb{E}[W \mathbf{1}_{\mathcal{R}}] \leq \text{Err} \leq 28N/\ell$ and $N\mathbb{P}(\mathcal{R}^c) \leq 1 \leq N/\ell$. Hence $\mathbb{E}|F_1 \triangle S| \leq 2K + 29N/\ell$ and $\mathbb{E}[\Phi_K^{(2)}] \leq K + 29N/\ell$ in this case.

Case $K > N^{1-\eta}$. Here $|F_1 \triangle S| \leq |F_1| + |S| \leq 2\Psi(N, B) \leq 6N/\ell$, using $|F_1| \leq \Psi(N, B)$ (Lemma 3.1(c)) and $|S| = \Phi_K \leq \Psi(N, B)$; and $\Phi_K^{(2)} \leq \Phi_K \leq \Psi(N, B) \leq 3N/\ell$. $\square$

Uses: Propositions 3.2, 4.6; Corollary 4.3; Lemma 3.1; Theorem 1.3.

Remark 5.2 (The term K in Corollary 1.5(c) is needed). [PROVEN]

Lean: remark_5_2, secondSCC_ge, expect_secondSCC_ge, prob_PhiK_le_K

Fix $\rho \in (0, 1)$ and let $K = K(N)$ satisfy $K \rightarrow \infty$ and $K \leq \sqrt{N}/2$. Then, for any rule choosing a largest SCC, $\liminf_{N \rightarrow \infty} \mathbb{E}[\Phi_K^{(2)}]/K \geq \frac{1}{2}$. (All $o(1)$ terms below are as $N \rightarrow \infty$ along this sequence and depend on ρ and on the sequence.)

Proof. Let $N \geq 16$. Then $(K+1)^2 \leq N$, and $B \geq \sqrt{N}$ because $N < (K+1)(B+1)$ and $K+1 \leq \sqrt{N}/2 + 1$. Let p be a prime in $(B, 2B]$, which exists by Bertrand's postulate. For $1 \leq s \leq N/p$ we have $s < N/B \leq \sqrt{N} \leq B < p$. So s is B -smooth and $p \nmid s$, which gives $R_B(ps) = p$. Also $ps \geq p > K$. Together with Lemma 3.1(c) this shows $V_p = \{ps : 1 \leq s \leq N'\}$ with $N' := \lfloor N/p \rfloor$. Since $ps \mid ps'$ if and only if $s \mid s'$, the map $s \mapsto ps$ is an isomorphism from $G_{N'}$ onto $G_N[V_p]$ that preserves the default orientation. Hence the restriction of $\mathcal{D}_\rho(N, K)$ to V_p has the law of $\mathcal{D}_\rho(N')$. As V_p is a union of connected components of $G_{N,K}$, its SCCs are SCCs of $\mathcal{D}_\rho(N, K)$. We have $K/2 \leq N' \leq K$, because $N < p(N'+1) \leq 2B(N'+1)$, $B(K+1) \leq N$ and $pN' \leq N < (K+1)(B+1) \leq (K+1)p$. So $N' \rightarrow \infty$. So by [9, Corollary 2] (or by Theorem 1.2 with $K=0$, since $\Psi(N', N') = N'$) the largest SCC inside V_p has expected size $(1 - o(1))N'$.

Next, $\log B \geq \frac{1}{2} \log N$, so the estimate in the proof of Remark 5.1 gives $\Psi(N, B)/N \geq 1 - \log 2 - 77/\log N \geq \frac{1}{4}$ for large N . Also $K \leq \sqrt{N}/2 \leq N/8$. By Theorem 1.3 and Markov's inequality,

$$\mathbb{P}(\Phi_K \leq K) \leq \frac{\mathbb{E}[\Psi(N, B) - \Phi_K]}{\Psi(N, B) - K} \leq \frac{30N/\log \log N}{N/8} = \frac{240}{\log \log N} \rightarrow 0.$$

On the event $\{\Phi_K > K\}$ the largest SCC lies in $F_1 = V_1$ (Lemma 3.1(c)), which is disjoint from V_p , so $\Phi_K^{(2)} \geq \#\Phi(\mathcal{D}_\rho(N, K)[V_p])$. Therefore

$$\mathbb{E}[\Phi_K^{(2)}] \geq \mathbb{E}[\#\Phi(\mathcal{D}_\rho(N, K)[V_p])] - N' \mathbb{P}(\Phi_K \leq K) = (1 - o(1))N' \geq (\frac{1}{2} - o(1))K. \quad \square$$

Remark 5.3 (Uniformity in ρ ; slowly vanishing ρ). [PROVEN]

Lean: `remark_5_3_uniform`, `remark_5_3_uniform_main`, `GoodT.mono`, `remark_5_3_slow`, `slow_rho_bound_explicit`

The proof depends on ρ only through $q = \rho(1 - \rho)$ and $\lambda = \max(\rho, 1 - \rho)$. Let $0 < \rho_0 \leq \frac{1}{2}$. For $\rho \in [\rho_0, 1 - \rho_0]$ we have $\lambda \leq \lambda_0 := 1 - \rho_0$ and $q \geq q_0 := \rho_0(1 - \rho_0)$. Then $s_0(\lambda) \leq s_0(\lambda_0)$. Conditions (R1)–(R4) and (R6) involve only N and s_0 , and they become weaker as s_0 decreases. Condition (R5) becomes weaker as q increases, because its left side is decreasing in q when $N^{2\delta} \geq 2$. So (R1)–(R6) at ρ_0 imply (R1)–(R6) at every $\rho \in [\rho_0, 1 - \rho_0]$. Hence Theorems 1.2 and 1.3 hold uniformly for $\rho \in [\rho_0, 1 - \rho_0]$, with $N_0 = N_0(\rho_0)$. More generally, Theorem 1.2 remains true for $\rho = \rho_N$ provided $\min(\rho_N, 1 - \rho_N) \log \log N \rightarrow \infty$. This follows from a quantitative form: for every $\varepsilon \in (0, 1]$ there is $N_0(\varepsilon)$ such that $\Psi(N, N/(K+1)) - \mathbb{E}[\Phi_K] \leq \varepsilon N$ for all $N \geq N_0(\varepsilon)$, all $0 \leq K < N$ and all $\rho \in (0, 1)$ with $\min(\rho, 1 - \rho) \log \log N \geq 8 \log(16/\varepsilon) + 8$. To prove it, write $\ell = \log \log N$ and apply Proposition 4.6 with $\eta = \varepsilon/4$, $\delta = \varepsilon\eta/32 \leq \eta/8$ and $s_0 := \lfloor \ell/4 \rfloor$, so that $\ell/8 \leq s_0 \leq \ell/4$. Each of the seven error terms is at most $\varepsilon N/8$ for large N . In particular, Lemma 2.1(c) gives $L \geq \log \delta + \ell - \log s_0 - 13 \geq \ell/2$, so the ω_z term is at most $12N/L \leq 24N/\ell$. Next, $2\lambda^{s_0+1} \leq 2 \exp(-\min(\rho, 1 - \rho)(s_0 + 1)) \leq \varepsilon/8$. Finally $N\mathbb{P}(\mathcal{R}^c) \leq N^3 \exp(-q(N^{2\delta} - 2)) \leq \varepsilon N/8$, since $q \geq \frac{1}{2} \min(\rho, 1 - \rho) \geq 1/(2\ell)$. The case $K > N^{1-\eta}$ is handled by Lemma 2.3 as before.

6. REMOVING THE VERTICES OF HIGHEST DEGREE

For $T \subset [N]$ write $\deg_T(v)$ for the degree of v in $G_N - T$. In G_N ,

$$\deg(m) = (\tau(m) - 1) + (\lfloor N/m \rfloor - 1).$$

Put $\tau_N := \max_{n \leq N} \tau(n)$. Then $\deg_T(m) \leq \deg(m) \leq \tau_N + N/m$ for every T and $m \notin T$. The proofs below use only this upper bound and the following lower bound: if every element of T is at most A and $1 \leq v \leq A \leq N$, then the multiples of v in $(A, N]$ are neighbours of v in $G_N - T$, so $\deg_T(v) \geq N/v - A/v - 1$.

Lean: `deg_le`, `degT_le_deg`, `degT_ge`

Lemma 6.1 (Monotonicity). [PROVEN]

Lean: `maxSCC_mono`, `maxSCC_Ioc_antitone`

If D is a directed graph and $U \subset V(D)$, then $\#\Phi(D[U]) \leq \#\Phi(D)$.

Proof. An SCC of $D[U]$ is strongly connected in D , so it is contained in an SCC of D . $\square$

Lemma 6.2 (Sandwich). [PROVEN]

Lean: `sandwich_static`, `sandwich_adaptive`

Let $0 < \varepsilon \leq \frac{1}{2}$ and $1 \leq K < N$ with $\frac{N}{K} \cdot \frac{\varepsilon}{2} \geq \hat{\tau} + 5$, where $\hat{\tau} \geq \tau(n)$ for all $n \leq N$ (for instance $\hat{\tau} = \tau_N$). Let T be a static or an adaptive top- K set as in Corollary 1.6. Then

$$\{1, \dots, \lfloor (1 - \varepsilon)K \rfloor\} \subset T \subset \{1, \dots, \lfloor (1 + \varepsilon)K \rfloor\}.$$

Proof. Static, upper inclusion. Suppose $w \in T$ with $w > (1 + \varepsilon)K$. Then $|T \cap [1, K]| \leq K - 1$, so some $v \leq K$ has $v \notin T$. Now

$$\deg(v) \geq \lfloor N/v \rfloor - 1 \geq N/K - 2 \quad \text{and} \quad \deg(w) < \tau_N + N/((1 + \varepsilon)K).$$

The static condition gives $\deg(w) \geq \deg(v)$, hence $\frac{N}{K} \cdot \frac{\varepsilon}{1 + \varepsilon} < \tau_N + 2$. This contradicts the hypothesis, since $\frac{\varepsilon}{1 + \varepsilon} \geq \frac{\varepsilon}{2}$.

Static, lower inclusion. Suppose $v \leq (1 - \varepsilon)K$ and $v \notin T$. Every $t \in T$ satisfies $\tau_N + N/t \geq \deg(t) \geq \deg(v) \geq N/v - 2 \geq \frac{N}{(1 - \varepsilon)K} - 2$. If $t > (1 - \frac{\varepsilon}{2})K$, this gives $\frac{N}{K} \left(\frac{1}{1 - \varepsilon} - \frac{1}{1 - \varepsilon/2} \right) < \tau_N + 2$. The left side is at least $\frac{N}{K} \cdot \frac{\varepsilon}{2}$, a contradiction. So $T \subset [1, (1 - \frac{\varepsilon}{2})K]$. But this set has fewer than K elements, which contradicts $|T| = K$.

Adaptive, upper inclusion. We show by induction on $j < K$ that $T_j := \{u_1, \dots, u_j\} \subset [1, (1 + \varepsilon)K]$ implies $u_{j+1} \leq (1 + \varepsilon)K$. Since $|T_j| < K$, some $v \leq K$ has $v \notin T_j$. No multiple of v in $((1 + \varepsilon)K, N]$ has been removed, so

$$\deg_{T_j}(v) \geq \frac{N - (1 + \varepsilon)K}{v} - 1 \geq \frac{N}{K} - 3.$$

Every $w > (1 + \varepsilon)K$ has $\deg_{T_j}(w) < \tau_N + N/((1 + \varepsilon)K)$, and this is at most $N/K - 3$ by the hypothesis. Hence the maximiser u_{j+1} satisfies $u_{j+1} \leq (1 + \varepsilon)K$.

Adaptive, lower inclusion. Suppose $v \leq (1 - \varepsilon)K$ and $v \notin T_K$. For every $j < K$, using $T_j \subset [1, (1 + \varepsilon)K]$,

$$\deg_{T_j}(v) \geq \frac{N - (1 + \varepsilon)K}{v} - 1 \geq \frac{N - 2K}{(1 - \varepsilon)K} - 1.$$

Also $\tau_N + N/u_{j+1} \geq \deg(u_{j+1}) \geq \deg_{T_j}(u_{j+1}) \geq \deg_{T_j}(v)$. If $u_{j+1} > (1 - \frac{\varepsilon}{2})K$, then

$$\frac{N}{K} \left(\frac{1}{1 - \varepsilon} - \frac{1}{1 - \varepsilon/2} \right) < \tau_N + 1 + \frac{2}{1 - \varepsilon} \leq \tau_N + 5.$$

The left side is at least $\frac{N}{K} \cdot \frac{\varepsilon}{2}$, a contradiction. So all K vertices $u_1, \dots, u_K$ lie in $[1, (1 - \frac{\varepsilon}{2})K]$, which has fewer than K elements. This is impossible. $\square$

Uses: the two degree bounds above, with $\hat{\tau}$ in place of τ_N .

Proof of Corollary 1.6. Let $\varepsilon = \frac{1}{4}$ and $K_{\pm} := \lfloor (1 \pm \varepsilon)K \rfloor$.

Step 1: the coupling. $\mathcal{D}_{\rho}(N) - T$ is the subgraph of $\mathcal{D}_{\rho}(N)$ induced on $[N] \setminus T$. For $K' < N$, the subgraph induced on $V_{N, K'}$ uses only the edges of $G_{N, K'}$, and their orientations are independent with the same law as in $\mathcal{D}_{\rho}(N, K')$: the product measure on the edges of G_N pushes forward to the product measure on the edges of $G_{N, K'}$. Hence $\mathbb{E}[\#\Phi(\mathcal{D}_{\rho}(N)[V_{N, K'}])] = \mathbb{E}[\Phi_{K'}]$.

Lean: `expect_maxSCC_Ioc`, `expectP_comp_injective`

Step 2: the sandwich. Put $\alpha := (1 - \theta)/2$ and $\beta := (1 + 3\theta)/4$, so $\theta < \beta < 1$ and $1 - \beta = \alpha + \frac{1 - \theta}{4}$. By Lemma 2.6 with exponent α , $\tau(n) \leq C_{\alpha} N^{\alpha} =: \hat{\tau}$ for all $n \leq N$. For large N we have $\log(K + 1)/\log N < \beta$, that is, $K + 1 < N^{\beta}$, so $N/K \geq N^{1 - \beta} = N^{\alpha} N^{(1 - \theta)/4}$. For large N also $N^{(1 - \theta)/4} \geq 16C_{\alpha}$ and $N^{1 - \beta} \geq 80$. Then $\hat{\tau} + 5 \leq N^{1 - \beta}/16 + N^{1 - \beta}/16 \leq \frac{N}{K} \cdot \frac{\varepsilon}{2}$, which is the hypothesis of Lemma 6.2 when $K \geq 1$. It gives $\{1, \dots, K_{-}\} \subset T \subset \{1, \dots, K_{+}\}$, that is, $V_{N, K_{+}} \subset [N] \setminus T \subset V_{N, K_{-}}$. If $K = 0$, then $T = \emptyset$ and this is trivial.

Step 3. Since T does not depend on the orientation, Lemma 6.1 applied to the single random graph $\mathcal{D}_\rho(N)$ gives $\#\Phi(\mathcal{D}_\rho(N)[V_{N,K_+}]) \leq \#\Phi(\mathcal{D}_\rho(N) - T) \leq \#\Phi(\mathcal{D}_\rho(N)[V_{N,K_-}])$ pointwise. By Step 1, $\mathbb{E}[\Phi_{K_+}] \leq \mathbb{E}[\#\Phi(\mathcal{D}_\rho(N) - T)] \leq \mathbb{E}[\Phi_{K_-}]$.

Step 4. We have $(K+1)/4 \leq K_- + 1 \leq K+1 \leq K_+ + 1 \leq 2(K+1)$, so $|\log(K_\pm + 1) - \log(K+1)| \leq \log 4$ and $\log(K_\pm + 1)/\log N \rightarrow \theta$. Since $\theta < 1$, $K_- \leq K_+ < N$ for large N . Corollary 1.4 applies to $K_\pm$, and both bounds divided by N tend to $\rho_{\text{Dick}}(1/(1-\theta))$.

The last claim. Let $C := C_\kappa$ be the constant of Lemma 2.6 with exponent κ , and let N be so large that $x := N^\kappa \geq 2C + 4$. Let $1 \leq K \leq N^{1/2-\kappa}$ and let T be a static set. Then $K^2 x^2 \leq N^{1-2\kappa} N^{2\kappa} = N$, so $K < N$. Suppose some $t \in T$ has $t > K$. As $|T| = K$, some $v \in \{1, \dots, K\}$ is not in T . Then $\deg(v) \geq N/v - 2 \geq N/K - 2$, and $\deg(t) \leq \tau(t) + N/t \leq Cx + N/(K+1)$. The static condition $\deg(v) \leq \deg(t)$ gives $N/(K(K+1)) \leq Cx + 2$. But $K(K+1) \leq 2K^2$ and $K^2 x^2 \leq N$ give $N/(K(K+1)) \geq x^2/2$, and $x^2/2 - Cx - 2 = x(x/2 - C) - 2 \geq 2x - 2 > 0$. This contradiction shows $T \subset \{1, \dots, K\}$, and $|T| = K$ gives $T = \{1, \dots, K\}$. $\square$

Uses: Lemmas 2.6, 6.1, 6.2; Corollary 1.4.

Referee check. The attack set T must not depend on the orientation. Otherwise $\mathcal{D}_\rho(N) - T$ is not an independently oriented copy of $G_N - T$, and the coupling argument breaks down. This is why in- and out-degree attacks are excluded.

7. FURTHER REMARKS

Remark 7.1 (Edge share of the hubs). [PROVEN]

Lean: `card_hubEdges`, `card_edges_zero`, `abs_card_hubEdges_sub_le`, `abs_card_edges_sub_le`, `hub_edge_share`

An edge meets $\{1, \dots, K\}$ if and only if its smaller endpoint d is at most K . So $\{1, \dots, K\}$ meets exactly $\sum_{d \leq K} (\lfloor N/d \rfloor - 1) = N \log K + O(N)$ of the $\sum_{d \leq N} (\lfloor N/d \rfloor - 1) = N \log N + O(N)$ edges, for $1 \leq K \leq N$; both $O(N)$ terms are at most $2N$ in absolute value. Hence, if $\log(K+1)/\log N \rightarrow \theta$, the hubs meet a fraction $\theta + o(1)$ of the edges.

Remark 7.2 (Why a hub clique does not suffice for $\frac{1}{3} < \theta < \frac{1}{2}$). [PROVEN; not used]

Lean: `remark_7_2_hubs`, `card_common_multiples_ge`, `prob_not_hub`, `remark_7_2_small`, `exists_divisor_of_subsum`, `three_primes_mem`, `three_primes_goodSet`, `eventually_attach_cond`, `remark_7_2`, `card_obst_ge`

A natural first approach takes the hub set $H = (K, N^{1/2-\delta}]$ for a fixed $\delta > 0$. Two hubs $a < b$ have $\text{lcm}(a, b) \leq ab \leq N^{1-2\delta}$, hence at least $\lfloor N^{2\delta} \rfloor$ common multiples in $[1, N]$. So H is strongly connected w.h.p., uniformly in K . Indeed, for each common multiple $y \in (b, N]$ the event $\{a \rightarrow y \rightarrow b\}$ has probability q and depends only on the edges $\{a, y\}$ and $\{b, y\}$, which are disjoint for distinct y . As in Lemma 4.1, the probability that some pair of hubs is not joined in both directions is at most $2N^{1-2\delta}(1-q)^{N^{2\delta}-2} \rightarrow 0$. One then attaches every m with many divisors in H . This captures only the integers having a divisor in $(K, N^{1/2-\delta}]$.

For $\theta < 1/3$, all but $o(N)$ elements of F_1 have a divisor in $(K, \sqrt{N}]$, so the obstruction below does not occur there. To see this, fix a small $\varepsilon > 0$ and put $\theta' = \theta + \varepsilon/2$ and $\eta = \varepsilon/8$. Let $m \in F_1$ with $m \geq N^{1-\eta}$. If every prime factor of m is $< m^{1-\theta'}$, apply the sub-sum lemma (Appendix A) to the numbers $\log p / \log m$, where p runs over the prime factors of m counted with multiplicity. This gives a divisor d of m with $m^{\theta'} \leq d \leq m^{1/2}$, and $d > K$ for large N . Otherwise m has a prime factor in $(N^a, N^b]$, where $a = (1-\eta)(1-\theta-\varepsilon)$ and $b = 1-\theta+\varepsilon$, because $B \leq N^b$ for large N . By Lemma 2.1(b), at most $N^{1-\eta} + N(\log \frac{b}{a} + \frac{18}{a \log N}) \leq (10\varepsilon + o(1))N$ elements of F_1 remain.

For $1/3 < \theta < 1/2$, however, let p_1, p_2, p_3 be distinct primes in $(N^{1/4}, N^{1/3}]$, let $1 \leq s \leq N^\gamma$ with $\gamma < \theta - \frac{1}{3}$, and let $m = p_1 p_2 p_3 s \leq N$. For large N we have $N^{1/3+\gamma} \leq K < N^{1/2}$ and $B \geq N^{1/3}$. Then m lies in F_1 and has no divisor in $(K, \sqrt{N}]$, so none in $(K, N^{1/2-\delta}]$. Indeed, a divisor of $p_1 p_2 p_3 s$ is either a multiple of some $p_i p_j$ ($i \neq j$), hence $> \sqrt{N}$, or a divisor of some

$p_k s$, hence $\leq N^{1/3+\gamma} \leq K$. The number of such m is $\gg N$. To see this, take $\gamma = (\theta - \frac{1}{3})/3$ and put $\ell_j = \frac{1-\gamma}{3} + \frac{j\gamma}{9}$, so that $(N^{\ell_0}, N^{\ell_3}] = (N^{(1-\gamma)/3}, N^{1/3}] \subset (N^{1/4}, N^{1/3}]$. Take p_{j+1} prime in $(N^{\ell_j}, N^{\ell_{j+1}}]$ for $j = 0, 1, 2$. Then $p_1 < p_2 < p_3$ and $p_1 p_2 p_3 > N^{1-\gamma}$, so every $s \leq N/(p_1 p_2 p_3)$ satisfies $s < N^\gamma$, and $m = p_1 p_2 p_3 s$ determines (p_1, p_2, p_3, s) . By Lemma 2.1(b), the number of these m is, for large N ,

$$\sum_{p_1, p_2, p_3} \left\lfloor \frac{N}{p_1 p_2 p_3} \right\rfloor \geq \frac{N}{2} \prod_{j=0}^2 \sum_{N^{\ell_j} < p \leq N^{\ell_{j+1}}} \frac{1}{p} \geq \frac{N}{2} \prod_{j=0}^2 \frac{1}{2} \log \frac{\ell_{j+1}}{\ell_j}.$$

The method above avoids this obstruction. Suppose that $\delta \leq \frac{1}{8}$ and $2(K+1)N^{1/3} \leq M$, which holds for large N when $\theta + \frac{1}{3} < 1 - 2\delta$. Then such an m lies in S_M itself if $m \leq M$. In any case its three divisors $m/p_i = p_j p_k s$ are distinct and lie in S_M , since $K < N^{1/2} < p_j p_k s < N^{3/4} \leq M$ and all their prime factors are at most $N^{1/3} \leq Y$. So $s(m) \geq 3$, and m attaches as in Lemma 4.4. The connectivity of the lower layer is supplied deterministically by Lemma 4.2, not by a clique.

Remark 7.3 (Relation to [9]). [PROVEN]

Lean: `kp_corollary_2`

The lower bound of [9] uses the triangles $\{1, d, m\}$ and needs the vertex 1. The present argument uses 2-paths through common *multiples* of divisibility pairs in the lower layer. It gives a new proof of [9, Corollary 2], the case $K = 0$ of Corollary 1.4, and extends it to all K . The case $K = 0$ is also checked formally (`kp_corollary_2`, and `Checks/ProfileCheck.lean`).

APPENDIX A. THE SUB-SUM LEMMA

The preliminary notes suggested proving the case $\theta \leq 1/3$ through the Poisson–Dirichlet description of the large prime factors together with the following deterministic lemma. Our proof of Theorem 1.2 makes this route unnecessary, but we record the lemma with a proof.

Lemma A.1 (Sub-sum lemma). [PROVEN; elementary]

Lean: `subsum`, `subsum_of_mem_Icc`, `subsum_sharp`, `subsum_infinite`

Let $x_1, \dots, x_n$ be real numbers with $\sum_i x_i = 1$, and let $0 < \theta \leq 1/3$. If $x_i < 1 - \theta$ for every i , then there is a set $I \subset \{1, \dots, n\}$ with $\sum_{i \in I} x_i \in [\theta, \frac{1}{2}]$. The same holds for an infinite sequence $x_1, x_2, \dots$ of real numbers with $\sum_i x_i = 1$ absolutely convergent and every $x_i < 1 - \theta$, with a set $I \subset \{1, 2, \dots\}$. This fails for every $\theta \in (\frac{1}{3}, \frac{1}{2}]$: take $(x_1, x_2, x_3) = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$.

In the setting of the preliminary notes, $x_1 \geq x_2 \geq \dots > 0$, and the hypothesis is $x_1 < 1 - \theta$. Neither monotonicity nor positivity is needed.

Proof. We use that if $S = \sum_{i \in I} x_i \in [\theta, 1 - \theta]$, then either $S \leq \frac{1}{2}$, or $1 - S = \sum_{i \notin I} x_i \in [\theta, \frac{1}{2}]$.

If some $x_i \geq \theta$, take $I = \{i\}$; then $x_i \in [\theta, 1 - \theta]$.

Otherwise all $x_i < \theta$. The partial sums $S_k = x_1 + \dots + x_k$ start at $S_0 = 0 < \theta$ and end at $S_n = 1 \geq \theta$. Let k be minimal with $S_k \geq \theta$. Then $S_{k-1} < \theta$, so $S_k = S_{k-1} + x_k < 2\theta \leq 1 - \theta$. So $S_k \in [\theta, 1 - \theta]$, and we take $I = \{1, \dots, k\}$ or its complement.

For an infinite sequence, choose n with $x_1 + \dots + x_n > \theta$. Then the tail $r = \sum_{i > n} x_i = 1 - (x_1 + \dots + x_n)$ satisfies $r < 1 - \theta$. Apply the finite case to $(x_1, \dots, x_n, r)$, and if the resulting set contains r , replace r by all indices $i > n$.

For the counterexample with $\theta \in (\frac{1}{3}, \frac{1}{2}]$: every $x_i = \frac{1}{3} < 1 - \theta$, and the sub-sums are $0, \frac{1}{3}, \frac{2}{3}, 1$, none of which lies in $[\theta, \frac{1}{2}]$. $\square$

APPENDIX B. STATUS OF ALL CLAIMS

The formal model. The formalisation is in the directory `lean/` of the repository (Lean and Mathlib v4.34.1), and the command `lake build HubRemoval` checks all of it. For each module, a

file in the directory **Checks/** prints the axioms of its results and applies them to concrete instances, so that no hypothesis is vacuous. The formal objects match the definitions of this paper as follows.

- For integers, $\Psi(x, y)$ counts the n with $1 \leq n \leq x$ all of whose prime factors are at most y . For real arguments, $\Psi(x, y) = \Psi(\lfloor x \rfloor, \lfloor y \rfloor)$. So $\Psi(N, N/(K+1)) = \Psi(N, B)$.
- An outcome of $\mathcal{D}_\rho(N, K)$ is a map from the edges of $G_{N,K}$ to $\{\text{reversed, not reversed}\}$, weighted by the product of the Bernoulli(ρ) probabilities; expectations and probabilities are finite sums. $\mathcal{D}_\rho(N)$ is the case $K = 0$. The law of $\mathcal{D}_\rho(N, K)$ is that of $\mathcal{D}_\rho(N)[V_{N,K}]$ (Step 1 of the proof of Corollary 1.6).
- $\#\Phi(D[V])$ is the largest size of a set $S \subset V$ any two of whose elements are joined by directed paths inside $D[V]$. SCCs are the classes of mutual reachability.
- Limits are Mathlib's **Tendsto** along **atTop**, and “for $N \geq N_0$ ” is $\exists N_0 \forall N \geq N_0$.

Status.

Claim	Status	Lean names / notes
Lemma 2.1 (Mertens, Chebyshev)	PROVEN	Lean: lemma_2_1 $c_0 = \log 4 + 2$, $C_1 = 18$; proved from scratch (Appendix C)
Lemma 2.2 (Dickman, uniform on $[1, U]$)	PROVEN	Lean: dickman_unique, dickman_asymptotic Proved from scratch (Appendix C)
Lemma 2.3 ($\Psi \leq \sqrt{x} + 2x(\log y + c_0)/\log x$)	PROVEN	Lean: psi_upper', dickman_le_two_div Integers $x \geq 2$, $y \geq 1$
Lemma 2.4 (variance of ω_z)	PROVEN	Lean: turan_variance, turan_count
Remark 2.5 (Turán–Kubilius)	PROVEN	Lean: remark_2_5, turan_kubilius $C = 130$; not needed for the main results
Lemma 2.6 ($\tau(n) \leq C_\varepsilon n^\varepsilon$)	PROVEN	Lean: divisor_bound_explicit Explicit C_ε
Lemma 2.7 (limit of $\Psi(N, N/(K+1))/N$)	PROVEN	Lean: psi_limit
Lemma 3.1 (fibres)	PROVEN	Lean: roughPart_eq_of_reachable, card_fibre_le, card_fibre_le_K Essentially known (cf. [10])
Proposition 3.2 ($\Phi_K \leq \Psi(N, B)$, $\Phi_K \leq \max(F_1 , K)$)	PROVEN	Lean: PhiK_le_psi, PhiK_le_max Deterministic
Corollary 3.3 (limsup)	PROVEN	Lean: cor_upper_profile Any $\rho \in [0, 1]$
Remark 3.4 (equivalent forms)	PROVEN	Lean: remark_3_4, remark_3_4_holds Markov; the equivalence holds for any $\rho \in [0, 1]$
Lemma 4.1 (robust edges)	PROVEN	Lean: prob_not_robust_exp, robust_reachable Union bound, no cross-pair independence
Lemma 4.2 (doubling)	PROVEN	Lean: doubling_connected Deterministic
Corollary 4.3	PROVEN	Lean: sigmaSet_mutual, sigmaSet_closed, sigmaSet_subset

Claim	Status	Lean names / notes
Lemma 4.4 (attachment)	PROVEN	Lean: prob_attachEv_eq, prob_attachEv_ge, attach_mutual
Lemma 4.5 (arithmetic core)	PROVEN	Lean: core_a, core_b, core_c, core_d'
Proposition 4.6, (2) and (3)	PROVEN	Deterministic, explicit exceptional set Lean: lower_bound_err', fixedN_sandwich', eq_4_2 Non-asymptotic, $C_1 = 18$
Theorem 1.3 (rate $30N/\log\log N$)	PROVEN	Lean: thm_rate, rate_cases, eventually_goodN
Theorem 1.2 (uniform asymptotic)	PROVEN	Lean: thm_main Answers [9, Q. 20]
Remark 5.1 (explicit loss for $K < \sqrt{N}$)	PROVEN	Lean: remark_5_1, smallK_psi Ψ -estimate with error $77/\log N$
Corollary 1.4 (profile $\rho_{\text{Dick}}(1/(1-\theta))$)	PROVEN	Lean: cor_profile, dickLim_continuous, dickLim_strictAntiOn
Corollary 1.5 (giant $\approx F_1$, second $\text{SCC} \leq K + o(N)$)	PROVEN	Lean: cor_structure, cor_structure_prob, cor_structure_whp
Remark 5.2 (term K needed)	PROVEN	Lean: remark_5_2, secondSCC_ge, prob_PhiK_le_K Uses [9, Cor. 2] (Remark 7.3)
Remark 5.3 (uniform in ρ ; $\rho_N \gg 1/\log\log N$)	PROVEN	Lean: remark_5_3_uniform, remark_5_3_slow, slow_rho_bound_explicit (R1)–(R6) are monotone in s_0 and q
Lemma 6.1, Lemma 6.2	PROVEN	Lean: maxSCC_mono, sandwich_static, sandwich_adaptive
Corollary 1.6 (static/adaptive degree attacks)	PROVEN	Lean: cor_degree_static, cor_degree_adaptive, static_eq_Icc Degrees taken in G_N
Remark 7.1 (edge share θ)	PROVEN	Lean: card_hubEdges, hub_edge_share $ O(N) \leq 2N$
Remark 7.2 (hub clique loses for $\frac{1}{3} < \theta < \frac{1}{2}$)	PROVEN; not used	Lean: remark_7_2_hubs, remark_7_2_small, three_primes_mem, three_primes_goodSet, eventually_attach_cond, remark_7_2 Triple sum written out; hubs strongly connected w.h.p.; the case $\theta < 1/3$
Remark 7.3 (Kim–Phillips' Corollary 2)	PROVEN	Lean: kp_corollary_2 The case $K = 0$ of Corollary 1.4
Lemma A.1 (sub-sum lemma, sharp)	PROVEN	Lean: subsum, subsum_infinite, subsum_sharp Finite and infinite sequences; not needed for the main results
Lemmas C.1 and C.2 (Buchstab, prime sums)	PROVEN	Lean: buchstab_real, primeSum_integral

APPENDIX C. PROOFS OF LEMMAS 2.1 AND 2.2

The proofs follow the formal proofs in the Lean files `Mertens`, `Mertens2`, `Dickman`, `Buchstab`, `PrimeSum` and `DickmanThm` of the directory `lean/HubRemoval/`. Throughout, $n \geq 1$ is an integer, $\vartheta(n) := \sum_{p \leq n} \log p$ and $S(n) := \sum_{p \leq n} \frac{\log p}{p}$.

Proof of Lemma 2.1. (i) *Chebyshev.* $\vartheta(n) \leq n \log 4$, because $\prod_{p \leq n} p \leq 4^n$ (the classical bound through the central binomial coefficient; it is in Mathlib).

(ii) $S(n) \leq \log n + \log 4$. By Legendre's formula, $v_p(n!) \geq \lfloor n/p \rfloor \geq n/p - 1$ for $p \leq n$. Every prime factor of $n!$ is at most n , and $n! \leq n^n$. Hence

$$\sum_{p \leq n} \left(\frac{n}{p} - 1 \right) \log p \leq \sum_{p \leq n} v_p(n!) \log p = \log n! \leq n \log n,$$

so $nS(n) \leq n \log n + \vartheta(n) \leq n \log n + n \log 4$.

(iii) $\sum_{p \leq n} \frac{\log p}{p(p-1)} \leq 2$. Put $h(k) := (\log k + 2)/k$. For $k \geq 2$,

$$h(k-1) - h(k) = \frac{\log k - k \log \frac{k}{k-1} + 2}{k(k-1)} \geq \frac{\log k}{k(k-1)},$$

because $k \log \frac{k}{k-1} \leq \frac{k}{k-1} \leq 2$. Summing over $2 \leq k \leq n$ gives $\sum_{p \leq n} \frac{\log p}{p(p-1)} \leq h(1) = 2$.

(iv) *Part (a).* Since $\frac{\log p}{p-1} = \frac{\log p}{p} + \frac{\log p}{p(p-1)}$, (ii) and (iii) give $\sum_{p \leq n} \frac{\log p}{p-1} \leq \log n + \log 4 + 2 = \log n + c_0$. For real $y \geq 2$ apply this to $n = \lfloor y \rfloor$.

(v) *Mertens' first theorem:* $|S(n) - \log n| \leq 3$. The upper bound is (ii), as $\log 4 < 3$. For the lower bound, Legendre's formula gives $(p-1)v_p(n!) \leq n$. So $\log n! \leq n \sum_{p \leq n} \frac{\log p}{p-1} \leq n(S(n) + 2)$ by (iii). Also $\log n! \geq n \log n - n$, because $n^n/n! \leq e^n$. Hence $S(n) \geq \log n - 3$.

(vi) *Part (b) for integers $2 \leq y \leq w$.* Since $S(k) - S(k-1) = \frac{\log k}{k}$ if k is prime and 0 otherwise,

$$\sum_{y < p \leq w} \frac{1}{p} = \sum_{y < k \leq w} \frac{S(k) - S(k-1)}{\log k}.$$

Write $S(k) = \log k + E(k)$, so $|E(k)| \leq 3$ by (v). Put $f(k) := 1/\log k$, which is positive and non-increasing for $k \geq 2$. By Abel summation,

$$\sum_{y < k \leq w} (E(k) - E(k-1))f(k) = E(w)f(w) - E(y)f(y) + \sum_{y < k \leq w} E(k-1)(f(k-1) - f(k)),$$

whose absolute value is at most $3f(w) + 3f(y) + 3(f(y) - f(w)) = 6/\log y$. The remaining part is $\sum_{y < k \leq w} (v_k - u_k)/v_k$ with $u_k := \log(k-1)$ and $v_k := \log k$. Since $\frac{v-u}{v} \leq \log \frac{v}{u} \leq \frac{v-u}{u}$, $v_k - u_k = \log \frac{k}{k-1} \leq \frac{1}{k-1}$ and $u_k, v_k \geq \log y$,

$$0 \leq \log \frac{v_k}{u_k} - \frac{v_k - u_k}{v_k} \leq \frac{(v_k - u_k)^2}{u_k v_k} \leq \frac{1}{(k-1)^2 \log^2 y}.$$

As $\sum_{y < k \leq w} \log \frac{v_k}{u_k} = \log \log w - \log \log y$ and $\sum_{j \geq 2} j^{-2} \leq 1$, we get

$$\left| \sum_{y < p \leq w} \frac{1}{p} - \log \frac{\log w}{\log y} \right| \leq \frac{6}{\log y} + \frac{1}{\log^2 y} \leq \frac{8}{\log y},$$

using $\log y \geq \log 2 > \frac{1}{2}$.

(vii) *Part (b) for reals.* For real $x \geq 2$ and $n = \lfloor x \rfloor$ we have $\log x - \log n \leq \log \frac{n+1}{n} \leq \frac{1}{n}$, so

$$0 \leq \log \log x - \log \log n \leq \frac{\log x - \log n}{\log n} \leq \frac{1}{n \log n} \leq \frac{1}{2 \log n},$$

and $\frac{1}{\log n} \leq \frac{2}{\log x}$, since $x < n + 1 \leq n^2$. Let $Y = \lfloor y \rfloor$ and $W = \lfloor w \rfloor$. The primes in $(y, w]$ are those in $(Y, W]$. By (vi), the error is at most $\frac{8}{\log Y} + \frac{1}{2\log W} + \frac{1}{2\log Y} \leq \frac{9}{\log Y} \leq \frac{18}{\log y}$.

(viii) *Part (c)*. Let $Z = \lfloor z \rfloor \geq 2$. By (vi) with $y = 2$ and $w = Z$, $\sum_{p \leq Z} \frac{1}{p} \geq \sum_{2 < p \leq Z} \frac{1}{p} \geq \log \log Z - \log \log 2 - \frac{8}{\log 2} \geq \log \log Z - 12$, as $\log \log 2 < 0$ and $8/\log 2 < 12$. By (vii), $\log \log z \leq \log \log Z + \frac{1}{2\log Z} \leq \log \log Z + 1$. Hence $\sum_{p \leq z} \frac{1}{p} \geq \log \log z - 13$.

(ix) *Part (d)*. At most $\sqrt{x}$ primes are $\leq \sqrt{x}$. Each prime $p \in (\sqrt{x}, x]$ has $\log p > \frac{1}{2} \log x$, so there are at most $\vartheta(x)/(\frac{1}{2} \log x) \leq 2x \log 4 / \log x$ of them, by (i). Also $\sqrt{x} \log x = 2\sqrt{x} \log \sqrt{x} \leq 2\sqrt{x}(\sqrt{x} - 1) \leq 2x$, so $\sqrt{x} \leq 2x / \log x$. Hence $\pi(x) \leq (2 + 2 \log 4)x / \log x \leq 5x / \log x$, as $\log 4 < 1.39$. $\square$

Proof of Lemma 2.2: the function. (i) *Construction*. Let $\rho_0 \equiv 1$ on $\mathbb{R}$, and for $n \geq 0$ let

$$\rho_{n+1}(u) := \begin{cases} \rho_n(u), & u \leq n+1, \\ \rho_n(n+1) - \int_{n+1}^u \rho_n(w-1) \frac{dw}{w}, & u \geq n+1. \end{cases}$$

By induction each ρ_n is continuous, and $\rho_m = \rho_n$ on $(-\infty, m+1]$ for $m \leq n$. Put $\rho_{\text{Dick}}(u) := \rho_n(u)$ for any n with $u \leq n+1$. Then ρ_{Dick} is continuous, $\rho_{\text{Dick}} = 1$ on $(-\infty, 1]$, and for $n+1 \leq u \leq n+2$ we have $\rho_{\text{Dick}}(u) = \rho_{\text{Dick}}(n+1) - \int_{n+1}^u \rho_{\text{Dick}}(w-1) \frac{dw}{w}$, since $w-1 \leq n+1$ there.

(ii) *The integral equation*. Adding the relations of (i) over $[1, 2], [2, 3], \dots$ gives $\rho_{\text{Dick}}(u) = 1 - \int_1^u \rho_{\text{Dick}}(w-1) \frac{dw}{w}$ for $u \geq 1$, and $\rho_{\text{Dick}}(u) = \rho_{\text{Dick}}(v) - \int_v^u \rho_{\text{Dick}}(w-1) \frac{dw}{w}$ for $1 \leq v \leq u$. By the fundamental theorem of calculus, ρ_{Dick} is differentiable at every $u > 1$ with $\rho'_{\text{Dick}}(u) = -\rho_{\text{Dick}}(u-1)/u$. On $[1, 2]$ the integrand is $1/w$, so $\rho_{\text{Dick}}(u) = 1 - \log u$.

(iii) *The integrated form*. $G(v) := v\rho_{\text{Dick}}(v) - \int_{v-1}^v \rho_{\text{Dick}}$ is continuous, and for $v > 1$, $G'(v) = \rho_{\text{Dick}}(v) + v\rho'_{\text{Dick}}(v) - (\rho_{\text{Dick}}(v) - \rho_{\text{Dick}}(v-1)) = 0$. By the mean value theorem, $G(u) = G(1) = 1 - \int_0^1 1 = 0$ for $u \geq 1$.

(iv) *Positivity*. Suppose $\rho_{\text{Dick}}(u) \leq 0$ for some u . The set $\{v : \rho_{\text{Dick}}(v) \leq 0\}$ is closed, non-empty and bounded below by 1; let u_0 be its least element. Then $u_0 > 1$ and $\rho_{\text{Dick}} > 0$ on $(-\infty, u_0)$. By (iii), $u_0\rho_{\text{Dick}}(u_0) = \int_{u_0-1}^{u_0} \rho_{\text{Dick}} > 0$, which contradicts $\rho_{\text{Dick}}(u_0) \leq 0 < u_0$.

(v) *Monotonicity*. For $u > 1$, $\rho'_{\text{Dick}}(u) = -\rho_{\text{Dick}}(u-1)/u < 0$, so ρ_{Dick} is strictly decreasing on $[1, \infty)$. Hence ρ_{Dick} is non-increasing on $\mathbb{R}$ and $\rho_{\text{Dick}} \leq \rho_{\text{Dick}}(1) = 1$.

(vi) *Uniqueness*. Let g be continuous on $[0, \infty)$, with $g = 1$ on $[0, 1]$ and $g'(u) = -g(u-1)/u$ for $u > 1$. We show $g = \rho_{\text{Dick}}$ on $[0, n+1]$ by induction on n ; the case $n = 0$ is given. If $g = \rho_{\text{Dick}}$ on $[0, n+1]$, then on $(n+1, n+2)$ the derivative of $g - \rho_{\text{Dick}}$ is $-(g(u-1) - \rho_{\text{Dick}}(u-1))/u = 0$. By the mean value theorem and continuity, $g - \rho_{\text{Dick}}$ is constant on $[n+1, n+2]$, and it vanishes at $n+1$. $\square$

Buchstab's identity and a prime-sum estimate.

Lemma C.1 (Buchstab's identity). [PROVEN]

Lean: `buchstab, buchstab_real, psi_prime_step, psi_nonprime_step, psi_eq_self_of_le`

For real x and $0 \leq y \leq z$,

$$\Psi(x, z) = \Psi(x, y) + \sum_{y < p \leq z} \Psi(x/p, p).$$

Moreover $\Psi(X, Y) = X$ for integers $0 \leq X \leq Y$.

Proof. First let $X = \lfloor x \rfloor$ and let $Y \leq Z$ be integers; we induct on $Z \geq Y$. If $Z+1$ is not prime, the $(Z+1)$ -smooth numbers are the Z -smooth ones. If $p = Z+1$ is prime, the $n \leq X$ that are p -smooth but not $(p-1)$ -smooth are exactly the $n = pm$ with $m \leq \lfloor X/p \rfloor$ and m p -smooth, and $m \mapsto pm$ is injective. So $\Psi(X, p) = \Psi(X, p-1) + \Psi(\lfloor X/p \rfloor, p)$. For real arguments use $\lfloor x/p \rfloor = \lfloor \lfloor x \rfloor / p \rfloor$ and $y < p \iff \lfloor y \rfloor < p$. Finally, if $X \leq Y$, every $n \in [1, X]$ has all prime factors $\leq n \leq Y$. $\square$

Lemma C.2 (Prime sums as Riemann sums). [PROVEN]

Lean: `primeSum_integral, step_bound, sum_primesBetween_inv, log_div_log_mem`

Let $0 < a \leq b$, and let g be continuous and non-increasing on $[a, b]$ with $0 \leq g \leq G$ there. For $x > 1$ and a prime p put $w_p := \log x / \log p$. Then for every $\varepsilon > 0$ there is x_0 such that for all $x \geq x_0$ and all $u \in [a, b]$,

$$\left| \sum_{x^{1/u} < p \leq x^{1/a}} \frac{g(w_p)}{p} - \int_a^u g(w) \frac{dw}{w} \right| \leq \varepsilon.$$

Proof. For $a \leq s \leq t \leq b$ let $P(s, t)$ be the set of primes p with $x^{1/t} < p \leq x^{1/s}$; for these, $s \leq w_p < t$. Assume $x^{1/b} \geq 2$. Lemma 2.1(b) with $y = x^{1/t}$ and $w = x^{1/s}$, where $\log(\log w / \log y) = \log(t/s)$ and $C_1 / \log y = 18t / \log x$, gives $|A - \log(t/s)| \leq \eta := 18b / \log x$ for $A := \sum_{p \in P(s, t)} \frac{1}{p}$. As g is non-increasing, $\sum_{p \in P(s, t)} g(w_p)/p$ lies in $[g(t)A, g(s)A]$, and $\int_s^t g(w) \frac{dw}{w}$ lies in $[g(t)\ell, g(s)\ell]$ with $\ell := \log(t/s) \in [0, (t-s)/a]$. Hence the two differ by at most

$$(g(s) - g(t)) \frac{t-s}{a} + G\eta.$$

Fix an integer $m \geq 1$ and put $t_i := a + i(u-a)/m$ for $0 \leq i \leq m$. The sets $P(t_i, t_{i+1})$ partition $P(a, u)$, and the integrals over $[t_i, t_{i+1}]$ add up to the integral over $[a, u]$. Summing the step bounds, the first terms telescope, and $t_{i+1} - t_i \leq (b-a)/m$, so the difference in the lemma is at most

$$(g(a) - g(u)) \frac{b-a}{ma} + mG\eta \leq \frac{G(b-a)}{ma} + \frac{18bmG}{\log x}.$$

Choose $m > 2G(b-a)/(a\varepsilon)$, so the first term is $\leq \varepsilon/2$. The second term is $\leq \varepsilon/2$ once $\log x \geq 36bmG/\varepsilon$, and $x^{1/b} \geq 2$ for large x . None of these conditions depends on u . $\square$

Proof of Lemma 2.2: the asymptotic. Write $\varphi(x, u) := \Psi(x, x^{1/u})/x$. We show by induction on $n \geq 2$: for every $\varepsilon > 0$, for all large x , $|\varphi(x, u) - \rho_{\text{Dick}}(u)| \leq \varepsilon$ for all $u \in [1, n]$. For $U \geq 1$ this gives the lemma with $n = \lceil U \rceil + 2$.

Lean: `psi_dickman_base, psi_dickman_step, psi_dickman_nat`

Base case $n = 2$. Let $x \geq 4$ and $u \in [1, 2]$, so $x^{1/u} \geq \sqrt{x} \geq 2$. Lemma C.1 with $y = x^{1/u}$ and $z = x$ gives $\Psi(x, x^{1/u}) = [x] - \sum_{x^{1/u} < p \leq x} \Psi(x/p, p)$. For these p we have $p > \sqrt{x}$, so $x/p < p$ and $\Psi(x/p, p) = [x/p] \in (x/p - 1, x/p]$. With $A := \sum_{x^{1/u} < p \leq x} \frac{1}{p}$ and Lemma 2.1(b),(d),

$$|\Psi(x, x^{1/u}) - x(1 - \log u)| \leq 1 + \pi(x) + x|A - \log u| \leq 1 + \frac{5x}{\log x} + \frac{36x}{\log x},$$

since $C_1 / \log x^{1/u} = 18u / \log x \leq 36 / \log x$. As $\rho_{\text{Dick}}(u) = 1 - \log u$ on $[1, 2]$, we get $|\varphi(x, u) - \rho_{\text{Dick}}(u)| \leq 1/x + 41 / \log x \leq \varepsilon$ for $x \geq \max(4, 2/\varepsilon)$ and $\log x \geq 82/\varepsilon$.

Step from n to $n+1$, $n \geq 2$. Let $\varepsilon > 0$. By the induction hypothesis there is T with $|\varphi(t, v) - \rho_{\text{Dick}}(v)| \leq \varepsilon/4$ for all $t \geq T$ and $v \in [1, n]$. Apply Lemma C.2 with $a = n$, $b = n+1$, $g(w) = \rho_{\text{Dick}}(w-1)$ (continuous, non-increasing, $0 < g \leq 1 = G$) and $\varepsilon/4$. Let x be so large that this applies, $\sqrt{x} \geq \max(T, 1)$, $x^{1/(n+1)} \geq 2$ and $\log x \geq 18(n+1)$. For $u \in [1, n]$ the induction hypothesis applies, so let $u \in [n, n+1]$. Lemma C.1 with $y = x^{1/u}$ and $z = x^{1/n}$ gives

$$\varphi(x, u) = \varphi(x, n) - \sum_{p \in P} \frac{\Psi(x/p, p)}{x}, \quad P := \{p : x^{1/u} < p \leq x^{1/n}\}.$$

For $p \in P$ we have $w_p = \log x / \log p \in [n, u)$, so $v_p := w_p - 1 \in [n-1, n) \subset [1, n]$, and $(x/p)^{1/v_p} = \exp((\log x - \log p)/v_p) = p$. Also $p \leq x^{1/n} \leq \sqrt{x}$, so $x/p \geq \sqrt{x} \geq T$. The induction hypothesis at

$t = x/p$ and $v = v_p$ gives $|\Psi(x/p, p)/(x/p) - \rho_{\text{Dick}}(w_p - 1)| \leq \varepsilon/4$, that is, $|\Psi(x/p, p)/x - \rho_{\text{Dick}}(w_p - 1)/p| \leq \varepsilon/(4p)$. By Lemma 2.1(b), $\sum_{p \in P} \frac{1}{p} \leq \log \frac{u}{n} + \frac{18u}{\log x} \leq 1 + 1$. Hence

$$\left| \sum_{p \in P} \frac{\Psi(x/p, p)}{x} - \sum_{p \in P} \frac{\rho_{\text{Dick}}(w_p - 1)}{p} \right| \leq \frac{\varepsilon}{2}.$$

By Lemma C.2, $|\sum_{p \in P} \rho_{\text{Dick}}(w_p - 1)/p - \int_n^u \rho_{\text{Dick}}(w - 1) \frac{dw}{w}| \leq \varepsilon/4$. By (ii) of the construction above, $\rho_{\text{Dick}}(u) = \rho_{\text{Dick}}(n) - \int_n^u \rho_{\text{Dick}}(w - 1) \frac{dw}{w}$. Finally $|\varphi(x, n) - \rho_{\text{Dick}}(n)| \leq \varepsilon/4$ by the induction hypothesis at $t = x \geq T$ and $v = n$. Together, $|\varphi(x, u) - \rho_{\text{Dick}}(u)| \leq \frac{\varepsilon}{4} + \frac{\varepsilon}{2} + \frac{\varepsilon}{4} = \varepsilon$. $\square$

REFERENCES

- [1] R. Albert, H. Jeong, A.-L. Barabási, Error and attack tolerance of complex networks, *Nature* **406** (2000), 378–382.
- [2] B. Bollobás, O. Riordan, Robustness and vulnerability of scale-free random graphs, *Internet Math.* **1** (2003), 1–35.
- [3] N. G. de Bruijn, On the number of positive integers $\leq x$ and free of prime factors $> y$, *Indag. Math.* **13** (1951), 50–60.
- [4] K. Dickman, On the frequency of numbers containing prime factors of a certain relative magnitude, *Ark. Mat. Astr. Fys.* **22A** (1930), no. 10, 1–14.
- [5] A. Granville, Smooth numbers: computational number theory and beyond, in: *Algorithmic Number Theory*, MSRI Publ. 44, Cambridge Univ. Press, 2008, 267–323.
- [6] G. H. Hardy, E. M. Wright, *An Introduction to the Theory of Numbers*, 6th ed., Oxford Univ. Press, 2008.
- [7] A. Hildebrand, On the number of positive integers $\leq x$ and free of prime factors $> y$, *J. Number Theory* **22** (1986), 289–307.
- [8] A. Hildebrand, G. Tenenbaum, Integers without large prime factors, *J. Théor. Nombres Bordeaux* **5** (1993), 411–484.
- [9] J. Kim, T. Phillips, On the largest strongly connected component of randomly oriented divisor graphs, arXiv:2604.05176v2 (2026).
- [10] N. McNew, Counting primitive subsets and other statistics of the divisor graph of $\{1, 2, \dots, n\}$, *European J. Combin.* **92** (2021), 103237.
- [11] F. Mertens, Ein Beitrag zur analytischen Zahlentheorie, *J. Reine Angew. Math.* **78** (1874), 46–62.
- [12] G. Tenenbaum, *Introduction to Analytic and Probabilistic Number Theory*, 3rd ed., Grad. Stud. Math. 163, Amer. Math. Soc., 2015.
- [13] P. Turán, On a theorem of Hardy and Ramanujan, *J. London Math. Soc.* **9** (1934), 274–276.