



# LaplacesDemon Examples

Statisticat, LLC

---

## Abstract

The **LaplacesDemon** package is a complete environment for Bayesian inference within R. Virtually any probability model may be specified. This vignette is a compendium of examples of how to specify different model forms.

*Keywords:* Bayesian, Bayesian Inference, Laplace's Demon, LaplacesDemon, R, Statisticat.

---

**LaplacesDemon** (Statisticat LLC. 2014), usually referred to as Laplace's Demon, is an R package that is available at <http://www.bayesian-inference.com/software>. A formal introduction to Laplace's Demon is provided in an accompanying vignette entitled "**LaplacesDemon** Tutorial", and an introduction to Bayesian inference is provided in the "Bayesian Inference" vignette.

The purpose of this document is to provide users of the **LaplacesDemon** package with examples of a variety of Bayesian methods. It is also a testament to the diverse applicability of **LaplacesDemon** to Bayesian inference.

To conserve space, the examples are not worked out in detail, and only the minimum of necessary materials is provided for using the various methodologies. Necessary materials include the form expressed in notation, data (which is often simulated), the **Model** function, and initial values. The provided data, model specification, and initial values may be copy/pasted into an R file and updated with the **LaplacesDemon** or (usually) **LaplaceApproximation** functions. Although many of these examples update quickly, some examples are computationally intensive.

Initial values are usually hard-coded in the examples, though the Parameter-Generating Function (PGF) is also specified. It is recommended to generate initial values with the **GIV** function according to the user-specified PGF.

Notation in this vignette follows these standards: Greek letters represent parameters, lower case letters represent indices, lower case bold face letters represent scalars or vectors, probability distributions are represented with calligraphic font, upper case letters represent index limits, and upper case bold face letters represent matrices. More information on notation is available at <http://www.bayesian-inference.com/notation>.

This vignette will grow over time as examples of more methods become included. Contributed examples are welcome. Please send contributed examples or discovered errors in a similar for-

mat in an email to [software@bayesian-inference.com](mailto:software@bayesian-inference.com) for review and testing. All accepted contributions are, of course, credited.

## Contents

- ANCOVA [1](#)
- ANOVA, One-Way [2](#)
- ANOVA, Two-Way [3](#)
- Approximate Bayesian Computation (ABC) [4](#)
- ARCH-M(1,1) [5](#)
- Autoregression, AR(1) [6](#)
- Autoregressive Conditional Heteroskedasticity, ARCH(1,1) [7](#)
- Autoregressive Moving Average, ARMA(1,1) [8](#)
- Beta Regression [9](#)
- Beta-Binomial [10](#)
- Binary Logit [11](#)
- Binary Log-Log Link Mixture [12](#)
- Binary Probit [13](#)
- Binary Robit [14](#)
- Binomial Logit [15](#)
- Binomial Probit [16](#)
- Binomial Robit [17](#)
- Change Point Regression [18](#)
- Cluster Analysis, Confirmatory (CCA) [19](#)
- Cluster Analysis, Exploratory (ECA) [20](#)
- Collaborative Filtering (CF) [34](#)
- Conditional Autoregression (CAR), Poisson [21](#)
- Conditional Predictive Ordinate (CPO) [22](#)
- Contingency Table [23](#)
- Covariance Separation Strategy [24](#)

- Discrete Choice, Conditional Logit [25](#)
- Discrete Choice, Mixed Logit [26](#)
- Discrete Choice, Multinomial Probit [27](#)
- Distributed Lag, Koyck [28](#)
- Dynamic Linear Model (DLM) [32](#) [88](#) [89](#) [90](#)
- Exponential Smoothing [29](#)
- Factor Analysis, Approximate Dynamic (ADFA) [30](#)
- Factor Analysis, Confirmatory (CFA) [31](#)
- Factor Analysis, Dynamic (DFA) [32](#)
- Factor Analysis, Exploratory (EFA) [33](#)
- Factor Analysis, Exploratory Ordinal (EOFA) [34](#)
- Factor Regression [35](#)
- Gamma Regression [36](#)
- GARCH(1,1) [37](#)
- GARCH-M(1,1) [38](#)
- Geographically Weighted Regression [39](#)
- Hidden Markov Model [40](#)
- Hierarchical Bayes [48](#)
- Horseshoe Regression [96](#)
- Inverse Gaussian Regression [41](#)
- Kriging [42](#)
- Kriging, Predictive Process [43](#)
- Laplace Regression [44](#)
- LASSO [95](#)
- Latent Dirichlet Allocation (LDA) [45](#)
- Linear Regression [46](#)
- Linear Regression, Frequentist [47](#)
- Linear Regression, Hierarchical Bayesian [48](#)
- Linear Regression, Multilevel [49](#)

- Linear Regression with Full Missingness 50
- Linear Regression with Missing Response 51
- Linear Regression with Missing Response via ABB 52
- Linear Regression with Power Priors 53
- Linear Regression with Zellner's g-Prior 54
- LSTAR 55
- MANCOVA 56
- MANOVA 57
- Missing Values 50 51 52
- Mixture Model, Dirichlet Process 20
- Mixture Model, Finite 19 58
- Mixture Model, Infinite 20 59
- Mixture Model, Poisson-Gamma 60
- Model Averaging 98 97
- Multilevel Model 49
- Multinomial Logit 61
- Multinomial Logit, Nested 62
- Multinomial Probit 63
- Multiple Discrete-Continuous Choice 64
- Multivariate Binary Probit 65
- Multivariate Laplace Regression 66
- Multivariate Regression 67
- Negative Binomial Regression 68
- Normal, Multilevel 69
- Ordinal Logit 70
- Ordinal Probit 71
- Panel, Autoregressive Poisson 72
- Penalized Spline Regression 73
- Poisson Regression 74

- Poisson Regression, Overdispersed [60](#) [68](#)
- Poisson-Gamma Regression [60](#)
- Polynomial Regression [75](#)
- Power Priors [53](#)
- Proportional Hazards Regression, Weibull [76](#)
- Quantile Regression [77](#)
- Revision, Normal [78](#)
- Ridge Regression [79](#)
- Robust Regression [80](#)
- Seemingly Unrelated Regression (SUR) [81](#)
- Simultaneous Equations [82](#)
- Space-Time, Dynamic [83](#)
- Space-Time, Nonseparable [84](#)
- Space-Time, Separable [85](#)
- Spatial Autoregression (SAR) [86](#)
- STARMA(1,1) [87](#)
- State Space Model (SSM), Dynamic Factor Analysis (DFA) [32](#)
- State Space Model (SSM), Linear Regression [88](#)
- State Space Model (SSM), Local Level [89](#)
- State Space Model (SSM), Local Linear Trend [90](#)
- State Space Model (SSM), Stochastic Volatility (SV) [91](#)
- Stochastic Volatility (SV) [91](#)
- Survival Model [76](#)
- T-test [2](#)
- TARCH(1) [92](#)
- Threshold Autoregression (TAR) [93](#)
- Topic Model [45](#)
- Time Varying AR(1) with Chebyshev Series [94](#)
- Variable Selection, BAL [95](#)

- Variable Selection, Horseshoe [96](#)
- Variable Selection, RJ [97](#)
- Variable Selection, SSVS [98](#)
- Variety Model [64](#)
- Vector Autoregression, VAR(1) [99](#)
- Weighted Regression [100](#)
- Zellner's g-Prior [54](#)
- Zero-Inflated Poisson (ZIP) [101](#)

## 1. ANCOVA

This example is essentially the same as the two-way ANOVA (see section [3](#)), except that a covariate  $\mathbf{X}_{,3}$  has been added, and its parameter is  $\delta$ .

### 1.1. Form

$$\begin{aligned}
 \mathbf{y}_i &\sim \mathcal{N}(\mu_i, \sigma_1^2) \\
 \mu_i &= \alpha + \beta[\mathbf{X}_{i,1}] + \gamma[\mathbf{X}_{i,2}] + \delta\mathbf{X}_{i,2}, \quad i = 1, \dots, N \\
 \epsilon_i &= \mathbf{y}_i - \mu_i \\
 \alpha &\sim \mathcal{N}(0, 1000) \\
 \beta_j &\sim \mathcal{N}(0, \sigma_2^2), \quad j = 1, \dots, J \\
 \beta_J &= -\sum_{j=1}^{J-1} \beta_j \\
 \gamma_k &\sim \mathcal{N}(0, \sigma_3^2), \quad k = 1, \dots, K \\
 \gamma_K &= -\sum_{k=1}^{K-1} \gamma_k \\
 \delta &\sim \mathcal{N}(0, 1000) \\
 \sigma_m &\sim \mathcal{HC}(25), \quad m = 1, \dots, 3
 \end{aligned}$$

### 1.2. Data

```

N <- 100
J <- 5 #Number of levels in factor (treatment) 1
K <- 3 #Number of levels in factor (treatment) 2
X <- cbind(rcat(N,rep(1/J,J)), rcat(N,rep(1/K,K)), runif(N,-2,2))

```

```

alpha <- runif(1,-1,1)
beta <- runif(J-1,-2,2)
beta <- c(beta, -sum(beta))
gamma <- runif(K-1,-2,2)
gamma <- c(gamma, -sum(gamma))
delta <- runif(1,-2,2)
y <- alpha + beta[X[,1]] + gamma[X[,2]] + delta*X[,3] + rnorm(N,0,0.1)
mon.names <- c("LP","beta[5]","gamma[3]","s.beta","s.gamma","s.epsilon")
parm.names <- as.parm.names(list(alpha=0, beta=rep(0,J-1), gamma=rep(0,K-1),
  delta=0, sigma=rep(0,3)))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
pos.gamma <- grep("gamma", parm.names)
pos.delta <- grep("delta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnorm(1,0,1), rnorm(Data$J-1,0,1),
  rnorm(Data$K-1,0,1), rnorm(1,0,1), rhalfcauchy(3,5)))
MyData <- list(J=J, K=K, N=N, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.alpha=pos.alpha,
  pos.beta=pos.beta, pos.gamma=pos.gamma, pos.delta=pos.delta,
  pos.sigma=pos.sigma, y=y)

```

### 1.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[Data$pos.alpha]
  beta <- parm[Data$pos.beta]
  beta <- c(beta, -sum(beta)) #Sum-to-zero constraint
  gamma <- parm[Data$pos.gamma]
  gamma <- c(gamma, -sum(gamma)) #Sum-to-zero constraint
  delta <- parm[Data$pos.delta]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  alpha.prior <- dnorm(alpha, 0, 1000, log=TRUE)
  beta.prior <- sum(dnorm(beta, 0, sigma[2], log=TRUE))
  gamma.prior <- sum(dnorm(gamma, 0, sigma[3], log=TRUE))
  delta.prior <- dnorm(delta, 0, 1000, log=TRUE)
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
  ### Log-Likelihood
  mu <- alpha + beta[Data$X[,1]] + gamma[Data$X[,2]] +
    delta*Data$X[,3]
  LL <- sum(dnorm(Data$y, mu, sigma[1], log=TRUE))
  ### Variance Components

```

```

s.beta <- sd(beta)
s.gamma <- sd(gamma)
s.epsilon <- sd(Data$y - mu)
### Log-Posterior
LP <- LL + alpha.prior + beta.prior + gamma.prior + delta.prior +
  sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP, beta[Data$J],
  gamma[Data$K], s.beta, s.gamma, s.epsilon),
  yhat=rnorm(length(mu), mu, sigma[1]), parm=parm)
return(Modelout)
}

```

#### 1.4. Initial Values

```
Initial.Values <- c(0, rep(0,(J-1)), rep(0,(K-1)), 0, rep(1,3))
```

## 2. ANOVA, One-Way

When  $J = 2$ , this is a Bayesian form of a t-test.

### 2.1. Form

$$\begin{aligned}
 \mathbf{y} &\sim \mathcal{N}(\mu, \sigma_1^2) \\
 \mu_i &= \alpha + \beta[\mathbf{x}_i], \quad i = 1, \dots, N \\
 \alpha &\sim \mathcal{N}(0, 1000) \\
 \beta_j &\sim \mathcal{N}(0, \sigma_2^2), \quad j = 1, \dots, J \\
 \beta_J &= -\sum_{j=1}^{J-1} \beta_j \\
 \sigma_{1:2} &\sim \mathcal{HC}(25)
 \end{aligned}$$

### 2.2. Data

```

N <- 100
J <- 3
x <- rcat(N, rep(1/J, J))
alpha <- runif(1,-1,1)
beta <- runif(J-1,-2,2)
beta <- c(beta, -sum(beta))
y <- alpha + beta[x] + rnorm(1,0,0.2)
mon.names <- c("LP","beta[3]")
parm.names <- as.parm.names(list(alpha=0, beta=rep(0,J-1), sigma=rep(0,2)))
pos.alpha <- grep("alpha", parm.names)

```



```

pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnorm(1,0,1), rnorm(Data$J-1,0,1),
  rhalfcauchy(2,5)))
MyData <- list(J=J, N=N, PGF=PGF, mon.names=mon.names,
  parm.names=parm.names, pos.alpha=pos.alpha, pos.beta=pos.beta,
  pos.sigma=pos.sigma, x=x, y=y)

```

## 2.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[Data$pos.alpha]
  beta <- parm[Data$pos.beta]
  beta <- c(beta, -sum(beta)) #Sum-to-zero constraint
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  alpha.prior <- dnorm(alpha, 0, 1000, log=TRUE)
  beta.prior <- sum(dnorm(beta, 0, sigma[2], log=TRUE))
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
  ### Log-Likelihood
  mu <- alpha + beta[Data$x]
  LL <- sum(dnorm(Data$y, mu, sigma[1], log=TRUE))
  ### Log-Posterior
  LP <- LL + alpha.prior + beta.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,beta[Data$J]),
    yhat=rnorm(length(mu), mu, sigma[1]), parm=parm)
  return(Modelout)
}

```

## 2.4. Initial Values

```
Initial.Values <- c(0, rep(0,(J-1)), rep(1,2))
```

# 3. ANOVA, Two-Way

In this representation,  $\sigma^m$  are the superpopulation variance components, **s.beta** and **s.gamma** are the finite-population within-variance components of the factors or treatments, and **s.epsilon** is the finite-population between-variance component.

### 3.1. Form

$$\begin{aligned}
\mathbf{y}_i &\sim \mathcal{N}(\mu_i, \sigma_1^2) \\
\mu_i &= \alpha + \beta[\mathbf{X}_{i,1}] + \gamma[\mathbf{X}_{i,2}], \quad i = 1, \dots, N \\
\epsilon_i &= \mathbf{y}_i - \mu_i \\
\alpha &\sim \mathcal{N}(0, 1000) \\
\beta_j &\sim \mathcal{N}(0, \sigma_2^2), \quad j = 1, \dots, J \\
\beta_J &= - \sum_{j=1}^{J-1} \beta_j \\
\gamma_k &\sim \mathcal{N}(0, \sigma_3^2), \quad k = 1, \dots, K \\
\gamma_K &= - \sum_{k=1}^{K-1} \gamma_k \\
\sigma_m &\sim \mathcal{HC}(25), \quad m = 1, \dots, 3
\end{aligned}$$

### 3.2. Data

```

N <- 100
J <- 5 #Number of levels in factor (treatment) 1
K <- 3 #Number of levels in factor (treatment) 2
X <- cbind(rcat(N,rep(1/J,J)), rcat(N,rep(1/K,K)))
alpha <- runif(1,-1,1)
beta <- runif(J-1,-2,2)
beta <- -sum(beta)
gamma <- runif(K-1,-2,2)
gamma <- -sum(gamma)
y <- alpha + beta[X[,1]] + gamma[X[,2]] + rnorm(1,0,0.1)
mon.names <- c("LP","beta[5]","gamma[3]","s.beta","s.gamma","s.epsilon")
parm.names <- as.parm.names(list(alpha=0, beta=rep(0,J-1), gamma=rep(0,K-1),
  sigma=rep(0,3)))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
pos.gamma <- grep("gamma", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnorm(1,0,1), rnorm(Data$J-1,0,1),
  rnorm(Data$K-1,0,1), rhalfcauchy(3,5)))
MyData <- list(J=J, K=K, N=N, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.alpha=pos.alpha, pos.beta=pos.beta,
  pos.gamma=pos.gamma, pos.sigma=pos.sigma, y=y)

```

### 3.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[Data$pos.alpha]
  beta <- parm[Data$pos.beta]
  beta <- c(beta, -sum(beta)) #Sum-to-zero constraint
  gamma <- parm[Data$pos.gamma]
  gamma <- c(gamma, -sum(gamma)) #Sum-to-zero constraint
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  alpha.prior <- dnorm(alpha, 0, 1000, log=TRUE)
  beta.prior <- sum(dnorm(beta, 0, sigma[2], log=TRUE))
  gamma.prior <- sum(dnorm(gamma, 0, sigma[3], log=TRUE))
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
  ### Log-Likelihood
  mu <- alpha + beta[Data$X[,1]] + gamma[Data$X[,2]]
  LL <- sum(dnorm(Data$y, mu, sigma[1], log=TRUE))
  ### Variance Components
  s.beta <- sd(beta)
  s.gamma <- sd(gamma)
  s.epsilon <- sd(Data$y - mu)
  ### Log-Posterior
  LP <- LL + alpha.prior + beta.prior + gamma.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP, beta[Data$J],
    gamma[Data$K], s.beta, s.gamma, s.epsilon),
    yhat=rnorm(length(mu), mu, sigma[1]), parm=parm)
  return(Modelout)
}
```

### 3.4. Initial Values

```
Initial.Values <- c(0, rep(0,(J-1)), rep(0,(K-1)), rep(1,3))
```

## 4. Approximate Bayesian Computation (ABC)

Approximate Bayesian Computation (ABC), also called likelihood-free estimation, is not a statistical method, but a family of numerical approximation techniques in Bayesian inference. ABC is especially useful when evaluation of the likelihood,  $p(\mathbf{y}|\Theta)$  is computationally prohibitive, or when suitable likelihoods are unavailable. The current example is the application of ABC in the context of linear regression. The log-likelihood is replaced with the negative sum of the distance between  $\mathbf{y}$  and  $\mathbf{y}^{rep}$  as the approximation of the log-likelihood. Distance reduces to the absolute difference. Although linear regression has an easily calculated likelihood, it is used as an example due to its generality. This example demonstrates how

ABC may be estimated either with MCMC via the `LaplacesDemon` function or with Laplace Approximation via the `LaplaceApproximation` function. In this method, a tolerance (which is found often in ABC) does not need to be specified, and the logarithm of the unnormalized joint posterior density is maximized, as usual. The negative and summed distance, above, may be replaced with the negative and summed distance between summaries of the data, rather than the data itself, but this has not been desirable in testing.

#### 4.1. Form

$$\begin{aligned} \mathbf{y} &= \boldsymbol{\mu} + \boldsymbol{\epsilon} \\ \boldsymbol{\mu} &= \mathbf{X}\boldsymbol{\beta} \\ \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \end{aligned}$$

#### 4.2. Data

```
data(demonsnacks)
y <- log(demonsnacks$Calories)
X <- cbind(1, as.matrix(log(demonsnacks[,c(1,4,10)]+1)))
J <- ncol(X)
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
mon.names <- c("LP", "sigma")
parm.names <- as.parm.names(list(beta=rep(0,J)))
pos.beta <- grep("beta", parm.names)
PGF <- function(Data) return(rnormv(Data$J, 0, 1000))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, y=y)
```

#### 4.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  ### Log-Likelihood Approximation
  mu <- as.vector(tcrossprod(Data$X, t(beta)))
  epsilon <- Data$y - mu
  sigma <- sd(epsilon)
  LL <- -sum(abs(epsilon))
  ### Log-Posterior Approximation
  LP <- LL + beta.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP, sigma),
    yhat=rnorm(length(mu), mu, sigma), parm=parm)
```

```
return(Modelout)
}
```

#### 4.4. Initial Values

```
Initial.Values <- c(rep(0,J))
```

### 5. ARCH-M(1,1)

#### 5.1. Form

$$\begin{aligned}
\mathbf{y}_t &\sim \mathcal{N}(\mu_t, \sigma_t^2), \quad t = 1, \dots, T \\
\mathbf{y}^{new} &\sim \mathcal{N}(\mu_{T+1}, \sigma_{new}^2) \\
\mu_t &= \alpha + \phi \mathbf{y}_{t-1} + \delta \sigma_{t-1}^2, \quad t = 1, \dots, (T+1) \\
\epsilon_t &= \mathbf{y}_t - \mu_t \\
\alpha &\sim \mathcal{N}(0, 1000) \\
\phi &\sim \mathcal{N}(0, 1000) \\
\delta &\sim \mathcal{N}(0, 1000) \\
\sigma_{new}^2 &= \omega + \theta \epsilon_T^2 \\
\sigma_t^2 &= \omega + \theta \epsilon_{t-1}^2 \\
\omega &\sim \mathcal{HC}(25) \\
\theta &\sim \mathcal{U}(0, 1)
\end{aligned}$$

#### 5.2. Data

```
y <- c(0.02, -0.51, -0.30, 1.46, -1.26, -2.15, -0.91, -0.53, -1.91,
2.64, 1.64, 0.15, 1.46, 1.61, 1.96, -2.67, -0.19, -3.28,
1.89, 0.91, -0.71, 0.74, -0.10, 3.20, -0.80, -5.25, 1.03,
-0.40, -1.62, -0.80, 0.77, 0.17, -1.39, -1.28, 0.48, -1.02,
0.09, -1.09, 0.86, 0.36, 1.51, -0.02, 0.47, 0.62, -1.36,
1.12, 0.42, -4.39, -0.87, 0.05, -5.41, -7.38, -1.01, -1.70,
0.64, 1.16, 0.87, 0.28, -1.69, -0.29, 0.13, -0.65, 0.83,
0.62, 0.05, -0.14, 0.01, -0.36, -0.32, -0.80, -0.06, 0.24,
0.23, -0.37, 0.00, -0.33, 0.21, -0.10, -0.10, -0.01, -0.40,
-0.35, 0.48, -0.28, 0.08, 0.28, 0.23, 0.27, -0.35, -0.19,
0.24, 0.17, -0.02, -0.23, 0.03, 0.02, -0.17, 0.04, -0.39,
-0.12, 0.16, 0.17, 0.00, 0.18, 0.06, -0.36, 0.22, 0.14,
-0.17, 0.10, -0.01, 0.00, -0.18, -0.02, 0.07, -0.06, 0.06,
-0.05, -0.08, -0.07, 0.01, -0.06, 0.01, 0.01, -0.02, 0.01,
```

```

0.01, 0.12, -0.03, 0.08, -0.10, 0.01, -0.03, -0.08, 0.04,
-0.09, -0.08, 0.01, -0.05, 0.08, -0.14, 0.06, -0.11, 0.09,
0.06, -0.12, -0.01, -0.05, -0.15, -0.05, -0.03, 0.04, 0.00,
-0.12, 0.04, -0.06, -0.05, -0.07, -0.05, -0.14, -0.05, -0.01,
-0.12, 0.05, 0.06, -0.10, 0.00, 0.01, 0.00, -0.08, 0.00,
0.00, 0.07, -0.01, 0.00, 0.09, 0.33, 0.13, 0.42, 0.24,
-0.36, 0.22, -0.09, -0.19, -0.10, -0.08, -0.07, 0.05, 0.07,
0.07, 0.00, -0.04, -0.05, 0.03, 0.08, 0.26, 0.10, 0.08,
0.09, -0.07, -0.33, 0.17, -0.03, 0.07, -0.04, -0.06, -0.06,
0.07, -0.03, 0.00, 0.08, 0.27, 0.11, 0.11, 0.06, -0.11,
-0.09, -0.21, 0.24, -0.12, 0.11, -0.02, -0.03, 0.02, -0.10,
0.00, -0.04, 0.01, 0.02, -0.03, -0.10, -0.09, 0.17, 0.07,
-0.05, -0.01, -0.05, 0.01, 0.00, -0.08, -0.05, -0.08, 0.07,
0.06, -0.14, 0.02, 0.01, 0.04, 0.00, -0.13, -0.17)
T <- length(y)
mon.names <- c("LP", "ynew", "sigma2.new")
parm.names <- c("alpha", "phi", "delta", "omega", "theta")
pos.alpha <- grep("alpha", parm.names)
pos.phi <- grep("phi", parm.names)
pos.delta <- grep("delta", parm.names)
pos.omega <- grep("omega", parm.names)
pos.theta <- grep("theta", parm.names)
PGF <- function(Data) return(c(rnorm(3,0,1), rhalfcauchy(1,5), runif(1)))
MyData <- list(PGF=PGF, T=T, mon.names=mon.names, parm.names=parm.names,
  pos.alpha=pos.alpha, pos.phi=pos.phi, pos.delta=pos.delta,
  pos.omega=pos.omega, pos.theta=pos.theta, y=y)

```

### 5.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[Data$pos.alpha]; phi <- parm[Data$pos.phi]
  delta <- parm[Data$pos.delta]
  omega <- interval(parm[Data$pos.omega], 1e-100, Inf)
  parm[Data$pos.omega] <- omega
  theta <- interval(parm[Data$pos.theta], 1e-10, 1-1e-5)
  parm[Data$pos.theta] <- theta
  ### Log(Prior Densities)
  alpha.prior <- dnormv(alpha, 0, 1000, log=TRUE)
  phi.prior <- dnormv(phi, 0, 1000, log=TRUE)
  delta.prior <- dnormv(delta, 0, 1000, log=TRUE)
  omega.prior <- dhalfcauchy(omega, 25, log=TRUE)
  theta.prior <- dunif(theta, 0, 1, log=TRUE)
  ### Log-Likelihood
  mu <- c(alpha, alpha + phi*Data$y[-Data$T])
}

```

```

epsilon <- Data$y - mu
sigma2 <- c(omega, omega + theta*epsilon[-Data$T]^2)
mu <- mu + delta*sigma2
sigma2.new <- omega + theta*epsilon[Data$T]^2
ynew <- rnorm(1, alpha + phi*Data$y[Data$T] + delta*sigma2[Data$T],
             sigma2.new)
LL <- sum(dnormv(Data$y, mu, sigma2, log=TRUE))
### Log-Posterior
LP <- LL + alpha.prior + phi.prior + delta.prior + omega.prior + theta.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP, ynew, sigma2.new),
                yhat=rnormv(length(mu), mu, sigma2), parm=parm)
return(Modelout)
}

```

#### 5.4. Initial Values

```
Initial.Values <- c(rep(0,3), rep(0.5,2))
```

## 6. Autoregression, AR(1)

### 6.1. Form

$$\begin{aligned}
 \mathbf{y}_t &\sim \mathcal{N}(\mu_t, \sigma^2), \quad t = 1, \dots, T \\
 \mathbf{y}^{new} &= \alpha + \mu_{T+1} \\
 \mu_t &= \alpha + \phi \mathbf{y}_{t-1}, \quad t = 1, \dots, (T+1) \\
 \alpha &\sim \mathcal{N}(0, 1000) \\
 \phi &\sim \mathcal{N}(0, 1000) \\
 \sigma &\sim \mathcal{HC}(25)
 \end{aligned}$$

### 6.2. Data

```

y <- c(0.02, -0.51, -0.30, 1.46, -1.26, -2.15, -0.91, -0.53, -1.91,
      2.64, 1.64, 0.15, 1.46, 1.61, 1.96, -2.67, -0.19, -3.28,
      1.89, 0.91, -0.71, 0.74, -0.10, 3.20, -0.80, -5.25, 1.03,
      -0.40, -1.62, -0.80, 0.77, 0.17, -1.39, -1.28, 0.48, -1.02,
      0.09, -1.09, 0.86, 0.36, 1.51, -0.02, 0.47, 0.62, -1.36,
      1.12, 0.42, -4.39, -0.87, 0.05, -5.41, -7.38, -1.01, -1.70,
      0.64, 1.16, 0.87, 0.28, -1.69, -0.29, 0.13, -0.65, 0.83,
      0.62, 0.05, -0.14, 0.01, -0.36, -0.32, -0.80, -0.06, 0.24,
      0.23, -0.37, 0.00, -0.33, 0.21, -0.10, -0.10, -0.01, -0.40,
      -0.35, 0.48, -0.28, 0.08, 0.28, 0.23, 0.27, -0.35, -0.19,

```

```

0.24, 0.17, -0.02, -0.23, 0.03, 0.02, -0.17, 0.04, -0.39,
-0.12, 0.16, 0.17, 0.00, 0.18, 0.06, -0.36, 0.22, 0.14,
-0.17, 0.10, -0.01, 0.00, -0.18, -0.02, 0.07, -0.06, 0.06,
-0.05, -0.08, -0.07, 0.01, -0.06, 0.01, 0.01, -0.02, 0.01,
0.01, 0.12, -0.03, 0.08, -0.10, 0.01, -0.03, -0.08, 0.04,
-0.09, -0.08, 0.01, -0.05, 0.08, -0.14, 0.06, -0.11, 0.09,
0.06, -0.12, -0.01, -0.05, -0.15, -0.05, -0.03, 0.04, 0.00,
-0.12, 0.04, -0.06, -0.05, -0.07, -0.05, -0.14, -0.05, -0.01,
-0.12, 0.05, 0.06, -0.10, 0.00, 0.01, 0.00, -0.08, 0.00,
0.00, 0.07, -0.01, 0.00, 0.09, 0.33, 0.13, 0.42, 0.24,
-0.36, 0.22, -0.09, -0.19, -0.10, -0.08, -0.07, 0.05, 0.07,
0.07, 0.00, -0.04, -0.05, 0.03, 0.08, 0.26, 0.10, 0.08,
0.09, -0.07, -0.33, 0.17, -0.03, 0.07, -0.04, -0.06, -0.06,
0.07, -0.03, 0.00, 0.08, 0.27, 0.11, 0.11, 0.06, -0.11,
-0.09, -0.21, 0.24, -0.12, 0.11, -0.02, -0.03, 0.02, -0.10,
0.00, -0.04, 0.01, 0.02, -0.03, -0.10, -0.09, 0.17, 0.07,
-0.05, -0.01, -0.05, 0.01, 0.00, -0.08, -0.05, -0.08, 0.07,
0.06, -0.14, 0.02, 0.01, 0.04, 0.00, -0.13, -0.17)
T <- length(y)
mon.names <- c("LP", "ynew")
parm.names <- c("alpha", "phi", "sigma")
PGF <- function(Data) return(c(rnorm(2,0,1), rhalfcauchy(1,5)))
MyData <- list(PGF=PGF, T=T, mon.names=mon.names, parm.names=parm.names,
               y=y)

```

### 6.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[1]; phi <- parm[2]      sigma <- interval(parm[3], 1e-100, Inf)
  parm[3] <- sigma
  ### Log(Prior Densities)
  alpha.prior <- dnormv(alpha, 0, 1000, log=TRUE)
  phi.prior <- dnormv(phi, 0, 1000, log=TRUE)
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
  ### Log-Likelihood
  mu <- c(alpha, alpha + phi*Data$y[-Data$T])
  ynew <- rnorm(1, alpha + phi*Data$y[Data$T], sigma)
  LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + alpha.prior + phi.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,ynew),
                  yhat=rnorm(length(mu), mu, sigma), parm=parm)
  return(Modelout)
}

```



## 6.4. Initial Values

```
Initial.Values <- c(rep(0,2), 1)
```

# 7. Autoregressive Conditional Heteroskedasticity, ARCH(1,1)

## 7.1. Form

$$\begin{aligned}
 \mathbf{y}_t &\sim \mathcal{N}(\mu_t, \sigma_t^2), \quad t = 1, \dots, T \\
 \mathbf{y}^{new} &\sim \mathcal{N}(\mu_{T+1}, \sigma_{new}^2) \\
 \mu_t &= \alpha + \phi \mathbf{y}_{t-1}, \quad t = 1, \dots, (T+1) \\
 \epsilon_t &= \mathbf{y}_t - \mu_t \\
 \alpha &\sim \mathcal{N}(0, 1000) \\
 \phi &\sim \mathcal{N}(0, 1000) \\
 \sigma_{new}^2 &= \omega + \theta \epsilon_T^2 \\
 \sigma_t^2 &= \omega + \theta \epsilon_{t-1}^2, \\
 \omega &\sim \mathcal{HC}(25) \\
 \theta &\sim \mathcal{U}(0, 1)
 \end{aligned}$$

## 7.2. Data

```
y <- c(0.02, -0.51, -0.30, 1.46, -1.26, -2.15, -0.91, -0.53, -1.91,
      2.64, 1.64, 0.15, 1.46, 1.61, 1.96, -2.67, -0.19, -3.28,
      1.89, 0.91, -0.71, 0.74, -0.10, 3.20, -0.80, -5.25, 1.03,
      -0.40, -1.62, -0.80, 0.77, 0.17, -1.39, -1.28, 0.48, -1.02,
      0.09, -1.09, 0.86, 0.36, 1.51, -0.02, 0.47, 0.62, -1.36,
      1.12, 0.42, -4.39, -0.87, 0.05, -5.41, -7.38, -1.01, -1.70,
      0.64, 1.16, 0.87, 0.28, -1.69, -0.29, 0.13, -0.65, 0.83,
      0.62, 0.05, -0.14, 0.01, -0.36, -0.32, -0.80, -0.06, 0.24,
      0.23, -0.37, 0.00, -0.33, 0.21, -0.10, -0.10, -0.01, -0.40,
      -0.35, 0.48, -0.28, 0.08, 0.28, 0.23, 0.27, -0.35, -0.19,
      0.24, 0.17, -0.02, -0.23, 0.03, 0.02, -0.17, 0.04, -0.39,
      -0.12, 0.16, 0.17, 0.00, 0.18, 0.06, -0.36, 0.22, 0.14,
      -0.17, 0.10, -0.01, 0.00, -0.18, -0.02, 0.07, -0.06, 0.06,
      -0.05, -0.08, -0.07, 0.01, -0.06, 0.01, 0.01, -0.02, 0.01,
      0.01, 0.12, -0.03, 0.08, -0.10, 0.01, -0.03, -0.08, 0.04,
      -0.09, -0.08, 0.01, -0.05, 0.08, -0.14, 0.06, -0.11, 0.09,
      0.06, -0.12, -0.01, -0.05, -0.15, -0.05, -0.03, 0.04, 0.00,
      -0.12, 0.04, -0.06, -0.05, -0.07, -0.05, -0.14, -0.05, -0.01,
```

```

-0.12, 0.05, 0.06, -0.10, 0.00, 0.01, 0.00, -0.08, 0.00,
0.00, 0.07, -0.01, 0.00, 0.09, 0.33, 0.13, 0.42, 0.24,
-0.36, 0.22, -0.09, -0.19, -0.10, -0.08, -0.07, 0.05, 0.07,
0.07, 0.00, -0.04, -0.05, 0.03, 0.08, 0.26, 0.10, 0.08,
0.09, -0.07, -0.33, 0.17, -0.03, 0.07, -0.04, -0.06, -0.06,
0.07, -0.03, 0.00, 0.08, 0.27, 0.11, 0.11, 0.06, -0.11,
-0.09, -0.21, 0.24, -0.12, 0.11, -0.02, -0.03, 0.02, -0.10,
0.00, -0.04, 0.01, 0.02, -0.03, -0.10, -0.09, 0.17, 0.07,
-0.05, -0.01, -0.05, 0.01, 0.00, -0.08, -0.05, -0.08, 0.07,
0.06, -0.14, 0.02, 0.01, 0.04, 0.00, -0.13, -0.17)
T <- length(y)
mon.names <- c("LP", "ynew", "sigma2.new")
parm.names <- c("alpha", "phi", "omega", "theta")
PGF <- function(Data) return(c(rnorm(2,0,1), rhalfcauchy(1,5), runif(1)))
MyData <- list(PGF=PGF, T=T, mon.names=mon.names, parm.names=parm.names,
              y=y)

```

### 7.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[1]; phi <- parm[2]
  parm[3] <- omega <- interval(parm[3], 1e-100, Inf)
  parm[4] <- theta <- interval(parm[4], 1e-10, 1-1e-5)
  ### Log(Prior Densities)
  alpha.prior <- dnormv(alpha, 0, 1000, log=TRUE)
  phi.prior <- dnormv(phi, 0, 1000, log=TRUE)
  omega.prior <- dhalfcauchy(omega, 25, log=TRUE)
  theta.prior <- dunif(theta, 0, 1, log=TRUE)
  ### Log-Likelihood
  mu <- c(alpha, alpha + phi*Data$y[-Data$T])
  epsilon <- Data$y - mu
  sigma2 <- c(omega, omega + theta*epsilon[-Data$T]^2)
  sigma2.new <- omega + theta*epsilon[Data$T]^2
  ynew <- rnormv(1, alpha + phi*Data$y[Data$T], sigma2.new)
  LL <- sum(dnormv(Data$y, mu, sigma2, log=TRUE))
  ### Log-Posterior
  LP <- LL + alpha.prior + phi.prior + omega.prior + theta.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP, ynew,
        sigma2.new), yhat=rnormv(length(mu), mu, sigma2), parm=parm)
  return(Modelout)
}

```

## 7.4. Initial Values

```
Initial.Values <- c(rep(0,2), rep(0.5,2))
```

# 8. Autoregressive Moving Average, ARMA(1,1)

## 8.1. Form

$$\mathbf{y}_t \sim \mathcal{N}(\mu_t, \sigma^2), \quad t = 1, \dots, T$$

$$\mathbf{y}^{new} = \alpha + \phi \mathbf{y}_T + \theta \epsilon_T$$

$$\mu_t = \alpha + \phi \mathbf{y}_{t-1} + \theta \epsilon_{t-1}$$

$$\epsilon_t = \mathbf{y}_t - \mu_t$$

$$\alpha \sim \mathcal{N}(0, 1000)$$

$$\phi \sim \mathcal{N}(0, 1000)$$

$$\sigma \sim \mathcal{HC}(25)$$

$$\theta \sim \mathcal{N}(0, 1000)$$

## 8.2. Data

```
y <- c(0.02, -0.51, -0.30, 1.46, -1.26, -2.15, -0.91, -0.53, -1.91,
      2.64, 1.64, 0.15, 1.46, 1.61, 1.96, -2.67, -0.19, -3.28,
      1.89, 0.91, -0.71, 0.74, -0.10, 3.20, -0.80, -5.25, 1.03,
      -0.40, -1.62, -0.80, 0.77, 0.17, -1.39, -1.28, 0.48, -1.02,
      0.09, -1.09, 0.86, 0.36, 1.51, -0.02, 0.47, 0.62, -1.36,
      1.12, 0.42, -4.39, -0.87, 0.05, -5.41, -7.38, -1.01, -1.70,
      0.64, 1.16, 0.87, 0.28, -1.69, -0.29, 0.13, -0.65, 0.83,
      0.62, 0.05, -0.14, 0.01, -0.36, -0.32, -0.80, -0.06, 0.24,
      0.23, -0.37, 0.00, -0.33, 0.21, -0.10, -0.10, -0.01, -0.40,
      -0.35, 0.48, -0.28, 0.08, 0.28, 0.23, 0.27, -0.35, -0.19,
      0.24, 0.17, -0.02, -0.23, 0.03, 0.02, -0.17, 0.04, -0.39,
      -0.12, 0.16, 0.17, 0.00, 0.18, 0.06, -0.36, 0.22, 0.14,
      -0.17, 0.10, -0.01, 0.00, -0.18, -0.02, 0.07, -0.06, 0.06,
      -0.05, -0.08, -0.07, 0.01, -0.06, 0.01, 0.01, -0.02, 0.01,
      0.01, 0.12, -0.03, 0.08, -0.10, 0.01, -0.03, -0.08, 0.04,
      -0.09, -0.08, 0.01, -0.05, 0.08, -0.14, 0.06, -0.11, 0.09,
      0.06, -0.12, -0.01, -0.05, -0.15, -0.05, -0.03, 0.04, 0.00,
      -0.12, 0.04, -0.06, -0.05, -0.07, -0.05, -0.14, -0.05, -0.01,
      -0.12, 0.05, 0.06, -0.10, 0.00, 0.01, 0.00, -0.08, 0.00,
      0.00, 0.07, -0.01, 0.00, 0.09, 0.33, 0.13, 0.42, 0.24,
      -0.36, 0.22, -0.09, -0.19, -0.10, -0.08, -0.07, 0.05, 0.07,
      0.07, 0.00, -0.04, -0.05, 0.03, 0.08, 0.26, 0.10, 0.08,
      0.09, -0.07, -0.33, 0.17, -0.03, 0.07, -0.04, -0.06, -0.06,
```

```

0.07, -0.03, 0.00, 0.08, 0.27, 0.11, 0.11, 0.06, -0.11,
-0.09, -0.21, 0.24, -0.12, 0.11, -0.02, -0.03, 0.02, -0.10,
0.00, -0.04, 0.01, 0.02, -0.03, -0.10, -0.09, 0.17, 0.07,
-0.05, -0.01, -0.05, 0.01, 0.00, -0.08, -0.05, -0.08, 0.07,
0.06, -0.14, 0.02, 0.01, 0.04, 0.00, -0.13, -0.17)
T <- length(y)
mon.names <- c("LP", "ynew")
parm.names <- c("alpha", "phi", "sigma", "theta")
PGF <- function(Data) return(c(rnorm(2,0,1), rhalfcauchy(1,5),
  rnorm(1,0,1)))
MyData <- list(PGF=PGF, T=T, mon.names=mon.names, parm.names=parm.names,
  y=y)

```

### 8.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[1]; phi <- parm[2]; theta <- parm[3]
  parm[4] <- sigma <- interval(parm[4], 1e-100, Inf)
  ### Log(Prior Densities)
  alpha.prior <- dnorm(alpha, 0, 1000, log=TRUE)
  phi.prior <- dnorm(phi, 0, 1000, log=TRUE)
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
  theta.prior <- dnorm(theta, 0, 1000, log=TRUE)
  ### Log-Likelihood
  mu <- c(alpha, alpha + phi*Data$y[-Data$T])
  epsilon <- Data$y - mu
  mu <- c(mu[1], mu[-1] + theta * epsilon[-Data$T])
  ynew <- rnorm(1, alpha + phi*Data$y[Data$T] + theta*epsilon[Data$T],
    sigma)
  LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + alpha.prior + phi.prior + sigma.prior + theta.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP, ynew),
    yhat=rnorm(length(mu), mu, sigma), parm=parm)
  return(Modelout)
}

```

### 8.4. Initial Values

```
Initial.Values <- c(rep(0,2), 0, 1)
```

## 9. Beta Regression

### 9.1. Form

$$\begin{aligned} \mathbf{y} &\sim \mathcal{BETA}(a, b) \\ a &= \mu\phi \\ b &= (1 - \mu)\phi \\ \mu &= \Phi(\beta_1 + \beta_2\mathbf{x}), \quad \mu \in (0, 1) \\ \beta_j &\sim \mathcal{N}(0, 10), \quad j = 1, \dots, J \\ \phi &\sim \mathcal{HC}(25) \end{aligned}$$

where  $\Phi$  is the normal CDF.

### 9.2. Data

```
N <- 100
x <- runif(N)
y <- rbeta(N, (0.5-0.2*x)*3, (1-(0.5-0.2*x))*3) mon.names <- "LP"
parm.names <- c("beta[1]", "beta[2]", "phi")
pos.beta <- grep("beta", parm.names)
pos.phi <- grep("phi", parm.names)
PGF <- function(Data) return(c(rnormv(2,0,10), rhalfcauchy(1,5)))
MyData <- list(PGF=PGF, mon.names=mon.names, parm.names=parm.names,
               pos.beta=pos.beta, pos.phi=pos.phi, x=x, y=y)
```

### 9.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  parm[Data$pos.phi] <- phi <- interval(parm[Data$pos.phi], 1e-100, Inf)
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 10, log=TRUE))
  phi.prior <- dhalfcauchy(phi, 25, log=TRUE)
  ### Log-Likelihood
  mu <- interval(pnorm(beta[1] + beta[2]*Data$x), 0.001, 0.999,
                 reflect=FALSE)
  a <- mu * phi
  b <- (1 - mu) * phi
  LL <- sum(dbeta(Data$y, a, b, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + phi.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
                  yhat=rbeta(length(mu), a, b), parm=parm)
  return(Modelout)
```

```
}
```

#### 9.4. Initial Values

```
Initial.Values <- c(rep(0,2), 0.01)
```

## 10. Beta-Binomial

### 10.1. Form

$$y_i \sim \mathcal{BIN}(\mathbf{n}_i, \pi_i), \quad i = 1, \dots, N$$

$$\pi_i \sim \mathcal{BETA}(\alpha, \beta) \in [0.001, 0.999]$$

### 10.2. Data

```
N <- 20
n <- round(runif(N, 50, 100))
y <- round(runif(N, 1, 10))
mon.names <- "LP"
parm.names <- as.parm.names(list(pi=rep(0,N)))
PGF <- function(Data) return(rbeta(Data$N,1,1))
MyData <- list(N=N, PGF=PGF, mon.names=mon.names, n=n,
  parm.names=parm.names, y=y)
```

### 10.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  parm[1:Data$N] <- pi <- interval(parm[1:Data$N], 0.001, 0.999)
  ### Log(Prior Densities)
  pi.prior <- sum(dbeta(pi, 1, 1, log=TRUE))
  ### Log-Likelihood
  LL <- sum(dbinom(Data$y, Data$n, pi, log=TRUE))
  ### Log-Posterior
  LP <- LL + pi.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rbinom(Data$N, Data$n, pi), parm=parm)
  return(Modelout)
}
```

## 10.4. Initial Values

```
Initial.Values <- c(rep(0.5,N))
```

# 11. Binary Logit

## 11.1. Form

$$\begin{aligned} \mathbf{y} &\sim \mathcal{BERN}(\eta) \\ \eta &= \frac{1}{1 + \exp(-\mu)} \\ \mu &= \mathbf{X}\beta \\ \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \end{aligned}$$

## 11.2. Data

```
data(demonsnacks)
J <- 3
y <- ifelse(demonsnacks$Calories <= 137, 0, 1)
X <- cbind(1, as.matrix(demonsnacks[,c(7,8)]))
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J)))
PGF <- function(Data) return(rnormv(Data$J,0,1000))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
               parm.names=parm.names, y=y)
```

## 11.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[1:Data$J]
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, t(beta))
  eta <- invlogit(mu)
  LL <- sum(dbern(Data$y, eta, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
                  yhat=rbern(length(eta), eta), parm=parm)
```

```
return(Modelout)
}
```

### 11.4. Initial Values

```
Initial.Values <- rep(0,J)
```

## 12. Binary Log-Log Link Mixture

A weighted mixture of the log-log and complementary log-log link functions is used, where  $\alpha$  is the weight. Since the log-log and complementary log-log link functions are asymmetric (as opposed to the symmetric logit and probit link functions), it may be unknown *a priori* whether the log-log or complementary log-log will perform better.

### 12.1. Form

$$\begin{aligned} \mathbf{y} &\sim \mathcal{BERN}(\eta) \\ \eta &= \alpha \exp(-\exp(\mu)) + (1 - \alpha)(1 - \exp(-\exp(\mu))) \\ \mu &= \mathbf{X}\beta \\ \alpha &\sim \mathcal{U}(0, 1) \\ \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \end{aligned}$$

### 12.2. Data

```
data(demonsnacks)
J <- 3
y <- ifelse(demonsnacks$Calories <= 30, 0, 1)
X <- cbind(1, as.matrix(demonsnacks[,c(7,8)]))
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
mon.names <- c("LP", "alpha")
parm.names <- as.parm.names(list(beta=rep(0,J), logit.alpha=0))
PGF <- function(Data) return(c(rnormv(Data$J,0,1000), runif(1)))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, y=y)
```

### 12.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  parm[Data$J+1] <- alpha <- interval(parm[Data$J+1], -700, 700)
  beta <- parm[1:Data$J]
```



```
### Log(Prior Densities)
alpha.prior <- dunif(alpha, 0, 1, log=TRUE)
beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
### Log-Likelihood
mu <- tcrossprod(Data$X, t(beta))
eta <- alpha*invloglog(mu) + (1-alpha)*invloglog(mu)
LL <- sum(dbern(Data$y, eta, log=TRUE))
### Log-Posterior
LP <- LL + alpha.prior + beta.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,alpha),
  yhat=rbern(length(eta), eta), parm=parm)
return(Modelout)
}
```

## 12.4. Initial Values

```
Initial.Values <- c(rep(0,J), 0)
```

# 13. Binary Probit

## 13.1. Form

$$\mathbf{y} \sim \text{BERN}(\mathbf{p})$$

$$\mathbf{p} = \phi(\mu)$$

$$\mu = \mathbf{X}\beta \in [-10, 10]$$

$$\beta_j \sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J$$

where  $\phi$  is the CDF of the standard normal distribution, and  $J=3$ .

## 13.2. Data

```
data(demonsnacks)
J <- 3
y <- ifelse(demonsnacks$Calories <= 137, 0, 1)
X <- cbind(1, as.matrix(demonsnacks[,c(7,8)]))
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J)))
PGF <- function(Data) return(rnormv(Data$J,0,1000))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, y=y)
```

### 13.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[1:Data$J]
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, t(beta))
  mu <- interval(mu, -10, 10, reflect=FALSE)
  p <- pnorm(mu)
  LL <- sum(dbern(Data$y, p, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rbern(length(p), p), parm=parm)
  return(Modelout)
}
```

### 13.4. Initial Values

```
Initial.Values <- rep(0,J)
```

## 14. Binary Robit

### 14.1. Form

$$\begin{aligned} \mathbf{y} &\sim \mathcal{BERN}(\mathbf{p}) \\ \mathbf{p} &= \mathbf{T}_\nu(\boldsymbol{\mu}) \\ \boldsymbol{\mu} &= \mathbf{X}\boldsymbol{\beta} \in [-10, 10] \\ \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\ \nu &\sim \mathcal{U}(5, 10) \end{aligned}$$

where  $\mathbf{T}_\nu$  is the CDF of the standard t-distribution with  $\nu$  degrees of freedom.

### 14.2. Data

```
data(demonsnacks)
y <- ifelse(demonsnacks$Calories <= 137, 0, 1)
X <- cbind(1, as.matrix(demonsnacks[,c(7,8)]))
J <- ncol(X)
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
```

```

mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J), nu=0))
PGF <- function(Data) return(c(rnormv(Data$J,0,1),
  runif(1,5,10)))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, y=y)

```

### 14.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[1:Data$J]
  parm[Data$J+1] <- nu <- interval(parm[Data$J+1], 5, 10)
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  nu.prior <- dunif(nu, 5, 10, log=TRUE)
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, t(beta))
  mu <- interval(mu, -10, 10, reflect=FALSE)
  p <- pst(mu, nu=nu)
  LL <- sum(dbern(Data$y, p, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + nu.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rbern(length(p), p), parm=parm)
  return(Modelout)
}

```

### 14.4. Initial Values

```
Initial.Values <- c(rep(0,J), 5)
```

## 15. Binomial Logit

### 15.1. Form

$$\mathbf{y} \sim \mathcal{BIN}(\mathbf{p}, \mathbf{n})$$

$$\mathbf{p} = \frac{1}{1 + \exp(-\boldsymbol{\mu})}$$

$$\boldsymbol{\mu} = \beta_1 + \beta_2 \mathbf{x}$$

$$\beta_j \sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J$$

## 15.2. Data

```
#10 Trials
exposed <- c(100,100,100,100,100,100,100,100,100,100)
deaths <- c(10,20,30,40,50,60,70,80,90,100)
dose <- c(1,2,3,4,5,6,7,8,9,10)
J <- 2 #Number of parameters
mon.names <- "LP"
parm.names <- c("beta[1]","beta[2]")
PGF <- function(Data) return(rnormv(Data$J,0,1000))
MyData <- list(J=J, PGF=PGF, n=exposed, mon.names=mon.names,
  parm.names=parm.names, x=dose, y=deaths)
```

## 15.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[1:Data$J]
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  ### Log-Likelihood
  mu <- beta[1] + beta[2]*Data$x
  p <- invlogit(mu)
  LL <- sum(dbinom(Data$y, Data$n, p, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rbinom(length(p), Data$n, p), parm=parm)
  return(Modelout)
}
```

## 15.4. Initial Values

```
Initial.Values <- rep(0,J)
```

# 16. Binomial Probit

## 16.1. Form

$$\mathbf{y} \sim \mathcal{BIN}(\mathbf{p}, \mathbf{n})$$

$$\mathbf{p} = \phi(\boldsymbol{\mu})$$

$$\boldsymbol{\mu} = \beta_1 + \beta_2 \mathbf{x} \in [-10, 10]$$

$$\beta_j \sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J$$

where  $\phi$  is the CDF of the standard normal distribution, and  $J=2$ .

## 16.2. Data

```
#10 Trials
exposed <- c(100,100,100,100,100,100,100,100,100,100)
deaths <- c(10,20,30,40,50,60,70,80,90,100)
dose <- c(1,2,3,4,5,6,7,8,9,10)
J <- 2 #Number of parameters
mon.names <- "LP"
parm.names <- c("beta[1]", "beta[2]")
PGF <- function(Data) return(rnormv(Data$J, 0, 1000))
MyData <- list(J=J, PGF=PGF, n=exposed, mon.names=mon.names,
               parm.names=parm.names, x=dose, y=deaths)
```

## 16.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[1:Data$J]
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  ### Log-Likelihood
  mu <- beta[1] + beta[2]*Data$x
  mu <- interval(mu, -10, 10, reflect=FALSE)
  p <- pnorm(mu)
  LL <- sum(dbinom(Data$y, Data$n, p, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
                  yhat=rbinom(length(p), Data$n, p), parm=parm)
  return(Modelout)
}
```

## 16.4. Initial Values

```
Initial.Values <- rep(0, J)
```

# 17. Binomial Robit

### 17.1. Form

$$\begin{aligned} \mathbf{y} &\sim \mathcal{BIN}(\mathbf{p}, \mathbf{n}) \\ \mathbf{p} &= \mathbf{T}_\nu(\boldsymbol{\mu}) \\ \boldsymbol{\mu} &= \beta_1 + \beta_2 \mathbf{x} \in [-10, 10] \\ \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\ \nu &\sim \mathcal{U}(5, 10) \end{aligned}$$

where  $\mathbf{T}_\nu$  is the CDF of the standard t-distribution with  $\nu$  degrees of freedom.

### 17.2. Data

```
#10 Trials
exposed <- c(100,100,100,100,100,100,100,100,100,100)
deaths <- c(10,20,30,40,50,60,70,80,90,100)
dose <- c(1,2,3,4,5,6,7,8,9,10)
J <- 2 #Number of parameters
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,2), nu=0))
PGF <- function(Data) return(c(rnormv(Data$J,0,1000), runif(1,5,10)))
MyData <- list(J=J, PGF=PGF, n=exposed, mon.names=mon.names,
               parm.names=parm.names, x=dose, y=deaths)
```

### 17.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[1:Data$J]
  parm[Data$J+1] <- nu <- interval(parm[Data$J+1], 5, 10)
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  nu.prior <- dunif(nu, 5, 10, log=TRUE)
  ### Log-Likelihood
  mu <- beta[1] + beta[2]*Data$x
  mu <- interval(mu, -10, 10, reflect=FALSE)
  p <- pst(mu, nu=nu)
  LL <- sum(dbinom(Data$y, Data$n, p, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + nu.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
                  yhat=rbinom(length(p), Data$n, p), parm=parm)
  return(Modelout)
}
```

## 17.4. Initial Values

```
Initial.Values <- c(rep(0,J), 5)
```

# 18. Change Point Regression

This example uses a popular variant of the stagnant water data set.

## 18.1. Form

$$\begin{aligned} \mathbf{y} &\sim \mathcal{N}(\mu, \sigma^2) \\ \mu &= \alpha + \beta_1 \mathbf{x} + \beta_2 (\mathbf{x} - \theta) [(\mathbf{x} - \theta) > 0] \\ \alpha &\sim \mathcal{N}(0, 1000) \\ \beta &\sim \mathcal{N}(0, 1000) \\ \sigma &\sim \mathcal{HC}(25) \\ \theta &\sim \mathcal{U}(-1.3, 1.1) \end{aligned}$$

## 18.2. Data

```
N <- 29
y <- c(1.12, 1.12, 0.99, 1.03, 0.92, 0.90, 0.81, 0.83, 0.65, 0.67, 0.60,
      0.59, 0.51, 0.44, 0.43, 0.43, 0.33, 0.30, 0.25, 0.24, 0.13, -0.01,
      -0.13, -0.14, -0.30, -0.33, -0.46, -0.43, -0.65)
x <- c(-1.39, -1.39, -1.08, -1.08, -0.94, -0.80, -0.63, -0.63, -0.25, -0.25,
      -0.12, -0.12, 0.01, 0.11, 0.11, 0.11, 0.25, 0.25, 0.34, 0.34, 0.44,
      0.59, 0.70, 0.70, 0.85, 0.85, 0.99, 0.99, 1.19)
mon.names <- "LP"
parm.names <- as.parm.names(list(alpha=0, beta=rep(0,2), sigma=0, theta=0))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)
pos.theta <- grep("theta", parm.names)
PGF <- function(Data) return(c(rnorm(1), rnorm(2), runif(1), runif(1)))
MyData <- list(N=N, PGF=PGF, mon.names=mon.names, parm.names=parm.names,
  pos.alpha=pos.alpha, pos.beta=pos.beta, pos.sigma=pos.sigma,
  pos.theta=pos.theta, x=x, y=y)
```

## 18.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[Data$pos.alpha]
```

```

beta <- parm[Data$pos.beta]
sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
parm[Data$pos.sigma] <- sigma
theta <- interval(parm[Data$pos.theta], -1.3, 1.1)
parm[Data$pos.theta] <- theta
### Log(Prior Densities)
alpha.prior <- dnorm(alpha, 0, 1000, log=TRUE)
beta.prior <- sum(dnorm(beta, 0, 1000, log=TRUE))
sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
theta.prior <- dunif(theta, -1.3, 1.1, log=TRUE)
### Log-Likelihood
mu <- alpha + beta[1]*x + beta[2]*(x - theta)*(x - theta) > 0
LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
### Log-Posterior
LP <- LL + alpha.prior + beta.prior + sigma.prior + theta.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
yhat=rnorm(length(mu), mu, sigma), parm=parm)
return(Modelout)
}

```

#### 18.4. Initial Values

```
Initial.Values <- c(0.2, -0.45, 0, 0.2, 0)
```

### 19. Cluster Analysis, Confirmatory (CCA)

This is a parametric, model-based, cluster analysis, also called a finite mixture model or latent class cluster analysis, where the number of clusters  $C$  is fixed. When the number of clusters is unknown, exploratory cluster analysis should be used (see section 20). The record-level cluster membership parameter vector,  $\theta$ , is a vector of discrete parameters. Discrete parameters are not supported in all algorithms. The example below is updated with the Griddy-Gibbs sampler.

#### 19.1. Form

$$\mathbf{Y}_{i,j} \sim \mathcal{N}(\mu_{\theta[i],j}, \sigma_{\theta[i]}^2), \quad i = 1, \dots, N, \quad j = 1, \dots, J$$

$$\theta_i \sim \mathcal{CAT}(\pi_{1:C}), \quad i = 1, \dots, N$$

$$\pi_{1:C} \sim \mathcal{D}(\alpha_{1:C})$$

$$\alpha_c = 1$$

$$\mu_{c,j} \sim \mathcal{N}(0, \nu_j^2)$$

$$\sigma_c \sim \mathcal{HC}(25)$$

$$\nu_j \sim \mathcal{HC}(25)$$



## 19.2. Data

```

data(demonsnacks)
Y <- as.matrix(log(demonsnacks + 1))
N <- nrow(Y)
J <- ncol(Y)
for (j in 1:J) {Y[,j] <- CenterScale(Y[,j])}
C <- 3 #Number of clusters
alpha <- rep(1,C) #Prior probability of cluster proportion
mon.names <- c("LP", paste("pi[", 1:C, "]", sep=""))
parm.names <- as.parm.names(list(theta=rep(0,N), mu=matrix(0,C,J),
  nu=rep(0,J), sigma=rep(0,C)))
pos.theta <- grep("theta", parm.names)
pos.mu <- grep("mu", parm.names)
pos.nu <- grep("nu", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rcat(N, p=rep(1/C, C)),
  rnorm(Data$C*Data$J,0,1), rhalfcauchy(Data$J+Data$C,5)))
MyData <- list(C=C, J=J, N=N, PGF=PGF, Y=Y, alpha=alpha,
  mon.names=mon.names, parm.names=parm.names, pos.theta=pos.theta,
  pos.mu=pos.mu, pos.nu=pos.nu, pos.sigma=pos.sigma)

```

## 19.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  theta <- parm[Data$pos.theta]
  mu <- matrix(parm[Data$pos.mu], Data$C, Data$J)
  parm[Data$pos.nu] <- nu <- interval(parm[Data$pos.nu], 1e-100, Inf)
  pi <- rep(0, Data$C)
  tab <- table(theta)
  pi[as.numeric(names(tab))] <- as.vector(tab)
  pi <- pi / sum(pi)
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  theta.prior <- sum(dcat(theta, pi, log=TRUE))
  mu.prior <- sum(dnorm(mu, 0, matrix(rep(nu,Data$C), Data$C,
    Data$J, byrow=TRUE), log=TRUE))
  nu.prior <- sum(dhalfcauchy(nu, 25, log=TRUE))
  pi.prior <- ddirichlet(pi, Data$alpha, log=TRUE)
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
  ### Log-Likelihood
  LL <- sum(dnorm(Data$Y, mu[theta,], sigma[theta], log=TRUE))
  ### Log-Posterior

```

```

LP <- LL + theta.prior + mu.prior + nu.prior + pi.prior +
  sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,pi),
  yhat=rnorm(prod(dim(mu[theta,])), mu[theta,], sigma[theta]),
  parm=parm)
return(Modelout)
}

```

## 19.4. Initial Values

```
Initial.Values <- c(rcat(N,rep(1/C,C)), rep(0,C*J), rep(1,J), rep(1,C))
```

# 20. Cluster Analysis, Exploratory (ECA)

In “exploratory cluster analysis”, the optimal number of clusters  $C$  is unknown before the model update. This is a nonparametric, model-based, infinite mixture model that uses truncated stick-breaking within a truncated Dirichlet process. The user must specify the maximum number of clusters (mixture components),  $C$  to explore, where  $C$  is discrete, greater than one, and less than the number of records,  $N$ . The records in the  $N \times J$  matrix  $\mathbf{Y}$  are clustered, where  $J$  is the number of predictors. The record-level cluster membership parameter vector,  $\theta$ , is a vector of discrete parameters. Discrete parameters are not supported in all algorithms. The example below is updated with the Griddy-Gibbs sampler.

## 20.1. Form

$$\begin{aligned}
 \mathbf{Y}_{i,j} &\sim \mathcal{N}(\mu_{\theta[i],j}, \sigma_{\theta[i]}^2), \quad i = 1, \dots, N, \quad j = 1, \dots, J \\
 \theta_i &\sim \mathcal{CAT}(\pi_{1:C}), \quad i = 1, \dots, N \\
 \pi_{1:C} &\sim \text{Stick}(\gamma) \\
 \mu_{c,j} &\sim \mathcal{N}(0, \nu_j^2) \\
 \alpha &\sim \mathcal{HC}(25) \\
 \beta &\sim \mathcal{HC}(25) \\
 \gamma &\sim \mathcal{G}(\alpha, \beta) \\
 \sigma_c &\sim \mathcal{HC}(25) \\
 \nu_j &\sim \mathcal{HC}(25)
 \end{aligned}$$

## 20.2. Data

```

data(demonsnacks)
Y <- as.matrix(log(demonsnacks + 1))
N <- nrow(Y)
J <- ncol(Y)

```

```

for (j in 1:J) {Y[,j] <- CenterScale(Y[,j])}
C <- 3 #Number of clusters to explore
mon.names <- c("LP", paste("pi[", 1:C, "]", sep=""))
parm.names <- as.parm.names(list(theta=rep(0,N), mu=matrix(0,C,J),
  nu=rep(0,J), sigma=rep(0,C), alpha=0, beta=0, gamma=0))
pos.theta <- grep("theta", parm.names)
pos.mu <- grep("mu", parm.names)
pos.nu <- grep("nu", parm.names)
pos.sigma <- grep("sigma", parm.names)
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
pos.gamma <- grep("gamma", parm.names)
PGF <- function(Data) return(c(rcat(Data$N, p=rep(1/Data$C,Data$C)),
  rnorm(Data$C*Data$J,0,1), rhalfcauchy(Data$J+Data$C,5),
  rhalfcauchy(2,5), rgamma(1,rhalfcauchy(2,5))))
MyData <- list(C=C, J=J, N=N, PGF=PGF, Y=Y, mon.names=mon.names,
  parm.names=parm.names, pos.theta=pos.theta, pos.mu=pos.mu,
  pos.nu=pos.nu, pos.sigma=pos.sigma, pos.alpha=pos.alpha,
  pos.beta=pos.beta, pos.gamma=pos.gamma)

```

## 20.3. Model

```

Model <- function(parm, Data)
{
  ### Hyperhyperparameters
  alpha <- interval(parm[Data$pos.alpha], 1e-100, Inf)
  parm[Data$pos.alpha] <- alpha
  beta <- interval(parm[Data$pos.beta], 1e-100, Inf)
  parm[Data$pos.beta] <- beta
  ### Hyperparameters
  gamma <- interval(parm[Data$pos.gamma], 1e-100, Inf)
  parm[Data$pos.nu] <- nu <- interval(parm[Data$pos.nu], 1e-100, Inf)
  ### Parameters
  theta <- parm[Data$pos.theta]
  mu <- matrix(parm[Data$pos.mu], Data$C, Data$J)
  pi <- rStick(Data$C-1, gamma)
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Hyperhyperprior Densities)
  alpha.prior <- dhalfcauchy(alpha, 25, log=TRUE)
  beta.prior <- dhalfcauchy(beta, 25, log=TRUE)
  ### Log(Hyperprior Densities)
  gamma.prior <- dgamma(gamma, alpha, beta, log=TRUE)
  nu.prior <- sum(dhalfcauchy(nu, 25, log=TRUE))
  ### Log(Prior Densities)
  theta.prior <- sum(dcat(theta, pi, log=TRUE))
}

```

```

mu.prior <- sum(dnorm(mu, 0, matrix(rep(nu,Data$C), Data$C,
  Data$J, byrow=TRUE), log=TRUE))
pi.prior <- dStick(pi, gamma, log=TRUE)
sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
### Log-Likelihood
LL <- sum(dnorm(Data$Y, mu[theta,], sigma[theta], log=TRUE))
### Log-Posterior
LP <- LL + theta.prior + mu.prior + nu.prior + pi.prior +
  alpha.prior + beta.prior + gamma.prior + sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,pi),
  yhat=rnorm(prod(dim(mu[theta,])), mu[theta,], sigma[theta]),
  parm=parm)
return(Modelout)
}

```

## 20.4. Initial Values

```

Initial.Values <- c(rcat(N,p=rep(1/C,C)), rep(0,C*J), rep(1,J), rep(1,C),
  rep(1,3))

```

## 21. Conditional Autoregression (CAR), Poisson

This CAR example is a slightly modified form of example 7.3 (Model A) in [Congdon \(2003\)](#). The Scottish lip cancer data also appears in the WinBUGS ([Spiegelhalter, Thomas, Best, and Lunn 2003](#)) examples and is a widely analyzed example. The data  $\mathbf{y}$  consists of counts for  $i = 1, \dots, 56$  counties in Scotland. A single predictor  $\mathbf{x}$  is provided. The errors,  $\epsilon$ , are allowed to include spatial effects as smoothing by spatial effects from areal neighbors. The vector  $\epsilon_\mu$  is the mean of each area's error, and is a weighted average of errors in contiguous areas. Areal neighbors are indicated in adjacency matrix  $\mathbf{A}$ .

### 21.1. Form

$$\begin{aligned}
 \mathbf{y} &\sim \mathcal{P}(\lambda) \\
 \lambda &= \exp(\log(\mathbf{E}) + \beta_1 + \beta_2 \mathbf{x} + \epsilon) \\
 \epsilon &\sim \mathcal{N}(\epsilon_\mu, \sigma^2) \\
 \epsilon_{\mu[i]} &= \rho \sum_{j=1}^J \mathbf{A}_{i,j} \epsilon_j, \quad i = 1, \dots, N \\
 \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\
 \rho &\sim \mathcal{U}(-1, 1) \\
 \sigma &\sim \mathcal{HC}(25)
 \end{aligned}$$

## 21.2. Data

```

N <- 56 #Number of areas
NN <- 264 #Number of adjacent areas
y <- c(9,39,11,9,15,8,26,7,6,20,13,5,3,8,17,9,2,7,9,7,16,31,11,7,19,15,7,
      10,16,11,5,3,7,8,11,9,11,8,6,4,10,8,2,6,19,3,2,3,28,6,1,1,1,1,0,0)
E <- c( 1.4,8.7,3.0,2.5,4.3,2.4,8.1,2.3,2.0,6.6,4.4,1.8,1.1,3.3,7.8,4.6,
      1.1,4.2,5.5,4.4,10.5,22.7,8.8,5.6,15.5,12.5,6.0,9.0,14.4,10.2,4.8,
      2.9,7.0,8.5,12.3,10.1,12.7,9.4,7.2,5.3,18.8,15.8,4.3,14.6,50.7,8.2,
      5.6,9.3,88.7,19.6,3.4,3.6,5.7,7.0,4.2,1.8) #Expected
x <- c(16,16,10,24,10,24,10,7,7,16,7,16,10,24,7,16,10,7,7,10,7,16,10,7,1,1,
      7,7,10,10,7,24,10,7,7,0,10,1,16,0,1,16,16,0,1,7,1,1,0,1,1,0,1,1,16,10)
A <- matrix(0, N, N)
A[1,c(5,9,11,19)] <- 1 #Area 1 is adjacent to areas 5, 9, 11, and 19
A[2,c(7,10)] <- 1 #Area 2 is adjacent to areas 7 and 10
A[3,c(6,12)] <- 1; A[4,c(18,20,28)] <- 1; A[5,c(1,11,12,13,19)] <- 1
A[6,c(3,8)] <- 1; A[7,c(2,10,13,16,17)] <- 1; A[8,6] <- 1
A[9,c(1,11,17,19,23,29)] <- 1; A[10,c(2,7,16,22)] <- 1
A[11,c(1,5,9,12)] <- 1; A[12,c(3,5,11)] <- 1; A[13,c(5,7,17,19)] <- 1
A[14,c(31,32,35)] <- 1; A[15,c(25,29,50)] <- 1
A[16,c(7,10,17,21,22,29)] <- 1; A[17,c(7,9,13,16,19,29)] <- 1
A[18,c(4,20,28,33,55,56)] <- 1; A[19,c(1,5,9,13,17)] <- 1
A[20,c(4,18,55)] <- 1; A[21,c(16,29,50)] <- 1; A[22,c(10,16)] <- 1
A[23,c(9,29,34,36,37,39)] <- 1; A[24,c(27,30,31,44,47,48,55,56)] <- 1
A[25,c(15,26,29)] <- 1; A[26,c(25,29,42,43)] <- 1
A[27,c(24,31,32,55)] <- 1; A[28,c(4,18,33,45)] <- 1
A[29,c(9,15,16,17,21,23,25,26,34,43,50)] <- 1
A[30,c(24,38,42,44,45,56)] <- 1; A[31,c(14,24,27,32,35,46,47)] <- 1
A[32,c(14,27,31,35)] <- 1; A[33,c(18,28,45,56)] <- 1
A[34,c(23,29,39,40,42,43,51,52,54)] <- 1; A[35,c(14,31,32,37,46)] <- 1
A[36,c(23,37,39,41)] <- 1; A[37,c(23,35,36,41,46)] <- 1
A[38,c(30,42,44,49,51,54)] <- 1; A[39,c(23,34,36,40,41)] <- 1
A[40,c(34,39,41,49,52)] <- 1; A[41,c(36,37,39,40,46,49,53)] <- 1
A[42,c(26,30,34,38,43,51)] <- 1; A[43,c(26,29,34,42)] <- 1
A[44,c(24,30,38,48,49)] <- 1; A[45,c(28,30,33,56)] <- 1
A[46,c(31,35,37,41,47,53)] <- 1; A[47,c(24,31,46,48,49,53)] <- 1
A[48,c(24,44,47,49)] <- 1; A[49,c(38,40,41,44,47,48,52,53,54)] <- 1
A[50,c(15,21,29)] <- 1; A[51,c(34,38,42,54)] <- 1
A[52,c(34,40,49,54)] <- 1; A[53,c(41,46,47,49)] <- 1
A[54,c(34,38,49,51,52)] <- 1; A[55,c(18,20,24,27,56)] <- 1
A[56,c(18,24,30,33,45,55)] <- 1
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,2), epsilon=rep(0,N), rho=0,
  sigma=0))
pos.beta <- grep("beta", parm.names)
pos.epsilon <- grep("epsilon", parm.names)
pos.rho <- grep("rho", parm.names)

```

```
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(2,0,1000), rnorm(Data$N,0,1),
  runif(1,-1,1), rhalfcauchy(1,5)))
MyData <- list(A=A, E=E, N=N, PGF=PGF, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.epsilon=pos.epsilon,
  pos.rho=pos.rho, pos.sigma=pos.sigma, x=x, y=y)
```

### 21.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  epsilon <- parm[Data$pos.epsilon]
  parm[Data$pos.rho] <- rho <- interval(parm[Data$pos.rho], -1, 1)
  epsilon.mu <- rho * rowSums(epsilon * Data$A)
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  epsilon.prior <- sum(dnorm(epsilon, epsilon.mu, sigma, log=TRUE))
  rho.prior <- dunif(rho, -1, 1, log=TRUE)
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
  ### Log-Likelihood
  lambda <- exp(log(Data$E) + beta[1] + beta[2]*Data$x/10 + epsilon)
  LL <- sum(dpois(Data$y, lambda, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + epsilon.prior + rho.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rpois(length(lambda), lambda), parm=parm)
  return(Modelout)
}
```

### 21.4. Initial Values

```
Initial.Values <- c(rep(0,2), rep(0,N), 0, 1)
```

## 22. Conditional Predictive Ordinate

For a more complete introduction to the conditional predictive ordinate (CPO), see the vignette entitled “Bayesian Inference”. Following is a brief guide to the applied use of CPO.

To include CPO in any model that is to be updated with MCMC, calculate and monitor the record-level inverse of the likelihood,  $\text{Inv}L_i$  for records  $i = 1, \dots, N$ .  $\text{CPO}_i$  is the inverse of the posterior mean of  $\text{Inv}L_i$ . The inverse  $\text{CPO}_i$ , or  $\text{ICPO}_i$ , is the posterior mean of  $\text{Inv}L_i$ . ICPOs larger than 40 can be considered as possible outliers, and higher than 70 as extreme

values.

Here, CPO is added to the linear regression example in section 46. In this data, record 6 is a possible outlier, and record 8 is an extreme value.

### 22.1. Form

$$\begin{aligned} \mathbf{y} &\sim \mathcal{N}(\mu, \sigma^2) \\ \mu &= \mathbf{X}\beta \\ \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\ \sigma &\sim \mathcal{HC}(25) \end{aligned}$$

### 22.2. Data

```
data(demonsnacks)
y <- log(demonsnacks$Calories)
X <- cbind(1, as.matrix(log(demonsnacks[,c(1,4,10)]+1)))
J <- ncol(X)
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
mon.names <- c("LP", as.parm.names(list(InvL=rep(0,N))))
parm.names <- as.parm.names(list(beta=rep(0,J), sigma=0))
pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,1000), rhalfcauchy(1,5)))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.sigma=pos.sigma, y=y)
```

### 22.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, t(beta))
  LL <- dnorm(Data$y, mu, sigma, log=TRUE)
  InvL <- 1 / exp(LL)
  LL <- sum(LL)
  ### Log-Posterior
```

```

LP <- LL + beta.prior + sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,InvL),
  yhat=rnorm(length(mu), mu, sigma), parm=parm)
return(Modelout)
}

```

## 22.4. Initial Values

```
Initial.Values <- c(rep(0,J), 1)
```

# 23. Contingency Table

The two-way contingency table, matrix  $\mathbf{Y}$ , can easily be extended to more dimensions. Contingency table  $\mathbf{Y}$  has  $J$  rows and  $K$  columns. The cell counts are fit with Poisson regression, according to intercept  $\alpha$ , main effects  $\beta_j$  for each row, main effects  $\gamma_k$  for each column, and interaction effects  $\delta_{j,k}$  for dependence effects. An omnibus (all cells) test of independence is done by estimating two models (one with  $\delta$ , and one without), and a large enough Bayes Factor indicates a violation of independence when the model with  $\delta$  fits better than the model without  $\delta$ . In an ANOVA-like style, main effects contrasts can be used to distinguish rows or groups of rows from each other, as well as with columns. Likewise, interaction effects contrasts can be used to test independence in groups of  $\delta_{j,k}$  elements. Finally, single-cell interactions can be used to indicate violations of independence for a given cell, such as when zero is not within its 95% probability interval.

## 23.1. Form

$$\begin{aligned}
 \mathbf{Y}_{j,k} &\sim \mathcal{P}(\lambda_{j,k}), \quad j = 1, \dots, J, \quad k = 1, \dots, K \\
 \lambda_{j,k} &= \exp(\alpha + \beta_j + \gamma_k + \delta_{j,k}), \quad j = 1, \dots, J, \quad k = 1, \dots, K \\
 \alpha &\sim \mathcal{N}(0, 1000) \\
 \beta_j &\sim \mathcal{N}(0, \beta_\sigma^2), \quad j = 1, \dots, J \\
 \beta_J &= -\sum_{j=1}^{J-1} \beta_j \\
 \beta_\sigma &\sim \mathcal{HC}(25) \\
 \gamma_k &\sim \mathcal{N}(0, \gamma_\sigma^2), \quad k = 1, \dots, K \\
 \gamma_K &= -\sum_{k=1}^{K-1} \gamma_k \\
 \gamma_\sigma &\sim \mathcal{HC}(25) \\
 \delta_{j,k} &\sim \mathcal{N}(0, \delta_\sigma^2) \\
 \delta_{J,K} &= -\sum \delta_{-J,-K}
 \end{aligned}$$



$$\delta_\sigma \sim \mathcal{HC}(25)$$

## 23.2. Data

```

J <- 3 #Rows
K <- 3 #Columns
alpha <- runif(1,10,20)
beta <- rnorm(J-1, 0, 2); beta <- c(beta, -sum(beta))
gamma <- rnorm(K-1, 0, 2); gamma <- c(gamma, -sum(gamma))
delta <- rnorm(J*K-1, 0, 0.5); delta <- c(delta, -sum(delta))
Y <- matrix(alpha + matrix(beta, J, K) + matrix(gamma, J, K, byrow=TRUE) +
            matrix(delta, J, K), J, K)
Y <- round(Y)
mon.names <- "LP"
parm.names <- as.parm.names(list(alpha=0, beta=rep(0,J-1),
                                gamma=rep(0,K-1), delta=rep(0,J*K-1), b.sigma=0, g.sigma=0,
                                d.sigma=0))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
pos.gamma <- grep("gamma", parm.names)
pos.delta <- grep("delta", parm.names)
pos.b.sigma <- grep("b.sigma", parm.names)
pos.g.sigma <- grep("g.sigma", parm.names)
pos.d.sigma <- grep("d.sigma", parm.names)
PGF <- function(Data) return(c(rnorm(1,log(mean(Y)),1), rnorm(Data$J-1),
                               rnorm(Data$K-1), rnorm(Data$J*Data$K-1), rhalfcauchy(3,5)))
MyData <- list(J=J, K=K, PGF=PGF, Y=Y, mon.names=mon.names,
              parm.names=parm.names, pos.alpha=pos.alpha, pos.beta=pos.beta,
              pos.gamma=pos.gamma, pos.delta=pos.delta, pos.b.sigma=pos.b.sigma,
              pos.g.sigma=pos.g.sigma, pos.d.sigma=pos.d.sigma)

```

## 23.3. Model

```

Model <- function(parm, Data)
{
  ### Hyperparameters
  beta.sigma <- interval(parm[Data$pos.b.sigma], 1e-100, Inf)
  parm[Data$pos.b.sigma] <- beta.sigma
  gamma.sigma <- interval(parm[Data$pos.g.sigma], 1e-100, Inf)
  parm[Data$pos.g.sigma] <- gamma.sigma
  delta.sigma <- interval(parm[Data$pos.d.sigma], 1e-100, Inf)
  parm[Data$pos.d.sigma] <- delta.sigma
  ### Parameters
  alpha <- parm[Data$pos.alpha]
  beta <- parm[Data$pos.beta]
  beta <- c(beta, -sum(beta))

```

```

gamma <- parm[Data$pos.gamma]
gamma <- c(gamma, -sum(gamma))
delta <- parm[Data$pos.delta]
delta <- c(delta, -sum(delta))
delta <- matrix(delta, Data$J, Data$K)
### Log(Hyperprior Densities)
beta.sigma.prior <- dhalfcauchy(beta.sigma, 25, log=TRUE)
gamma.sigma.prior <- dhalfcauchy(gamma.sigma, 25, log=TRUE)
delta.sigma.prior <- dhalfcauchy(delta.sigma, 25, log=TRUE)
### Log(Prior Densities)
alpha.prior <- dnorm(alpha, 0, 1000, log=TRUE)
beta.prior <- sum(dnorm(beta, 0, beta.sigma, log=TRUE))
gamma.prior <- sum(dnorm(gamma, 0, gamma.sigma, log=TRUE))
delta.prior <- sum(dnorm(delta, 0, delta.sigma, log=TRUE))
### Log-Likelihood
beta <- matrix(beta, Data$J, Data$K)
gamma <- matrix(gamma, Data$J, Data$K, byrow=TRUE)
lambda <- exp(alpha + beta + gamma + delta)
LL <- sum(dpois(Data$Y, lambda, log=TRUE))
### Log-Posterior
LP <- LL + alpha.prior + beta.prior + beta.sigma.prior +
      gamma.prior + gamma.sigma.prior + delta.prior +
      delta.sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
                 yhat=rpois(length(lambda), lambda),
                 parm=parm)
return(Modelout)
}

```

### 23.4. Initial Values

```

Initial.Values <- c(log(mean(Y)), rep(0,J-1), rep(0,K-1), rep(0,J*K-1),
                    rep(1,3))

```

## 24. Covariance Separation Strategy

A Seemingly Unrelated Regression (SUR) model is used to provide an example of a flexible way to estimate covariance or precision matrices with the “separation strategy” decomposition of [Barnard, McCulloch, and Meng \(2000\)](#). For more information on SUR models, see section 81.

The most common way of specifying a covariance matrix, such as for the multivariate normal distribution, may be with the conjugate inverse Wishart distribution. Alternatively, the conjugate Wishart distribution is often used for a precision matrix. The Wishart and inverse Wishart distributions, however, do not always perform well, due to only one parameter for

variability, and usually in the case of small sample sizes or when its dimension approaches the sample size. There are several alternatives. This example decomposes a covariance matrix into a standard deviation vector and a correlation matrix, each of which are easy to understand (as opposed to setting priors on eigenvalues). A precision matrix may be decomposed similarly, though the separated components are interpreted differently.

Barnard *et al.* (2000) prefer to update the covariance separation strategy with Gibbs sampling rather than Metropolis-Hastings, though the form presented here works well in testing with Adaptive MCMC.

### 24.1. Form

$$\begin{aligned}
\mathbf{Y}_{t,j} &\sim \mathcal{N}(\mu_{t,j}, \Sigma), \quad t = 1, \dots, T; \quad j = 1, \dots, J \\
\mu_{t,1} &= \alpha_1 + \alpha_2 \mathbf{X}_{t-1,1} + \alpha_3 \mathbf{X}_{t-1,2}, \quad t = 2, \dots, T \\
\mu_{t,2} &= \beta_1 + \beta_2 \mathbf{X}_{t-1,3} + \beta_3 \mathbf{X}_{t-1,4}, \quad t = 2, \dots, T \\
\Sigma &= \mathbf{SRS} \\
\alpha_k &\sim \mathcal{N}(0, 1000), \quad k = 1, \dots, K \\
\beta_k &\sim \mathcal{N}(0, 1000), \quad k = 1, \dots, K \\
\mathbf{R}_{i,j} &\sim \mathcal{N}(\rho_\mu, \rho_\sigma^2), \quad \mathbf{R}_{i,j} \in [-1, 1], \quad i = 1, \dots, J \\
\mathbf{S} &= \sigma \mathbf{I}_J \\
\rho_\mu &\sim \mathcal{N}(0, 2), \quad \in [-1, 1] \\
\rho_\sigma &\sim \mathcal{HC}(25), \quad \in (0, 1000] \\
\sigma_j &\sim \mathcal{N}(\sigma_\mu, \sigma_\sigma) \\
\sigma_\mu &\sim \mathcal{HN}(1000), \quad \in (0, 1000] \\
\sigma_\sigma &\sim \mathcal{HC}(25)
\end{aligned}$$

### 24.2. Data

```

T <- 20 #Time-periods
year <- c(1935,1936,1937,1938,1939,1940,1941,1942,1943,1944,1945,1946,
1947,1948,1949,1950,1951,1952,1953,1954)
IG <- c(33.1,45.0,77.2,44.6,48.1,74.4,113.0,91.9,61.3,56.8,93.6,159.9,
147.2,146.3,98.3,93.5,135.2,157.3,179.5,189.6)
VG <- c(1170.6,2015.8,2803.3,2039.7,2256.2,2132.2,1834.1,1588.0,1749.4,
1687.2,2007.7,2208.3,1656.7,1604.4,1431.8,1610.5,1819.4,2079.7,
2371.6,2759.9)
CG <- c(97.8,104.4,118.0,156.2,172.6,186.6,220.9,287.8,319.9,321.3,319.6,
346.0,456.4,543.4,618.3,647.4,671.3,726.1,800.3,888.9)
IW <- c(12.93,25.90,35.05,22.89,18.84,28.57,48.51,43.34,37.02,37.81,
39.27,53.46,55.56,49.56,32.04,32.24,54.38,71.78,90.08,68.60)
VW <- c(191.5,516.0,729.0,560.4,519.9,628.5,537.1,561.2,617.2,626.7,

```

```

737.2,760.5,581.4,662.3,583.8,635.2,723.8,864.1,1193.5,1188.9)
CW <- c(1.8,0.8,7.4,18.1,23.5,26.5,36.2,60.8,84.4,91.2,92.4,86.0,111.1,
130.6,141.8,136.7,129.7,145.5,174.8,213.5)
J <- 2 #Number of dependent variables
Y <- matrix(c(IG,IW), T, J)
R <- diag(J)
mon.names <- "LP"
parm.names <- as.parm.names(list(alpha=rep(0,3), beta=rep(0,3),
R=diag(J), rho.mu=0, rho.sigma=0, log.sigma=rep(0,J), sigma.mu=0,
log.sig.sigma=0), uppertri=c(0,0,1,0,0,0,0,0))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
pos.R <- grep("R", parm.names)
pos.rho.mu <- grep("rho.mu", parm.names)
pos.rho.sigma <- grep("rho.sigma", parm.names)
pos.log.sigma <- grep("log.sigma", parm.names)
pos.sigma.mu <- grep("sigma.mu", parm.names)
pos.log.sig.sigma <- grep("log.sig.sigma", parm.names)
PGF <- function(Data) return(c(rnormv(3,0,10), rnormv(3,0,10),
runif(length(upper.triangle(diag(Data$J), diag=TRUE))), -1, 1),
rtrunc(1, "norm", a=-1, b=1, mean=0, sd=2),
log(rhalfcauchy(Data$J+1,5)), rhalfnorm(1, 10),
log(rhalfcauchy(1,5))))
MyData <- list(J=J, PGF=PGF, T=T, Y=Y, CG=CG, CW=CW, IG=IG, IW=IW, VG=VG,
VW=VW, mon.names=mon.names, parm.names=parm.names, )
pos.alpha=pos.alpha, pos.beta=pos.beta, pos.R=pos.R,
pos.rho.mu=pos.rho.mu, pos.rho.sigma=pos.rho.sigma,
pos.log.sigma=pos.log.sigma, pos.sigma.mu=pos.sigma.mu,
pos.log.sig.sigma=pos.log.sig.sigma)

```

### 24.3. Model

```

Model <- function(parm, Data)
{
  ### Hyperparameters
  rho.mu <- interval(parm[Data$pos.rho.mu], -1, 1)
  parm[Data$pos.rho.mu] <- rho.mu
  rho.sigma <- interval(parm[Data$pos.rho.sigma], 1e-100, Inf)
  parm[Data$pos.rho.sigma] <- rho.sigma
  sigma.mu <- interval(parm[Data$pos.sigma.mu], 1e-100, Inf)
  parm[Data$pos.sigma.mu] <- sigma.mu
  sigma.sigma <- sigma.sigma <- exp(parm[Data$pos.log.sig.sigma])
  ### Parameters
  alpha <- parm[Data$pos.alpha]
  beta <- parm[Data$pos.beta]
  R <- as.parm.matrix(R, Data$J, parm, Data, a=-1, b=1)

```

```

parm[Data$pos.R] <- upper.triangle(R, diag=TRUE)
sigma <- exp(parm[Data$pos.log.sigma])
S <- diag(sigma)
Sigma <- as.symmetric.matrix(S %*% R %*% S)
### Log(Hyperprior Densities)
rho.mu.prior <- dtrunc(rho.mu, "norm", a=-1, b=1, mean=0, sd=2,
  log=TRUE)
rho.sigma.prior <- dhalfcauchy(rho.sigma, 25, log=TRUE)
sigma.mu.prior <- dhalfnorm(sigma.mu, 1000, log=TRUE)
sigma.sigma.prior <- dhalfcauchy(sigma.sigma, 25, log=TRUE)
### Log(Prior Densities)
alpha.prior <- sum(dnorm(alpha, 0, 1000, log=TRUE))
beta.prior <- sum(dnorm(beta, 0, 1000, log=TRUE))
R.prior <- sum(dtrunc(upper.triangle(R, diag=TRUE), "norm",
  a=-1, b=1, mean=rho.mu, sd=rho.sigma, log=TRUE))
sigma.prior <- sum(dnorm(sigma, sigma.mu, sigma.sigma, log=TRUE))
### Log-Likelihood
mu <- Data$Y
mu[-1,1] <- alpha[1] + alpha[2]*Data$CG[-Data$T] +
  alpha[3]*Data$VG[-Data$T]
mu[-1,2] <- beta[1] + beta[2]*Data$CW[-Data$T] +
  beta[3]*Data$VW[-Data$T]
LL <- sum(dmvn(Data$Y[-1,], mu[-1,], Sigma, log=TRUE))
### Log-Posterior
LP <- LL + alpha.prior + beta.prior + R.prior + rho.mu.prior +
  rho.sigma.prior + sigma.prior + sigma.mu.prior +
  sigma.sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=rmvn(nrow(mu), mu, Sigma), parm=parm)
return(Modelout)
}

```

## 24.4. Initial Values

```

Initial.Values <- c(rep(0,3), rep(0,3), upper.triangle(R, diag=TRUE),      rep(0,2),
rep(0,J), rep(1,2))

```

# 25. Discrete Choice, Conditional Logit

## 25.1. Form

$$\mathbf{y}_i \sim \mathcal{CAT}(\mathbf{p}_{i,1:J}), \quad i = 1, \dots, N, \quad j = 1, \dots, J$$

$$\mathbf{p}_{i,j} = \frac{\phi_{i,j}}{\sum_{j=1}^J \phi_{i,j}}$$

$$\phi = \exp(\mu)$$

$$\mu_{i,j} = \beta_{j,1:K} \mathbf{X}_{i,1:K} + \gamma \mathbf{Z}_{i,1:C} \in [-700, 700], \quad j = 1, \dots, (J-1)$$

$$\mu_{i,J} = \gamma \mathbf{Z}_{i,1:C}$$

$$\beta_{j,k} \sim \mathcal{N}(0, 1000), \quad j = 1, \dots, (J-1)$$

$$\gamma_c \sim \mathcal{N}(0, 1000)$$

## 25.2. Data

```

y <- x01 <- x02 <- z01 <- z02 <- c(1:300)
y[1:100] <- 1
y[101:200] <- 2
y[201:300] <- 3
x01[1:100] <- rnorm(100, 25, 2.5)
x01[101:200] <- rnorm(100, 40, 4.0)
x01[201:300] <- rnorm(100, 35, 3.5)
x02[1:100] <- rnorm(100, 2.51, 0.25)
x02[101:200] <- rnorm(100, 2.01, 0.20)
x02[201:300] <- rnorm(100, 2.70, 0.27)
z01[1:100] <- 1
z01[101:200] <- 2
z01[201:300] <- 3
z02[1:100] <- 40
z02[101:200] <- 50
z02[201:300] <- 100
N <- length(y)
J <- 3 #Number of categories in y
K <- 3 #Number of individual attributes (including the intercept)
C <- 2 #Number of choice-based attributes (intercept is not included)
X <- matrix(c(rep(1,N),x01,x02),N,K) #Design matrix of individual attrib.
Z <- matrix(c(z01,z02),N,C) #Design matrix of choice-based attributes
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=matrix(0,J-1,K), gamma=rep(0,C)))
pos.beta <- grep("beta", parm.names)
pos.gamma <- grep("gamma", parm.names)
PGF <- function(Data) return(c(rnormv((Data$J-1)*Data$K,0,10),
  rnormv(Data$C,0,10)))
MyData <- list(C=C, J=J, K=K, N=N, PGF=PGF, X=X, Z=Z, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.gamma=pos.gamma, y=y)

```

## 25.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters

```

```

beta <- matrix(parm[Data$pos.beta], Data$J-1, Data$K)
gamma <- parm[Data$pos.gamma]
### Log(Prior Densities)
beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
gamma.prior <- sum(dnormv(gamma, 0, 1000, log=TRUE))
### Log-Likelihood
mu <- matrix(rep(tcrossprod(gamma, Data$Z), Data$J), Data$N, Data$J)
mu[, -Data$J] <- mu[, -Data$J] + tcrossprod(Data$X, beta)
mu <- interval(mu, -700, 700, reflect=FALSE)
phi <- exp(mu)
p <- phi / rowSums(phi)
LL <- sum(dcat(Data$y, p, log=TRUE))
### Log-Posterior
LP <- LL + beta.prior + gamma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP, yhat=rcat(nrow(p), p),
  parm=parm)
return(Modelout)
}

```

## 25.4. Initial Values

```
Initial.Values <- c(rep(0, (J-1)*K), rep(0, C))
```

# 26. Discrete Choice, Mixed Logit

## 26.1. Form

$$\mathbf{y}_i \sim \mathcal{CAT}(\mathbf{p}_{i,1:J}), \quad i = 1, \dots, N$$

$$\mathbf{p}_{i,j} = \frac{\phi_{i,j}}{\sum_{j=1}^J \phi_{i,j}}$$

$$\phi = \exp(\mu)$$

$$\mu_{i,j} = \beta_{j,1:K} \mathbf{X}_{i,1:K} + \gamma \mathbf{Z}_{i,1:C} \in [-700, 700], \quad j = 1, \dots, (J-1)$$

$$\mu_{i,J} = \gamma \mathbf{Z}_{i,1:C}$$

$$\beta_{j,k} \sim \mathcal{N}(0, 1000), \quad j = 1, \dots, (J-1)$$

$$\gamma_c \sim \mathcal{N}(\zeta_{\mu[c]}, \zeta_{\sigma[c]}^2)$$

$$\zeta_{\mu[c]} \sim \mathcal{N}(0, 1000)$$

$$\zeta_{\sigma[c]} \sim \mathcal{HC}(25)$$

## 26.2. Data

```

y <- x01 <- x02 <- z01 <- z02 <- c(1:300)
y[1:100] <- 1
y[101:200] <- 2
y[201:300] <- 3
x01[1:100] <- rnorm(100, 25, 2.5)
x01[101:200] <- rnorm(100, 40, 4.0)
x01[201:300] <- rnorm(100, 35, 3.5)
x02[1:100] <- rnorm(100, 2.51, 0.25)
x02[101:200] <- rnorm(100, 2.01, 0.20)
x02[201:300] <- rnorm(100, 2.70, 0.27)
z01[1:100] <- 1
z01[101:200] <- 2
z01[201:300] <- 3
z02[1:100] <- 40
z02[101:200] <- 50
z02[201:300] <- 100
N <- length(y)
J <- 3 #Number of categories in y
K <- 3 #Number of individual attributes (including the intercept)
C <- 2 #Number of choice-based attributes (intercept is not included)
X <- matrix(c(rep(1,N),x01,x02),N,K) #Design matrix of individual attrib.
Z <- matrix(c(z01,z02),N,C) #Design matrix of choice-based attributes
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=matrix(0,J-1,K), gamma=rep(0,C),
  zeta.mu=rep(0,C), zeta.sigma=rep(0,C)))
pos.beta <- grep("beta", parm.names)
pos.gamma <- grep("gamma", parm.names)
pos.zeta.mu <- grep("zeta.mu", parm.names)
pos.zeta.sigma <- grep("zeta.sigma", parm.names)
PGF <- function(Data) return(c(rnormv((Data$J-1)*Data$K,0,1000),
  rnorm(Data$N*Data$C,
  matrix(rnormv(Data$C,0,1000), Data$N, Data$C, byrow=TRUE),
  matrix(rhalfcauchy(Data$C,5), Data$N, Data$C, byrow=TRUE)),
  rnormv(Data$C,0,1000), rhalfcauchy(Data$C,5)))
MyData <- list(C=C, J=J, K=K, N=N, X=X, Z=Z, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.gamma=pos.gamma,
  pos.zeta.mu=pos.zeta.mu, pos.zeta.sigma=pos.zeta.sigma, y=y)

```

## 26.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  beta <- matrix(parm[Data$pos.beta], Data$J-1, Data$K)
  gamma <- parm[Data$pos.gamma]

```



```

zeta.mu <- parm[Data$pos.zeta.mu]
zeta.sigma <- interval(parm[Data$pos.zeta.sigma], 1e-100, Inf)
parm[Data$pos.zeta.sigma] <- zeta.sigma
### Log(Prior Densities)
beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
gamma.prior <- sum(dnorm(gamma, matrix(zeta.mu, Data$N, Data$C,
    byrow=TRUE), matrix(zeta.sigma, Data$N, Data$C, byrow=TRUE),
log=TRUE))
zeta.mu.prior <- sum(dnormv(zeta.mu, 0, 1000, log=TRUE))
zeta.sigma.prior <- sum(dhalfcauchy(zeta.sigma, 25, log=TRUE))
### Log-Likelihood
mu <- matrix(rep(rowSums(gamma * Data$Z), Data$J), Data$N, Data$J)
mu[, -Data$J] <- tcrossprod(Data$X, beta) + gamma * Data$Z
mu <- interval(mu, -700, 700, reflect=FALSE)
phi <- exp(mu)
p <- phi / rowSums(phi)
LL <- sum(dcat(Data$y, p, log=TRUE))
### Log-Posterior
LP <- LL + beta.prior + gamma.prior + zeta.mu.prior + zeta.sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP, yhat=rcat(nrow(p), p),
    parm=parm)
return(Modelout)
}

```

## 26.4. Initial Values

```
Initial.Values <- c(rep(0, (J-1)*K), rep(0, N*C), rep(0, C), rep(1, C))
```

# 27. Discrete Choice, Multinomial Probit

## 27.1. Form

$$\mathbf{W}_{i,1:(J-1)} \sim \mathcal{N}_{J-1}(\mu_{i,1:(J-1)}, \Sigma), \quad i = 1, \dots, N$$

$$\mathbf{W}_{i,j} \in \begin{cases} [0,10] & \text{if } \mathbf{y}_i = j \\ [-10,0] & \end{cases}$$

$$\mu_{1:N,j} = \mathbf{X}\beta_{j,1:K} + \mathbf{Z}\gamma$$

$$\Sigma = \mathbf{U}^T \mathbf{U}$$

$$\beta_{j,k} \sim \mathcal{N}(0, 10), \quad j = 1, \dots, (J-1), \quad k = 1, \dots, K$$

$$\gamma_c \sim \mathcal{N}(0, 10), \quad c = 1, \dots, C$$

$$\mathbf{U}_{j,k} \sim \mathcal{N}(0, 1), \quad j = 1, \dots, (J-1), \quad k = 1, \dots, (J-1), \quad j \geq k, \quad j \neq k = 1$$

## 27.2. Data

```

N <- 50
J <- 5 #Categories of y
K <- 8 #Number of columns in design matrix X
C <- 2 #Number of choice-based attributes
X <- matrix(runif(N*K,-2,2), N, K)
X[,1] <- 1
beta <- matrix(runif((J-1)*K), J-1, K)
gamma <- runif(C)
Z <- matrix(runif(N*C), N, C) #Design matrix of choice-based attributes
Z[,1] <- 1
mu <- tcrossprod(X, beta) + as.vector(tcrossprod(Z, t(gamma)))
S <- diag(J-1)
u <- c(0, rnorm((J-2) + (factorial(J-1) /
  (factorial(J-1-2)*factorial(2))),0,1))
U <- diag(J-1)
U[upper.tri(U, diag=TRUE)] <- u
diag(U) <- exp(diag(U))
Sigma <- t(U) %*% U
Sigma[1,] <- Sigma[,1] <- U[1,]
mu <- tcrossprod(X, beta)
W <- rmvN(N, mu, Sigma) + matrix(rnorm(N*(J-1),0,0.1), N, J-1)
y <- max.col(cbind(W,0))
table(y)
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=matrix(0,(J-1),K), gamma=rep(0,C),
  U=U, W=matrix(0,N,J-1)), uppertri=c(0,0,1,0))
parm.names <- parm.names[-which(parm.names == "U[1,1]")]
pos.beta <- grep("beta", parm.names)
pos.gamma <- grep("gamma", parm.names)
pos.U <- grep("U", parm.names)
pos.W <- grep("W", parm.names)
PGF <- function(Data) {
  beta <- rnormv((Data$J-1)*Data$K,0,1)
  gamma <- rnormv(Data$C,0,1)
  U <- rnorm((Data$J-2) + (factorial(Data$J-1) /
    (factorial(Data$J-1-2)*factorial(2))),0,1)
  W <- matrix(runif(Data$N*(Data$J-1),-10,0), Data$N, Data$J-1)
  Y <- as.indicator.matrix(Data$y)
  W <- ifelse(Y[, -Data$J] == 1, abs(W), W)
  return(c(beta, gamma, U, as.vector(W)))}
MyData <- list(C=C, J=J, K=K, N=N, PGF=PGF, S=S, X=X, Z=Z,
  mon.names=mon.names, parm.names=parm.names, pos.beta=pos.beta,
  pos.gamma=pos.gamma, pos.U=pos.U, pos.W=pos.W, y=y)

```

### 27.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  beta <- matrix(parm[Data$pos.beta], Data$J-1, Data$K)
  gamma <- parm[Data$pos.gamma]
  u <- c(0, parm[Data$pos.U])
  U <- diag(Data$J-1)
  U[upper.tri(U, diag=TRUE)] <- u
  diag(U) <- exp(diag(U))
  Sigma <- t(U) %*% U
  Sigma[1,] <- Sigma[,1] <- U[1,]
  W <- matrix(parm[Data$pos.W], Data$N, Data$J-1)
  Y <- as.indicator.matrix(Data$y)
  temp <- which(Y[, -c(Data$J)] == 1)
  W[temp] <- interval(W[temp], 0, 10)
  temp <- which(Y[, -c(Data$J)] == 0)
  W[temp] <- interval(W[temp], -10, 0)
  parm[Data$pos.W] <- as.vector(W)
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 10, log=TRUE))
  gamma.prior <- sum(dnormv(gamma, 0, 10, log=TRUE))
  U.prior <- sum(dnorm(u[-length(u)], 0, 1, log=TRUE))
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, beta) +
    as.vector(tcrossprod(Data$Z, t(gamma)))
  #eta <- exp(cbind(mu,0))
  #p <- eta / rowSums(eta)
  LL <- sum(dmvn(W, mu, Sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + gamma.prior + U.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=max.col(cbind(rmvn(nrow(mu), mu, Sigma),0)), parm=parm)
  return(Modelout)
}

```

### 27.4. Initial Values

```
Initial.Values <- GIV(Model, MyData, PGF=TRUE)
```

## 28. Distributed Lag, Koyck

This example applies Koyck or geometric distributed lags to  $k = 1, \dots, K$  discrete events in covariate  $\mathbf{x}$ , transforming the covariate into a  $N \times K$  matrix  $\mathbf{X}$  and creates a  $N \times K$  lag matrix  $\mathbf{L}$ .

### 28.1. Form

$$\begin{aligned}
\mathbf{y} &\sim \mathcal{N}(\mu, \sigma^2) \\
\mu_t &= \alpha + \phi \mathbf{y}_{t-1} + \sum_{k=1}^K \mathbf{X}_{t,k} \beta \lambda^{\mathbf{L}[t,k]}, \quad k = 1, \dots, K, \quad t = 2, \dots, T \\
\mu_1 &= \alpha + \sum_{k=1}^K \mathbf{X}_{1,k} \beta \lambda^{\mathbf{L}[1,k]}, \quad k = 1, \dots, K \\
\alpha &\sim \mathcal{N}(0, 1000) \\
\beta &\sim \mathcal{N}(0, 1000) \\
\lambda &\sim \mathcal{U}(0, 1) \\
\phi &\sim \mathcal{N}(0, 1000) \\
\sigma &\sim \mathcal{HC}(25)
\end{aligned}$$

### 28.2. Data

```

y <- c(0.02, -0.51, -0.30, 1.46, -1.26, -2.15, -0.91, -0.53, -1.91,
      2.64, 1.64, 0.15, 1.46, 1.61, 1.96, -2.67, -0.19, -3.28,
      1.89, 0.91, -0.71, 0.74, -0.10, 3.20, -0.80, -5.25, 1.03,
      -0.40, -1.62, -0.80, 0.77, 0.17, -1.39, -1.28, 0.48, -1.02,
      0.09, -1.09, 0.86, 0.36, 1.51, -0.02, 0.47, 0.62, -1.36,
      1.12, 0.42, -4.39, -0.87, 0.05, -5.41, -7.38, -1.01, -1.70,
      0.64, 1.16, 0.87, 0.28, -1.69, -0.29, 0.13, -0.65, 0.83,
      0.62, 0.05, -0.14, 0.01, -0.36, -0.32, -0.80, -0.06, 0.24,
      0.23, -0.37, 0.00, -0.33, 0.21, -0.10, -0.10, -0.01, -0.40,
      -0.35, 0.48, -0.28, 0.08, 0.28, 0.23, 0.27, -0.35, -0.19,
      0.24, 0.17, -0.02, -0.23, 0.03, 0.02, -0.17, 0.04, -0.39,
      -0.12, 0.16, 0.17, 0.00, 0.18, 0.06, -0.36, 0.22, 0.14,
      -0.17, 0.10, -0.01, 0.00, -0.18, -0.02, 0.07, -0.06, 0.06,
      -0.05, -0.08, -0.07, 0.01, -0.06, 0.01, 0.01, -0.02, 0.01,
      0.01, 0.12, -0.03, 0.08, -0.10, 0.01, -0.03, -0.08, 0.04,
      -0.09, -0.08, 0.01, -0.05, 0.08, -0.14, 0.06, -0.11, 0.09,
      0.06, -0.12, -0.01, -0.05, -0.15, -0.05, -0.03, 0.04, 0.00,
      -0.12, 0.04, -0.06, -0.05, -0.07, -0.05, -0.14, -0.05, -0.01,
      -0.12, 0.05, 0.06, -0.10, 0.00, 0.01, 0.00, -0.08, 0.00,
      0.00, 0.07, -0.01, 0.00, 0.09, 0.33, 0.13, 0.42, 0.24,
      -0.36, 0.22, -0.09, -0.19, -0.10, -0.08, -0.07, 0.05, 0.07,
      0.07, 0.00, -0.04, -0.05, 0.03, 0.08, 0.26, 0.10, 0.08,
      0.09, -0.07, -0.33, 0.17, -0.03, 0.07, -0.04, -0.06, -0.06,
      0.07, -0.03, 0.00, 0.08, 0.27, 0.11, 0.11, 0.06, -0.11,
      -0.09, -0.21, 0.24, -0.12, 0.11, -0.02, -0.03, 0.02, -0.10,
      0.00, -0.04, 0.01, 0.02, -0.03, -0.10, -0.09, 0.17, 0.07,
      -0.05, -0.01, -0.05, 0.01, 0.00, -0.08, -0.05, -0.08, 0.07,

```

```

0.06, -0.14, 0.02, 0.01, 0.04, 0.00, -0.13, -0.17)
x <- c(0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0,
      1, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
      0, 1, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
      0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
      0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
      0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
      0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
      0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
      0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
      0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
      0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
      0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0)
T <- length(y)
K <- length(which(x != 0))
L <- X <- matrix(0, T, K)
for (i in 1:K) {
  X[which(x != 0)[i]:T,i] <- x[which(x != 0)[i]]
  L[(which(x != 0)[i]):T,i] <- 0:(T - which(x != 0)[i])}
mon.names <- "LP"
parm.names <- c("alpha","beta","lambda","phi","sigma")
PGF <- function(Data) return(c(rnormv(2,0,1000), runif(1),
  rnormv(1,0,1000), rhalfcauchy(1,5)))
MyData <- list(L=L, PGF=PGF, T=T, X=X, mon.names=mon.names,
  parm.names=parm.names, y=y)

```

### 28.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[1]; beta <- parm[2]
  parm[3] <- lambda <- interval(parm[3], 0, 1)
  phi <- parm[4]
  parm[5] <- sigma <- interval(parm[5], 1e-100, Inf)
  ### Log(Prior Densities)
  alpha.prior <- dnormv(alpha, 0, 1000, log=TRUE)
  beta.prior <- dnormv(beta, 0, 1000, log=TRUE)
  lambda.prior <- dunif(lambda, 0, 1, log=TRUE)
  phi.prior <- dnormv(phi, 0, 1000, log=TRUE)
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
  ### Log-Likelihood
  mu <- c(alpha, alpha + phi*Data$y[-Data$T]) +
    rowSums(Data$X * beta * lambda^Data$L)
  LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + alpha.prior + beta.prior + lambda.prior + phi.prior +

```

```

    sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rnorm(length(mu), mu, sigma), parm=parm)
  return(Modelout)
}

```

## 28.4. Initial Values

```
Initial.Values <- c(rep(0,2), 0.5, 0, 1)
```

# 29. Exponential Smoothing

## 29.1. Form

$$\begin{aligned}
 \mathbf{y} &\sim \mathcal{N}(\mu, \sigma^2) \\
 \mu_t &= \alpha \mathbf{y}_{t-1} + (1 - \alpha) \mu_{t-1}, \quad t = 2, \dots, T \\
 \alpha &\sim \mathcal{U}(0, 1) \\
 \sigma &\sim \mathcal{HC}
 \end{aligned}$$

## 29.2. Data

```

T <- 10
y <- rep(0,T)
y[1] <- 0
for (t in 2:T) {y[t] <- y[t-1] + rnorm(1,0,0.1)}
mon.names <- "LP"
parm.names <- c("alpha","sigma")
PGF <- function(Data) return(c(runif(1), rhalfcauchy(1,5)))
MyData <- list(PGF=PGF, T=T, mon.names=mon.names, parm.names=parm.names,
  y=y)

```

## 29.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  parm[1] <- alpha <- interval(parm[1], 0, 1)
  parm[2] <- sigma <- interval(parm[2], 1e-100, Inf)
  ### Log(Prior Densities)
  alpha.prior <- dunif(alpha, 0, 1, log=TRUE)
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
}

```

```

### Log-Likelihood
mu <- y
mu[-1] <- alpha*Data$y[-1]
mu[-1] <- mu[-1] + (1 - alpha) * mu[-Data$T]
LL <- sum(dnorm(Data$y[-1], mu[-Data$T], sigma, log=TRUE))
### Log-Posterior
LP <- LL + alpha.prior + sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=rnorm(length(mu), mu, sigma), parm=parm)
return(Modelout)
}

```

## 29.4. Initial Values

```
Initial.Values <- c(0.5, 1)
```

# 30. Factor Analysis, Approximate Dynamic

The Approximate Dynamic Factor Analysis (ADFA) model has many names, including the approximate factor model and approximate dynamic factor model. An ADFA is a Dynamic Factor Analysis (DFA) in which the factor scores of the dynamic factors are approximated with principal components. This is a combination of principal components and common factor analysis, in which the factor loadings of common factors are estimated from the data and factor scores are estimated from principal components. This is a two-stage model: principal components are estimated in the first stage and a decision is made regarding how many principal components to retain, and ADFA is estimated in the second stage. For more information on DFA, see section 32.

## 30.1. Form

$$\begin{aligned}
 \mathbf{Y}_{t,j} &\sim \mathcal{N}(\mu_{t,j}, \sigma_j^2), \quad t = 2, \dots, T, \quad j = 1, \dots, J \\
 \mu_t &= \mathbf{F}_{t-1} \Lambda \\
 \Lambda_{p,j} &\sim \mathcal{N}(0, 1000), \quad p = 1, \dots, P, \quad j = 1, \dots, J \\
 \sigma_j &\sim \mathcal{HC}(25), \quad j = 1, \dots, J
 \end{aligned}$$

## 30.2. Data

```

T <- 10 #Number of time-periods
J <- 20 #Number of variables
P <- 5 #Number of approximate dynamic factors
Lambda <- matrix(runif(J*P,-1,1), P, J)
Sigma <- matrix(runif(P*P), P, P); diag(Sigma) <- runif(P)*5
Sigma <- as.symmetric.matrix(Sigma); Sigma <- as.positive.definite(Sigma)

```

```

F <- rmvn(T, rep(0,P), Sigma)
Y <- tcrossprod(F, t(Lambda))
PCA <- prcomp(Y, scale=TRUE)
F <- PCA$x[,1:P]
mon.names <- c("LP", paste("ynew[", 1:J, "]", sep=""))
parm.names <- as.parm.names(list(Lambda=matrix(0,P,J), sigma=rep(0,J)))
pos.Lambda <- grep("Lambda", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$P*Data$J,0,1000),
  rhalfcauchy(Data$J,5)))
MyData <- list(F=F, J=J, P=P, PGF=PGF, T=T, Y=Y, mon.names=mon.names,
  parm.names=parm.names, pos.Lambda=pos.Lambda, pos.sigma=pos.sigma)

```

### 30.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  Lambda <- matrix(parm[Data$pos.Lambda], Data$P, Data$J)
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  Lambda.prior <- sum(dnormv(Lambda, 0, 1000, log=TRUE))
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
  ### Log-Likelihood
  mu <- rbind(rep(0, Data$J), tcrossprod(F[-Data$T,], t(Lambda)))
  Sigma <- matrix(sigma, Data$T, Data$J, byrow=TRUE)
  ynew <- rnorm(Data$J, tcrossprod(F[Data$T,], t(Lambda)), sigma)
  LL <- sum(dnorm(Data$Y, mu, Sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + Lambda.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,ynew),
    yhat=rnorm(prod(dim(mu)), mu, Sigma), parm=parm)
  return(Modelout)
}

```

### 30.4. Initial Values

```
Initial.Values <- c(rep(0,P*J), rep(1,J))
```

## 31. Factor Analysis, Confirmatory

Factor scores are in matrix  $\mathbf{F}$ , factor loadings for each variable are in vector  $\lambda$ , and  $\mathbf{f}$  is a vector that indicates which variable loads on which factor.



### 31.1. Form

$$\begin{aligned}
\mathbf{Y}_{i,m} &\sim \mathcal{N}(\mu_{i,m}, \sigma_m^2), \quad i = 1, \dots, N, \quad m = 1, \dots, M \\
\mu &= \alpha^T + \mathbf{F}_{1:N, \mathbf{f}} \lambda^T \\
\mathbf{F}_{i, 1:P} &\sim \mathcal{N}_P(\gamma, \Omega^{-1}), \quad i = 1, \dots, N \\
\alpha_m &\sim \mathcal{N}(0, 1000), \quad m = 1, \dots, M \\
\lambda_m &\sim \mathcal{N}(0, 1000), \quad m = 1, \dots, M \\
\sigma_m &\sim \mathcal{HC}(25), \quad m = 1, \dots, M \\
\Omega &\sim \mathcal{W}_N(\mathbf{S}), \quad \mathbf{S} = \mathbf{I}_P
\end{aligned}$$

### 31.2. Data

```

data(swiss)
Y <- cbind(swiss$Agriculture, swiss$Examination, swiss$Education,
           swiss$Catholic, swiss$Infant.Mortality)
M <- ncol(Y) #Number of variables
N <- nrow(Y) #Number of records
P <- 3 #Number of factors
f <- c(1,3,2,2,1) #Indicator f for the factor for each variable m
gamma <- rep(0,P)
S <- diag(P)
mon.names <- "LP"
parm.names <- as.parm.names(list(F=matrix(0,N,P), lambda=rep(0,M),
    U=diag(P), alpha=rep(0,M), sigma=rep(0,M)), uppertri=c(0,0,1,0,0))
pos.F <- grep("F", parm.names)
pos.lambda <- grep("lambda", parm.names)
pos.alpha <- grep("alpha", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rmvnpc(Data$N, Data$gamma,
    rwishartc(Data$N,Data$S)), rnormv(Data$M,0,1000),
    upper.triangle(rwishartc(Data$N,Data$S), diag=TRUE),
    rnormv(Data$M,0,1000), rhalfcauchy(Data$M,5)))
MyData <- list(M=M, N=N, P=P, PGF=PGF, S=S, Y=Y, f=f, gamma=gamma,
    mon.names=mon.names, parm.names=parm.names, pos.F=pos.F,
    pos.lambda=pos.lambda, pos.alpha=pos.alpha, pos.sigma=pos.sigma)

```

### 31.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[Data$pos.alpha]

```

```

lambda <- parm[Data$pos.lambda]
sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
parm[Data$pos.sigma] <- sigma
F <- matrix(parm[Data$pos.F], Data$N, Data$P)
U <- as.parm.matrix(U, Data$P, parm, Data, chol=TRUE)
diag(U) <- exp(diag(U))
### Log(Prior Densities)
alpha.prior <- sum(dnormv(alpha, 0, 1000, log=TRUE))
lambda.prior <- sum(dnormv(lambda, 0, 1000, log=TRUE))
sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
U.prior <- dwishartc(U, Data$N, Data$S, log=TRUE)
F.prior <- sum(dmvnpc(F, Data$gamma, U, log=TRUE))
### Log-Likelihood
mu <- matrix(alpha, Data$N, Data$M, byrow=TRUE) + F[,Data$f] *
  matrix(lambda, Data$N, Data$M, byrow=TRUE)
Sigma <- matrix(sigma, Data$N, Data$M, byrow=TRUE)
LL <- sum(dnorm(Data$Y, mu, Sigma, log=TRUE))
### Log-Posterior
LP <- LL + alpha.prior + lambda.prior + sigma.prior + F.prior +
  U.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=rnorm(prod(dim(mu)), mu, Sigma), parm=parm)
return(Modelout)
}

```

### 31.4. Initial Values

```

Initial.Values <- c(rep(0,N*P), rep(0,M), upper.triangle(S, diag=TRUE),
  rep(0,M), rep(1,M))

```

## 32. Factor Analysis, Dynamic

The factor scores in  $\mathbf{F}$  are dynamic with respect to time, and are estimated as in a state space model (SSM) or dynamic linear model (DLM) with constant variance in the state vector. For more information on SSMs, see section 88. For more information on exploratory factor analysis, see section 33.

### 32.1. Form

$$\begin{aligned}
\mathbf{Y}_{t,j} &\sim \mathcal{N}(\mu_{t,j}, \sigma_j^2), \quad t = 2, \dots, T, \quad j = 1, \dots, J \\
\mu_{2:T} &= \mathbf{F}_{1:(T-1)} \Lambda \\
\mathbf{F}_{1,1:P} &\sim \mathcal{N}_P(0, \Omega^{-1}) \\
\mathbf{F}_{t,1:P} &\sim \mathcal{N}_P(\mathbf{F}_{t-1,1:P}, \Omega^{-1}), \quad t = 2, \dots, T
\end{aligned}$$

$$\Lambda_{p,j} \sim \mathcal{N}(0, 1000), \quad p = 1, \dots, P, \quad j = 1, \dots, J$$

$$\Omega \sim \mathcal{W}_N(\mathbf{S}), \quad \mathbf{S} = \mathbf{I}_P$$

$$\sigma_j \sim \mathcal{HC}(25), \quad j = 1, \dots, J$$

### 32.2. Data

```

T <- 10 #Number of time-periods
J <- 20 #Number of time-series
P <- 3 #Number of dynamic factors
Lambda <- matrix(runif(J*P,-1,1), P, J)
Sigma <- matrix(runif(P*P), P, P); diag(Sigma) <- runif(P)*5
Sigma <- as.symmetric.matrix(Sigma); Sigma <- as.positive.definite(Sigma)
F <- rmvn(T, rep(0,P), Sigma)
Y <- tcrossprod(F, t(Lambda))
S <- diag(P)
mon.names <- c("LP", paste("ynew[", 1:J, "]", sep=""))
parm.names <- as.parm.names(list(F=matrix(0,T,P), U=diag(P),
    Lambda=matrix(0,P,J), sigma=rep(0,J)), uppertri=c(0,1,0,0))
pos.F <- grep("F", parm.names)
pos.Lambda <- grep("Lambda", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rmvnpc(Data$T, rep(0,Data$P),
    rwishartc(Data$P+1,Data$S)),
    upper.triangle(rwishartc(Data$P+1,Data$S), diag=TRUE),
    rnormv(Data$P*Data$J,0,1000), rhalfcauchy(Data$J,5)))
MyData <- list(J=J, P=P, PGF=PGF, S=S, T=T, Y=Y, mon.names=mon.names,
    parm.names=parm.names, pos.F=pos.F, pos.Lambda=pos.Lambda,
    pos.sigma=pos.sigma)
Dyn <- matrix(".", T, P)
for (t in 1:T) {for (p in 1:P) {
    Dyn[t,p] <- paste("F[",t,"",p,"]", sep="")}}

```

### 32.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  F <- matrix(parm[Data$pos.F], Data$T, Data$P)
  U <- as.parm.matrix(U, Data$P, parm, Data, chol=TRUE)
  diag(U) <- exp(diag(U))
  Lambda <- matrix(parm[Data$pos.Lambda], Data$P, Data$J)
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  F.prior <- sum(dmvnpc(F, rbind(rep(0, Data$P), F[-Data$T,]), U,

```

```

log=TRUE))
U.prior <- dwishartc(U, Data$P+1, Data$S, log=TRUE)
Lambda.prior <- sum(dnormv(Lambda, 0, 1000, log=TRUE))
sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
### Log-Likelihood
mu <- rbind(rep(0, Data$J), tcrossprod(F[-Data$T,], t(Lambda)))
Sigma <- matrix(sigma, Data$T, Data$J, byrow=TRUE)
ynew <- rnorm(Data$J, tcrossprod(F[Data$T,], t(Lambda)), sigma)
LL <- sum(dnorm(Data$Y, mu, Sigma, log=TRUE))
### Log-Posterior
LP <- LL + F.prior + U.prior + Lambda.prior + sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,ynew),
  yhat=rnorm(prod(dim(mu)), mu, Sigma), parm=parm)
return(Modelout)
}

```

### 32.4. Initial Values

```

Initial.Values <- c(rep(0,T*P), S[upper.tri(S, diag=TRUE)], rep(0,P*J),
  rep(1,J))

```

## 33. Factor Analysis, Exploratory

Factor scores are in matrix  $\mathbf{F}$  and factor loadings are in matrix  $\Lambda$ .

### 33.1. Form

$$\begin{aligned}
\mathbf{Y}_{i,m} &\sim \mathcal{N}(\mu_{i,m}, \sigma_m^2), \quad i = 1, \dots, N, \quad m = 1, \dots, M \\
\mu &= \alpha^T + \mathbf{F}\Lambda \\
\mathbf{F}_{i,1:P} &\sim \mathcal{N}_P(\gamma, \Omega^{-1}), \quad i = 1, \dots, N \\
\alpha_m &\sim \mathcal{N}(0, 1000), \quad m = 1, \dots, M \\
\gamma_p &= 0, \quad p = 1, \dots, P \\
\Lambda_{p,m} &\sim \mathcal{N}(0, 1000), \quad p = 1, \dots, P, \quad m = 1, \dots, M \\
\Omega &\sim \mathcal{W}_N(\mathbf{S}), \quad \mathbf{S} = \mathbf{I}_P \\
\sigma_m &\sim \mathcal{HC}(25), \quad m = 1, \dots, M
\end{aligned}$$

### 33.2. Data

```

M <- 10 #Number of variables
N <- 20 #Number of records

```

```

P <- 3 #Number of factors
alpha <- runif(M)*10
Lambda <- matrix(runif(M*P,-1,1), P, M)
U <- diag(P) U[upper.tri(U, diag=TRUE)] <- runif(length(upper.triangle(U, diag=TRUE)))
F <- rmvnc(N, rep(0,P), U)
Y <- matrix(alpha, N, M, byrow=TRUE) + tcrossprod(F, t(Lambda))
gamma <- rep(0,P)
S <- diag(P)
mon.names <- "LP"
parm.names <- as.parm.names(list(F=matrix(0,N,P), Lambda=matrix(0,P,M),
  U=diag(P), alpha=rep(0,M), sigma=rep(0,M)), uppertri=c(0,0,1,0,0))
pos.F <- grep("F", parm.names)
pos.Lambda <- grep("Lambda", parm.names)
pos.alpha <- grep("alpha", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rmvnnc(Data$N, Data$gamma,
  rwishartc(Data$N, Data$S)), rnormv(Data$P*Data$M,0,1000),
  upper.triangle(rwishartc(Data$N, Data$S), diag=TRUE),
  rnormv(Data$M,0,1000), rhalfcauchy(Data$M,5)))
MyData <- list(M=M, N=N, P=P, PGF=PGF, S=S, Y=Y, gamma=gamma,
  mon.names=mon.names, parm.names=parm.names, pos.F=pos.F,
  pos.Lambda=pos.Lambda, pos.alpha=pos.alpha, pos.sigma=pos.sigma)

```

### 33.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  F <- matrix(parm[Data$pos.F], Data$N, Data$P)
  Lambda <- matrix(parm[Data$pos.Lambda], Data$P, Data$M)
  U <- as.parm.matrix(U, Data$P, parm, Data, chol=TRUE)
  diag(U) <- exp(diag(U))
  alpha <- parm[Data$pos.alpha]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  F.prior <- sum(dmvnnc(F, Data$gamma, U, log=TRUE))
  Lambda.prior <- sum(dnormv(Lambda, 0, 1000, log=TRUE))
  U.prior <- dwishartc(U, Data$N, Data$S, log=TRUE)
  alpha.prior <- sum(dnormv(alpha, 0, 1000, log=TRUE))
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
  ### Log-Likelihood
  mu <- matrix(alpha, Data$N, Data$M, byrow=TRUE) +
    tcrossprod(F, t(Lambda))
  Sigma <- matrix(sigma, Data$N, Data$M, byrow=TRUE)
  LL <- sum(dnorm(Data$Y, mu, Sigma, log=TRUE))
}

```

```

### Log-Posterior
LP <- LL + F.prior + Lambda.prior + U.prior + alpha.prior +
  sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=rnorm(prod(dim(mu)), mu, Sigma), parm=parm)
return(Modelout)
}

```

### 33.4. Initial Values

```

Initial.Values <- c(rep(0,N*P), rep(0,P*M), upper.triangle(S, diag=TRUE),
  rep(0,M), rep(1,M))

```

## 34. Factor Analysis, Exploratory Ordinal

This exploratory ordinal factor analysis (EOFA) model form is also suitable for collaborative filtering, and automatically handles missing values.

### 34.1. Form

$$\begin{aligned}
\mathbf{Y}_{i,m} &\sim \mathcal{CAT}(\mathbf{P}_{i,m,1:K}), \quad i = 1, \dots, N, \quad m = 1, \dots, M \\
\mathbf{P}_{:,K} &= 1 - \mathbf{Q}_{:,(K-1)} \\
\mathbf{P}_{:,k} &= |\mathbf{Q}_{:,k} - \mathbf{Q}_{:,(k-1)}|, \quad k = 2, \dots, (K-1) \\
\mathbf{P}_{:,1} &= \mathbf{Q}_{:,1} \\
\mathbf{Q} &= \phi(\boldsymbol{\mu}) \\
\boldsymbol{\mu}_{:,k} &= \boldsymbol{\alpha}_k - \mathbf{F}\boldsymbol{\Lambda}, \quad k = 1, \dots, (K-1) \\
\mathbf{F}_{i,1:P} &\sim \mathcal{N}_P(\boldsymbol{\gamma}, \boldsymbol{\Omega}^{-1}), \quad i = 1, \dots, N \\
\boldsymbol{\gamma}_p &= 0, \quad p = 1, \dots, P \\
\boldsymbol{\Lambda}_{p,m} &\sim \mathcal{N}(0, 1000), \quad p = 1, \dots, P, \quad m = 1, \dots, M \\
\boldsymbol{\Omega} &\sim \mathcal{W}_N(\mathbf{S}), \quad \mathbf{S} = \mathbf{I}_P \\
\alpha_k &\sim \mathcal{N}(0, 1) \in [(k-1), k] \in [-5, 5], \quad k = 2, \dots, (K-1)
\end{aligned}$$

### 34.2. Data

```

M <- 10 #Number of variables
N <- 20 #Number of records
K <- 3 #Number of discrete values
P <- 3 #Number of factors
alpha <- sort(rnorm(K-1))
Lambda <- matrix(runif(M*P, -1, 1), P, M)

```

```

U <- diag(P)
U[upper.tri(U, diag=TRUE)] <- runif(length(upper.triangle(U, diag=TRUE)))
F <- rmvnc(N, rep(0,P), U)
mu <- aperm(array(alpha, dim=c(K-1, M, N)), perm=c(3,2,1))
mu <- mu - array(tcrossprod(F, t(Lambda)), dim=c(N, M, K-1))
Pr <- Q <- pnorm(mu)
Pr[, , -1] <- abs(Q[, , -1] - Q[, , -(K-1)])
Pr <- array(Pr, dim=c(N, M, K))
Pr[, , K] <- 1 - Q[, , (K-1)]
dim(Pr) <- c(N*M, K)
Y <- matrix(rcat(nrow(Pr), Pr), N, M) #Make sure Y has all values
gamma <- rep(0,P)
S <- diag(P)
mon.names <- "LP"
parm.names <- as.parm.names(list(F=matrix(0,N,P), Lambda=matrix(0,P,M),
  U=diag(P), alpha=rep(0,K-1)), uppertri=c(0,0,1,0))
pos.F <- grep("F", parm.names)
pos.Lambda <- grep("Lambda", parm.names)
pos.alpha <- grep("alpha", parm.names)
PGF <- function(Data) return(c(rmvnnc(Data$N, Data$gamma,
  rwishartc(Data$N, Data$S)), rnorm(Data$P*Data$M),
  upper.triangle(rwishartc(Data$N, Data$S), diag=TRUE),
  sort(rnorm(Data$K-1))))
MyData <- list(K=K, M=M, N=N, P=P, PGF=PGF, S=S, Y=Y, gamma=gamma,
  mon.names=mon.names, parm.names=parm.names, pos.F=pos.F,
  pos.Lambda=pos.Lambda, pos.alpha=pos.alpha)

```

### 34.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  F <- matrix(parm[Data$pos.F], Data$N, Data$P)
  Lambda <- matrix(parm[Data$pos.Lambda], Data$P, Data$M)
  U <- as.parm.matrix(U, Data$P, parm, Data, chol=TRUE)
  diag(U) <- exp(diag(U))
  alpha <- sort(interval(parm[Data$pos.alpha], -5, 5))
  parm[Data$pos.alpha] <- alpha
  ### Log(Prior Densities)
  F.prior <- sum(dmvnnc(F, Data$gamma, U, log=TRUE))
  Lambda.prior <- sum(dnormv(Lambda, 0, 1000, log=TRUE))
  U.prior <- dwishartc(U, Data$N, Data$S, log=TRUE)
  alpha.prior <- sum(dnormv(alpha, 0, 1000, log=TRUE))
  ### Log-Likelihood
  mu <- aperm(array(alpha, dim=c(Data$K-1, Data$M, Data$N)),
    perm=c(3,2,1))

```

```

mu <- mu - array(tcrossprod(F, t(Lambda)),
  dim=c(Data$N, Data$M, Data$K-1))
P <- Q <- pnorm(mu)
P[ , , -1] <- abs(Q[ , , -1] - Q[ , , -(Data$K-1)])
P <- array(P, dim=c(Data$N, Data$M, Data$K))
P[ , , Data$K] <- 1 - Q[ , , (Data$K-1)]
y <- as.vector(Data$Y)
dim(P) <- c(Data$N*Data$M, Data$K)
LL <- sum(dcat(y, P, log=TRUE))
### Log-Posterior
LP <- LL + F.prior + Lambda.prior + U.prior + alpha.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=matrix(rcat(nrow(P), P), Data$N, Data$M), parm=parm)
return(Modelout)
}

```

### 34.4. Initial Values

```

Initial.Values <- c(rep(0,N*P), rep(0,P*M), upper.triangle(S, diag=TRUE),
  seq(from=-1, to=1, len=K-1))

```

## 35. Factor Regression

This example of factor regression is constrained to the case where the number of factors is equal to the number of independent variables (IVs) less the intercept. The purpose of this form of factor regression is to orthogonalize the IVs with respect to  $\mathbf{y}$ , rather than variable reduction. This method is the combination of confirmatory factor analysis in section 31 and linear regression in section 46.

### 35.1. Form

$$\begin{aligned}
 \mathbf{y} &\sim \mathcal{N}(\nu, \sigma_{J+1}^2) \\
 \nu &= \mu\beta \\
 \mu_{i,1} &= 1 \\
 \mu_{i,j+1} &= \mu_{i,j}, \quad j = 1, \dots, J \\
 \mathbf{X}_{i,j} &\sim \mathcal{N}(\mu_{i,j}, \sigma_j^2), \quad i = 1, \dots, N, \quad j = 2, \dots, J \\
 \mu_{i,j} &= \alpha_j + \lambda_j \mathbf{F}_{i,j}, \quad i = 1, \dots, N, \quad j = 2, \dots, J \\
 \mathbf{F}_{i,1:J} &\sim \mathcal{N}_{J-1}(0, \Omega^{-1}), \quad i = 1, \dots, N \\
 \alpha_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, (J-1) \\
 \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\
 \lambda_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, (J-1)
 \end{aligned}$$



$$\sigma_j \sim \mathcal{HC}(25), \quad j = 1, \dots, (J+1)$$

$$\Omega \sim \mathcal{W}_N(\mathbf{S}), \quad \mathbf{S} = \mathbf{I}_J$$

### 35.2. Data

```
data(demonsnacks)
N <- nrow(demonsnacks)
y <- log(demonsnacks$Calories)
X <- as.matrix(log(demonsnacks[,c(1,4,10)]+1))
J <- ncol(X)
for (j in 1:J) {X[,j] <- CenterScale(X[,j])}
S <- diag(J)
mon.names <- "LP"
parm.names <- as.parm.names(list(alpha=rep(0,J), beta=rep(0,J+1),
  lambda=rep(0,J), sigma=rep(0,J+1), F=matrix(0,N,J), U=diag(J)),
  uppertri=c(0,0,0,0,0,1))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
pos.lambda <- grep("lambda", parm.names)
pos.sigma <- grep("sigma", parm.names)
pos.F <- grep("F", parm.names)
PGF <- function(Data) return(c(rnormmv(Data$J,0,1000),
  rnormmv(Data$J+1,0,1000), rnormmv(Data$J,0,1000),
  rhalfcauchy(Data$J+1,5), rmvnpnc(Data$N, rep(0,Data$J), Data$S),
  diag=TRUE)))
MyData <- list(J=J, N=N, PGF=PGF, S=S, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.alpha=pos.alpha, pos.beta=pos.beta,
  pos.lambda=pos.lambda, pos.sigma=pos.sigma, pos.F=pos.F, y=y)
```

### 35.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[Data$pos.alpha]
  beta <- parm[Data$pos.beta]
  lambda <- parm[Data$pos.lambda]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  F <- matrix(Data$pos.F, Data$N, Data$J)
  U <- as.parm.matrix(U, Data$J, parm, Data, chol=TRUE)
  diag(U) <- exp(diag(U))
  ### Log(Prior Densities)
  alpha.prior <- sum(dnormmv(alpha, 0, 1000, log=TRUE))
  beta.prior <- sum(dnormmv(beta, 0, 1000, log=TRUE))
```

```

lambda.prior <- sum(dnormv(lambda, 0, 1000, log=TRUE))
sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
F.prior <- sum(dmvnnc(F, rep(0,Data$J), U, log=TRUE))
U.prior <- dwishartc(U, Data$J+1, Data$S, log=TRUE)
### Log-Likelihood
mu <- matrix(alpha, Data$N, Data$J, byrow=TRUE) + F *
  matrix(lambda, Data$N, Data$J, byrow=TRUE)
nu <- tcrossprod(beta, cbind(rep(1,Data$N),mu))
LL <- sum(dnorm(Data$X, mu, matrix(sigma[1:Data$J], Data$N, Data$J,
  byrow=TRUE)), dnorm(Data$y, nu, sigma[Data$J+1], log=TRUE))
### Log-Posterior
LP <- LL + alpha.prior + beta.prior + lambda.prior + sigma.prior +
  F.prior + U.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=rnorm(Data$N, nu, sigma[Data$J+1]), parm=parm)
return(Modelout)
}

```

### 35.4. Initial Values

```

Initial.Values <- c(rep(0,J), rep(0,J+1), rep(0,J), rep(0,J+1),      rep(0,N*J),
S[upper.tri(S, diag=TRUE)])

```

## 36. Gamma Regression

### 36.1. Form

$$\begin{aligned}
 \mathbf{y} &\sim \mathcal{G}(\lambda\tau, \tau) \\
 \lambda &= \exp(\mathbf{X}\beta) \\
 \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\
 \tau &\sim \mathcal{HC}(25)
 \end{aligned}$$

### 36.2. Data

```

N <- 20
J <- 3
X <- matrix(runif(N*J,-2,2),N,J); X[,1] <- 1
beta <- runif(J,-2,2)
y <- round(exp(tcrossprod(X, t(beta)))) + 0.1 #Must be > 0
mon.names <- c("LP","sigma2")
parm.names <- as.parm.names(list(beta=rep(0,J), tau=0))
pos.beta <- grep("beta", parm.names)

```

```
pos.tau <- grep("tau", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,1000), rhalfcauchy(1,5)))
MyData <- list(J=J, N=N, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.tau=pos.tau, y=y)
```

### 36.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  tau <- interval(parm[Data$pos.tau], 1e-100, Inf)
  parm[Data$pos.tau] <- tau
  sigma2 <- 1/tau
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  tau.prior <- dhalfcauchy(tau, 25, log=TRUE)
  ### Log-Likelihood
  lambda <- exp(tcrossprod(Data$X, t(beta)))
  LL <- sum(dgamma(Data$y, tau*lambda, tau, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + tau.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,sigma2),
    yhat=rgamma(nrow(lambda), tau*lambda, tau), parm=parm)
  return(Modelout)
}
```

### 36.4. Initial Values

```
Initial.Values <- c(rep(0,J), 1)
```

## 37. GARCH(1,1)

### 37.1. Form

$$\begin{aligned}
 \mathbf{y}_t &\sim \mathcal{N}(\mu_t, \sigma_t^2), \quad t = 1, \dots, T \\
 \mathbf{y}^{new} &\sim \mathcal{N}(\mu_{T+1}, \sigma_{new}^2) \\
 \mu_t &= \alpha + \phi \mathbf{y}_{t-1}, \quad t = 1, \dots, (T+1) \\
 \epsilon_t &= \mathbf{y}_t - \mu_t \\
 \alpha &\sim \mathcal{N}(0, 1000) \\
 \phi &\sim \mathcal{N}(0, 1000)
 \end{aligned}$$

$$\sigma_{new}^2 = \theta_1 + \theta_2 \epsilon_T^2 + \theta_3 \sigma_T^2$$

$$\sigma_t^2 = \theta_1 + \theta_2 \epsilon_{t-1}^2 + \theta_3 \sigma_{t-1}^2$$

$$\theta_k = \frac{1}{1 + \exp(-\theta_k)}, \quad k = 1, \dots, 3$$

$$\theta_k \sim \mathcal{N}(0, 1000) \in [-10, 10], \quad k = 1, \dots, 3$$

### 37.2. Data

```

y <- c(0.02, -0.51, -0.30, 1.46, -1.26, -2.15, -0.91, -0.53, -1.91,
      2.64, 1.64, 0.15, 1.46, 1.61, 1.96, -2.67, -0.19, -3.28,
      1.89, 0.91, -0.71, 0.74, -0.10, 3.20, -0.80, -5.25, 1.03,
      -0.40, -1.62, -0.80, 0.77, 0.17, -1.39, -1.28, 0.48, -1.02,
      0.09, -1.09, 0.86, 0.36, 1.51, -0.02, 0.47, 0.62, -1.36,
      1.12, 0.42, -4.39, -0.87, 0.05, -5.41, -7.38, -1.01, -1.70,
      0.64, 1.16, 0.87, 0.28, -1.69, -0.29, 0.13, -0.65, 0.83,
      0.62, 0.05, -0.14, 0.01, -0.36, -0.32, -0.80, -0.06, 0.24,
      0.23, -0.37, 0.00, -0.33, 0.21, -0.10, -0.10, -0.01, -0.40,
      -0.35, 0.48, -0.28, 0.08, 0.28, 0.23, 0.27, -0.35, -0.19,
      0.24, 0.17, -0.02, -0.23, 0.03, 0.02, -0.17, 0.04, -0.39,
      -0.12, 0.16, 0.17, 0.00, 0.18, 0.06, -0.36, 0.22, 0.14,
      -0.17, 0.10, -0.01, 0.00, -0.18, -0.02, 0.07, -0.06, 0.06,
      -0.05, -0.08, -0.07, 0.01, -0.06, 0.01, 0.01, -0.02, 0.01,
      0.01, 0.12, -0.03, 0.08, -0.10, 0.01, -0.03, -0.08, 0.04,
      -0.09, -0.08, 0.01, -0.05, 0.08, -0.14, 0.06, -0.11, 0.09,
      0.06, -0.12, -0.01, -0.05, -0.15, -0.05, -0.03, 0.04, 0.00,
      -0.12, 0.04, -0.06, -0.05, -0.07, -0.05, -0.14, -0.05, -0.01,
      -0.12, 0.05, 0.06, -0.10, 0.00, 0.01, 0.00, -0.08, 0.00,
      0.00, 0.07, -0.01, 0.00, 0.09, 0.33, 0.13, 0.42, 0.24,
      -0.36, 0.22, -0.09, -0.19, -0.10, -0.08, -0.07, 0.05, 0.07,
      0.07, 0.00, -0.04, -0.05, 0.03, 0.08, 0.26, 0.10, 0.08,
      0.09, -0.07, -0.33, 0.17, -0.03, 0.07, -0.04, -0.06, -0.06,
      0.07, -0.03, 0.00, 0.08, 0.27, 0.11, 0.11, 0.06, -0.11,
      -0.09, -0.21, 0.24, -0.12, 0.11, -0.02, -0.03, 0.02, -0.10,
      0.00, -0.04, 0.01, 0.02, -0.03, -0.10, -0.09, 0.17, 0.07,
      -0.05, -0.01, -0.05, 0.01, 0.00, -0.08, -0.05, -0.08, 0.07,
      0.06, -0.14, 0.02, 0.01, 0.04, 0.00, -0.13, -0.17)
T <- length(y)
mon.names <- c("LP", "ynew", "sigma2.new")
parm.names <- c("alpha", "phi", "omega", "theta[1]", "theta[2]")
PGF <- function(Data) return(c(rnormv(2,0,1000), rhalfcauchy(1,5),
  runif(2)))
MyData <- list(PGF=PGF, T=T, mon.names=mon.names, parm.names=parm.names,
  y=y)

```

### 37.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[1]; phi <- parm[2]
  parm[3] <- omega <- interval(parm[3], 1e-100, Inf)
  theta <- interval(parm[4:5], 1e-5, 1-1e-5)
  if(sum(theta) >= 1) theta[2] <- 1 - 1e-5 - theta[1]
  parm[4:5] <- theta      ### Log(Prior Densities)
  alpha.prior <- dnormv(alpha, 0, 1000, log=TRUE)
  phi.prior <- dnormv(phi, 0, 1000, log=TRUE)
  omega.prior <- dhalfcauchy(omega, 25, log=TRUE)
  theta.prior <- sum(dunif(theta, 0, 1, log=TRUE))
  ### Log-Likelihood
  mu <- c(alpha, alpha + phi*Data$y[-Data$T])
  epsilon <- Data$y - mu
  sigma2 <- c(omega, omega + theta[1]*epsilon[-Data$T]^2)
  sigma2[-1] <- sigma2[-1] + theta[2]*sigma2[-Data$T]
  sigma2.new <- omega + theta[1]*epsilon[Data$T]^2 +
    theta[2]*sigma2[Data$T]
  ynew <- rnormv(1, alpha + phi*Data$y[Data$T], sigma2.new)
  LL <- sum(dnormv(Data$y, mu, sigma2, log=TRUE))
  ### Log-Posterior
  LP <- LL + alpha.prior + phi.prior + omega.prior + theta.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP, ynew, sigma2.new),
    yhat=rnormv(length(mu), mu, sigma2), parm=parm)
  return(Modelout)
}
```

### 37.4. Initial Values

```
Initial.Values <- c(rep(0,2), rep(0.4,3))
```

## 38. GARCH-M(1,1)

### 38.1. Form

$$\begin{aligned}
 \mathbf{y}_t &\sim \mathcal{N}(\mu_t, \sigma_t^2), \quad t = 1, \dots, T \\
 \mathbf{y}^{new} &\sim \mathcal{N}(\mu_{T+1}, \sigma_{new}^2) \\
 \mu_t &= \alpha + \phi \mathbf{y}_{t-1} + \delta \sigma_{t-1}^2, \quad t = 1, \dots, (T+1) \\
 \epsilon_t &= \mathbf{y}_t - \mu_t \\
 \alpha &\sim \mathcal{N}(0, 1000)
 \end{aligned}$$

$$\phi \sim \mathcal{N}(0, 1000)$$

$$\sigma_{new}^2 = \omega + \theta_1 \epsilon_T^2 + \theta_2 \sigma_T^2$$

$$\sigma_t^2 = \omega + \theta_1 \epsilon_{t-1}^2 + \theta_2 \sigma_{t-1}^2$$

$$\omega \sim \mathcal{HC}(25)$$

$$\theta_k \sim \mathcal{U}(0, 1), \quad k = 1, \dots, 2$$

### 38.2. Data

```

y <- c(0.02, -0.51, -0.30, 1.46, -1.26, -2.15, -0.91, -0.53, -1.91,
      2.64, 1.64, 0.15, 1.46, 1.61, 1.96, -2.67, -0.19, -3.28,
      1.89, 0.91, -0.71, 0.74, -0.10, 3.20, -0.80, -5.25, 1.03,
      -0.40, -1.62, -0.80, 0.77, 0.17, -1.39, -1.28, 0.48, -1.02,
      0.09, -1.09, 0.86, 0.36, 1.51, -0.02, 0.47, 0.62, -1.36,
      1.12, 0.42, -4.39, -0.87, 0.05, -5.41, -7.38, -1.01, -1.70,
      0.64, 1.16, 0.87, 0.28, -1.69, -0.29, 0.13, -0.65, 0.83,
      0.62, 0.05, -0.14, 0.01, -0.36, -0.32, -0.80, -0.06, 0.24,
      0.23, -0.37, 0.00, -0.33, 0.21, -0.10, -0.10, -0.01, -0.40,
      -0.35, 0.48, -0.28, 0.08, 0.28, 0.23, 0.27, -0.35, -0.19,
      0.24, 0.17, -0.02, -0.23, 0.03, 0.02, -0.17, 0.04, -0.39,
      -0.12, 0.16, 0.17, 0.00, 0.18, 0.06, -0.36, 0.22, 0.14,
      -0.17, 0.10, -0.01, 0.00, -0.18, -0.02, 0.07, -0.06, 0.06,
      -0.05, -0.08, -0.07, 0.01, -0.06, 0.01, 0.01, -0.02, 0.01,
      0.01, 0.12, -0.03, 0.08, -0.10, 0.01, -0.03, -0.08, 0.04,
      -0.09, -0.08, 0.01, -0.05, 0.08, -0.14, 0.06, -0.11, 0.09,
      0.06, -0.12, -0.01, -0.05, -0.15, -0.05, -0.03, 0.04, 0.00,
      -0.12, 0.04, -0.06, -0.05, -0.07, -0.05, -0.14, -0.05, -0.01,
      -0.12, 0.05, 0.06, -0.10, 0.00, 0.01, 0.00, -0.08, 0.00,
      0.00, 0.07, -0.01, 0.00, 0.09, 0.33, 0.13, 0.42, 0.24,
      -0.36, 0.22, -0.09, -0.19, -0.10, -0.08, -0.07, 0.05, 0.07,
      0.07, 0.00, -0.04, -0.05, 0.03, 0.08, 0.26, 0.10, 0.08,
      0.09, -0.07, -0.33, 0.17, -0.03, 0.07, -0.04, -0.06, -0.06,
      0.07, -0.03, 0.00, 0.08, 0.27, 0.11, 0.11, 0.06, -0.11,
      -0.09, -0.21, 0.24, -0.12, 0.11, -0.02, -0.03, 0.02, -0.10,
      0.00, -0.04, 0.01, 0.02, -0.03, -0.10, -0.09, 0.17, 0.07,
      -0.05, -0.01, -0.05, 0.01, 0.00, -0.08, -0.05, -0.08, 0.07,
      0.06, -0.14, 0.02, 0.01, 0.04, 0.00, -0.13, -0.17)
T <- length(y)
mon.names <- c("LP", "ynew", "sigma2.new")
parm.names <- c("alpha", "phi", "delta", "omega", "theta[1]", "theta[2]")
PGF <- function(Data) return(c(rnormv(3, 0, 1000), rhalfcauchy(1, 5),
  runif(2)))
MyData <- list(PGF=PGF, T=T, mon.names=mon.names, parm.names=parm.names,
  y=y)

```

### 38.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[1]; phi <- parm[2]; delta <- parm[3]
  parm[4] <- omega <- interval(parm[4], 1e-100, Inf)
  parm[5:6] <- theta <- interval(parm[5:6], 1e-10, 1-1e-5)
  ### Log(Prior Densities)
  alpha.prior <- dnormv(alpha, 0, 1000, log=TRUE)
  phi.prior <- dnormv(phi, 0, 1000, log=TRUE)
  delta.prior <- dnormv(delta, 0, 1000, log=TRUE)
  omega.prior <- dhalfcauchy(omega, 25, log=TRUE)
  theta.prior <- sum(dunif(theta, 0, 1, log=TRUE))
  ### Log-Likelihood
  mu <- c(alpha, alpha + phi*Data$y[-Data$T])
  epsilon <- Data$y - mu
  sigma2 <- c(omega, omega + theta[1]*epsilon[-Data$T]^2)
  sigma2[-1] <- sigma2[-1] + theta[2]*sigma2[-Data$T]
  sigma2.new <- omega + theta[1]*epsilon[Data$T]^2 +
    theta[2]*sigma2[Data$T]
  mu <- mu + delta*sigma2
  ynew <- rnormv(1, alpha + phi*Data$y[Data$T] + delta*sigma2[Data$T],
    sigma2.new)
  LL <- sum(dnormv(Data$y, mu, sigma2, log=TRUE))
  ### Log-Posterior
  LP <- LL + alpha.prior + phi.prior + delta.prior + omega.prior +
    theta.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP, ynew, sigma2.new),
    yhat=rnormv(length(mu), mu, sigma2), parm=parm)
  return(Modelout)
}
```

### 38.4. Initial Values

```
Initial.Values <- c(rep(0,3), rep(0.3,3))
```

## 39. Geographically Weighted Regression

### 39.1. Form

$$\mathbf{y}_{i,k} \sim \mathcal{N}(\mu_{i,k}, \tau_{i,k}^{-1}), \quad i = 1, \dots, N, \quad k = 1, \dots, N$$

$$\mu_{i,1:N} = \mathbf{X}\beta_{i,1:J}$$

$$\begin{aligned}
\tau &= \frac{1}{\sigma^2} \mathbf{w} \nu \\
\mathbf{w} &= \frac{\exp(-0.5 \mathbf{Z}^2)}{\mathbf{h}} \\
\alpha &\sim \mathcal{U}(1.5, 100) \\
\beta_{i,j} &\sim \mathcal{N}(0, 1000), \quad i = 1, \dots, N, \quad j = 1, \dots, J \\
\mathbf{h} &\sim \mathcal{N}(0.1, 1000) \in [0.1, \infty] \\
\nu_{i,k} &\sim \mathcal{G}(\alpha, 2), \quad i = 1, \dots, N, \quad k = 1, \dots, N \\
\sigma_i &\sim \mathcal{HC}(25), \quad i = 1, \dots, N
\end{aligned}$$

### 39.2. Data

```

crime <- c(18.802, 32.388, 38.426, 0.178, 15.726, 30.627, 50.732,
          26.067, 48.585, 34.001, 36.869, 20.049, 19.146, 18.905, 27.823,
          16.241, 0.224, 30.516, 33.705, 40.970, 52.794, 41.968, 39.175,
          53.711, 25.962, 22.541, 26.645, 29.028, 36.664, 42.445, 56.920,
          61.299, 60.750, 68.892, 38.298, 54.839, 56.706, 62.275, 46.716,
          57.066, 54.522, 43.962, 40.074, 23.974, 17.677, 14.306, 19.101,
          16.531, 16.492)
income <- c(21.232, 4.477, 11.337, 8.438, 19.531, 15.956, 11.252,
           16.029, 9.873, 13.598, 9.798, 21.155, 18.942, 22.207, 18.950,
           29.833, 31.070, 17.586, 11.709, 8.085, 10.822, 9.918, 12.814,
           11.107, 16.961, 18.796, 11.813, 14.135, 13.380, 17.017, 7.856,
           8.461, 8.681, 13.906, 14.236, 7.625, 10.048, 7.467, 9.549,
           9.963, 11.618, 13.185, 10.655, 14.948, 16.940, 18.739, 18.477,
           18.324, 25.873)
housing <- c(44.567, 33.200, 37.125, 75.000, 80.467, 26.350, 23.225,
            28.750, 18.000, 96.400, 41.750, 47.733, 40.300, 42.100, 42.500,
            61.950, 81.267, 52.600, 30.450, 20.300, 34.100, 23.600, 27.000,
            22.700, 33.500, 35.800, 26.800, 27.733, 25.700, 43.300, 22.850,
            17.900, 32.500, 22.500, 53.200, 18.800, 19.900, 19.700, 41.700,
            42.900, 30.600, 60.000, 19.975, 28.450, 31.800, 36.300, 39.600,
            76.100, 44.333)
easting <- c(35.62, 36.50, 36.71, 33.36, 38.80, 39.82, 40.01, 43.75,
            39.61, 47.61, 48.58, 49.61, 50.11, 51.24, 50.89, 48.44, 46.73,
            43.44, 43.37, 41.13, 43.95, 44.10, 43.70, 41.04, 43.23, 42.67,
            41.21, 39.32, 41.09, 38.3, 41.31, 39.36, 39.72, 38.29, 36.60,
            37.60, 37.13, 37.85, 35.95, 35.72, 35.76, 36.15, 34.08, 30.32,
            27.94, 27.27, 24.25, 25.47, 29.02)
northing <- c(42.38, 40.52, 38.71, 38.41, 44.07, 41.18, 38.00, 39.28,
             34.91, 36.42, 34.46, 32.65, 29.91, 27.80, 25.24, 27.93, 31.91,
             35.92, 33.46, 33.14, 31.61, 30.40, 29.18, 28.78, 27.31, 24.96,
             25.90, 25.85, 27.49, 28.82, 30.90, 32.88, 30.64, 30.35, 32.09,
             34.08, 36.12, 36.30, 36.40, 35.60, 34.66, 33.92, 30.42, 28.26,

```



```

      29.85, 28.21, 26.69, 25.71, 26.58)
N <- length(crime)
J <- 3 #Number of predictors, including the intercept
X <- matrix(c(rep(1,N), income, housing),N,J)
D <- as.matrix(dist(cbind(northing,easting), diag=TRUE, upper=TRUE))
Z <- D / sd(as.vector(D))
y <- matrix(0,N,N); for (i in 1:N) {for (k in 1:N) {y[i,k] <- crime[k]}}
mon.names <- "LP"
parm.names <- as.parm.names(list(alpha=0, beta=matrix(0,N,J), H=0,
  nu=matrix(0,N,N), sigma=rep(0,N)))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
pos.H <- grep("H", parm.names)
pos.nu <- grep("nu", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(runif(1,1.5,100),
  rnormmv(Data$N*Data$J,0,1000), runif(1, 0.1, 1000),
  rgamma(Data$N*Data$N,runif(1,1.5,100),2), rhalfcauchy(Data$N,5)))
MyData <- list(J=J, N=N, PGF=PGF, X=X, Z=Z, latitude=northing,
  longitude=easting, mon.names=mon.names, parm.names=parm.names,
  pos.alpha=pos.alpha, pos.beta=pos.beta, pos.H=pos.H, pos.nu=pos.nu,
  pos.sigma=pos.sigma, y=y)

```

### 39.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  alpha <- interval(parm[Data$pos.alpha], 1.5, 100)
  parm[Data$pos.alpha] <- alpha
  beta <- matrix(parm[Data$pos.beta], Data$N, Data$J)
  parm[Data$pos.H] <- H <- interval(parm[Data$pos.H], 0.1, Inf)
  parm[Data$pos.nu] <- nu <- interval(parm[Data$pos.nu], 1e-100, Inf)
  nu <- matrix(nu, Data$N, Data$N)      sigma <- interval(parm[Data$pos.sigma],
1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  alpha.prior <- dunif(alpha, 1.5, 100, log=TRUE)
  beta.prior <- sum(dnormmv(beta, 0, 1000, log=TRUE))
  h.prior <- dhalfnorm(H-0.1, 1000, log=TRUE)
  nu.prior <- sum(dgamma(nu, alpha, 2, log=TRUE))
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
  ### Log-Likelihood
  w <- exp(-0.5 * Data$Z^2) / H
  tau <- (1/sigma^2) * w * nu
  mu <- tcrossprod(Data$X, beta)

```

```

LL <- sum(dnormp(Data$y, mu, tau, log=TRUE))
#WSE <- w * nu * (Data$y - mu)^2; w.y <- w * nu * Data$y
#WMSE <- rowMeans(WSE); y.w <- rowSums(w.y) / rowSums(w)
#LAR2 <- 1 - WMSE / sd(y.w)^2
### Log-Posterior
LP <- LL + alpha.prior + beta.prior + h.prior + nu.prior + sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=rnormp(prod(dim(mu)), mu, tau), parm=parm)
return(Modelout)
}

```

### 39.4. Initial Values

```
Initial.Values <- c(runif(1,1.5,100), rep(0,N*J), 1, rep(1,N*N), rep(1,N))
```

## 40. Hidden Markov Model

### 40.1. Form

This introductory hidden Markov model (HMM) includes  $N$  discrete states. Discrete draws of the states of  $\theta$  are updated with the Griddy-Gibbs sampler.

$$\mathbf{y}_t \sim \mathcal{N}(\mu_\theta, \sigma_\theta^2), \quad t = 1, \dots, T$$

$$\mu \sim \mathcal{N}(\mu_0, \lambda \sigma^2)$$

$$\sigma^2 \sim \mathcal{G}^{-1}(\nu, \sigma_0^2)$$

$$\theta \sim \mathcal{CAT}(\phi)$$

$$\phi_{1:N} \sim \mathcal{D}(\alpha_{1:N})$$

$$\lambda \sim \mathcal{HC}(25)$$

$$\mu_0 \sim \mathcal{N}(0, 1000)$$

$$\nu \sim \mathcal{HC}(25)$$

$$\sigma_0^2 \sim \mathcal{HC}(25)$$

### 40.2. Data

```

T <- 10 #Number of time-periods
N <- 2 #Number of discrete (hidden) states
y <- cumsum(rnorm(T))
alpha <- rep(1,N) #Concentration hyperparameter
mon.names <- "LP"
parm.names <- as.parm.names(list(lambda=rep(0,N), mu0=rep(0,N),
  mu1=rep(0,N), nu=rep(0,N), phi=rep(0,N), sigma0=rep(0,N),

```

```

sigma2=rep(0,N), theta=rep(0,T)))
pos.lambda <- grep("lambda", parm.names)
pos.mu0 <- grep("mu0", parm.names)
pos.mu1 <- grep("mu1", parm.names)
pos.nu <- grep("nu", parm.names)
pos.phi <- grep("phi", parm.names)
pos.sigma0 <- grep("sigma0", parm.names)
pos.sigma2 <- grep("sigma2", parm.names)
pos.theta <- grep("theta", parm.names)
PGF <- function(Data) {      phi <- runif(Data$N)
  phi <- phi / sum(phi)
  return(c(runif(Data$N), rnorm(Data$N),
    rnorm(Data$N), runif(Data$N), phi, runif(Data$N), runif(Data$N),
    rcat(Data$T, rep(1/Data$N,Data$N))))})
MyData <- list(N=N, PGF=PGF, T=T, alpha=alpha, mon.names=mon.names,
  parm.names=parm.names, pos.lambda=pos.lambda, pos.mu0=pos.mu0,
  pos.mu1=pos.mu1, pos.nu=pos.nu, pos.phi=pos.phi,
  pos.sigma0=pos.sigma0, pos.sigma2=pos.sigma2, pos.theta=pos.theta,
  y=y)

```

### 40.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  lambda <- interval(parm[Data$pos.lambda], 1e-100, Inf)
  parm[Data$pos.lambda] <- lambda
  mu0 <- parm[Data$pos.mu0]
  mu <- parm[Data$pos.mu1]
  parm[Data$pos.mu1] <- mu <- mu[order(mu)]
  parm[Data$pos.nu] <- nu <- interval(parm[Data$pos.nu], 1e-100, Inf)
  phi <- abs(parm[Data$pos.phi])
  parm[Data$pos.phi] <- phi <- phi / sum(phi)
  sigma0 <- interval(parm[Data$pos.sigma0], 1e-100, Inf)
  parm[Data$pos.sigma0] <- sigma0
  sigma2 <- interval(parm[Data$pos.sigma2], 1e-100, Inf)
  parm[Data$pos.sigma2] <- sigma2
  theta <- parm[Data$pos.theta]
  ### Log(Prior Densities)
  lambda.prior <- sum(dhalfcauchy(lambda, 25, log=TRUE))
  mu0.prior <- sum(dnormv(mu0, 0, 1000, log=TRUE))
  mu.prior <- sum(dnormv(mu, mu0, lambda*sigma2, log=TRUE))
  nu.prior <- sum(dhalfcauchy(nu, 25, log=TRUE))
  phi.prior <- sum(ddirichlet(phi, Data$alpha, log=TRUE))
  sigma0.prior <- sum(dhalfcauchy(sigma0, 25, log=TRUE))
  sigma2.prior <- sum(dinvgamma(sigma2, nu, sigma0, log=TRUE))

```

```

theta.prior <- sum(dcat(theta, phi, log=TRUE))
### Log-Likelihood
LL <- sum(dnormv(Data$y, mu[theta], sigma2[theta], log=TRUE))
### Log-Posterior
LP <- LL + lambda.prior + mu0.prior + mu.prior + nu.prior +
      phi.prior + sigma0.prior + sigma2.prior + theta.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
      yhat=rnormv(length(theta), mu[theta], sigma2[theta]), parm=parm)
return(Modelout)
}

```

#### 40.4. Initial Values

```

Initial.Values <- c(runif(N), rnorm(N), rnorm(N), runif(N), runif(N),
      runif(N), runif(N), rcat(T, rep(1/N,N)))

```

### 41. Inverse Gaussian Regression

#### 41.1. Form

$$\begin{aligned}
 \mathbf{y} &\sim \mathcal{N}^{-1}(\mu, \lambda) \\
 \mu &= \exp(\mathbf{X}\beta) + C \\
 \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\
 \lambda &\sim \mathcal{HC}(25)
 \end{aligned}$$

where  $C$  is a small constant, such as  $1.0\text{E-}10$ .

#### 41.2. Data

```

N <- 100
J <- 3 #Number of predictors, including the intercept
X <- matrix(1,N,J)
for (j in 2:J) {X[,j] <- rnorm(N,runif(1,-3,3),runif(1,0.1,1))}
beta.orig <- runif(J,-3,3)
e <- rnorm(N,0,0.1)
y <- exp(tcrossprod(X, t(beta.orig)) + e)
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J), lambda=0))
pos.beta <- grep("beta", parm.names)
pos.lambda <- grep("lambda", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,1000), rhalfcauchy(1,5)))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
      parm.names=parm.names, pos.beta=pos.beta, pos.lambda=pos.lambda, y=y)

```

### 41.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  lambda <- interval(parm[Data$pos.lambda], 1e-100, Inf)
  parm[Data$pos.lambda] <- lambda
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  lambda.prior <- dhalfcauchy(lambda, 25, log=TRUE)
  ### Log-Likelihood
  mu <- exp(tcrossprod(Data$X, t(beta))) + 1.0E-10
  LL <- sum(dinvgaussian(Data$y, mu, lambda, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + lambda.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rinvgaussian(length(mu), mu, lambda), parm=parm)
  return(Modelout)
}
```

### 41.4. Initial Values

```
Initial.Values <- c(rep(0,J), 1)
```

## 42. Kriging

This is an example of universal kriging of  $\mathbf{y}$  given  $\mathbf{X}$ , regression effects  $\beta$ , and spatial effects  $\zeta$ . Euclidean distance between spatial coordinates (longitude and latitude) is used for each of  $i = 1, \dots, N$  records of  $\mathbf{y}$ . An additional record is created from the same data-generating process to compare the accuracy of interpolation. For the spatial component,  $\phi$  is the rate of spatial decay and  $\kappa$  is the scale.  $\kappa$  is often difficult to identify, so it is set to 1 (Gaussian), but may be allowed to vary up to 2 (Exponential). In practice,  $\phi$  is also often difficult to identify. While  $\Sigma$  is spatial covariance, spatial correlation is  $\rho = \exp(-\phi\mathbf{D})$ . To extend this to a large data set, consider the predictive process kriging example in section 43.

### 42.1. Form

$$\begin{aligned}\mathbf{y} &\sim \mathcal{N}(\mu, \sigma_1^2) \\ \mu &= \mathbf{X}\beta + \zeta \\ \mathbf{y}^{new} &= \mathbf{X}\beta + \sum_{i=1}^N \left( \frac{\rho_i}{\sum \rho} \zeta_i \right) \\ \rho &= \exp(-\phi\mathbf{D}^{new})^\kappa\end{aligned}$$

$$\zeta \sim \mathcal{N}_N(\zeta_\mu, \Sigma)$$

$$\Sigma = \sigma_2^2 \exp(-\phi \mathbf{D})^\kappa$$

$$\beta_j \sim \mathcal{N}(0, 1000), \quad j = 1, \dots, 2$$

$$\sigma_j \sim \mathcal{HC}(25) \in [0.1, 10], \quad j = 1, \dots, 2$$

$$\phi \sim \mathcal{U}(1, 5)$$

$$\zeta_\mu = 0$$

$$\kappa = 1$$

## 42.2. Data

```

N <- 20
longitude <- runif(N+1,0,100)
latitude <- runif(N+1,0,100)
D <- as.matrix(dist(cbind(longitude,latitude), diag=TRUE, upper=TRUE))
Sigma <- 10000 * exp(-1.5 * D)
zeta <- as.vector(apply(rmvn(1000, rep(0,N+1), Sigma), 2, mean))
beta <- c(50,2)
X <- matrix(runif((N+1)*2,-2,2),(N+1),2); X[,1] <- 1
mu <- as.vector(tcrossprod(X, t(beta)))
y <- mu + zeta
longitude.new <- longitude[N+1]; latitude.new <- latitude[N+1]
Xnew <- X[N+1,]; ynew <- y[N+1]
longitude <- longitude[1:N]; latitude <- latitude[1:N]
X <- X[1:N,]; y <- y[1:N]
D <- as.matrix(dist(cbind(longitude,latitude), diag=TRUE, upper=TRUE))
D.new <- sqrt((longitude - longitude.new)^2 + (latitude - latitude.new)^2)
mon.names <- c("LP","ynew")
parm.names <- as.parm.names(list(zeta=rep(0,N), beta=rep(0,2),
                                sigma=rep(0,2), phi=0))
pos.zeta <- grep("zeta", parm.names)
pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)
pos.phi <- grep("phi", parm.names)
PGF <- function(Data) return(c(rmvn(1, rep(0, Data$N),
                                rhalfcauchy(1,25)^2 *exp(-runif(1,1,5)*Data$D)),
                                rnormv(2,0,1000), rhalfcauchy(2,5), runif(1,1,5)))
MyData <- list(D=D, D.new=D.new, latitude=latitude, longitude=longitude,
              N=N, PGF=PGF, X=X, Xnew=Xnew, mon.names=mon.names,
              parm.names=parm.names, pos.zeta=pos.zeta, pos.beta=pos.beta,
              pos.sigma=pos.sigma, pos.phi=pos.phi, y=y)

```

### 42.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  zeta <- parm[Data$pos.zeta]
  kappa <- 1
  sigma <- interval(parm[Data$pos.sigma], 0.1, 10)
  parm[Data$pos.sigma] <- sigma
  parm[Data$pos.phi] <- phi <- interval(parm[Data$pos.phi], 1, 5)
  Sigma <- sigma[2]*sigma[2] * exp(-phi * Data$D)^kappa
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  zeta.prior <- dmvn(zeta, rep(0, Data$N), Sigma, log=TRUE)
  sigma.prior <- sum(dhalfcauchy(sigma - 1, 25, log=TRUE))
  phi.prior <- dunif(phi, 1, 5, log=TRUE)
  ### Interpolation
  rho <- exp(-phi * Data$D.new)^kappa
  ynew <- rnorm(1, sum(beta * Data$Xnew) + sum(rho / sum(rho) * zeta),
    sigma[1])
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, t(beta)) + zeta
  LL <- sum(dnorm(Data$y, mu, sigma[1], log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + zeta.prior + sigma.prior + phi.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,ynew),
    yhat=rnorm(length(mu), mu, sigma[1]), parm=parm)
  return(Modelout)
}
```

### 42.4. Initial Values

```
Initial.Values <- c(rep(0,N), rep(0,2), rep(1,2), 1)
```

## 43. Kriging, Predictive Process

The first  $K$  of  $N$  records in  $\mathbf{y}$  are used as knots for the parent process, and the predictive process involves records  $(K + 1), \dots, N$ . For more information on kriging, see section [42](#).

### 43.1. Form

$$\mathbf{y} \sim \mathcal{N}(\mu, \sigma_1^2)$$

$$\mu_{1:K} = \mathbf{X}_{1:K,1:J}\beta + \zeta$$

$$\begin{aligned}\mu_{(K+1):N} &= \mathbf{X}_{(K+1):N,1:J}\beta + \sum_{p=1}^{N-K} \frac{\lambda_{p,1:K}}{\sum_{q=1}^{N-K} \lambda_{q,1:K}} \zeta^T \\ \lambda &= \exp(-\phi \mathbf{D}_P)^\kappa \\ \mathbf{y}^{new} &= \mathbf{X}\beta + \sum_{k=1}^K \left( \frac{\rho_k}{\sum \rho} \zeta_k \right) \\ \rho &= \exp(-\phi \mathbf{D}^{new})^\kappa \\ \zeta &\sim \mathcal{N}_K(0, \Sigma) \\ \Sigma &= \sigma_2^2 \exp(-\phi \mathbf{D})^\kappa \\ \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, 2 \\ \sigma_j &\sim \mathcal{HC}(25), \quad j = 1, \dots, 2 \\ \phi &\sim \mathcal{N}(0, 1000) \in [1, 5] \\ \kappa &= 1\end{aligned}$$

### 43.2. Data

```

N <- 100
K <- 30 #Number of knots
longitude <- runif(N+1,0,100)
latitude <- runif(N+1,0,100)
D <- as.matrix(dist(cbind(longitude,latitude), diag=TRUE, upper=TRUE))
Sigma <- 10000 * exp(-1.5 * D)
zeta <- as.vector(apply(rmvn(1000, rep(0,N+1), Sigma), 2, mean))
beta <- c(50,2)
X <- matrix(runif((N+1)*2,-2,2),(N+1),2); X[,1] <- 1
mu <- as.vector(tcrossprod(X, t(beta)))
y <- mu + zeta
longitude.new <- longitude[N+1]; latitude.new <- latitude[N+1]
Xnew <- X[N+1,]; ynew <- y[N+1]
longitude <- longitude[1:N]; latitude <- latitude[1:N]
X <- X[1:N,]; y <- y[1:N]
D <- as.matrix(dist(cbind(longitude[1:K],latitude[1:K]), diag=TRUE,
  upper=TRUE))
D.P <- matrix(0, N-K, K)
for (i in (K+1):N) {
  D.P[K+1-i,] <- sqrt((longitude[1:K] - longitude[i])^2 +
    (latitude[1:K] - latitude[i])^2)}
D.new <- sqrt((longitude[1:K] - longitude.new)^2 +
  (latitude[1:K] - latitude.new)^2)
mon.names <- c("LP","ynew")
parm.names <- as.parm.names(list(zeta=rep(0,K), beta=rep(0,2),
  sigma=rep(0,2), phi=0))

```



```

pos.zeta <- grep("zeta", parm.names)
pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)
pos.phi <- grep("phi", parm.names)
PGF <- function(Data) return(c(rmvn(1, rep(0, Data$K),
  rhalfcauchy(1,5)^2 *exp(-runif(1,1,5)*Data$D)),
  rnormmv(2,0,1000), rhalfcauchy(2,5), runif(1,1,5)))
MyData <- list(D=D, D.new=D.new, D.P=D.P, K=K, N=N, PGF=PGF, X=X,
  Xnew=Xnew, latitude=latitude, longitude=longitude,
  mon.names=mon.names, parm.names=parm.names, pos.zeta=pos.zeta,
  pos.beta=pos.beta, pos.sigma=pos.sigma, pos.phi=pos.phi, y=y)

```

### 43.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  zeta <- parm[Data$pos.zeta]
  kappa <- 1
  sigma <- interval(parm[Data$pos.sigma], 1, Inf)
  parm[Data$pos.sigma] <- sigma
  parm[Data$pos.phi] <- phi <- interval(parm[Data$pos.phi], 1, 5)
  Sigma <- sigma[2]*sigma[2] * exp(-phi * Data$D)^kappa
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  zeta.prior <- dmvn(zeta, rep(0, Data$K), Sigma, log=TRUE)
  sigma.prior <- sum(dhalfcauchy(sigma - 1, 25, log=TRUE))
  phi.prior <- dunif(phi, 1, 5, log=TRUE)
  ### Interpolation
  rho <- exp(-phi * Data$D.new)^kappa
  ynew <- rnorm(1, sum(beta * Data$Xnew) + sum(rho / sum(rho) * zeta),
    sigma)
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, t(beta))
  mu[1:Data$K] <- mu[1:Data$K] + zeta
  lambda <- exp(-phi * Data$D.P)^kappa
  mu[(Data$K+1):Data$N] <- mu[(Data$K+1):Data$N] +
    rowSums(lambda / rowSums(lambda) *
      matrix(zeta, Data$N - Data$K, Data$K, byrow=TRUE))
  LL <- sum(dnorm(Data$y, mu, sigma[1], log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + zeta.prior + sigma.prior + phi.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,ynew),
    yhat=rnorm(length(mu), mu, sigma[1]), parm=parm)
  return(Modelout)
}

```

```
}
```

#### 43.4. Initial Values

```
Initial.Values <- c(rep(0,K), c(mean(y), 0), rep(1,2), 3)
```

### 44. Laplace Regression

This linear regression specifies that  $\mathbf{y}$  is Laplace-distributed, where it is usually Gaussian or normally-distributed. It has been claimed that it should be surprising that the normal distribution became the standard, when the Laplace distribution usually fits better and has wider tails (Kotz, Kozubowski, and Podgorski 2001). Another popular alternative is to use the t-distribution (see Robust Regression in section 80), though it is more computationally expensive to estimate, because it has three parameters. The Laplace distribution has only two parameters, location and scale like the normal distribution, and is computationally easier to fit. This example could be taken one step further, and the parameter vector  $\beta$  could be Laplace-distributed. Laplace's Demon recommends that users experiment with replacing the normal distribution with the Laplace distribution.

#### 44.1. Form

$$\begin{aligned}\mathbf{y} &\sim \mathcal{L}(\mu, \sigma^2) \\ \mu &= \mathbf{X}\beta \\ \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\ \sigma &\sim \mathcal{HC}(25)\end{aligned}$$

#### 44.2. Data

```
N <- 10000
J <- 5
X <- matrix(1,N,J)
for (j in 2:J) {X[,j] <- rnorm(N,runif(1,-3,3),runif(1,0.1,1))}
beta <- runif(J,-3,3)
e <- rlaplace(N,0,0.1)
y <- tcrossprod(X, t(beta)) + e
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J), sigma=0))
pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,1000), rhalfcauchy(1,5)))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.sigma=pos.sigma, y=y)
```

### 44.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, t(beta))
  LL <- sum(dlaplace(Data$y, mu, sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rlaplace(length(mu), mu, sigma), parm=parm)
  return(Modelout)
}
```

### 44.4. Initial Values

```
Initial.Values <- c(rep(0,J), 1)
```

## 45. Latent Dirichlet Allocation

### 45.1. Form

$$\begin{aligned}
 \mathbf{Y}_{m,n} &\sim \mathcal{CAT}(\phi[\mathbf{Z}_{m,n},]), \quad m = 1, \dots, M, \quad n = 1, \dots, N \\
 \mathbf{Z}_{m,n} &\sim \mathcal{CAT}(\theta_{m,1:K}) \\
 \phi_{k,v} &\sim \mathcal{D}(\beta) \\
 \theta_{m,k} &\sim \mathcal{D}(\alpha) \\
 \alpha_k &= 1, \quad k = 1, \dots, K \\
 \beta_v &= 1, \quad v = 1, \dots, V
 \end{aligned}$$

### 45.2. Data

```
K <- 2 #Number of (latent) topics
M <- 4 #Number of documents in corpus
N <- 15 #Maximum number of (used) words per document
V <- 5 #Maximum number of occurrences of any word (Vocabulary size)
```

```

Y <- matrix(rcat(M*N,rep(1/V,V)), M, N)
rownames(Y) <- paste("doc", 1:nrow(Y), sep="")
colnames(Y) <- paste("word", 1:ncol(Y), sep="")
#Note: Y is usually represented as w, a matrix of word counts.
if(min(Y) == 0) Y <- Y + 1 #A zero cannot occur, Y must be 1,2,...,V.
V <- max(Y) #Maximum number of occurrences of any word (Vocabulary size)
alpha <- rep(1,K) # hyperparameters (constant)
beta <- rep(1,V)
mon.names <- "LP"
parm.names <- as.parm.names(list(phi=matrix(0,K,V), theta=matrix(0,M,K),
  Z=matrix(0,M,N)))
pos.phi <- grep("phi", parm.names)
pos.theta <- grep("theta", parm.names)
pos.Z <- grep("Z", parm.names)
PGF <- function(Data) {
  phi <- matrix(runif(Data$J*Data$V), Data$K, Data$V)
  phi <- phi / rowSums(phi)
  theta <- matrix(runif(Data$M*Data$K), Data$M, Data$K)
  theta <- theta / rowSums(theta)
  z <- rcat(Data$M*Data$N, rep(1/Data$K,Data$K))
  return(c(as.vector(phi), as.vector(theta), z))}
MyData <- list(K=K, M=M, N=N, PGF=PGF, V=V, Y=Y, alpha=alpha, beta=beta,
  mon.names=mon.names, parm.names=parm.names, pos.phi=pos.phi,
  pos.theta=pos.theta, pos.Z=pos.Z)

```

### 45.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  phi <- matrix(interval(parm[Data$pos.phi], 0, 1), Data$K, Data$V)
  phi <- phi / rowSums(phi)
  parm[Data$pos.phi] <- as.vector(phi)
  theta <- matrix(interval(parm[Data$pos.theta], 0, 1), Data$M, Data$K)
  theta <- theta / rowSums(theta)
  parm[Data$pos.theta] <- as.vector(theta)
  Z <- matrix(parm[Data$pos.Z], Data$M, Data$N)
  ### Log(Prior Densities)
  phi.prior <- sum(dldirichlet(phi, beta, log=TRUE))
  theta.prior <- sum(dldirichlet(theta, alpha, log=TRUE))
  ### Log-Likelihood
  LL <- Z.prior <- 0
  Yhat <- Data$Y
  for (m in 1:Data$M) {for (n in 1:Data$N) {
    Z.prior + Z.prior + dcat(Z[m,n], theta[m,], log=TRUE)
    LL <- LL + dcat(Data$Y[m,n], as.vector(phi[Z[m,n],]), log=TRUE)
  }
}

```

```

      Yhat[m,n] <- rcat(1, as.vector(phi[Z[m,n],]))}}
### Log-Posterior
LP <- LL + phi.prior + theta.prior + Z.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP, yhat=Yhat, parm=parm)
return(Modelout)
}

```

#### 45.4. Initial Values

```
Initial.Values <- c(rep(1/V,K*V), rep(1/K,M*K), rcat(M*N,rep(1/K,K)))
```

## 46. Linear Regression

### 46.1. Form

$$\begin{aligned}
 \mathbf{y} &\sim \mathcal{N}(\boldsymbol{\mu}, \sigma^2) \\
 \boldsymbol{\mu} &= \mathbf{X}\boldsymbol{\beta} \\
 \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\
 \sigma &\sim \mathcal{HC}(25)
 \end{aligned}$$

### 46.2. Data

```

N <- 10000
J <- 5
X <- matrix(1,N,J)
for (j in 2:J) {X[,j] <- rnorm(N,runif(1,-3,3),runif(1,0.1,1))}
beta <- runif(J,-3,3)
e <- rnorm(N,0,0.1)
y <- tcrossprod(X, t(beta)) + e
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J), sigma=0))
pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,1000), rhalfcauchy(1,5)))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
              parm.names=parm.names, pos.beta=pos.beta, pos.sigma=pos.sigma, y=y)

```

### 46.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters

```

```

beta <- parm[Data$pos.beta]
sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
parm[Data$pos.sigma] <- sigma
### Log(Prior Densities)
beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
sigma.prior <- dgamma(sigma, 25, log=TRUE)
### Log-Likelihood
mu <- tcrossprod(Data$X, t(beta))
LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
### Log-Posterior
LP <- LL + beta.prior + sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=rnorm(length(mu), mu, sigma), parm=parm)
return(Modelout)
}

```

#### 46.4. Initial Values

```
Initial.Values <- c(rep(0,J), 1)
```

## 47. Linear Regression, Frequentist

By eliminating prior probabilities, a frequentist linear regression example is presented. Although frequentism is not endorsed here, the purpose of this example is to illustrate how the **LaplaceDemon** package can be used for Bayesian or frequentist inference.

### 47.1. Form

$$\begin{aligned}
 \mathbf{y} &\sim \mathcal{N}(\boldsymbol{\mu}, \sigma^2) \\
 \boldsymbol{\mu} &= \mathbf{X}\boldsymbol{\beta}
 \end{aligned}$$

### 47.2. Data

```

N <- 10000
J <- 5
X <- matrix(1,N,J)
for (j in 2:J) {X[,j] <- rnorm(N,runif(1,-3,3),runif(1,0.1,1))}
beta <- runif(J,-3,3)
e <- rnorm(N,0,0.1)
y <- tcrossprod(X, t(beta)) + e
mon.names <- "LL"
parm.names <- as.parm.names(list(beta=rep(0,J), sigma=0))
pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)

```

```
PGF <- function(Data) return(c(rnormv(Data$J,0,1000), rhalfcauchy(1,5)))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.sigma=pos.sigma, y=y)
```

### 47.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  sigma <- interval(parm[Data$pos.sigma, 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, t(beta))
  LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
  Modelout <- list(LP=LL, Dev=-2*LL, Monitor=LL,
    yhat=rnorm(length(mu), mu, sigma), parm=parm)
  return(Modelout)
}
```

### 47.4. Initial Values

```
Initial.Values <- c(rep(0,J), 1)
```

## 48. Linear Regression, Hierarchical Bayesian

### 48.1. Form

$$\begin{aligned}
 \mathbf{y} &\sim \mathcal{N}(\boldsymbol{\mu}, \sigma^2) \\
 \boldsymbol{\mu} &= \mathbf{X}\boldsymbol{\beta} \\
 \beta_j &\sim \mathcal{N}(\gamma, \delta), \quad j = 1, \dots, J \\
 \gamma &\sim \mathcal{N}(0, 1000) \\
 \delta &\sim \mathcal{HC}(25) \\
 \sigma &\sim \mathcal{HC}(\tau) \\
 \tau &\sim \mathcal{HC}(25)
 \end{aligned}$$

### 48.2. Data

```
data(demonsnacks)
y <- log(demonsnacks$Calories)
```

```

X <- cbind(1, as.matrix(log(demonsnacks[,c(1,4,10)]+1)))
J <- ncol(X)
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J), gamma=0, delta=0, sigma=0,
    tau=0))
pos.beta <- grep("beta", parm.names)
pos.gamma <- grep("gamma", parm.names)
pos.delta <- grep("delta", parm.names)
pos.sigma <- grep("sigma", parm.names)
pos.tau <- grep("tau", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,1000), rnormv(1,0,1000),
    rhalfcauchy(3,5)))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
    parm.names=parm.names, pos.beta=pos.beta, pos.gamma=pos.gamma,
    pos.delta=pos.delta, pos.sigma=pos.sigma, pos.tau=pos.tau, y)

```

### 48.3. Model

```

Model <- function(parm, Data)
{
  ### Hyperparameters
  gamma <- parm[Data$pos.gamma]
  delta <- interval(parm[Data$pos.delta], 1e-100, Inf)
  parm[Data$pos.delta] <- delta
  parm[Data$pos.tau] <- tau <- interval(parm[Data$pos.tau], 1e-100, Inf)
  ### Parameters
  beta <- parm[Data$pos.beta]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Hyperprior Densities)
  gamma.prior <- dnormv(gamma, 0, 1000, log=TRUE)
  delta.prior <- dhalfcauchy(delta, 25, log=TRUE)
  tau.prior <- dhalfcauchy(tau, 25, log=TRUE)
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, gamma, delta, log=TRUE))
  sigma.prior <- dhalfcauchy(sigma, tau, log=TRUE)
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, t(beta))
  LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + gamma.prior + delta.prior + sigma.prior +
    tau.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rnorm(length(mu), mu, sigma), parm=parm)

```



```
return(Modelout)
}
```

#### 48.4. Initial Values

```
Initial.Values <- c(rep(0,J), 0, rep(1,3))
```

### 49. Linear Regression, Multilevel

#### 49.1. Form

$$\begin{aligned} \mathbf{y} &\sim \mathcal{N}(\mu, \sigma^2) \\ \mu_i &= \mathbf{X}\beta_{\mathbf{m}[i],1:J} \\ \beta_{g,1:J} &\sim \mathcal{N}_J(\gamma, \Omega^{-1}), \quad g = 1, \dots, G \\ \Omega &\sim \mathcal{W}_{J+1}(\mathbf{S}), \quad \mathbf{S} = \mathbf{I}_J \\ \gamma_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\ \sigma &\sim \mathcal{HC}(25) \end{aligned}$$

where  $\mathbf{m}$  is a vector of length  $N$ , and each element indicates the multilevel group ( $g = 1, \dots, G$ ) for the associated record.

#### 49.2. Data

```
N <- 30
J <- 2 ### Number of predictors (including intercept)
G <- 2 ### Number of Multilevel Groups
X <- matrix(rnorm(N,0,1),N,J); X[,1] <- 1
Sigma <- matrix(runif(J*J,-1,1),J,J)
diag(Sigma) <- runif(J,1,5)
Sigma <- as.positive.definite(Sigma)
gamma <- runif(J,-1,1)
beta <- matrix(NA,G,J)
for (g in 1:G) {beta[g,] <- rmvn(1, gamma, Sigma)}
m <- rcat(N, rep(1/G,G)) ### Multilevel group indicator
y <- rowSums(beta[m,] * X) + rnorm(N,0,0.1)
S <- diag(J)
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=matrix(0,G,J), gamma=rep(0,J),
  sigma=0, U=S), uppertri=c(0,0,0,1))
pos.beta <- grep("beta", parm.names)
pos.gamma <- grep("gamma", parm.names)
pos.sigma <- grep("sigma", parm.names)
```

```
PGF <- function(Data) return(c(rmvnpc(Data$G, rnormv(Data$J,0,100),
  rwishartc(Data$J+1, Data$S)), rnormv(Data$J,0,100), rhalfcauchy(1,5),
  upper.triangle(rwishartc(Data$J+1, Data$S), diag=TRUE)))
MyData <- list(G=G, J=J, N=N, PGF=PGF, S=S, X=X, m=m, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.gamma=pos.gamma,
  pos.sigma=pos.sigma, y=y)
```

### 49.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- matrix(parm[Data$pos.beta], Data$G, Data$J)
  gamma <- parm[Data$pos.gamma]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  U <- as.parm.matrix(U, Data$J, parm, Data, chol=TRUE)
  diag(U) <- exp(diag(U))
  ### Log(Prior Densities)
  U.prior <- dwishartc(U, Data$J+1, Data$S, log=TRUE)
  beta.prior <- sum(dmvnpc(beta, gamma, U, log=TRUE))
  gamma.prior <- sum(dnormv(gamma, 0, 100, log=TRUE))
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
  ### Log-Likelihood
  mu <- rowSums(beta[Data$m,] * Data$X)
  LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + U.prior + beta.prior + gamma.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rnorm(length(mu), mu, sigma), parm=parm)
  return(Modelout)
}
```

### 49.4. Initial.Values

```
Initial.Values <- c(rep(0,G*J), rep(0,J), 1,
  upper.triangle(S, diag=TRUE))
```

## 50. Linear Regression with Full Missingness

With ‘full missingness’, there are missing values for both the dependent variable (DV) and at least one independent variable (IV). The ‘full likelihood’ approach to full missingness is excellent as long as the model is identifiable. When it is not identifiable, imputation may be done in a previous stage, such as with the MISS function. In this example, matrix  $\alpha$  is for regression effects for IVs, vector  $\beta$  is for regression effects for the DV, vector  $\gamma$  is for missing

values for IVs, and  $\delta$  is for missing values for the DV.

### 50.1. Form

$$\begin{aligned}
 \mathbf{y}^{imp} &\sim \mathcal{N}(\nu, \sigma_J^2) \\
 \mathbf{X}^{imp} &\sim \mathcal{N}(\mu, \sigma_{-J}^2) \\
 \nu &= \mathbf{X}^{imp} \beta \\
 \mu &= \mathbf{X}^{imp} \alpha \\
 \mathbf{y}^{imp} &= \begin{cases} \delta & \text{if } \mathbf{y}^{mis} \\ \mathbf{y}^{obs} & \end{cases} \\
 \mathbf{X}^{imp} &= \begin{cases} \gamma & \text{if } \mathbf{X}^{mis} \\ \mathbf{X}^{obs} & \end{cases} \\
 \alpha_{j,l} &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, (J-1), \quad l = 1, \dots, (J-1) \\
 \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\
 \gamma_m &\sim \mathcal{N}(0, 1000), \quad m = 1, \dots, M \\
 \delta_p &\sim \mathcal{N}(0, 1000), \quad p = 1, \dots, P \\
 \sigma_j &\sim \mathcal{HC}(25), \quad j = 1, \dots, J
 \end{aligned}$$

### 50.2. Data

```

N <- 100
J <- 5
X <- matrix(runif(N*J,-2,2),N,J); X[,1] <- 1 #Design matrix X
M <- matrix(round(runif(N*J)-0.45),N,J); M[,1] <- 0 #Missing indicators
X <- ifelse(M == 1, NA, X) #Simulated X gets missings according to M
beta.orig <- runif(J,-2,2)
y <- as.vector(tcrossprod(X, t(beta.orig)) + rnorm(N,0,0.1))
y[sample(1:N, round(N*.05))] <- NA
m <- ifelse(is.na(y), 1, 0) #Missing indicator for vector y
mon.names <- "LP"
parm.names <- as.parm.names(list(alpha=matrix(0,J-1,J-1),
  beta=rep(0,J),
  gamma=rep(0,sum(is.na(X))),
  delta=rep(0,sum(is.na(y))),
  sigma=rep(0,J)))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
pos.gamma <- grep("gamma", parm.names)
pos.delta <- grep("delta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv((Data$J-1)*(Data$J-1),0,10),

```

```

rnormmv(Data$J,0,10),
rnormmv(sum(is.na(Data$X)),0,10),
rnormmv(sum(is.na(Data$y)),mean(Data$y, na.rm=TRUE),1),
rhalfcauchy(Data$J,5)))
MyData <- list(J=J, N=N, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.alpha=pos.alpha, pos.beta=pos.beta,
  pos.gamma=pos.gamma, pos.delta=pos.delta, pos.sigma=pos.sigma, y=y)

```

### 50.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  alpha <- matrix(parm[Data$pos.alpha], Data$J-1, Data$J-1)
  beta <- parm[Data$pos.beta]
  gamma <- parm[Data$pos.gamma]
  delta <- parm[Data$pos.delta]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  alpha.prior <- sum(dnormmv(alpha, 0, 1000, log=TRUE))
  beta.prior <- sum(dnormmv(beta, 0, 1000, log=TRUE))
  gamma.prior <- sum(dnormmv(gamma, 0, 1000, log=TRUE))
  delta.prior <- sum(dnormmv(delta, 0, 1000, log=TRUE))
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
  ### Log-Likelihood
  mu <- X.imputed <- Data$X
  X.imputed[which(is.na(X.imputed))] <- gamma
  y.imputed <- Data$y
  y.imputed[which(is.na(y.imputed))] <- delta
  for (j in 2:Data$J) {mu[,j] <- tcrossprod(X.imputed[,-j],
    t(alpha[, (j-1)]))}
  nu <- tcrossprod(X.imputed, t(beta))
  LL <- sum(dnorm(X.imputed[,-1], mu[,-1],
    matrix(sigma[1:(Data$J-1)], Data$N, Data$J-1), log=TRUE),
    dnorm(y.imputed, nu, sigma[Data$J], log=TRUE))
  ### Log-Posterior
  LP <- LL + alpha.prior + beta.prior + gamma.prior + delta.prior +
    sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rnorm(length(nu), nu, sigma[Data$J]), parm=parm)
  return(Modelout)
}

```

## 50.4. Initial Values

```
Initial.Values <- c(rep(0, (J-1)^2), rep(0,J), rep(0, sum(is.na(X))),
  rep(0, sum(is.na(y))), rep(1,J))
```

# 51. Linear Regression with Missing Response

This is an introductory example to missing values using data augmentation with auxiliary variables. The dependent variable, or response, has both observed values,  $\mathbf{y}^{obs}$ , and missing values,  $\mathbf{y}^{mis}$ . The  $\alpha$  vector is for missing value imputation, and enables the use of the full-likelihood by augmenting the state with these auxiliary variables. In the model form,  $M$  is used to denote the number of missing values, though it is used as an indicator in the data.

## 51.1. Form

$$\begin{aligned}\mathbf{y}^{imp} &\sim \mathcal{N}(\mu, \sigma^2) \\ \mathbf{y}^{imp} &= \begin{cases} \alpha & \text{if } \mathbf{y}^{mis} \\ \mathbf{y}^{obs} & \end{cases} \\ \mu &= \mathbf{X}\beta \\ \alpha_m &\sim \mathcal{N}(0, 1000), \quad m = 1, \dots, M \\ \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\ \sigma &\sim \mathcal{HC}(25)\end{aligned}$$

## 51.2. Data

```
data(demonsnacks)
N <- nrow(demonsnacks)
J <- ncol(demonsnacks)
y <- log(demonsnacks$Calories)
y[sample(1:N, round(N*0.05))] <- NA
M <- ifelse(is.na(y), 1, 0)
X <- cbind(1, as.matrix(demonsnacks[,c(1,3:10)]))
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
mon.names <- "LP"
parm.names <- as.parm.names(list(alpha=rep(0,sum(M)), beta=rep(0,J),
  sigma=0))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnorm(sum(Data$M), mean(y, na.rm=TRUE), 1),
  rnormv(Data$J, 0, 1000), rhalfcauchy(1, 5)))
MyData <- list(J=J, M=M, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.alpha=pos.alpha, pos.beta=pos.beta,
```

```
pos.sigma=pos.sigma, y=y)
```

### 51.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[Data$pos.alpha]
  beta <- parm[Data$pos.beta]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  alpha.prior <- sum(dnormv(alpha, 0, 1000, log=TRUE))
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  sigma.prior <- dgamma(sigma, 25, log=TRUE)
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, t(beta))
  y.imputed <- Data$y
  y.imputed[which(is.na(Data$y))] <- alpha
  LL <- sum(dnorm(y.imputed, mu, sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + alpha.prior + beta.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rnorm(length(mu), mu, sigma), parm=parm)
  return(Modelout)
}
```

### 51.4. Initial Values

```
Initial.Values <- c(rep(0,sum(M)), rep(0,J), 1)
```

## 52. Linear Regression with Missing Response via ABB

The Approximate Bayesian Bootstrap (ABB), using the `ABB` function, is used to impute missing values in the dependent variable (DV), or response, given a propensity score. In this example, vector  $\alpha$  is used to estimate propensity score  $\eta$ , while vector  $\beta$  is for regression effects, and vector  $\gamma$  has the monitored missing values. For more information on ABB, see the `ABB` function.

### 52.1. Form

$$\mathbf{y}^{imp} \sim \mathcal{N}(\mu, \sigma^2)$$

$$\mathbf{y}^{imp} = \begin{cases} \gamma & \text{if } \mathbf{y}^{mis} \\ \mathbf{y}^{obs} & \end{cases}$$

$$\begin{aligned}\mu &= \mathbf{X}\beta \\ \gamma &\sim p(\mathbf{y}^{obs} | \mathbf{y}^{obs}, \mathbf{y}^{mis}, \eta) \\ \eta &= \frac{1}{1 + \exp(-\nu)} \\ \nu &= \mathbf{X}\alpha \\ \alpha_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\ \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\ \sigma &\sim \mathcal{HC}(25)\end{aligned}$$

## 52.2. Data

```
data(demonsnacks)
N <- nrow(demonsnacks)
J <- ncol(demonsnacks)
y <- log(demonsnacks$Calories)
y[sample(1:N, round(N*0.05))] <- NA
M <- ifelse(is.na(y), 1, 0)
X <- cbind(1, as.matrix(demonsnacks[,c(1,3:10)]))
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
mon.names <- c("LP", paste("gamma[", 1:sum(is.na(y)), "]", sep=""))
parm.names <- as.parm.names(list(alpha=rep(0,J), beta=rep(0,J), sigma=0))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,10), rnormv(Data$J,0,10),
  rhalfcauchy(1,5)))
MyData <- list(J=J, M=M, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.alpha=pos.alpha, pos.beta=pos.beta, pos.sigma=pos.sigma,
  y=y)
```

## 52.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[Data$pos.alpha]
  beta <- parm[Data$pos.beta]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  alpha.prior <- sum(dnormv(alpha, 0, 1000, log=TRUE))
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  sigma.prior <- dgamma(sigma, 25, log=TRUE)
```

```

### Log-Likelihood
y.imputed <- Data$y
mu <- tcrossprod(Data$X, t(beta))
nu <- as.vector(tcrossprod(Data$X, t(alpha)))
eta <- invlogit(nu)
breaks <- as.vector(quantile(eta, probs=c(0,0.2,0.4,0.6,0.8,1)))
B <- matrix(breaks[-length(breaks)], length(Data$y), 5, byrow=TRUE)
z <- rowSums(eta >= B)
for (i in 1:5) {
  if(any(is.na(Data$y[which(z == i)]))) {
    imp <- unlist(ABB(Data$y[which(z == i)]))
    y.imputed[which({z == i} & is.na(Data$y))] <- imp}}
gamma <- y.imputed[which(is.na(Data$y))]
LL <- sum(dbern(Data$M, eta, log=TRUE),
  dnorm(y.imputed, mu, sigma, log=TRUE))
### Log-Posterior
LP <- LL + alpha.prior + beta.prior + sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,gamma),
  yhat=rnorm(length(mu), mu, sigma), parm=parm)
return(Modelout)
}

```

## 52.4. Initial Values

```
Initial.Values <- c(rep(0,J), rep(0,J), 1)
```

## 53. Linear Regression with Power Priors

Power priors (Ibrahim and Chen 2000) are a class of informative priors when relevant historical data is available. Power priors may be used when it is desirable to take historical data into account while analyzing similar, current data. Both the current data,  $\mathbf{y}$  and  $\mathbf{X}$ , and historical data,  $\mathbf{y}_h$  and  $\mathbf{X}_h$ , are included in the power prior analysis, where  $h$  indicates historical data. Each data set receives its own likelihood function, though the likelihood of the historical data is raised to an exponential power,  $\alpha \in [0, 1]$ . In this example,  $\alpha$  is a constant.

### 53.1. Form

$$\begin{aligned}
 \mathbf{y} &\sim \mathcal{N}(\mu, \sigma^2) \\
 \mathbf{y}_h &\sim \mathcal{N}(\mu_h, \sigma^2)^\alpha \\
 \mu &= \mathbf{X}\beta \\
 \mu_h &= \mathbf{X}_h\beta \\
 \alpha &= 0.5 \\
 \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J
 \end{aligned}$$



$$\sigma \sim \mathcal{HC}(25)$$

### 53.2. Data

```

N <- 100
J <- 5 #Number of predictors, including the intercept
X <- Xh <- matrix(1,N,J)
for (j in 2:J) {
  X[,j] <- rnorm(N,runif(1,-3,3),runif(1,0.1,1))
  Xh[,j] <- rnorm(N,runif(1,-3,3),runif(1,0.1,1))}
beta.orig <- runif(J,-3,3)
e <- rnorm(N,0,0.1)
yh <- as.vector(tcrossprod(beta.orig, Xh) + e)
y <- as.vector(tcrossprod(beta.orig, X) + e)
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J), sigma=0))
pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,1000), rhalfcauchy(1,5)))
MyData <- list(alpha=0.5, J=J, PGF=PGF, X=X, Xh=Xh, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.sigma=pos.sigma, y=y,
  yh=yh)

```

### 53.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
  ### Log-Likelihood
  muh <- tcrossprod(Data$Xh, t(beta))
  mu <- tcrossprod(Data$X, t(beta))
  LL <- sum(Data$alpha*dnorm(Data$yh, muh, sigma, log=TRUE) +
    dnorm(Data$y, mu, sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rnorm(length(mu), mu, sigma), parm=parm)
  return(Modelout)
}

```

### 53.4. Initial Values

```
Initial.Values <- c(rep(0,J), 1)
```

## 54. Linear Regression with Zellner's g-Prior

For more information on Zellner's g-prior, see the documentation for the `dzellner` function.

### 54.1. Form

$$\begin{aligned} \mathbf{y} &\sim \mathcal{N}(\mu, \sigma^2) \\ \mu &= \mathbf{X}\beta \\ \beta &\sim \mathcal{N}_J(0, g\sigma^2(\mathbf{X}^T\mathbf{X})^{-1}) \\ g &\sim \mathcal{HG}(\alpha), \quad \alpha = 3 \\ \sigma &\sim \mathcal{HC}(25) \end{aligned}$$

### 54.2. Data

```
data(demonsnacks)
y <- log(demonsnacks$Calories)
X <- cbind(1, as.matrix(demonsnacks[,c(1,3:10)]))
J <- ncol(X)
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J), g0=0, sigma=0))
pos.beta <- grep("beta", parm.names)
pos.g <- grep("g0", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,10), rhalfcauchy(2,5)))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.g=pos.g,      pos.sigma=pos.sigma,
  y=y)
```

### 54.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  parm[Data$pos.g] <- g <- interval(parm[Data$pos.g], 1e-100, Inf)
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
```

```

parm[Data$pos.sigma] <- sigma
### Log(Hyperprior Densities)
g.prior <- dhyperg(g, alpha=3, log=TRUE)
### Log(Prior Densities)
beta.prior <- dzellner(beta, g, sigma, Data$X, log=TRUE)
sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
### Log-Likelihood
mu <- tcrossprod(Data$X, t(beta))
LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
### Log-Posterior
LP <- LL + beta.prior + g.prior + sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=rnorm(length(mu), mu, sigma), parm=parm)
return(Modelout)
}

```

#### 54.4. Initial Values

```
Initial.Values <- c(rep(1,J), rep(1,2))
```

## 55. LSTAR

This is a Logistic Smooth-Threshold Autoregression (LSTAR), and is specified with a transition function that includes  $\gamma$  as the shape parameter,  $\mathbf{y}$  as the transition variable,  $\theta$  as the location parameter, and  $d$  as the delay parameter.

### 55.1. Form

$$\begin{aligned}
 \mathbf{y}_t &\sim \mathcal{N}(\mu_t, \sigma^2), \quad t = 1, \dots, T \\
 \mu_t &= \pi_t(\alpha_1 + \phi_1 \mathbf{y}_{t-1}) + (1 - \pi_t)(\alpha_2 + \phi_2 \mathbf{y}_{t-1}), \quad t = 2, \dots, T \\
 \pi_t &= \frac{1}{1 + \exp(-(\gamma(\mathbf{y}_{t-d} - \theta)))} \\
 \alpha_j &\sim \mathcal{N}(0, 1000) \in [\mathbf{y}_{min}, \mathbf{y}_{max}], \quad j = 1, \dots, 2 \\
 \phi_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, 2 \\
 \gamma &\sim \mathcal{HC}(25) \\
 \theta &\sim \mathcal{U}(\mathbf{y}_{min}, \mathbf{y}_{max}) \\
 \pi_1 &\sim \mathcal{U}(0.001, 0.999) \\
 \sigma &\sim \mathcal{HC}(25)
 \end{aligned}$$

### 55.2. Data

```

y <- c(26.73, 26.75, 26.24, 25.94, 27.40, 26.14, 23.99, 23.08, 22.55,
  20.64, 23.28, 24.92, 25.07, 26.53, 28.14, 30.10, 27.43, 27.24,

```

```

23.96, 25.85, 26.76, 26.05, 26.79, 26.69, 29.89, 29.09, 23.84,
24.87, 24.47, 22.85, 22.05, 22.82, 22.99, 21.60, 20.32, 20.80,
19.78, 19.87, 18.78, 19.64, 20.00, 21.51, 21.49, 21.96, 22.58,
21.22, 22.34, 22.76, 18.37, 17.50, 17.55, 12.14, 4.76, 3.75,
2.05, 2.69, 3.85, 4.72, 5.00, 3.31, 3.02, 3.15, 2.50,
3.33, 3.95, 4.00, 3.86, 3.87, 3.51, 3.19, 2.39, 2.33,
2.57, 2.80, 2.43, 2.43, 2.10, 2.31, 2.21, 2.11, 2.10,
1.70, 1.35, 1.83, 1.55, 1.63, 1.91, 2.14, 2.41, 2.06,
1.87, 2.11, 2.28, 2.26, 2.03, 2.06, 2.08, 1.91, 1.95,
1.56, 1.44, 1.60, 1.77, 1.77, 1.95, 2.01, 1.65, 1.87,
2.01, 1.84, 1.94, 1.93, 1.93, 1.75, 1.73, 1.80, 1.74,
1.80, 1.75, 1.67, 1.60, 1.61, 1.55, 1.56, 1.57, 1.55,
1.56, 1.57, 1.69, 1.66, 1.74, 1.64, 1.65, 1.62, 1.54,
1.58, 1.49, 1.41, 1.42, 1.37, 1.45, 1.31, 1.37, 1.26,
1.35, 1.41, 1.29, 1.28, 1.23, 1.08, 1.03, 1.00, 1.04,
1.04, 0.92, 0.96, 0.90, 0.85, 0.78, 0.73, 0.59, 0.54,
0.53, 0.41, 0.46, 0.52, 0.42, 0.42, 0.43, 0.43, 0.35,
0.35, 0.35, 0.42, 0.41, 0.41, 0.50, 0.83, 0.96, 1.38,
1.62, 1.26, 1.48, 1.39, 1.20, 1.10, 1.02, 0.95, 1.00,
1.07, 1.14, 1.14, 1.10, 1.05, 1.08, 1.16, 1.42, 1.52,
1.60, 1.69, 1.62, 1.29, 1.46, 1.43, 1.50, 1.46, 1.40,
1.34, 1.41, 1.38, 1.38, 1.46, 1.73, 1.84, 1.95, 2.01,
1.90, 1.81, 1.60, 1.84, 1.72, 1.83, 1.81, 1.78, 1.80,
1.70, 1.70, 1.66, 1.67, 1.69, 1.66, 1.56, 1.47, 1.64,
1.71, 1.66, 1.65, 1.60, 1.61, 1.61, 1.53, 1.48, 1.40,
1.47, 1.53, 1.39, 1.41, 1.42, 1.46, 1.46, 1.33, 1.16)
T <- length(y)
mon.names <- c("LP", "ynew", "pi.new")
parm.names <- as.parm.names(list(alpha=rep(0,2), phi=rep(0,2), gamma=0,
  theta=0, pi=0, sigma=0))
pos.alpha <- grep("alpha", parm.names)
pos.phi <- grep("phi", parm.names)
pos.gamma <- grep("gamma", parm.names)
pos.theta <- grep("theta", parm.names)
pos.pi <- grep("pi", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(2,0,10), rnormv(2,0,10),
  rhalfcauchy(1,5), runif(1,min(Data$y),max(Data$y)),
  runif(1,0.001,0.999), rhalfcauchy(1,5)))
MyData <- list(PGF=PGF, T=T, mon.names=mon.names, parm.names=parm.names,
  pos.alpha=pos.alpha, pos.phi=pos.phi, pos.gamma=pos.gamma,
  pos.theta=pos.theta, pos.pi=pos.pi, pos.sigma=pos.sigma, y=y)

```

### 55.3. Model

```
Model <- function(parm, Data)
```

```

{
### Parameters
alpha <- interval(parm[Data$pos.alpha], min(Data$y), max(Data$y))
parm[Data$pos.alpha] <- alpha
parm[Data$pos.phi] <- phi <- interval(parm[Data$pos.phi], -1, 1)
gamma <- interval(parm[Data$pos.gamma], 1e-100, Inf)
parm[Data$pos.gamma] <- gamma
theta <- interval(parm[Data$pos.theta], min(Data$y), max(Data$y))
parm[Data$pos.theta] <- theta
parm[Data$pos.pi] <- pi <- interval(parm[Data$pos.pi], 0.001, 0.999)
sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
parm[Data$pos.sigma] <- sigma
### Log(Prior Densities)
alpha.prior <- sum(dnormv(alpha, 0, 1000, log=TRUE))
phi.prior <- sum(dnormv(phi, 0, 1000, log=TRUE))
gamma.prior <- dhalfcauchy(gamma, 25, log=TRUE)
theta.prior <- dunif(theta, min(Data$y), max(Data$y), log=TRUE)
pi.prior <- dunif(pi, 0.001, 0.999, log=TRUE)
sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
### Log-Likelihood
pi <- c(pi, 1 / (1 + exp(-(gamma*(Data$y[-Data$T]-theta)))))
pi.new <- 1 / (1 + exp(-(gamma*(Data$y[Data$T]-theta)))))
mu <- pi * c(alpha[1], alpha[1] + phi[1]*Data$y[-Data$T]) +
      (1-pi) * c(alpha[2], alpha[2] + phi[2]*Data$y[-Data$T])
ynew <- rnorm(1, pi.new * (alpha[1] + phi[1]*Data$y[Data$T]) +
              (1-pi.new) * (alpha[2] + phi[2]*Data$y[Data$T]), sigma)
LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
### Log-Posterior
LP <- LL + alpha.prior + phi.prior + gamma.prior + theta.prior +
      pi.prior + sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,sigma,ynew,pi.new),
                 yhat=rnorm(length(mu), mu, sigma), parm=parm)
return(Modelout)
}

```

#### 55.4. Initial Values

```
Initial.Values <- c(rep(min(y),4), 1, mean(y), 0.5, 1)
```

## 56. MANCOVA

Since this is a multivariate extension of ANCOVA, please see the ANCOVA example in section 1 for a univariate introduction.

### 56.1. Form

$$\mathbf{Y}_{i,1:J} \sim \mathcal{N}_K(\mu_{i,1:J}, \Sigma), \quad i = 1, \dots, N$$

$$\mu_{i,k} = \alpha_k + \beta_{k,\mathbf{X}[i,1]} + \gamma_{k,\mathbf{X}[i,1]} + \mathbf{X}_{1:N,3:(C+J)} \delta_{k,1:C}$$

$$\epsilon_{i,k} = \mathbf{Y}_{i,k} - \mu_{i,k}$$

$$\alpha_k \sim \mathcal{N}(0, 1000), \quad k = 1, \dots, K$$

$$\beta_{k,l} \sim \mathcal{N}(0, \sigma_1^2), \quad l = 1, \dots, (L-1)$$

$$\beta_{1:K,L} = - \sum_{l=1}^{L-1} \beta_{1:K,l}$$

$$\gamma_{k,m} \sim \mathcal{N}(0, \sigma_2^2), \quad m = 1, \dots, (M-1)$$

$$\gamma_{1:K,M} = - \sum_{m=1}^{M-1} \beta_{1:K,m}$$

$$\delta_{k,c} \sim \mathcal{N}(0, 1000)$$

$$\Omega \sim \mathcal{W}_{K+1}(\mathbf{S}), \quad \mathbf{S} = \mathbf{I}_K$$

$$\Sigma = \Omega^{-1}$$

$$\sigma_{1:J} \sim \mathcal{HC}(25)$$

### 56.2. Data

```

C <- 2 #Number of covariates
J <- 2 #Number of factors (treatments)
K <- 3 #Number of endogenous (dependent) variables
L <- 4 #Number of levels in factor (treatment) 1
M <- 5 #Number of levels in factor (treatment) 2
N <- 100
X <- matrix(cbind(rcat(N, rep(1/L,L)), rcat(N, rep(1/M,M))),
            runif(C*N,0,1)), N, J + C)
alpha <- runif(K,-1,1)
beta <- matrix(runif(K*L,-2,2), K, L)
beta[,L] <- -rowSums(beta[, -L])
gamma <- matrix(runif(K*M,-2,2), K, M)
gamma[,M] <- -rowSums(gamma[, -M])
delta <- matrix(runif(K*C), K, C)
Y <- matrix(NA,N,K)
for (k in 1:K) {
  Y[,k] <- alpha[k] + beta[k,X[,1]] + gamma[k,X[,2]] +
    tcrossprod(delta[k,], X[, -c(1,2)]) + rnorm(1,0,0.1)}
S <- diag(K)
mon.names <- c("LP", "s.o.beta", "s.o.gamma", "s.o.epsilon",
  as.parm.names(list(s.beta=rep(0,K), s.gamma=rep(0,K),

```

```

      s.epsilon=rep(0,K)))
parm.names <- as.parm.names(list(alpha=rep(0,K), beta=matrix(0,K,(L-1)),
  gamma=matrix(0,K,(M-1)), delta=matrix(0,K,C), U=diag(K),
  sigma=rep(0,2)), uppertri=c(0,0,0,0,1,0))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
pos.gamma <- grep("gamma", parm.names)
pos.delta <- grep("delta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$K,0,1000),
  rnorm(Data$K*(Data$L-1),0,rhalfcauchy(1,5)),
  rnorm(Data$K*(Data$M-1),0,rhalfcauchy(1,5)),
  rnormv(Data$K*Data$C,0,1000),
  upper.triangle(rwishartc(Data$K+1,Data$S), diag=TRUE),
  rhalfcauchy(2,5)))
MyData <- list(C=C, J=J, K=K, L=L, M=M, N=N, PGF=PGF, S=S, X=X, Y=Y,
  mon.names=mon.names, parm.names=parm.names, pos.alpha=pos.alpha,
  pos.beta=pos.beta, pos.gamma=pos.gamma, pos.delta=pos.delta,
  pos.sigma=pos.sigma)

```

### 56.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[Data$pos.alpha]
  beta <- matrix(c(parm[Data$pos.beta], rep(0,Data$K)), Data$K, Data$L)
  beta[,Data$L] <- -rowSums(beta[,Data$L])
  gamma <- matrix(c(parm[Data$pos.gamma],
    rep(0,Data$K)), Data$K, Data$M)
  gamma[,Data$M] <- -rowSums(gamma[,Data$M])
  delta <- matrix(parm[Data$pos.delta], Data$K, Data$C)
  U <- as.parm.matrix(U, Data$K, parm, Data, chol=TRUE)
  diag(U) <- exp(diag(U))
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  alpha.prior <- sum(dnormv(alpha, 0, 1000, log=TRUE))
  beta.prior <- sum(dnorm(beta, 0, sigma[1], log=TRUE))
  gamma.prior <- sum(dnorm(gamma, 0, sigma[2], log=TRUE))
  delta.prior <- sum(dnormv(delta, 0, 1000, log=TRUE))
  U.prior <- dwishartc(U, Data$K+1, Data$S, log=TRUE)
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
  ### Log-Likelihood
  mu <- matrix(0,Data$N,Data$K)
  for (k in 1:Data$K) {

```

```

      mu[,k] <- alpha[k] + beta[k,Data$X[,1]] + gamma[k,Data$X[,2]] +
      tcrossprod(Data$X[,-c(1,2)], t(delta[k,]))}
LL <- sum(dmvnpc(Data$Y, mu, U, log=TRUE))
### Variance Components, Omnibus
s.o.beta <- sd(as.vector(beta))
s.o.gamma <- sd(as.vector(gamma))
s.o.epsilon <- sd(as.vector(Data$Y - mu))
### Variance Components, Univariate
s.beta <- apply(beta,1,sd)
s.gamma <- apply(gamma,1,sd)
s.epsilon <- apply(Data$Y - mu,2,sd)
### Log-Posterior
LP <- LL + alpha.prior + beta.prior + gamma.prior + delta.prior +
      U.prior + sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP, s.o.beta, s.o.gamma,
      s.o.epsilon, s.beta, s.gamma, s.epsilon),
      yhat=rmvnp(nrow(mu), mu, U), parm=parm)
return(Modelout)
}

```

#### 56.4. Initial Values

```

Initial.Values <- c(rep(0,K), rep(0,K*(L-1)), rep(0,K*(M-1)),
      rep(0,C*K), upper.triangle(S, diag=TRUE), rep(1,2))

```

## 57. MANOVA

Since this is a multivariate extension of ANOVA, please see the two-way ANOVA example in section 3 for a univariate introduction.

### 57.1. Form

$$\mathbf{Y}_{i,1:J} \sim \mathcal{N}_K(\mu_{i,1:J}, \Omega^{-1}), \quad i = 1, \dots, N$$

$$\mu_{i,k} = \alpha_k + \beta_{k,\mathbf{X}[i,1]} + \gamma_{k,\mathbf{X}[i,1]}$$

$$\epsilon_{i,k} = \mathbf{Y}_{i,k} - \mu_{i,k}$$

$$\alpha_k \sim \mathcal{N}(0, 1000), \quad k = 1, \dots, K$$

$$\beta_{k,l} \sim \mathcal{N}(0, \sigma_1^2), \quad l = 1, \dots, (L-1)$$

$$\beta_{1:K,L} = - \sum_{l=1}^{L-1} \beta_{1:K,l}$$

$$\gamma_{k,m} \sim \mathcal{N}(0, \sigma_2^2), \quad m = 1, \dots, (M-1)$$



$$\gamma_{1:K,M} = - \sum_{m=1}^{M-1} \beta_{1:K,m}$$

$$\Omega \sim \mathcal{W}_{K+1}(\mathbf{S}), \quad \mathbf{S} = \mathbf{I}_K$$

$$\sigma_{1:J} \sim \mathcal{HC}(25)$$

## 57.2. Data

```

J <- 2 #Number of factors (treatments)
K <- 3 #Number of endogenous (dependent) variables
L <- 4 #Number of levels in factor (treatment) 1
M <- 5 #Number of levels in factor (treatment) 2
N <- 100
X <- cbind(rcat(N, rep(1/L,L)), rcat(N, rep(1/M,M)))
alpha <- runif(K,-1,1)
beta <- matrix(runif(K*L,-2,2), K, L)
beta[,L] <- -rowSums(beta[, -L])
gamma <- matrix(runif(K*M,-2,2), K, M)
gamma[,M] <- -rowSums(gamma[, -M])
Y <- matrix(NA,N,K)
for (k in 1:K) {
  Y[,k] <- alpha[k] + beta[k,X[,1]] + gamma[k,X[,2]] + rnorm(1,0,0.1)}
S <- diag(K)
mon.names <- c("LP", "s.o.beta", "s.o.gamma", "s.o.epsilon",
  as.parm.names(list(s.beta=rep(0,K), s.gamma=rep(0,K),
    s.epsilon=rep(0,K))))
parm.names <- as.parm.names(list(alpha=rep(0,K), beta=matrix(0,K,(L-1)),
  gamma=matrix(0,K,(M-1)), U=diag(K), sigma=rep(0,2)),
  uppertri=c(0,0,0,1,0))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
pos.gamma <- grep("gamma", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$K,0,1000),
  rnorm(Data$K*(Data$L-1),0,rhalfcauchy(1,5)),
  rnorm(Data$K*(Data$M-1),0,rhalfcauchy(1,5)),
  upper.triangle(rwishartc(Data$K+1,Data$S), diag=TRUE),
  rhalfcauchy(2,5)))
MyData <- list(J=J, K=K, L=L, M=M, N=N, PGF=PGF, S=S, X=X, Y=Y,
  mon.names=mon.names, parm.names=parm.names, pos.alpha=pos.alpha,
  pos.beta=pos.beta, pos.gamma=pos.gamma, pos.sigma=pos.sigma)

```

## 57.3. Model

```

Model <- function(parm, Data)
{

```

```

#### Parameters
alpha <- parm[Data$pos.alpha]
beta <- matrix(c(parm[Data$pos.beta], rep(0,Data$K)),
  beta[,Data$L] <- -rowSums(beta[,~Data$L])
gamma <- matrix(c(parm[Data$pos.gamma],
  rep(0,Data$K)), Data$K, Data$M)
gamma[,Data$M] <- -rowSums(gamma[,~Data$M])
U <- as.parm.matrix(U, Data$K, parm, Data, chol=TRUE)
diag(U) <- exp(diag(U))
sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
parm[Data$pos.sigma] <- sigma
#### Log(Prior Densities)
alpha.prior <- sum(dnorm(alpha, 0, 1000, log=TRUE))
beta.prior <- sum(dnorm(beta, 0, sigma[1], log=TRUE))
gamma.prior <- sum(dnorm(gamma, 0, sigma[2], log=TRUE))
U.prior <- dwishartc(U, Data$K+1, Data$S, log=TRUE)
sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
#### Log-Likelihood
mu <- matrix(0,Data$N,Data$K)
for (k in 1:Data$K) {
  mu[,k] <- alpha[k] + beta[k,Data$X[,1]] + gamma[k,Data$X[,2]]}
LL <- sum(dmvnpc(Data$Y, mu, U, log=TRUE))
#### Variance Components, Omnibus
s.o.beta <- sd(as.vector(beta))
s.o.gamma <- sd(as.vector(gamma))
s.o.epsilon <- sd(as.vector(Data$Y - mu))
#### Variance Components, Univariate
s.beta <- apply(beta,1,sd)
s.gamma <- apply(gamma,1,sd)
s.epsilon <- apply(Data$Y - mu,2,sd)
#### Log-Posterior
LP <- LL + alpha.prior + beta.prior + gamma.prior + U.prior +
  sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP, s.o.beta, s.o.gamma,
  s.o.epsilon, s.beta, s.gamma, s.epsilon),
  yhat=rmvnpc(nrow(mu), mu, U), parm=parm)
return(Modelout)
}

```

## 57.4. Initial Values

```

Initial.Values <- c(rep(0,K), rep(0,K*(L-1)), rep(0,K*(M-1)),
  upper.triangle(S, diag=TRUE), rep(1,2))

```

## 58. Mixture Model, Finite

This finite mixture model (FMM) imposes a multilevel structure on each of the  $J$  regression effects in  $\beta$ , so that mixture components share a common residual standard deviation,  $\nu_j$ . Identifiability is gained at the expense of some shrinkage. The record-level mixture membership parameter vector,  $\theta$ , is a vector of discrete parameters. Discrete parameters are not supported in all algorithms. The example below is updated with the Griddy-Gibbs sampler.

### 58.1. Form

$$\begin{aligned} \mathbf{y} &\sim \mathcal{N}(\mu_{1:N,\theta}, \sigma^2) \\ \theta_i &\sim \mathcal{CAT}(\pi_{1:M}), \quad i = 1, \dots, N \\ \mu_{1:N,\theta} &= \mathbf{X}\beta_{\theta,1:J} \\ \beta_{m,j} &\sim \mathcal{N}(0, \nu_j^2), \quad j = 1, \dots, J, \quad m = 1, \dots, M \\ \nu_j &\sim \mathcal{HC}(25) \\ \sigma &\sim \mathcal{HC}(25) \\ \pi_{1:M} &\sim \mathcal{D}(\alpha_{1:M}) \\ \alpha_m &= 1 \end{aligned}$$

### 58.2. Data

```
M <- 2 #Number of mixtures
alpha <- rep(1,M) #Prior probability of mixing probabilities
data(demonsnacks)
N <- nrow(demonsnacks)
y <- log(demonsnacks$Calories)
X <- cbind(1, as.matrix(log(demonsnacks[,c(1,4,10)]+1)))
J <- ncol(X)
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
mon.names <- "LP"
parm.names <- as.parm.names(list(theta=rep(0,N), beta=matrix(0,M,J),
  nu=rep(0,J), sigma=0))
pos.theta <- grep("theta", parm.names)
pos.beta <- grep("beta", parm.names)
pos.nu <- grep("nu", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rcat(Data$N, rep(1/Data$M, Data$M)),
  rnorm(Data$M*Data$J,0,1), rhalfcauchy(Data$J+1,5)))
MyData <- list(J=J, M=M, N=N, PGF=PGF, X=X, alpha=alpha,
  mon.names=mon.names, parm.names=parm.names, pos.theta=pos.theta,
  pos.beta=pos.beta, pos.nu=pos.nu, pos.sigma=pos.sigma, y=y)
```

### 58.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  beta <- matrix(parm[Data$pos.beta], Data$M, Data$J)
  theta <- parm[Data$pos.theta]
  parm[Data$pos.nu] <- nu <- interval(parm[Data$pos.nu], 1e-100, Inf)
  pi <- rep(0, Data$M)
  tab <- table(theta)
  pi[as.numeric(names(tab))] <- as.vector(tab)
  pi <- pi / sum(pi)
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  beta.prior <- sum(dnorm(beta, 0, matrix(rep(nu, Data$M), Data$M,
    Data$J, byrow=TRUE), log=TRUE))
  theta.prior <- sum(dcat(theta, p=pi, log=TRUE))
  pi.prior <- ddirichlet(pi, Data$alpha, log=TRUE)
  nu.prior <- sum(dhalfcauchy(nu, 25, log=TRUE))
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, beta)
  mu <- diag(mu[,theta])
  LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + theta.prior + pi.prior + nu.prior +
    sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rnorm(length(mu), mu, sigma), parm=parm)
  return(Modelout)
}

```

### 58.4. Initial Values

```
Initial.Values <- c(rcat(N,rep(1/M,M)), rep(0,M*J), rep(1,J), 1)
```

## 59. Mixture Model, Infinite

This infinite mixture model (IMM) imposes a multilevel structure on each of the  $J$  regression effects in  $\beta$ , so that mixture components share a common residual standard deviation,  $\nu_j$ . The infinite number of mixture components is truncated to a finite number, and the user specifies the maximum number to explore,  $M$ , where  $M$  is discrete, greater than one, and less than the number of records,  $N$ . A truncated stick-breaking process within a truncated Dirichlet process defines the nonparametric mixture component selection. The record-level mixture membership parameter vector,  $\theta$ , is a vector of discrete parameters. Discrete parameters

are not supported in all algorithms. The example below is updated with the Griddy-Gibbs sampler.

### 59.1. Form

$$\begin{aligned}
 \mathbf{y} &\sim \mathcal{N}(\mu_{1:N,\theta}, \sigma^2) \\
 \mu_{1:N,\theta} &= \mathbf{X}\beta_{\theta,1:J} \\
 \theta_i &\sim \mathcal{CAT}(\pi_{1:M}) \\
 \beta_{m,j} &\sim \mathcal{N}(0, \nu_j^2), \quad j = 1, \dots, J, \quad m = 1, \dots, M \\
 \nu_j &\sim \mathcal{HC}(25) \\
 \sigma &\sim \mathcal{HC}(25) \\
 \pi &\sim \text{Stick}(\gamma) \\
 \alpha &\sim \mathcal{HC}(25) \\
 \iota &\sim \mathcal{HC}(25) \\
 \gamma &\sim \mathcal{G}(\alpha, \iota)
 \end{aligned}$$

### 59.2. Data

```

M <- 3 #Maximum number of mixtures to explore
data(demonsnacks)
N <- nrow(demonsnacks)
y <- log(demonsnacks$Calories)
X <- cbind(1, as.matrix(log(demonsnacks[,c(1,4,10)]+1)))
J <- ncol(X)
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
mon.names <- c("LP", as.parm.names(list(pi=rep(0,M))))
parm.names <- as.parm.names(list(theta=rep(0,N), beta=matrix(0,M,J),
  nu=rep(0,J), sigma=0, alpha=0, iota=0, gamma=0))
pos.theta <- grep("theta", parm.names)
pos.beta <- grep("beta", parm.names)
pos.nu <- grep("nu", parm.names)
pos.sigma <- grep("sigma", parm.names)
pos.alpha <- grep("alpha", parm.names)
pos.iota <- grep("iota", parm.names)
pos.gamma <- grep("gamma", parm.names)
PGF <- function(Data) return(c(rcat(Data$N, rep(1/Data$M,Data$M)),
  rnorm(Data$M*Data$J,0,1), rhalfcauchy(Data$J,5), rhalfcauchy(1,5),
  rhalfcauchy(2,5), rgamma(1,rhalfcauchy(1,25),rhalfcauchy(1,5))))
MyData <- list(J=J, M=M, N=N, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.theta=pos.theta, pos.beta=pos.beta,
  pos.nu=pos.nu, pos.sigma=pos.sigma, pos.alpha=pos.alpha,

```

```
pos.iota=pos.iota, pos.gamma=pos.gamma, y=y)
```

### 59.3. Model

```
Model <- function(parm, Data)
{
  ### Hyperhyperparameters
  alpha <- interval(parm[Data$pos.alpha], 1e-100, Inf)
  parm[Data$pos.alpha] <- alpha
  iota <- interval(parm[Data$pos.iota], 1e-100, Inf)
  parm[Data$pos.iota] <- iota
  ### Hyperparameters
  gamma <- interval(parm[Data$pos.gamma], 1e-100, Inf)
  parm[Data$pos.gamma] <- gamma
  parm[Data$pos.nu] <- nu <- interval(parm[Data$pos.nu], 1e-100, Inf)
  ### Parameters
  beta <- matrix(parm[Data$pos.beta], Data$M, Data$J)
  theta <- parm[Data$pos.theta]
  pi <- rStick(Data$M-1, gamma)
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Hyperhyperprior Densities)
  alpha.prior <- dhalfcauchy(alpha, 25, log=TRUE)
  iota.prior <- dhalfcauchy(iota, 25, log=TRUE)
  ### Log(Hyperprior Densities)
  gamma.prior <- dgamma(gamma, alpha, iota, log=TRUE)
  nu.prior <- sum(dhalfcauchy(nu, 25, log=TRUE))
  ### Log(Prior Densities)
  beta.prior <- sum(dnorm(beta, 0, matrix(rep(nu, Data$M), Data$M,
    Data$J, byrow=TRUE), log=TRUE))
  theta.prior <- sum(dcat(theta, pi, log=TRUE))
  pi.prior <- dStick(pi, gamma, log=TRUE)
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, beta)
  mu <- diag(mu[,theta])
  LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + theta.prior + pi.prior + nu.prior +
    sigma.prior + alpha.prior + iota.prior + gamma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,pi),
    yhat=rnorm(length(mu), mu, sigma), parm=parm)
  return(Modelout)
}
```

#### 59.4. Initial Values

```
Initial.Values <- c(rcat(N, rep(1/M,M)), rep(0,M*J), rep(1,J), rep(1,4))
```

## 60. Mixture Model, Poisson-Gamma

### 60.1. Form

$$\begin{aligned} \mathbf{y} &\sim \mathcal{P}(\lambda) \\ \lambda &\sim \mathcal{G}(\alpha\mu, \alpha) \\ \mu &= \exp(\mathbf{X}\beta) \\ \alpha &\sim \mathcal{HC}(25) \\ \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \end{aligned}$$

### 60.2. Data

```
N <- 20
J <- 3
X <- matrix(runif(N*J,-2,2),N,J); X[,1] <- 1
beta <- runif(J,-2,2)
y <- as.vector(round(exp(tcrossprod(X, t(beta)))))
mon.names <- "LP"
parm.names <- as.parm.names(list(alpha=0, beta=rep(0,J), lambda=rep(0,N)))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
pos.lambda <- grep("lambda", parm.names)
PGF <- function(Data) return(c(rhalfcauchy(1,5),
  rnormv(Data$J,0,1), rgamma(Data$N,
  exp(tcrossprod(Data$X, t(rnormv(Data$J,0,1))))*rhalfcauchy(1,5),
  rhalfcauchy(1,5)))
MyData <- list(J=J, N=N, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.alpha=pos.alpha, pos.beta=pos.beta,
  pos.lambda=pos.lambda, y=y)
```

### 60.3. Model

```
Model <- function(parm, Data)
{
  ### Hyperparameters
  alpha <- interval(parm[Data$pos.alpha], 1e-100, Inf)
  parm[Data$pos.alpha] <- alpha
  ### Parameters
```

```

beta <- parm[Data$pos.beta]
lambda <- interval(parm[Data$pos.lambda], 1e-100, Inf)
parm[Data$pos.lambda] <- lambda
mu <- exp(tcrossprod(Data$X, t(beta)))
### Log(Hyperprior Densities)
alpha.prior <- dhalfcauchy(alpha, 25, log=TRUE)
### Log(Prior Densities)
beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
lambda.prior <- sum(dgamma(lambda, alpha*mu, alpha, log=TRUE))
### Log-Likelihood
LL <- sum(dpois(Data$y, lambda, log=TRUE))
### Log-Posterior
LP <- LL + alpha.prior + beta.prior + lambda.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=rpois(length(lambda), lambda), parm=parm)
return(Modelout)
}

```

#### 60.4. Initial Values

```
Initial.Values <- c(1, rep(0,J), rep(1,N))
```

## 61. Multinomial Logit

### 61.1. Form

$$\mathbf{y}_i \sim \mathcal{CAT}(\mathbf{p}_{i,1:J}), \quad i = 1, \dots, N$$

$$\mathbf{p}_{i,j} = \frac{\phi_{i,j}}{\sum_{j=1}^J \phi_{i,j}}, \quad \sum_{j=1}^J \mathbf{p}_{i,j} = 1$$

$$\phi = \exp(\mu)$$

$$\mu_{i,J} = 0, \quad i = 1, \dots, N$$

$$\mu_{i,j} = \mathbf{X}_{i,1:K} \beta_{j,1:K} \in [-700, 700], \quad j = 1, \dots, (J-1)$$

$$\beta_{j,k} \sim \mathcal{N}(0, 1000), \quad j = 1, \dots, (J-1), \quad k = 1, \dots, K$$

### 61.2. Data

```

y <- x01 <- x02 <- c(1:300)
y[1:100] <- 1
y[101:200] <- 2
y[201:300] <- 3
x01[1:100] <- rnorm(100, 25, 2.5)

```



```
x01[101:200] <- rnorm(100, 40, 4.0)
x01[201:300] <- rnorm(100, 35, 3.5)
x02[1:100] <- rnorm(100, 2.51, 0.25)
x02[101:200] <- rnorm(100, 2.01, 0.20)
x02[201:300] <- rnorm(100, 2.70, 0.27)
N <- length(y)
J <- 3 #Number of categories in y
K <- 3 #Number of predictors (including the intercept)
X <- matrix(c(rep(1,N),x01,x02),N,K)
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=matrix(0,J-1,K)))
PGF <- function(Data) return(rnormv((Data$J-1)*Data$K,0,1000))
MyData <- list(J=J, K=K, N=N, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, y=y)
```

### 61.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- matrix(parm, Data$J-1, Data$K)
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  ### Log-Likelihood
  mu <- matrix(0, Data$N, Data$J)
  mu[,-Data$J] <- tcrossprod(Data$X, beta)
  mu <- interval(mu, -700, 700, reflect=FALSE)
  phi <- exp(mu)
  p <- phi / rowSums(phi)
  LL <- sum(dcat(Data$y, p, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP, yhat=rcat(nrow(p), p),
    parm=parm)
  return(Modelout)
}
```

### 61.4. Initial Values

```
Initial.Values <- c(rep(0,(J-1)*K))
```

## 62. Multinomial Logit, Nested

### 62.1. Form

$$\mathbf{y}_i \sim \mathcal{CAT}(\mathbf{P}_{i,1:J}), \quad i = 1, \dots, N$$

$$\mathbf{P}_{1:N,1} = \frac{\mathbf{R}}{\mathbf{R} + \exp(\alpha \mathbf{I})}$$

$$\mathbf{P}_{1:N,2} = \frac{(1 - \mathbf{P}_{1:N,1})\mathbf{S}_{1:N,1}}{\mathbf{V}}$$

$$\mathbf{P}_{1:N,3} = \frac{(1 - \mathbf{P}_{1:N,1})\mathbf{S}_{1:N,2}}{\mathbf{V}}$$

$$\mathbf{R}_{1:N} = \exp(\mu_{1:N,1})$$

$$\mathbf{S}_{1:N,1:2} = \exp(\mu_{1:N,2:3})$$

$$\mathbf{I} = \log(\mathbf{V})$$

$$\mathbf{V}_i = \sum_{k=1}^K \mathbf{S}_{i,k}, \quad i = 1, \dots, N$$

$$\mu_{1:N,1} = \mathbf{X}\boldsymbol{\iota} \in [-700, 700]$$

$$\mu_{1:N,2} = \mathbf{X}\boldsymbol{\beta}_{2,1:K} \in [-700, 700]$$

$$\boldsymbol{\iota} = \alpha \boldsymbol{\beta}_{1,1:K}$$

$$\alpha \sim \mathcal{EKP}(1) \in [0, 2]$$

$$\beta_{j,k} \sim \mathcal{N}(0, 1000), \quad j = 1, \dots, (J-1) \quad k = 1, \dots, K$$

where there are  $J = 3$  categories of  $\mathbf{y}$ ,  $K = 3$  predictors,  $\mathbf{R}$  is the non-nested alternative,  $\mathbf{S}$  is the nested alternative,  $\mathbf{V}$  is the observed utility in the nest,  $\alpha$  is effectively 1 - correlation and has a truncated exponential distribution, and  $\boldsymbol{\iota}$  is a vector of regression effects for the isolated alternative after  $\alpha$  is taken into account. The third alternative is the reference category.

### 62.2. Data

```

y <- x01 <- x02 <- c(1:300)
y[1:100] <- 1
y[101:200] <- 2
y[201:300] <- 3
x01[1:100] <- rnorm(100, 25, 2.5)
x01[101:200] <- rnorm(100, 40, 4.0)
x01[201:300] <- rnorm(100, 35, 3.5)
x02[1:100] <- rnorm(100, 2.51, 0.25)
x02[101:200] <- rnorm(100, 2.01, 0.20)
x02[201:300] <- rnorm(100, 2.70, 0.27)
N <- length(y)
J <- 3 #Number of categories in y
K <- 3 #Number of predictors (including the intercept)
X <- matrix(c(rep(1,N),x01,x02),N,K)
mon.names <- c("LP", as.parm.names(list(iota=rep(0,K))))

```

```

parm.names <- as.parm.names(list(alpha=0, beta=matrix(0,J-1,K)))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
PGF <- function(Data) return(c(rtrunc(1,"exp",a=0,b=2,rate=1),
  rnormv((Data$J-1)*Data$K,0,1)))
MyData <- list(J=J, K=K, N=N, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.alpha=pos.alpha, pos.beta=pos.beta, y=y)

```

### 62.3. Model

```

Model <- function(parm, Data)
{
  ### Hyperparameters
  alpha.rate <- 1
  ### Parameters
  parm[Data$pos.alpha] <- alpha <- interval(parm[Data$pos.alpha],0,2)
  beta <- matrix(parm[Data$pos.beta], Data$J-1, Data$K)
  ### Log(Prior Densities)
  alpha.prior <- dtrunc(alpha, "exp", a=0, b=2, rate=alpha.rate,
    log=TRUE)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  ### Log-Likelihood
  mu <- P <- matrix(0, Data$N, Data$J)
  iota <- alpha * beta[1,]
  mu[,1] <- tcrossprod(Data$X, t(iota))
  mu[,2] <- tcrossprod(Data$X, t(beta[2,]))
  mu <- interval(mu, -700, 700, reflect=FALSE)
  R <- exp(mu[,1])
  S <- exp(mu[, -1])
  V <- rowSums(S)
  I <- log(V)
  P[,1] <- R / (R + exp(alpha*I))
  P[,2] <- (1 - P[,1]) * S[,1] / V
  P[,3] <- (1 - P[,1]) * S[,2] / V
  LL <- sum(dcat(Data$y, P, log=TRUE))
  ### Log-Posterior
  LP <- LL + alpha.prior + beta.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,iota),
    yhat=rcat(nrow(P), P), parm=parm)
  return(Modelout)
}

```

### 62.4. Initial Values

```

Initial.Values <- c(0.5, rep(0.1,(J-1)*K))

```

## 63. Multinomial Probit

### 63.1. Form

$$\begin{aligned}\mathbf{W}_{i,1:(J-1)} &\sim \mathcal{N}_{J-1}(\mu_{i,1:(J-1)}, \Sigma), \quad i = 1, \dots, N \\ \mathbf{W}_{i,j} &\in \begin{cases} [0,10] & \text{if } \mathbf{y}_i = j \\ [-10,0] & \end{cases} \\ \mu_{1:N,j} &= \mathbf{X}\beta_{j,1:K} \\ \Sigma &= \mathbf{U}^T \mathbf{U} \\ \beta_{j,k} &\sim \mathcal{N}(0,10), \quad j = 1, \dots, (J-1), \quad k = 1, \dots, K \\ \mathbf{U}_{j,k} &\sim \mathcal{N}(0,1), \quad j = 1, \dots, (J-1), \quad k = 1, \dots, (J-1), \quad j \geq k, \quad j \neq k = 1\end{aligned}$$

### 63.2. Data

```

N <- 50
J <- 5 #Categories of y
K <- 8 #Number of columns in design matrix X
X <- matrix(runif(N*K,-2,2), N, K)
X[,1] <- 1
beta <- matrix(runif((J-1)*K), J-1, K)
mu <- tcrossprod(X, beta)
S <- diag(J-1)
u <- c(0, rnorm((J-2) + (factorial(J-1) /
  (factorial(J-1-2)*factorial(2))),0,1))
U <- diag(J-1)
U[upper.tri(U, diag=TRUE)] <- u
diag(U) <- exp(diag(U))
Sigma <- t(U) %*% U
Sigma[1,] <- Sigma[,1] <- U[1,]
mu <- tcrossprod(X, beta)
W <- rmvN(N, mu, Sigma) + matrix(rnorm(N*(J-1),0,0.1), N, J-1)
y <- max.col(cbind(W,0))
table(y)
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=matrix(0,(J-1),K),
  U=U, W=matrix(0,N,J-1)), uppertri=c(0,1,0))
parm.names <- parm.names[-which(parm.names == "U[1,1]")]
pos.beta <- grep("beta", parm.names)
pos.U <- grep("U", parm.names)
pos.W <- grep("W", parm.names)
PGF <- function(Data) {
  beta <- rnormv((Data$J-1)*Data$K,0,1)
  U <- rnorm((Data$J-2) + (factorial(Data$J-1) /
```

```

      (factorial(Data$J-1-2)*factorial(2))),0,1)
W <- matrix(runif(Data$N*(Data$J-1),-10,0), Data$N, Data$J-1)
Y <- as.indicator.matrix(Data$y)
W <- ifelse(Y[, -Data$J] == 1, abs(W), W)
return(c(beta, U, as.vector(W)))}
MyData <- list(J=J, K=K, N=N, PGF=PGF, S=S, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.U=pos.U, pos.W=pos.W,
  y=y)

```

### 63.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  beta <- matrix(parm[Data$pos.beta], Data$J-1, Data$K)
  u <- c(0, parm[Data$pos.U])
  U <- diag(Data$J-1)
  U[upper.tri(U, diag=TRUE)] <- u
  diag(U) <- exp(diag(U))
  Sigma <- t(U) %*% U
  Sigma[1,] <- Sigma[,1] <- U[1,]
  W <- matrix(parm[Data$pos.W], Data$N, Data$J-1)
  Y <- as.indicator.matrix(Data$y)
  temp <- which(Y[, -c(Data$J)] == 1)
  W[temp] <- interval(W[temp], 0, 10)
  temp <- which(Y[, -c(Data$J)] == 0)
  W[temp] <- interval(W[temp], -10, 0)
  parm[Data$pos.W] <- as.vector(W)
  ### Log(Prior Densities)
  beta.prior <- sum(dnorm(beta, 0, 10, log=TRUE))
  U.prior <- sum(dnorm(u[-length(u)], 0, 1, log=TRUE))
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, beta)
  #eta <- exp(cbind(mu,0))
  #p <- eta / rowSums(eta)
  LL <- sum(dmvn(W, mu, Sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + U.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=max.col(cbind(rmvn(nrow(mu), mu, Sigma),0)), parm=parm)
  return(Modelout)
}

```

### 63.4. Initial Values

```
Initial.Values <- GIV(Model, MyData, PGF=TRUE)
```

## 64. Multiple Discrete-Continuous Choice

This form of a multivariate discrete-continuous choice model was introduced in [Kim, Allenby, and Rossi \(2002\)](#) and referred to as a variety model. The original version is presented with log-normally distributed errors, but a gamma regression form is used here instead, which has always mixed better in testing. Note that the  $\gamma$  parameters are fixed here, as recommended for identifiability in future articles by these authors.

### 64.1. Form

$$\begin{aligned} \mathbf{Y} &\sim \mathcal{G}(\lambda\tau, \tau) \\ \lambda_{i,j} &= \exp(\mathbf{Z}_{i,j} \log(\psi 1_{m[i],j}) + \mathbf{X} \mathbf{1}_{1:K} \log(\beta) + \mathbf{X} \mathbf{2}_{1:L} \log(\delta)) (\mathbf{Y}_{i,j} + \gamma_j)^\alpha_j, \quad i = 1, \dots, N, j = 1, \dots, J \\ \alpha_j &\sim \mathcal{U}(0, 1), \quad j = 1, \dots, J \\ \log(\beta_k) &\sim \mathcal{N}(0, 1000), \quad k = 1, \dots, K \\ \gamma_j &= 1, \quad j = 1, \dots, J \\ \log(\delta_{j,l}) &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, (J-1), \quad l = 1, \dots, L \\ \log(\psi 0_j) &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\ \log(\psi 1_{g,j}) &\sim \mathcal{N}_J(\log(\psi 0), \Omega^{-1}), \quad g = 1, \dots, G, \quad j = 1, \dots, J \\ \Omega &\sim \mathcal{W}_{J+1}(\mathbf{S}), \quad \mathbf{S} = \mathbf{I}_J \\ \tau_j &\sim \mathcal{HC}(25), \quad j = 1, \dots, J \end{aligned}$$

### 64.2. Data

```
G <- 6 #Number of Multilevel Groups (decision-makers, households, etc.)
J <- 3 #Number of products
K <- 4 #Number of product attributes
L <- 5 #Number of decision-maker attributes
N <- 30 #Number of records
X1 <- matrix(rnorm(N*K), N, K) #Product attributes
X2 <- matrix(rnorm(N*L), N, L) #Decision-maker attributes
Sigma <- matrix(runif((J-1)*(J-1), -1, 1), J-1, J-1)
diag(Sigma) <- runif(J-1, 1, 5)
Sigma <- as.positive.definite(Sigma) / 100
alpha <- runif(J)
log.beta <- rnorm(K, 0, 0.1)
log.delta <- matrix(rnorm((J-1)*L, 0, 0.1), J-1, L)
log.psi0 <- rnorm(J)
log.psi1 <- rmvn(G, log.psi0, Sigma)
```

```

m <- rcat(N, rep(1/G,G)) # Multilevel group indicator
Z <- as.indicator.matrix(m)
Y <- matrix(0, N, J)
Y <- round(exp(tcrossprod(Z, t(cbind(log.psi1,0)))) +
  matrix(tcrossprod(X1, t(log.beta)), N, J) +
  tcrossprod(X2, rbind(log.delta, colSums(log.delta)*-1))) *
  (Y + 1)^matrix(alpha,N,J,byrow=TRUE) +
  matrix(rnorm(N*J,0,0.1),N,J))
S <- diag(J)
mon.names <- "LP"
parm.names <- as.parm.names(list(alpha=rep(0,J), log.beta=rep(0,K),
  log.delta=matrix(0,J-1,L), log.psi0=rep(0,J),
  log.psi1=matrix(0,G,J), tau=rep(0,J), U=S),
  uppertri=c(0,0,0,0,0,0,1))
pos.alpha <- grep("alpha", parm.names)
pos.log.beta <- grep("log.beta", parm.names)
pos.log.delta <- grep("delta", parm.names)
pos.log.psi0 <- grep("log.psi0", parm.names)
pos.log.psi1 <- grep("log.psi1", parm.names)
pos.tau <- grep("tau", parm.names)
PGF <- function(Data) {
  log.psi0 <- rnorm(Data$J,0,0.1)
  U <- rwishartc(Data$J+1, Data$S)
  return(c(runif(Data$J,0.9,1), rnorm(Data$K,0,0.1),
    rnorm((Data$J-1)*Data$L,0,0.1), log.psi0, rmvnp(c(Data$G, log.psi0, U),
    runif(Data$J), upper.triangle(U, diag=TRUE))))}
MyData <- list(G=G, J=J, K=K, L=L, N=N, PGF=PGF, S=S, X1=X1, X2=X2, Y=Y,
  Z=Z, m=m, mon.names=mon.names, parm.names=parm.names,
  pos.alpha=pos.alpha, pos.log.beta=pos.log.beta,
  pos.log.delta=pos.log.delta, pos.log.psi0=pos.log.psi0,
  pos.log.psi1=pos.log.psi1, pos.tau=pos.tau)

```

### 64.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  parm[Data$pos.alpha] <- alpha <- interval(parm[Data$pos.alpha], 0, 1)
  log.beta <- parm[Data$pos.log.beta]
  log.delta <- matrix(parm[Data$pos.log.delta], Data$J-1, Data$L)
  log.psi0 <- parm[Data$pos.log.psi0]
  log.psi1 <- matrix(parm[Data$pos.log.psi1], Data$G, Data$J)
  parm[Data$pos.tau] <- tau <- interval(parm[Data$pos.tau], 1e-100, Inf)
  U <- as.parm.matrix(U, Data$J, parm, Data, chol=TRUE)
  diag(U) <- exp(diag(U))
  lambda <- tcrossprod(Data$Z, t(log.psi1)) +

```

```

      matrix(tcrossprod(Data$X1, t(log.beta)), Data$N, Data$J) +
      tcrossprod(Data$X2, rbind(log.delta, colSums(log.delta)*-1))
### Log(Prior Densities)
U.prior <- dwishartc(U, Data$J+1, Data$S, log=TRUE)
alpha.prior <- sum(dunif(alpha, 0, 1, log=TRUE))
log.beta.prior <- sum(dnormv(log.beta, 0, 1000, log=TRUE))
log.delta.prior <- sum(dnormv(log.delta, 0, 1000, log=TRUE))
log.psi0.prior <- sum(dnormv(log.psi0, 0, 1000, log=TRUE))
log.psi1.prior <- sum(dmvnpc(lambda,
      matrix(log.psi0, Data$N, Data$J, byrow=TRUE), U, log=TRUE))
tau.prior <- sum(dhalfcauchy(tau, 25, log=TRUE))
### Log-Likelihood
alpha <- matrix(alpha, Data$N, Data$J, byrow=TRUE)
lambda <- exp(lambda)*(Data$Y + 1)^alpha
tau <- matrix(tau, Data$N, Data$J, byrow=TRUE)
LL <- sum(dgamma(Data$Y+1, lambda*tau, tau, log=TRUE))
### Log-Posterior
LP <- LL + U.prior + alpha.prior + log.beta.prior + log.delta.prior +
      log.psi0.prior + log.psi1.prior + tau.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
      yhat=rgamma(prod(dim(lambda)), lambda*tau, tau)-1,
      parm=parm)
return(Modelout)
}

```

#### 64.4. Initial Values

```

Initial.Values <- c(runif(J,0.9,1), rnorm(K,0,0.1),
      rnorm((J-1)*L,0,0.1), rnorm(J,0,0.1),
      rmvnpc(G, rnorm(J,0,0.1), rwishartc(J+1,S)), runif(J),
      upper.triangle(rwishartc(J+1,S), diag=TRUE))

```

### 65. Multivariate Binary Probit

#### 65.1. Form

$$\begin{aligned}
 \mathbf{Z}_{i,1:J} &\sim \mathcal{N}_J(\mu_{i,1:J}, \Sigma), \quad i = 1, \dots, N \\
 \mathbf{Z}_{i,j} &\in \begin{cases} [0,10] & \text{if } \mathbf{y}_i = j \\ [-10,0] & \end{cases} \\
 \mu_{1:N,j} &= \mathbf{X}\beta_{j,1:K} \\
 \Sigma &\sim \mathcal{IW}_{J+1}(\mathbf{S}^{-1}), \quad \mathbf{S} = \mathbf{I}_J, \quad \Sigma[1,1] = 1 \\
 \beta_{j,k} &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, (J-1), \quad k = 1, \dots, K
 \end{aligned}$$



$$\beta_{J,k} = - \sum_{j=1}^{J-1} \beta_{j,k}$$

$$\mathbf{Z}_{i,j} \sim \mathcal{N}(0, 1000) \in [-10, 10]$$

## 65.2. Data

```

N <- 30
J <- 2 #Number of binary dependent variables
K <- 3 #Number of columns to be in design matrix X
X <- matrix(1, N, K)
for (k in 2:K) {X[,k] <- rnorm(N, runif(1,-3,3), runif(1,0.1,3))}
beta <- matrix(runif(J*K), J, K)
mu <- tcrossprod(X, beta)
S <- diag(J)
u <- c(0, rnorm((J-1) + (factorial(J) /
  (factorial(J-2)*factorial(2))),0,1))
U <- diag(J)
U[upper.tri(U, diag=TRUE)] <- u
diag(U) <- exp(diag(U))
Sigma <- t(U) %*% U
Sigma[1,] <- Sigma[,1] <- U[1,]
W <- rmvn(N, mu, Sigma) + matrix(rnorm(N*J,0,0.1), N, J)
Y <- 1 * (W >= 0)
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=matrix(0,J,K),
  U=U, W=matrix(0,N,J)), uppertri=c(0,1,0))
parm.names <- parm.names[-which(parm.names == "U[1,1]")]
pos.beta <- grep("beta", parm.names)
pos.U <- grep("U", parm.names)
pos.W <- grep("W", parm.names)
PGF <- function(Data) {
  beta <- rnormv(Data$J*Data$K,0,1)
  U <- rnorm((Data$J-1) + (factorial(Data$J) /
    (factorial(Data$J-2)*factorial(2))),0,1)
  W <- matrix(runif(Data$N*Data$J,-10,0), Data$N, Data$J)
  W <- ifelse(Y == 1, abs(W), W)
  return(c(beta, U, as.vector(W)))}
MyData <- list(J=J, K=K, N=N, PGF=PGF, S=S, X=X, Y=Y,
  mon.names=mon.names, parm.names=parm.names, pos.beta=pos.beta,
  pos.U=pos.U, pos.W=pos.W)

```

## 65.3. Model

```

Model <- function(parm, Data)
{

```

```

#### Parameters
beta <- matrix(parm[Data$pos.beta], Data$J, Data$K)
u <- c(0, parm[Data$pos.U])
U <- diag(Data$J)
U[upper.tri(U, diag=TRUE)] <- u
diag(U) <- exp(diag(U))
Sigma <- t(U) %*% U
Sigma[1,] <- Sigma[,1] <- U[1,]
W <- matrix(parm[Data$pos.W], Data$N, Data$J)
W[Data$Y == 0] <- interval(W[Data$Y == 0], -10, 0)
W[Data$Y == 1] <- interval(W[Data$Y == 1], 0, 10)
parm[Data$pos.W] <- as.vector(W)
#### Log(Prior Densities)
beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
U.prior <- sum(dnorm(u[-length(u)], 0, 1, log=TRUE))
#### Log-Likelihood
mu <- tcrossprod(Data$X, beta)
LL <- sum(dmvn(W, mu, Sigma, log=TRUE))
#### Log-Posterior
LP <- LL + beta.prior + U.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=1*(rmvn(prod(nrow(mu)), mu, Sigma) >= 0), parm=parm)
return(Modelout)
}

```

## 65.4. Initial Values

```
Initial.Values <- GIV(Model, MyData, PGF=TRUE)
```

# 66. Multivariate Laplace Regression

## 66.1. Form

$$\begin{aligned}
 \mathbf{Y}_{i,k} &\sim \mathcal{L}_K(\mu_{i,k}, \Sigma), \quad i = 1, \dots, N; \quad k = 1, \dots, K \\
 \mu_{i,k} &= \mathbf{X}_{1:N,k} \beta_{k,1:J} \\
 \Sigma &= \Omega^{-1} \\
 \Omega &\sim \mathcal{W}_{K+1}(\mathbf{S}), \quad \mathbf{S} = \mathbf{I}_K \\
 \beta_{k,j} &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J
 \end{aligned}$$

## 66.2. Data

```

N <- 100
J <- 6 #Number of columns in design matrix

```

```

K <- 3 #Number of DVs
X <- matrix(runif(N*J),N,J); X[,1] <- 1
Y <- mu <- tcrossprod(X, matrix(rnorm(J*K),K,J))
Sigma <- matrix(runif(K*K),K,K); diag(Sigma) <- runif(K,1,K)
Sigma <- as.symmetric.matrix(Sigma)
for (i in 1:N) {Y[i,] <- colMeans(rmvn(1000, mu[i,], Sigma))}
S <- diag(K)
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=matrix(0,K,J), U=diag(K)),
  uppertri=c(0,1))
pos.beta <- grep("beta", parm.names)
PGF <- function(Data) return(c(rnormv(Data$K*Data$J,0,1),
  upper.triangle(rwishartc(Data$K+1,Data$S), diag=TRUE)))
MyData <- list(J=J, K=K, N=N, PGF=PGF, S=S, X=X, Y=Y, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta)

```

### 66.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  beta <- matrix(parm[Data$pos.beta], Data$K, Data$J)
  U <- as.parm.matrix(U, Data$K, parm, Data, chol=TRUE)
  diag(U) <- exp(diag(U))
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  U.prior <- dwishart(U, Data$K+1, Data$S, log=TRUE)
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, beta)
  LL <- sum(dmvlc(Data$Y, mu, U, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + U.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rmvnc(nrow(mu), mu, U), parm=parm)
  return(Modelout)
}

```

### 66.4. Initial Values

```

Initial.Values <- c(rep(0,J*K), upper.triangle(S, diag=TRUE))

```

## 67. Multivariate Regression

### 67.1. Form

$$\begin{aligned} \mathbf{Y}_{i,k} &\sim \mathcal{N}_K(\mu_{i,k}, \Omega^{-1}), \quad i = 1, \dots, N; \quad k = 1, \dots, K \\ \mu_{i,k} &= \mathbf{X}_{1:N,k} \beta_{k,1:J} \\ \Omega &\sim \mathcal{W}_{K+1}(\mathbf{S}), \quad \mathbf{S} = \mathbf{I}_K \\ \beta_{k,j} &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \end{aligned}$$

### 67.2. Data

```

N <- 100
J <- 6 #Number of columns in design matrix
K <- 3 #Number of DVs
X <- matrix(runif(N*J),N,J); X[,1] <- 1
Y <- mu <- tcrossprod(X, matrix(rnorm(J*K),K,J))
Sigma <- matrix(runif(K*K),K,K); diag(Sigma) <- runif(K,1,K)
Sigma <- as.symmetric.matrix(Sigma)
for (i in 1:N) {Y[i,] <- colMeans(rmvn(1000, mu[i,], Sigma))}
S <- diag(K)
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=matrix(0,K,J), U=diag(K)),
  uppertri=c(0,1))
pos.beta <- grep("beta", parm.names)
PGF <- function(Data) return(c(rnormv(Data$K*Data$J,0,1),
  upper.triangle(rwishartc(Data$K+1,Data$S), diag=TRUE)))
MyData <- list(J=J, K=K, N=N, PGF=PGF, S=S, X=X, Y=Y, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta)

```

### 67.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  beta <- matrix(parm[Data$pos.beta], Data$K, Data$J)
  U <- as.parm.matrix(U, Data$K, parm, Data, chol=TRUE)
  diag(U) <- exp(diag(U))
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  U.prior <- dwishartc(U, Data$K+1, Data$S, log=TRUE)
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, beta)
  LL <- sum(dmvnpc(Data$Y, mu, U, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + U.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,

```

```

        yhat=rmvnp(nrow(mu), mu, U), parm=parm)
    return(Modelout)
}

```

#### 67.4. Initial Values

```
Initial.Values <- c(rep(0,J*K), upper.triangle(S, diag=TRUE))
```

## 68. Negative Binomial Regression

This example was contributed by Jim Robison-Cox.

### 68.1. Form

$$\begin{aligned}
 \mathbf{y} &\sim \mathcal{NB}(\mu, \kappa) \\
 p &= \frac{\kappa}{\kappa + \mu} \\
 \mu &= \exp(\mathbf{X}\beta) \\
 \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\
 \kappa &\sim \mathcal{HC}(25) \in (0, \infty]
 \end{aligned}$$

### 68.2. Data

```

N <- 100
J <- 5 #Number of predictors, including the intercept
kappa.orig <- 2
beta.orig <- runif(J,-2,2)
X <- matrix(runif(J*N,-2, 2), N, J); X[,1] <- 1
mu <- exp(tcrossprod(X, t(beta.orig)) + rnorm(N))
p <- kappa.orig / (kappa.orig + mu)
y <- rnbinom(N, size=kappa.orig, mu=mu)
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J), kappa=0))
pos.beta <- grep("beta", parm.names)
pos.kappa <- grep("kappa", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,1000), rhalfcauchy(1,5)))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
              parm.names=parm.names, pos.beta=pos.beta, pos.kappa=pos.kappa, y=y)

```

### 68.3. Model

```

Model <- function(parm, Data)
{

```

```

#### Parameters
beta <- parm[Data$pos.beta]
parm[Data$J + 1] <- kappa <- interval(parm[Data$pos.kappa],
  .Machine$double.xmin, Inf)
#### Log(Prior Densities)
beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
kappa.prior <- dhalfcauchy(kappa, 25, log=TRUE)
#### Log-Likelihood
mu <- as.vector(exp(tcrossprod(Data$X, t(beta))))
#p <- kappa / (kappa + mu)
LL <- sum(dnbinom(Data$y, size=kappa, mu=mu, log=TRUE))
#### Log-Posterior
LP <- LL + beta.prior + kappa.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=rnbinom(length(mu), size=kappa, mu=mu), parm=parm)
return(Modelout)
}

```

#### 68.4. Initial Values

```
Initial.Values <- c(rep(0,J), 1)
```

## 69. Normal, Multilevel

This is Gelman's school example (Gelman, Carlin, Stern, and Rubin 2004). Note that **LaplacesDemon** is slower to converge than WinBUGS through the **R2WinBUGS** package (Gelman 2013), an R package on CRAN. This example is very sensitive to the prior distributions. The recommended, default, half-Cauchy priors with scale 25 on scale parameters overwhelms the likelihood, so uniform priors are used.

### 69.1. Form

$$\begin{aligned}
 \mathbf{y}_j &\sim \mathcal{N}(\theta_j, \sigma_j^2), \quad j = 1, \dots, J \\
 \theta_j &\sim \mathcal{N}(\theta_\mu, \theta_\sigma^2) \\
 \theta_\mu &\sim \mathcal{N}(0, 1000000) \\
 \theta_{\sigma[j]} &\sim \mathcal{N}(0, 1000) \\
 \sigma &\sim \mathcal{U}(0, 1000)
 \end{aligned}$$

### 69.2. Data

```

J <- 8
y <- c(28.4, 7.9, -2.8, 6.8, -0.6, 0.6, 18.0, 12.2)

```

```

sd <- c(14.9, 10.2, 16.3, 11.0, 9.4, 11.4, 10.4, 17.6)
mon.names <- "LP"
parm.names <- as.parm.names(list(theta=rep(0,J), theta.mu=0,
  theta.sigma=0))
pos.theta <- 1:J
pos.theta.mu <- J+1
pos.theta.sigma <- J+2
PGF <- function(Data) return(c(rnorm(Data$J, rnormp(1,0,1E-6),
  runif(1,0,10)), rnormp(1,0,1E-6), runif(1,0,10)))
MyData <- list(J=J, PGF=PGF, mon.names=mon.names, parm.names=parm.names,
  pos.theta=pos.theta, pos.theta.mu=pos.theta.mu,
  pos.theta.sigma=pos.theta.sigma, sd=sd, y=y)

```

### 69.3. Model

```

Model <- function(parm, Data)
{
  ### Hyperparameters
  theta.mu <- parm[Data$pos.theta.mu]
  theta.sigma <- interval(parm[Data$pos.theta.sigma], 1e-100, Inf)
  parm[Data$pos.theta.sigma] <- theta.sigma
  ### Parameters
  theta <- parm[Data$pos.theta]
  ### Log(Hyperprior Densities)
  theta.mu.prior <- dnormp(theta.mu, 0, 1.0E-6, log=TRUE)
  theta.sigma.prior <- dunif(theta.sigma, 0, 1000, log=TRUE)
  ### Log(Prior Densities)
  theta.prior <- sum(dnorm(theta, theta.mu, theta.sigma, log=TRUE))
  sigma.prior <- sum(dunif(Data$sd, 0, 1000, log=TRUE))
  ### Log-Likelihood
  LL <- sum(dnorm(Data$y, theta, Data$sd, log=TRUE))
  ### Log-Posterior
  LP <- LL + theta.prior + theta.mu.prior + theta.sigma.prior +
    sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rnorm(length(theta), theta, Data$sd), parm=parm)
  return(Modelout)
}

```

### 69.4. Initial Values

```

Initial.Values <- c(rep(mean(y),J), mean(y), 1)

```

## 70. Ordinal Logit

### 70.1. Form

$$\begin{aligned}
 \mathbf{y}_i &\sim \mathcal{CAT}(P_{i,1:J}) \\
 P_{,J} &= 1 - Q_{,(J-1)} \\
 P_{,j} &= |Q_{,j} - Q_{,(j-1)}|, \quad j = 2, \dots, (J-1) \\
 P_{,1} &= Q_{,1} \\
 Q &= \frac{1}{1 + \exp(\mu)} \\
 \mu_{,j} &= \delta_j - \mathbf{X}\beta, \quad \in [-5, 5] \\
 \beta_k &\sim \mathcal{N}(0, 1000), \quad k = 1, \dots, K \\
 \delta_j &\sim \mathcal{N}(0, 1) \in [(j-1), j] \in [-5, 5], \quad j = 1, \dots, (J-1)
 \end{aligned}$$

### 70.2. Data

```

data(demonsnacks)
N <- nrow(demonsnacks)
J <- 3 #Number of categories in y
X <- as.matrix(demonsnacks[,c(1,3:10)])
K <- ncol(demonsnacks) #Number of columns in design matrix X
y <- log(demonsnacks$Calories)
y <- ifelse(y < 4.5669, 1, ifelse(y > 5.5268, 3, 2)) #Discretize
for (k in 1:K) {X[,k] <- CenterScale(X[,k])}
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,K), delta=rep(0,J-1)))
pos.beta <- grep("beta", parm.names)
pos.delta <- grep("delta", parm.names)
PGF <- function(Data)
{
  delta <- rnorm(Data$J-1,0,1)
  delta <- delta[order(delta)]
  return(c(rnormmv(Data$K,0,10), delta))
}
MyData <- list(J=J, K=K, N=N, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.delta=pos.delta, y=y)

```

### 70.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  delta <- interval(parm[Data$pos.delta], -5, 5)

```



```

delta <- delta[order(delta)]
parm[Data$pos.delta] <- delta
### Log(Prior Densities)
beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
delta.prior <- sum(dtrunc(delta, "norm", a=-5, b=5, log=TRUE,
  mean=0, sd=1)
### Log-Likelihood
mu <- matrix(delta, Data$N, Data$J-1, byrow=TRUE) -
  matrix(tcrossprod(Data$X, t(beta)), Data$N, Data$J-1)
P <- Q <- invlogit(mu)
P[, -1] <- abs(Q[, -1] - Q[, -(Data$J-1)])
P <- cbind(P, 1 - Q[, (Data$J-1)])
LL <- sum(dcat(Data$y, P, log=TRUE))
### Log-Posterior
LP <- LL + beta.prior + delta.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP, yhat=rcat(nrow(P), P)
  parm=parm)
return(Modelout)
}

```

## 70.4. Initial Values

```
Initial.Values <- c(rep(0,K), seq(from=-1, to=1, len=(J-1)))
```

## 71. Ordinal Probit

### 71.1. Form

$$\begin{aligned}
 \mathbf{y}_i &\sim \mathcal{CAT}(P_{i,1:J}) \\
 P_{i,J} &= 1 - Q_{i,(J-1)} \\
 P_{i,j} &= |Q_{i,j} - Q_{i,(j-1)}|, \quad j = 2, \dots, (J-1) \\
 P_{i,1} &= Q_{i,1} \\
 Q &= \phi(\mu) \\
 \mu_{i,j} &= \delta_j - \mathbf{X}_i \beta, \quad \in [-5, 5] \\
 \beta_k &\sim \mathcal{N}(0, 1000), \quad k = 1, \dots, K \\
 \delta_j &\sim \mathcal{N}(0, 1) \in [(j-1), j] \in [-5, 5], \quad j = 1, \dots, (J-1)
 \end{aligned}$$

### 71.2. Data

```

data(demonsnacks)
N <- nrow(demonsnacks)

```

```

J <- 3 #Number of categories in y
X <- as.matrix(demonsnacks[,c(1,3:10)])
K <- ncol(demonsnacks) #Number of columns in design matrix X
y <- log(demonsnacks$Calories)
y <- ifelse(y < 4.5669, 1, ifelse(y > 5.5268, 3, 2)) #Discretize
for (k in 1:K) {X[,k] <- CenterScale(X[,k])}
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,K), delta=rep(0,J-1)))
pos.beta <- grep("beta", parm.names)
pos.delta <- grep("delta", parm.names)
PGF <- function(Data)
{
  delta <- rnorm(Data$J-1,0,1)
  delta <- delta[order(delta)]
  return(c(rnormv(Data$K,0,10), delta))
}
MyData <- list(J=J, K=K, N=N, PGF=PGF, X=X, mon.names=mon.names,
parm.names=parm.names, pos.beta=pos.beta, pos.delta=pos.delta, y=y)

```

### 71.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  delta <- interval(parm[Data$pos.delta], -5, 5)
  delta <- delta[order(delta)]
  parm[Data$pos.delta] <- delta
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  delta.prior <- sum(dtrunc(delta, "norm", a=-5, b=5, log=TRUE,
    mean=0, sd=1))
  ### Log-Likelihood
  mu <- matrix(delta, Data$N, Data$J-1, byrow=TRUE) -
    matrix(tcrossprod(Data$X, t(beta)), Data$N, Data$J-1)
  P <- Q <- pnorm(mu)
  P[, -1] <- abs(Q[, -1] - Q[, -(Data$J-1)])
  P <- cbind(P, 1 - Q[, (Data$J-1)])
  LL <- sum(dcat(Data$y, P, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + delta.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP, yhat=rcat(nrow(P), P)
    parm=parm)
  return(Modelout)
}

```

### 71.4. Initial Values

```
Initial.Values <- c(rep(0,K), seq(from=-1, to=1, len=(J-1)))
```

## 72. Panel, Autoregressive Poisson

### 72.1. Form

$$\begin{aligned}
 \mathbf{Y} &\sim \mathcal{P}(\Lambda) \\
 \Lambda_{1:N,1} &= \exp(\alpha + \beta \mathbf{x}) \\
 \Lambda_{1:N,t} &= \exp(\alpha + \beta \mathbf{x} + \rho \log(\mathbf{Y}_{1:N,t-1})), \quad t = 2, \dots, T \\
 \alpha_i &\sim \mathcal{N}(\alpha_\mu, \alpha_\sigma^2), \quad i = 1, \dots, N \\
 \alpha_\mu &\sim \mathcal{N}(0, 1000) \\
 \alpha_\sigma &\sim \mathcal{HC}(25) \\
 \beta &\sim \mathcal{N}(0, 1000) \\
 \rho &\sim \mathcal{N}(0, 1000)
 \end{aligned}$$

### 72.2. Data

```

N <- 10
T <- 10
alpha <- rnorm(N,2,0.5)
rho <- 0.5
beta <- 0.5
x <- runif(N,0,1)
Y <- matrix(NA,N,T)
Y[,1] <- exp(alpha + beta*x)
for (t in 2:T) {Y[,t] <- exp(alpha + beta*x + rho*log(Y[,t-1]))}
Y <- round(Y)
mon.names <- "LP"
parm.names <- as.parm.names(list(alpha=rep(0,N), alpha.mu=0,
  alpha.sigma=0, beta=0, rho=0))
pos.alpha <- 1:N
pos.alpha.mu <- grep("alpha.mu", parm.names)
pos.alpha.sigma <- grep("alpha.sigma", parm.names)
pos.beta <- grep("beta", parm.names)
pos.rho <- grep("rho", parm.names)
PGF <- function(Data) return(c(rnorm(Data$N, rnormv(1,0,10),
  rhalfcauchy(1,5)), rnormv(1,0,10), rhalfcauchy(1,5),
  rnormv(2,0,10)))
MyData <- list(N=N, PGF=PGF, T=T, Y=Y, mon.names=mon.names,
  parm.names=parm.names, pos.alpha=pos.alpha, pos.alpha.mu=pos.alpha.mu,

```

```
pos.alpha.sigma=pos.alpha.sigma, pos.beta=pos.beta, pos.rho=pos.rho,
x=x)
```

### 72.3. Model

```
Model <- function(parm, Data)
{
  ### Hyperparameters
  alpha.mu <- parm[Data$pos.alpha.mu]
  alpha.sigma <- interval(parm[Data$pos.alpha.sigma], 1e-100, Inf)
  parm[Data$pos.alpha.sigma] <- alpha.sigma
  ### Parameters
  alpha <- parm[Data$pos.alpha]
  beta <- parm[Data$pos.beta]
  rho <- parm[Data$pos.rho]
  ### Log(Hyperprior Densities)
  alpha.mu.prior <- dnormv(alpha.mu, 0, 1000, log=TRUE)
  alpha.sigma.prior <- dhalfcauchy(alpha.sigma, 25, log=TRUE)
  ### Log(Prior Densities)
  alpha.prior <- sum(dnorm(alpha, alpha.mu, alpha.sigma, log=TRUE))
  beta.prior <- dnormv(beta, 0, 1000, log=TRUE)
  rho.prior <- dnormv(rho, 0, 1000, log=TRUE)
  ### Log-Likelihood
  Lambda <- Data$Y
  Lambda[,1] <- exp(alpha + beta*x)
  Lambda[,2:Data$T] <- exp(alpha + beta*Data$x +
    rho*log(Data$Y[,1:(Data$T-1)]))
  LL <- sum(dpois(Data$Y, Lambda, log=TRUE))
  ### Log-Posterior
  LP <- LL + alpha.prior + alpha.mu.prior + alpha.sigma.prior +
    beta.prior + rho.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rpois(prod(dim(Lambda)), parm=parm)
  return(Modelout)
}
```

### 72.4. Initial Values

```
Initial.Values <- c(rep(0,N), 0, 1, 0, 0)
```

## 73. Penalized Spline Regression

This example applies penalized splines to one predictor in a linear regression. The user selects the degree of the polynomial,  $D$ , and the number of knots,  $K$ .

### 73.1. Form

$$\begin{aligned} \mathbf{y} &\sim \mathcal{N}(\mu, \sigma_1^2) \\ \mu &= \mathbf{X}\beta + \mathbf{S} \\ \mathbf{S} &= \mathbf{Z}\gamma \\ \mathbf{Z}_{i,k} &= \begin{cases} (\mathbf{x}_i - k)^D & \text{if } \mathbf{Z}_{i,k} > 0 \\ 0 & \text{otherwise} \end{cases} \\ \beta_d &\sim \mathcal{N}(0, 1000), \quad d = 1, \dots, (D+1) \\ \gamma_k &\sim \mathcal{N}(0, \sigma_2^2), \quad k = 1, \dots, K \\ \sigma_j &\sim \mathcal{HC}(25), \quad j = 1, \dots, 2 \end{aligned}$$

### 73.2. Data

```

N <- 100
x <- 1:N
y <- sin(2*pi*x/N) + runif(N,-1,1)
K <- 10 #Number of knots
D <- 2 #Degree of polynomial
x <- CenterScale(x)
k <- as.vector(quantile(x, probs=(1:K / (K+1))))
X <- cbind(1, matrix(x, N, D))
for (d in 1:D) {X[,d+1] <- X[,d+1]^d}
Z <- matrix(x, N, K) - matrix(k, N, K, byrow=TRUE)
Z <- ifelse(Z > 0, Z, 0); Z <- Z^D
mon.names <- c("LP", paste("S[", 1:nrow(X) ,"]", sep=""))
parm.names <- as.parm.names(list(beta=rep(0,1+D), gamma=rep(0,K),
  log.sigma=rep(0,2)))
pos.beta <- grep("beta", parm.names)
pos.gamma <- grep("gamma", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(1+Data$D,0,10), rnorm(Data$K,0,10),
  rhalfcauchy(2,5)))
MyData <- list(D=D, K=K, N=N, PGF=PGF, Z=Z, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.gamma=pos.gamma,
  pos.sigma=pos.sigma, y=y)

```

### 73.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  gamma <- parm[Data$pos.gamma]

```

```

sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
parm[Data$pos.sigma] <- sigma
### Log(Prior Densities)
beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
gamma.prior <- sum(dnorm(gamma, 0, sigma[2], log=TRUE))
sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
### Log-Likelihood
S <- as.vector(tcrossprod(Data$Z, t(gamma)))
mu <- as.vector(tcrossprod(Data$X, t(beta))) + S
LL <- sum(dnorm(Data$y, mu, sigma[1], log=TRUE))
### Log-Posterior
LP <- LL + beta.prior + gamma.prior + sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,S),
  yhat=rnorm(length(mu), mu, sigma[1]), parm=parm)
return(Modelout)
}

```

### 73.4. Initial Values

```
Initial.Values <- c(rep(0,1+D), rep(0,K), c(1,1))
```

## 74. Poisson Regression

### 74.1. Form

$$\mathbf{y} \sim \mathcal{P}(\lambda)$$

$$\lambda = \exp(\mathbf{X}\beta)$$

$$\beta_j \sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J$$

### 74.2. Data

```

N <- 10000
J <- 5
X <- matrix(runif(N*J,-2,2),N,J); X[,1] <- 1
beta <- runif(J,-2,2)
y <- round(exp(tcrossprod(X, t(beta))))
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J)))
PGF <- function(Data) return(rnormv(Data$J,0,1000))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, y=y)

```

### 74.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  ### Log-Likelihood
  lambda <- exp(tcrossprod(Data$X, t(beta)))
  LL <- sum(dpois(Data$y, lambda, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rpois(length(lambda), lambda), parm=parm)
  return(Modelout)
}
```

### 74.4. Initial Values

```
Initial.Values <- rep(0,J)
```

## 75. Polynomial Regression

In this univariate example, the degree of the polynomial is specified as  $D$ . For a more robust extension to estimating nonlinear relationships between  $\mathbf{y}$  and  $\mathbf{x}$ , see penalized spline regression in section ??.

### 75.1. Form

$$\begin{aligned} \mathbf{y} &\sim \mathcal{N}(\mu, \sigma^2) \\ \mu &= \mathbf{X}\beta \\ \mathbf{X}_{i,d} &= \mathbf{x}_i^{d-1}, \quad d = 1, \dots, (D+1) \\ \mathbf{X}_{i,1} &= 1 \\ \beta_d &\sim \mathcal{N}(0, 1000), \quad d = 1, \dots, (D+1) \\ \sigma &\sim \mathcal{HC}(25) \end{aligned}$$

### 75.2. Data

```
data(demonsnacks)
N <- nrow(demonsnacks)
D <- 2 #Degree of polynomial
y <- log(demonsnacks$Calories)
```

```

x <- log(demonsnacks[,10]+1)
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,D+1), sigma=0))
pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$D+1,0,1000), rhalfcauchy(1,5)))
MyData <- list(D=D, N=N, PGF=PGF, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.sigma=pos.sigma, x=x,
  y=y)

```

### 75.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
  ### Log-Likelihood
  X <- matrix(Data$x, Data$N, Data$D)
  for (d in 2:Data$D) {X[,d] <- X[,d]^d}
  X <- cbind(1,X)
  mu <- tcrossprod(X, t(beta))
  LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rnorm(length(mu), mu, sigma), parm=parm)
  return(Modelout)
}

```

### 75.4. Initial Values

```
Initial.Values <- c(rep(0,D+1), 1)
```

## 76. Proportional Hazards Regression, Weibull

Although the dependent variable is usually denoted as **t** in survival analysis, it is denoted here as **y** so Laplace's Demon recognizes it as a dependent variable for posterior predictive checks. This example does not support censoring, but it will be included soon.



### 76.1. Form

$$\begin{aligned} \mathbf{y}_i &\sim \mathcal{WELB}(\gamma, \mu_i), \quad i = 1, \dots, N \\ \mu &= \exp(\mathbf{X}\beta) \\ \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\ \gamma &\sim \mathcal{G}(1, 0.001) \end{aligned}$$

### 76.2. Data

```
N <- 50
J <- 5
X <- matrix(runif(N*J,-2,2),N,J); X[,1] <- 1
beta <- c(1,runif(J-1,-1,1))
y <- round(exp(tcrossprod(X, t(beta)))) + 1 # Undefined at zero
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J), gamma=0))
pos.beta <- grep("beta", parm.names)
pos.gamma <- grep("gamma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,10), rgamma(1,1E-3)))
MyData <- list(J=J, N=N, PGF=PGF, X=X, mon.names=mon.names,
               parm.names=parm.names, pos.beta=pos.beta, pos.gamma=pos.gamma, y=y)
```

### 76.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  gamma <- interval(parm[Data$pos.gamma], 1e-100, Inf)
  parm[Data$pos.gamma] <- gamma
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  gamma.prior <- dgamma(gamma, 1, 1.0E-3, log=TRUE)
  ### Log-Likelihood
  mu <- exp(tcrossprod(Data$X, t(beta)))
  LL <- sum(dweibull(Data$y, gamma, mu, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + gamma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
                  yhat=rweibull(length(mu), gamma, mu), parm=parm)
  return(Modelout)
}
```

#### 76.4. Initial Values

```
Initial.Values <- c(rep(0,J), 1)
```

## 77. Quantile Regression

### 77.1. Form

$$\begin{aligned} \mathbf{y} &\sim \mathcal{N}(\phi, \sigma^2) \\ \phi &= \frac{(1-2P)}{P(1-P)}\zeta + \mu \\ \mu &= \mathbf{X}\beta \\ \sigma &= \frac{P(1-P)\tau}{2\zeta} \\ \beta &\sim \mathcal{N}(0, 1000) \\ \tau &\sim \mathcal{HC}(25) \\ \zeta &\sim \mathcal{ECP}(\tau) \end{aligned}$$

where  $P$  is the user-specified quantile in  $(0, 1)$ .

### 77.2. Data

```
data(demonsnacks)
y <- log(demonsnacks$Calories)
X <- cbind(1, as.matrix(log(demonsnacks[,c(1,4,10)]+1)))
N <- nrow(X)
J <- ncol(X)
P <- 0.5 #Quantile in (0,1)
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J), tau=0, zeta=rep(0,N)))
pos.beta <- grep("beta", parm.names)
pos.tau <- grep("tau", parm.names)
pos.zeta <- grep("zeta", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,10), rhalfcauchy(1,5),
  rexp(Data$N)))
MyData <- list(J=J, N=N, P=P, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.tau=pos.tau,
  pos.zeta=pos.zeta, y=y)
```

### 77.3. Model

```
Model <- function(parm, Data)
{
```

```

### Parameters
beta <- parm[Data$pos.beta]
parm[Data$pos.tau] <- tau <- interval(parm[Data$pos.tau], 1e-100, Inf)
zeta <- interval(parm[Data$pos.zeta], 1e-100, Inf)
parm[Data$pos.zeta] <- zeta
### Log(Prior Densities)
beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
tau.prior <- dhalfcauchy(tau, 25, log=TRUE)
zeta.prior <- sum(dexp(zeta, tau, log=TRUE))
### Log-Likelihood
mu <- tcrossprod(Data$X, t(beta))
phi <- (1 - 2*Data$P) / (Data$P*(1 - Data$P))*zeta + mu
sigma <- (Data$P*(1 - Data$P)*tau) / (2*zeta)
LL <- sum(dnorm(Data$y, phi, sigma, log=TRUE))
### Log-Posterior
LP <- LL + beta.prior + tau.prior + zeta.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=rnorm(length(phi), phi, sigma), parm=parm)
return(Modelout)
}

```

#### 77.4. Initial Values

```
Initial.Values <- c(rep(0,J), 1, rep(1,N))
```

## 78. Revision, Normal

This example provides both an analytic solution and numerical approximation of the revision of a normal distribution. Given a normal prior distribution ( $\alpha$ ) and data distribution ( $\beta$ ), the posterior ( $\gamma$ ) is the revised normal distribution. This is an introductory example of Bayesian inference, and allows the user to experiment with numerical approximation, such as with MCMC in *LaplacesDemon*. Note that, regardless of the data sample size  $N$  in this example, Laplace Approximation is inappropriate due to asymptotics since the data ( $\beta$ ) is perceived by the algorithm as a single datum rather than a collection of data. MCMC, on the other hand, is biased only by the effective number of samples taken of the posterior.

#### ### Analytic Solution

```

prior.mu <- 0
prior.sigma <- 10
N <- 10
data.mu <- 1
data.sigma <- 2
posterior.mu <- (prior.sigma^-2 * prior.mu + N * data.sigma^-2 * data.mu) /
  (prior.sigma^-2 + N * data.sigma^-2)
posterior.sigma <- sqrt(1/(prior.sigma^-2 + data.sigma^-2))
posterior.mu
posterior.sigma

```

### 78.1. Form

$$\alpha \sim \mathcal{N}(0, 10)$$

$$\beta \sim \mathcal{N}(1, 2)$$

$$\gamma = \frac{\alpha_{\sigma}^{-2}\alpha + N\beta_{\sigma}^{-2}\beta}{\alpha_{\sigma}^{-2} + N\beta_{\sigma}^{-2}}$$

### 78.2. Data

```
N <- 10
mon.names <- c("LP","gamma")
parm.names <- c("alpha","beta")
PGF <- function(Data) return(c(rnorm(1,0,10), rnorm(1,1,2)))
MyData <- list(N=N, PGF=PGF, mon.names=mon.names, parm.names=parm.names)
```

### 78.3. Model

```
Model <- function(parm, Data)
{
  ### Hyperparameters
  alpha.mu <- 0
  alpha.sigma <- 10
  beta.mu <- 1
  beta.sigma <- 2
  ### Parameters
  alpha <- parm[1]
  beta <- parm[2]
  ### Log(Prior Densities)
  alpha.prior <- dnorm(alpha, alpha.mu, alpha.sigma, log=TRUE)
  ### Log-Likelihood Density
  LL <- dnorm(beta, beta.mu, beta.sigma, log=TRUE)
  ### Posterior
  gamma <- (alpha.sigma^-2 * alpha + N * beta.sigma^-2 * beta) /
    (alpha.sigma^-2 + N * beta.sigma^-2)
  ### Log(Posterior Density)
  LP <- LL + alpha.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,gamma),
    yhat=rnorm(1, beta.mu, beta.sigma), parm=parm)
  return(Modelout)
}
```

## 78.4. Initial Values

```
Initial.Values <- c(0,0)
```

# 79. Ridge Regression

## 79.1. Form

$$\begin{aligned} \mathbf{y} &\sim \mathcal{N}(\mu, \sigma_1^2) \\ \mu &= \mathbf{X}\beta \\ \beta_1 &\sim \mathcal{N}(0, 1000) \\ \beta_j &\sim \mathcal{N}(0, \sigma_2^2), \quad j = 2, \dots, J \\ \sigma_k &\sim \mathcal{HC}(25), \quad k = 1, \dots, 2 \end{aligned}$$

## 79.2. Data

```
data(demonsnacks)
y <- log(demonsnacks$Calories)
X <- cbind(1, as.matrix(log(demonsnacks[,-2]+1)))
J <- ncol(X)
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J), sigma=rep(0,2)))
pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,10), rhalfcauchy(2,5)))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
               parm.names=parm.names, pos.beta=pos.beta, pos.sigma=pos.sigma, y=y)
```

## 79.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  beta.prior <- sum(dnorm(beta, 0, c(1000, rep(sigma[2], Data$J-1)),
                    log=TRUE))
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
  ### Log-Likelihood
```

```

mu <- tcrossprod(Data$X, t(beta))
LL <- sum(dnorm(Data$y, mu, sigma[1], log=TRUE))
### Log-Posterior
LP <- LL + beta.prior + sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=rnorm(length(mu), mu, sigma[1]), parm=parm)
return(Modelout)
}

```

#### 79.4. Initial Values

```
Initial.Values <- c(rep(1,J), rep(1,2))
```

## 80. Robust Regression

By replacing the normal distribution with the Student  $t$  distribution, linear regression is often called robust regression. As an alternative approach to robust regression, consider Laplace regression (see section 44).

### 80.1. Form

$$\begin{aligned}
 \mathbf{y} &\sim t(\mu, \sigma^2, \nu) \\
 \mu &= \mathbf{X}\beta \\
 \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\
 \sigma &\sim \mathcal{HC}(25) \\
 \nu &\sim \mathcal{HC}(25)
 \end{aligned}$$

### 80.2. Data

```

N <- 100
J <- 5
X <- matrix(1,N,J)
for (j in 2:J) {X[,j] <- rnorm(N,runif(1,-3,3),runif(1,0.1,1))}
beta <- runif(J,-3,3)
e <- rst(N,0,1,5)
y <- tcrossprod(X, t(beta)) + e
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J), sigma=0, nu=0))
pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)
pos.nu <- grep("nu", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,10), rhalfcauchy(2,5)))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,

```

```
parm.names=parm.names, pos.beta=pos.beta, pos.sigma=pos.sigma,
pos.nu=pos.nu, y=y)
```

### 80.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[1:Data$J]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  parm[Data$pos.nu] <- nu <- interval(parm[Data$pos.nu], 1e-100, Inf)
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
  nu.prior <- dhalfcauchy(nu, 25, log=TRUE)
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, t(beta))
  LL <- sum(dst(Data$y, mu, sigma, nu, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + sigma.prior + nu.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rst(length(mu), mu, sigma, nu), parm=parm)
  return(Modelout)
}
```

### 80.4. Initial Values

```
Initial.Values <- c(rep(0,J), 1, 5)
```

## 81. Seemingly Unrelated Regression (SUR)

The following data was used by Zellner (1962) when introducing the Seemingly Unrelated Regression methodology. This model uses the conjugate Wishart distribution for precision in a multivariate normal distribution. See section 24 for a non-Wishart alternative that is more flexible and converges much faster.

### 81.1. Form

$$\begin{aligned} \mathbf{Y}_{t,k} &\sim \mathcal{N}_K(\mu_{t,k}, \Omega^{-1}), \quad t = 1, \dots, T; \quad k = 1, \dots, K \\ \mu_{1,t} &= \alpha_1 + \alpha_2 \mathbf{X}_{t-1,1} + \alpha_3 \mathbf{X}_{t-1,2}, \quad t = 2, \dots, T \\ \mu_{2,t} &= \beta_1 + \beta_2 \mathbf{X}_{t-1,3} + \beta_3 \mathbf{X}_{t-1,4}, \quad t = 2, \dots, T \\ \Omega &\sim \mathcal{W}_{K+1}(\mathbf{S}), \quad \mathbf{S} = \mathbf{I}_K \end{aligned}$$

$$\alpha_j \sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J$$

$$\beta_j \sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J$$

where  $J=3$ ,  $K=2$ , and  $T=20$ .

## 81.2. Data

```
T <- 20 #Time-periods
year <- c(1935,1936,1937,1938,1939,1940,1941,1942,1943,1944,1945,1946,
         1947,1948,1949,1950,1951,1952,1953,1954)
IG <- c(33.1,45.0,77.2,44.6,48.1,74.4,113.0,91.9,61.3,56.8,93.6,159.9,
        147.2,146.3,98.3,93.5,135.2,157.3,179.5,189.6)
VG <- c(1170.6,2015.8,2803.3,2039.7,2256.2,2132.2,1834.1,1588.0,1749.4,
        1687.2,2007.7,2208.3,1656.7,1604.4,1431.8,1610.5,1819.4,2079.7,
        2371.6,2759.9)
CG <- c(97.8,104.4,118.0,156.2,172.6,186.6,220.9,287.8,319.9,321.3,319.6,
        346.0,456.4,543.4,618.3,647.4,671.3,726.1,800.3,888.9)
IW <- c(12.93,25.90,35.05,22.89,18.84,28.57,48.51,43.34,37.02,37.81,
        39.27,53.46,55.56,49.56,32.04,32.24,54.38,71.78,90.08,68.60)
VW <- c(191.5,516.0,729.0,560.4,519.9,628.5,537.1,561.2,617.2,626.7,
        737.2,760.5,581.4,662.3,583.8,635.2,723.8,864.1,1193.5,1188.9)
CW <- c(1.8,0.8,7.4,18.1,23.5,26.5,36.2,60.8,84.4,91.2,92.4,86.0,111.1,
        130.6,141.8,136.7,129.7,145.5,174.8,213.5)
J <- 2 #Number of dependent variables
Y <- matrix(c(IG,IW), T, J)
S <- diag(J)
mon.names <- "LP"
parm.names <- as.parm.names(list(alpha=rep(0,3), beta=rep(0,3),
                                U=diag(J)), uppertri=c(0,0,1))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
PGF <- function(Data) return(c(rnormv(3,0,10), rnormv(3,0,10),
                                upper.triangle(rwishartc(Data$J+1,Data$S), diag=TRUE)))
MyData <- list(J=J, PGF=PGF, S=S, T=T, Y=Y, CG=CG, CW=CW, IG=IG, IW=IW,
              VG=VG, VW=VW, mon.names=mon.names, parm.names=parm.names,
              pos.alpha=pos.alpha, pos.beta=pos.beta)
```

## 81.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[Data$pos.alpha]
  beta <- parm[Data$pos.beta]
  U <- as.parm.matrix(U, Data$J, parm, Data, chol=TRUE)
  diag(U) <- exp(diag(U))
```



```

### Log(Prior Densities)
alpha.prior <- sum(dnormv(alpha, 0, 1000, log=TRUE))
beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
U.prior <- dwishartc(U, Data$J+1, Data$S, log=TRUE)
### Log-Likelihood
mu <- Data$Y
mu[-1,1] <- alpha[1] + alpha[2]*Data$CG[-Data$T] +
  alpha[3]*Data$VG[-Data$T]
mu[-1,2] <- beta[1] + beta[2]*Data$CW[-Data$T] +
  beta[3]*Data$VW[-Data$T]
LL <- sum(dmvnpc(Data$Y[-1,], mu[-1,], U, log=TRUE))
### Log-Posterior
LP <- LL + alpha.prior + beta.prior + U.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=rmvnpc(nrow(mu), mu, U), parm=parm)
return(Modelout)
}

```

#### 81.4. Initial Values

```
Initial.Values <- c(rep(0,3), rep(0,3), upper.triangle(S, diag=TRUE))
```

## 82. Simultaneous Equations

This example of simultaneous equations uses Klein's Model I ([Kleine 1950](#)) regarding economic fluctuations in the United States in 1920-1941 ( $N=22$ ). Usually, this example is modeled with 3-stage least squares (3SLS), excluding the uncertainty from multiple stages. By constraining each element in the instrumental variables matrix  $\nu \in [-10, 10]$ , this example estimates the model without resorting to stages. The dependent variable is matrix  $\mathbf{Y}$ , in which  $\mathbf{Y}_{1,1:N}$  is  $\mathbf{C}$  or Consumption,  $\mathbf{Y}_{2,1:N}$  is  $\mathbf{I}$  or Investment, and  $\mathbf{Y}_{3,1:N}$  is  $\mathbf{Wp}$  or Private Wages. Here is a data dictionary:

```

A = Time Trend measured as years from 1931
C = Consumption
G = Government Nonwage Spending
I = Investment
K = Capital Stock
P = Private (Corporate) Profits
T = Indirect Business Taxes Plus Neg Exports
Wg = Government Wage Bill
Wp = Private Wages
X = Equilibrium Demand (GNP)

```

See [Kleine \(1950\)](#) for more information.

## 82.1. Form

$$\begin{aligned}
\mathbf{Y} &\sim \mathcal{N}_3(\mu, \Omega^{-1}) \\
\mu_{1,1} &= \alpha_1 + \alpha_2 \nu_{1,1} + \alpha_4 \nu_{2,1} \\
\mu_{1,i} &= \alpha_1 + \alpha_2 \nu_{1,i} + \alpha_3 \mathbf{P}_{i-1} + \alpha_4 \nu_{2,i}, \quad i = 2, \dots, N \\
\mu_{2,1} &= \beta_1 + \beta_2 \nu_{1,1} + \beta_4 \mathbf{K}_1 \\
\mu_{2,i} &= \beta_1 + \beta_2 \nu_{1,i} + \beta_3 \mathbf{P}_{i-1} + \beta_4 \mathbf{K}_i, \quad i = 2, \dots, N \\
\mu_{3,1} &= \gamma_1 + \gamma_2 \nu_{3,1} + \gamma_4 \mathbf{A}_1 \\
\mu_{3,i} &= \gamma_1 + \gamma_2 \nu_{3,i} + \gamma_3 \mathbf{X}_{i-1} + \gamma_4 \mathbf{A}_i, \quad i = 2, \dots, N \\
\mathbf{Z}_{j,i} &\sim \mathcal{N}(\nu_{j,i}, \sigma_j^2), \quad j = 1, \dots, 3 \\
\nu_{j,1} &= \pi_{j,1} + \pi_{j,3} \mathbf{K}_1 + \pi_{j,5} \mathbf{A}_1 + \pi_{j,6} \mathbf{T}_1 + \pi_{j,7} \mathbf{G}_1, \quad j = 1, \dots, 3 \\
\nu_{j,i} &= \pi_{j,1} + \pi_{j,2} \mathbf{P}_{i-1} + \pi_{j,3} \mathbf{K}_i + \pi_{j,4} \mathbf{X}_{i-1} + \pi_{j,5} \mathbf{A}_i + \pi_{j,6} \mathbf{T}_i + \pi_{j,7} \mathbf{G}_i, \quad i = 1, \dots, N, \quad j = 1, \dots, 3 \\
\alpha_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, 4 \\
\beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, 4 \\
\gamma_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, 4 \\
\pi_{j,i} &\sim \mathcal{N}(0, 1000) \in [-10, 10], \quad j = 1, \dots, 3, \quad i = 1, \dots, N \\
\sigma_j &\sim \mathcal{HC}(25), \quad j = 1, \dots, 3 \\
\Omega &\sim \mathcal{W}_4(\mathbf{S}), \quad \mathbf{S} = \mathbf{I}_3
\end{aligned}$$

## 82.2. Data

```

N <- 22
A <- c(-11, -10, -9, -8, -7, -6, -5, -4, -3, -2, -1, 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10)
C <- c(39.8, 41.9, 45, 49.2, 50.6, 52.6, 55.1, 56.2, 57.3, 57.8, 55, 50.9, 45.6, 46.5,
      48.7, 51.3, 57.7, 58.7, 57.5, 61.6, 65, 69.7)
G <- c(2.4, 3.9, 3.2, 2.8, 3.5, 3.3, 3.3, 4, 4.2, 4.1, 5.2, 5.9, 4.9, 3.7, 4, 4.4, 2.9, 4.3,
      5.3, 6.6, 7.4, 13.8)
I <- c(2.7, -0.2, 1.9, 5.2, 3, 5.1, 5.6, 4.2, 3, 5.1, 1, -3.4, -6.2, -5.1, -3, -1.3, 2.1, 2,
      -1.9, 1.3, 3.3, 4.9)
K <- c(180.1, 182.8, 182.6, 184.5, 189.7, 192.7, 197.8, 203.4, 207.6, 210.6, 215.7,
      216.7, 213.3, 207.1, 202, 199, 197.7, 199.8, 201.8, 199.9, 201.2, 204.5)
P <- c(12.7, 12.4, 16.9, 18.4, 19.4, 20.1, 19.6, 19.8, 21.1, 21.7, 15.6, 11.4, 7, 11.2,
      12.3, 14, 17.6, 17.3, 15.3, 19, 21.1, 23.5)
T <- c(3.4, 7.7, 3.9, 4.7, 3.8, 5.5, 7, 6.7, 4.2, 4, 7.7, 7.5, 8.3, 5.4, 6.8, 7.2, 8.3, 6.7,
      7.4, 8.9, 9.6, 11.6)
Wg <- c(2.2, 2.7, 2.9, 2.9, 3.1, 3.2, 3.3, 3.6, 3.7, 4, 4.2, 4.8, 5.3, 5.6, 6, 6.1, 7.4,
      6.7, 7.7, 7.8, 8, 8.5)
Wp <- c(28.8, 25.5, 29.3, 34.1, 33.9, 35.4, 37.4, 37.9, 39.2, 41.3, 37.9, 34.5, 29, 28.5,
      30.6, 33.2, 36.8, 41, 38.2, 41.6, 45, 53.3)

```

```

X <- c(44.9,45.6,50.1,57.2,57.1,61,64,64.4,64.5,67,61.2,53.4,44.3,45.1,
      49.7,54.4,62.7,65,60.9,69.5,75.7,88.4)
year <- c(1920,1921,1922,1923,1924,1925,1926,1927,1928,1929,1930,1931,1932,
        1933,1934,1935,1936,1937,1938,1939,1940,1941)
Y <- matrix(c(C,I,Wp),3,N, byrow=TRUE)
Z <- matrix(c(P, Wp+Wg, X), 3, N, byrow=TRUE)
S <- diag(nrow(Y))
mon.names <- "LP"
parm.names <- as.parm.names(list(alpha=rep(0,4), beta=rep(0,4),
      gamma=rep(0,4), pi=matrix(0,3,7), sigma=rep(0,3),
      U=diag(3)), uppertri=c(0,0,0,0,0,1))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
pos.gamma <- grep("gamma", parm.names)
pos.pi <- grep("pi", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormmv(4,0,10), rnormmv(4,0,10),
      rnormmv(4,0,10), rnormmv(3*7,0,10), rhalfcauchy(3,5),
      upper.triangle(rwishartc(nrow(Data$S)+1,Data$S), diag=TRUE)))
MyData <- list(A=A, C=C, G=G, I=I, K=K, N=N, P=P, PGF=PGF, S=S, T=T, Wg=Wg,
      Wp=Wp, X=X, Y=Y, Z=Z, mon.names=mon.names, parm.names=parm.names,
      pos.alpha=pos.alpha, pos.beta=pos.beta, pos.gamma=pos.gamma,
      pos.pi=pos.pi, pos.sigma=pos.sigma)

```

### 82.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[Data$pos.alpha]
  beta <- parm[Data$pos.beta]
  gamma <- parm[Data$pos.gamma]
  parm[Data$pos.pi] <- pi <- interval(parm[Data$pos.pi], -10, 10)
  pi <- matrix(pi, 3, 7)      sigma <- interval(parm[Data$pos.sigma], 1e-100,
Inf)
  parm[Data$pos.sigma] <- sigma      U <- as.parm.matrix(U, nrow(Data$S), parm,
Data, chol=TRUE)
  parm[grep("Omega", Data$parm.names)] <- upper.triangle(Omega,
    diag=TRUE)
  diag(U) <- exp(diag(U))
  ### Log(Prior Densities)
  alpha.prior <- sum(dnormmv(alpha, 0, 1000, log=TRUE))
  beta.prior <- sum(dnormmv(beta, 0, 1000, log=TRUE))
  gamma.prior <- sum(dnormmv(gamma, 0, 1000, log=TRUE))
  pi.prior <- sum(dnormmv(pi, 0, 1000, log=TRUE))
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))

```

```

U.prior <- dwishartc(U, nrow(Data$S)+1, Data$S, log=TRUE)
### Log-Likelihood
mu <- nu <- matrix(0,3,Data$N)
for (i in 1:3) {
  nu[i,1] <- pi[i,1] + pi[i,3]*Data$K[1] + pi[i,5]*Data$A[1] +
    pi[i,6]*Data$T[1] + pi[i,7]*Data$G[1]
  nu[i,-1] <- pi[i,1] + pi[i,2]*Data$P[-Data$N] +
    pi[i,3]*Data$K[-1] + pi[i,4]*Data$X[-Data$N] +
    pi[i,5]*Data$A[-1] + pi[i,6]*Data$T[-1] +
    pi[i,7]*Data$G[-1]}
LL <- sum(dnorm(Data$Z, nu, matrix(sigma, 3, Data$N), log=TRUE))
mu[1,1] <- alpha[1] + alpha[2]*nu[1,1] + alpha[4]*nu[2,1]
mu[1,-1] <- alpha[1] + alpha[2]*nu[1,-1] +
  alpha[3]*Data$P[-Data$N] + alpha[4]*nu[2,-1]
mu[2,1] <- beta[1] + beta[2]*nu[1,1] + beta[4]*Data$K[1]
mu[2,-1] <- beta[1] + beta[2]*nu[1,-1] +
  beta[3]*Data$P[-Data$N] + beta[4]*Data$K[-1]
mu[3,1] <- gamma[1] + gamma[2]*nu[3,1] + gamma[4]*Data$A[1]
mu[3,-1] <- gamma[1] + gamma[2]*nu[3,-1] +
  gamma[3]*Data$X[-Data$N] + gamma[4]*Data$A[-1]
LL <- LL + sum(dmvnpc(t(Data$Y), t(mu), U, log=TRUE))
### Log-Posterior
LP <- LL + alpha.prior + beta.prior + gamma.prior + pi.prior +
  sigma.prior + U.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=t(rmvnp(ncol(mu), t(mu), U)), parm=parm)
return(Modelout)
}

```

## 82.4. Initial Values

```

Initial.Values <- c(rep(0,4), rep(0,4), rep(0,4), rep(0,3*7), rep(1,3),
  upper.triangle(S, diag=TRUE))

```

## 83. Space-Time, Dynamic

This approach to space-time or spatiotemporal modeling applies kriging to a stationary spatial component for points in space  $s = 1, \dots, S$  first at time  $t = 1$ , where space is continuous and time is discrete. Vector  $\zeta$  contains these spatial effects. Next, SSM (State Space Model) or DLM (Dynamic Linear Model) components are applied to the spatial parameters ( $\phi$ ,  $\kappa$ , and  $\lambda$ ) and regression effects ( $\beta$ ). These parameters are allowed to vary dynamically with time  $t = 2, \dots, T$ , and the resulting spatial process is estimated for each of these time-periods. When time is discrete, a dynamic space-time process can be applied. The matrix  $\Theta$  contains the dynamically varying stationary spatial effects, or space-time effects. Spatial coordinates are given in longitude and latitude for  $s = 1, \dots, S$  points in space and measurements are

taken across discrete time-periods  $t = 1, \dots, T$  for  $\mathbf{Y}_{s,t}$ . The dependent variable is also a function of design matrix  $\mathbf{X}$  (which may also be dynamic, but is static in this example) and dynamic regression effects matrix  $\beta_{1:J,1:T}$ . For more information on kriging, see section 42. For more information on SSMS or DLMS, see section 88. To extend this to a large spatial data set, consider incorporating the predictive process kriging example in section 43.

### 83.1. Form

$$\begin{aligned}
\mathbf{Y}_{s,t} &\sim \mathcal{N}(\mu_{s,t}, \sigma_1^2), \quad s = 1, \dots, S, \quad t = 1, \dots, T \\
\mu_{s,t} &= \mathbf{X}_{s,1:J} \beta_{1:J,t} + \Theta_{s,t} \\
\Theta_{s,t} &= \frac{\Sigma_{s,s,t}}{\sum_{r=1}^S \Sigma_{r,s,t}} \Theta_{s,t-1}, \quad s = 1, \dots, S, \quad t = 2, \dots, T \\
\Theta_{s,1} &= \zeta_s \\
\zeta &\sim \mathcal{N}_S(0, \Sigma_{1:S,1:S,1}) \\
\Sigma_{1:S,1:S,t} &= \lambda_t^2 \exp(-\phi_t \mathbf{D})^{\kappa[t]} \\
\sigma_1 &\sim \mathcal{HC}(25) \\
\beta_{j,1} &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, 2 \\
\beta_{1,t} &\sim \mathcal{N}(\beta_{1,t-1}, \sigma_2^2), \quad t = 2, \dots, T \\
\beta_{2,t} &\sim \mathcal{N}(\beta_{2,t-1}, \sigma_3^2), \quad t = 2, \dots, T \\
\phi_1 &\sim \mathcal{HN}(1000) \\
\phi_t &\sim \mathcal{N}(\phi_{t-1}, \sigma_4^2) \in [0, \infty], \quad t = 2, \dots, T \\
\kappa_1 &\sim \mathcal{HN}(1000) \\
\kappa_t &\sim \mathcal{N}(\kappa_{t-1}, \sigma_5^2) \in [0, \infty], \quad t = 2, \dots, T \\
\lambda_1 &\sim \mathcal{HN}(1000) \\
\lambda_t &\sim \mathcal{N}(\lambda_{t-1}, \sigma_6^2) \in [0, \infty], \quad t = 2, \dots, T
\end{aligned}$$

### 83.2. Data

```

S <- 20
T <- 10
longitude <- runif(S,0,100)
latitude <- runif(S,0,100)
D <- as.matrix(dist(cbind(longitude,latitude), diag=TRUE, upper=TRUE))
beta <- matrix(c(50,2), 2, T)
phi <- rep(1,T); kappa <- rep(1.5,T); lambda <- rep(10000,T)
for (t in 2:T) {
  beta[1,t-1] <- beta[1,t-1] + rnorm(1,0,1)
  beta[2,t-1] <- beta[2,t-1] + rnorm(1,0,0.1)
  phi[t] <- phi[t-1] + rnorm(1,0,0.1)
}

```

```

    if(phi[t] < 0.001) phi[t] <- 0.001
    kappa[t] <- kappa[t-1] + rnorm(1,0,0.1)
    lambda[t] <- lambda[t-1] + rnorm(1,0,1000)}
Sigma <- array(0, dim=c(S,S,T))
for (t in 1:T) {
  Sigma[ , ,t] <- lambda[t] * exp(-phi[t] * D)^kappa[t]}
zeta <- as.vector(apply(rmvn(1000, rep(0,S), Sigma[ , ,1]), 2, mean))
Theta <- matrix(zeta,S,T)
for (t in 2:T) {for (s in 1:S) {
  Theta[,t] <- sum(Sigma[,s,t] / sum(Sigma[,s,t]) * Theta[,t-1])}}
X <- matrix(runif(S*2,-2,2),S,2); X[,1] <- 1
mu <- tcrossprod(X, t(beta))
Y <- mu + Theta + matrix(rnorm(S*T,0,0.1),S,T)
mon.names <- "LP"
parm.names <- as.parm.names(list(zeta=rep(0,S), beta=matrix(0,2,T),
  phi=rep(0,T), kappa=rep(0,T), lambda=rep(0,T), sigma=rep(0,6)))
pos.zeta <- grep("zeta", parm.names)
pos.beta <- grep("beta", parm.names)
pos.phi <- grep("phi", parm.names)
pos.kappa <- grep("kappa", parm.names)
pos.lambda <- grep("lambda", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rmvn(1, rep(0,Data$S),
  rhalfnorm(1,sqrt(1000))^2 *
  exp(-rhalfnorm(1,sqrt(1000))*Data$D)^rhalfnorm(1,sqrt(1000))),
  rnormv(2*Data$T,0,1000), rhalfnorm(Data$T,sqrt(1000)),
  rhalfnorm(Data$T,sqrt(1000)), rhalfnorm(Data$T,sqrt(1000)),
  rhalfcauchy(6,5)))
MyData <- list(D=D, PGF=PGF, S=S, T=T, X=X, Y=Y, latitude=latitude,
  longitude=longitude, mon.names=mon.names, parm.names=parm.names,
  pos.zeta=pos.zeta, pos.beta=pos.beta, pos.phi=pos.phi,
  pos.kappa=pos.kappa, pos.lambda=pos.lambda, pos.sigma=pos.sigma)

```

### 83.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  beta <- matrix(parm[Data$pos.beta], 2, Data$T)
  zeta <- parm[Data$pos.zeta]
  parm[Data$pos.phi] <- phi <- interval(parm[Data$pos.phi], 1e-100, Inf)
  kappa <- interval(parm[Data$pos.kappa], 1e-100, Inf)
  parm[Data$pos.kappa] <- kappa
  lambda <- interval(parm[Data$pos.lambda], 1e-100, Inf)
  parm[Data$pos.lambda] <- lambda
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)

```

```

parm[Data$pos.sigma] <- sigma
Sigma <- array(0, dim=c(Data$S, Data$S, Data$T))
for (t in 1:Data$T) {
  Sigma[ , ,t] <- lambda[t]^2 * exp(-phi[t] * Data$D)^kappa[t]}
### Log(Prior Densities)
beta.prior <- sum(dnorm(beta[,1], 0, 1000, log=TRUE),
  dnorm(beta[,-1], beta[,-Data$T], matrix(sigma[2:3], 2,
    Data$T-1), log=TRUE))
zeta.prior <- dmvn(zeta, rep(0,Data$S), Sigma[ , , 1], log=TRUE)
phi.prior <- sum(dhalfnorm(phi[1], sqrt(1000), log=TRUE),
  dtrunc(phi[-1], "norm", a=0, b=Inf, mean=phi[-Data$T],
    sd=sigma[4], log=TRUE))
kappa.prior <- sum(dhalfnorm(kappa[1], sqrt(1000), log=TRUE),
  dtrunc(kappa[-1], "norm", a=0, b=Inf, mean=kappa[-Data$T],
    sd=sigma[5], log=TRUE))
lambda.prior <- sum(dhalfnorm(lambda[1], sqrt(1000), log=TRUE),
  dtrunc(lambda[-1], "norm", a=0, b=Inf, mean=lambda[-Data$T],
    sd=sigma[6], log=TRUE))
sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
### Log-Likelihood
mu <- tcrossprod(Data$X, t(beta))
Theta <- matrix(zeta, Data$S, Data$T)
for (t in 2:Data$T) {
  for (s in 1:Data$S) {
    Theta[,t] <- Sigma[,s,t] / sum(Sigma[,s,t]) * Theta[,t-1]}
mu <- mu + Theta
LL <- sum(dnorm(Data$Y, mu, sigma[1], log=TRUE))
### Log-Posterior
LP <- LL + beta.prior + zeta.prior + sum(phi.prior) +
  sum(kappa.prior) + sum(lambda.prior) + sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=rnorm(prod(dim(mu)), mu, sigma[1]), parm=parm)
return(Modelout)
}

```

#### 83.4. Initial Values

```

Initial.Values <- c(rep(0,S), rep(c(mean(Y),0),T), rep(1,T), rep(1,T),
  rep(1,T), rep(1,6))

```

### 84. Space-Time, Nonseparable

This approach to space-time or spatiotemporal modeling applies kriging both to the stationary spatial and temporal components, where space is continuous and time is discrete. Matrix  $\Xi$  contains the space-time effects. Spatial coordinates are given in longitude and latitude for

$s = 1, \dots, S$  points in space and measurements are taken across time-periods  $t = 1, \dots, T$  for  $\mathbf{Y}_{s,t}$ . The dependent variable is also a function of design matrix  $\mathbf{X}$  and regression effects vector  $\beta$ . For more information on kriging, see section 42. This example uses a nonseparable, stationary covariance function in which space and time are separable only when  $\psi = 0$ . To extend this to a large space-time data set, consider incorporating the predictive process kriging example in section 43.

### 84.1. Form

$$\begin{aligned}
 \mathbf{Y}_{s,t} &\sim \mathcal{N}(\mu_{s,t}, \sigma_1^2), \quad s = 1, \dots, S, \quad t = 1, \dots, T \\
 \mu &= \mathbf{X}\beta + \Xi \\
 \Xi &\sim \mathcal{N}_{ST}(\Xi_\mu, \Sigma) \\
 \Sigma &= \sigma_2^2 \exp \left( -\frac{\mathbf{D}_S^\kappa}{\phi_1} - \frac{\mathbf{D}_T^\lambda}{\phi_2} - \psi \frac{\mathbf{D}_S^\kappa}{\phi_1} \frac{\mathbf{D}_T^\lambda}{\phi_2} \right) \\
 \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\
 \phi_k &\sim \mathcal{U}(1, 5), \quad k = 1, \dots, 2 \\
 \sigma_k &\sim \mathcal{HC}(25), \quad k = 1, \dots, 2 \\
 \psi &\sim \mathcal{HC}(25) \\
 \Xi_\mu &= 0 \\
 \kappa &= 1, \quad \lambda = 1
 \end{aligned}$$

### 84.2. Data

```

S <- 10
T <- 5
longitude <- runif(S,0,100)
latitude <- runif(S,0,100)
D.S <- as.matrix(dist(cbind(rep(longitude,T),rep(latitude,T)), diag=TRUE,
  upper=TRUE))
D.T <- as.matrix(dist(cbind(rep(1:T,each=S),rep(1:T,each=S)), diag=TRUE,
  upper=TRUE))
Sigma <- 10000 * exp(-D.S/3 - D.T/2 - 0.2*(D.S/3)*(D.T/2))
Xi <- as.vector(apply(rmvn(1000, rep(0,S*T), Sigma), 2, mean))
Xi <- matrix(Xi,S,T)
beta <- c(50,2)
X <- matrix(runif(S*2,-2,2),S,2); X[,1] <- 1
mu <- as.vector(tcrossprod(X, t(beta)))
Y <- mu + Xi
mon.names <- "LP"
parm.names <- as.parm.names(list(Xi=matrix(0,S,T), beta=rep(0,2),
  phi=rep(0,2), sigma=rep(0,2), psi=0))

```



```

pos.Xi <- grep("Xi", parm.names)
pos.beta <- grep("beta", parm.names)
pos.phi <- grep("phi", parm.names)
pos.sigma <- grep("sigma", parm.names)
pos.psi <- grep("psi", parm.names)
PGF <- function(Data) return(c(rmvn(1, rep(0,Data$S*Data$T),
  rhalfcauchy(1,25)^2 * exp(-(Data$D.S / runif(1,1,5)) -
    (Data$D.T / runif(1,1,5)) -
    rhalfcauchy(1,5)*(Data$D.S / rhalfcauchy(1,5)))),
  rnormv(2,0,1000), runif(2,1,5), rhalfcauchy(3,25)))
MyData <- list(D.S=D.S, D.T=D.T, PGF=PGF, S=S, T=T, X=X, Y=Y,
  latitude=latitude, longitude=longitude, mon.names=mon.names,
  parm.names=parm.names, pos.Xi=pos.Xi, pos.beta=pos.beta,
  pos.phi=pos.phi, pos.sigma=pos.sigma, pos.psi=pos.psi)

```

### 84.3. Model

```

Model <- function(parm, Data)
{
  ### Hyperparameters
  Xi.mu <- rep(0,Data$S*Data$T)
  ### Parameters
  beta <- parm[Data$pos.beta]
  Xi <- parm[Data$pos.Xi]
  kappa <- 1; lambda <- 1
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  parm[Data$pos.phi] <- phi <- interval(parm[Data$pos.phi], 1, 5)
  parm[Data$pos.psi] <- psi <- interval(parm[Data$pos.psi], 1e-100, Inf)
  Sigma <- sigma[2]*sigma[2] * exp(-(Data$D.S / phi[1])^kappa -
    (Data$D.T / phi[2])^lambda -
    psi*(Data$D.S / phi[1])^kappa * (Data$D.T / phi[2])^lambda)
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  Xi.prior <- dmvn(Xi, Xi.mu, Sigma, log=TRUE)
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
  phi.prior <- sum(dunif(phi, 1, 5, log=TRUE))
  psi.prior <- dhalfcauchy(psi, 25, log=TRUE)
  ### Log-Likelihood
  Xi <- matrix(Xi, Data$S, Data$T)
  mu <- as.vector(tcrossprod(Data$X, t(beta))) + Xi
  LL <- sum(dnorm(Data$Y, mu, sigma[1], log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + Xi.prior + sigma.prior + phi.prior + psi.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rnorm(prod(dim(mu)), mu, sigma[1]), parm=parm)
}

```

```
return(Modelout)
}
```

#### 84.4. Initial Values

```
Initial.Values <- c(rep(0,S*T), mean(Y), 1, rep(1,2), rep(1,2), 1)
```

### 85. Space-Time, Separable

This introductory approach to space-time or spatiotemporal modeling applies kriging both to the stationary spatial and temporal components, where space is continuous and time is discrete. Vector  $\zeta$  contains the spatial effects and vector  $\theta$  contains the temporal effects. Spatial coordinates are given in longitude and latitude for  $s = 1, \dots, S$  points in space and measurements are taken across time-periods  $t = 1, \dots, T$  for  $\mathbf{Y}_{s,t}$ . The dependent variable is also a function of design matrix  $\mathbf{X}$  and regression effects vector  $\beta$ . For more information on kriging, see section 42. This example uses separable space-time covariances, which is more convenient but usually less appropriate than a nonseparable covariance function. To extend this to a large space-time data set, consider incorporating the predictive process kriging example in section 43.

#### 85.1. Form

$$\begin{aligned}
\mathbf{Y}_{s,t} &\sim \mathcal{N}(\mu_{s,t}, \sigma_1^2), \quad s = 1, \dots, S, \quad t = 1, \dots, T \\
\mu_{s,t} &= \mathbf{X}_{s,1:J}\beta + \zeta_s + \Theta_{s,t} \\
\Theta_{s,1:T} &= \theta \\
\theta &\sim \mathcal{N}_N(\theta_\mu, \Sigma_T) \\
\Sigma_T &= \sigma_3^2 \exp(-\phi_2 \mathbf{D}_T)^\lambda \\
\zeta &\sim \mathcal{N}_N(\zeta_\mu, \Sigma_S) \\
\Sigma_S &= \sigma_2^2 \exp(-\phi_1 \mathbf{D}_S)^\kappa \\
\beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, 2 \\
\sigma_k &\sim \mathcal{HC}(25), \quad k = 1, \dots, 3 \\
\phi_k &\sim \mathcal{U}(1, 5), \quad k = 1, \dots, 2 \\
\zeta_\mu &= 0 \\
\theta_\mu &= 0 \\
\kappa &= 1, \quad \lambda = 1
\end{aligned}$$

#### 85.2. Data

```
S <- 20
T <- 10
```

```

longitude <- runif(S,0,100)
latitude <- runif(S,0,100)
D.S <- as.matrix(dist(cbind(longitude,latitude), diag=TRUE, upper=TRUE))
Sigma.S <- 10000 * exp(-1.5 * D.S)
zeta <- as.vector(apply(rmvn(1000, rep(0,S), Sigma.S), 2, mean))
D.T <- as.matrix(dist(cbind(c(1:T),c(1:T)), diag=TRUE, upper=TRUE))
Sigma.T <- 10000 * exp(-3 * D.T)
theta <- as.vector(apply(rmvn(1000, rep(0,T), Sigma.T), 2, mean))
Theta <- matrix(theta,S,T,byrow=TRUE)
beta <- c(50,2)
X <- matrix(runif(S*2,-2,2),S,2); X[,1] <- 1
mu <- as.vector(tcrossprod(X, t(beta)))
Y <- mu + zeta + Theta + matrix(rnorm(S*T,0,0.1),S,T)
mon.names <- "LP"
parm.names <- as.parm.names(list(zeta=rep(0,S), theta=rep(0,T),
    beta=rep(0,2), phi=rep(0,2), sigma=rep(0,3)))
pos.zeta <- grep("zeta", parm.names)
pos.theta <- grep("theta", parm.names)
pos.beta <- grep("beta", parm.names)
pos.phi <- grep("phi", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rmvn(1, rep(0,Data$S),
    rhalfcauchy(1,5)^2 * exp(-runif(1,1,5)*Data$D.S)),
    rmvn(1, rep(0,Data$T), rhalfcauchy(1,5)^2 *
    exp(-runif(1,1,5)*Data$D.T)), rnormv(2,0,1000), runif(2,1,5),
    rhalfcauchy(3,5)))
MyData <- list(D.S=D.S, D.T=D.T, PGF=PGF, S=S, T=T, X=X, Y=Y,
    latitude=latitude, longitude=longitude, mon.names=mon.names,
    parm.names=parm.names, pos.zeta=pos.zeta, pos.theta=pos.theta,
    pos.beta=pos.beta, pos.phi=pos.phi, pos.sigma=pos.sigma)

```

### 85.3. Model

```

Model <- function(parm, Data)
{
  ### Hyperparameters
  zeta.mu <- rep(0,Data$S)
  theta.mu <- rep(0,Data$T)
  ### Parameters
  beta <- parm[Data$pos.beta]
  zeta <- parm[Data$pos.zeta]
  theta <- parm[Data$pos.theta]
  kappa <- 1; lambda <- 1
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  parm[Data$pos.phi] <- phi <- interval(parm[Data$pos.phi], 1, 5)
}

```

```

Sigma.S <- sigma[2]^2 * exp(-phi[1] * Data$D.S)^kappa
Sigma.T <- sigma[3]^2 * exp(-phi[2] * Data$D.T)^lambda
### Log(Prior Densities)
beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
zeta.prior <- dmvn(zeta, zeta.mu, Sigma.S, log=TRUE)
theta.prior <- dmvn(theta, theta.mu, Sigma.T, log=TRUE)
sigma.prior <- sum(dhalfcauchy(25, log=TRUE))
phi.prior <- sum(dunif(phi, 1, 5, log=TRUE))
### Log-Likelihood
Theta <- matrix(theta, Data$S, Data$T, byrow=TRUE)
mu <- as.vector(tcrossprod(Data$X, t(beta))) + zeta + Theta
LL <- sum(dnorm(Data$Y, mu, sigma[1], log=TRUE))
### Log-Posterior
LP <- LL + beta.prior + zeta.prior + theta.prior + sigma.prior +
  phi.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=rnorm(prod(dim(mu)), mu, sigma[1]), parm=parm)
return(Modelout)
}

```

#### 85.4. Initial Values

```
Initial.Values <- c(rep(0,S), rep(0,T), rep(0,2), rep(1,2), rep(1,3))
```

## 86. Spatial Autoregression (SAR)

The spatial autoregressive (SAR) model in this example uses areal data that consists of first-order neighbors that were specified and converted from point-based data with longitude and latitude coordinates.

### 86.1. Form

$$\begin{aligned}
 \mathbf{y} &\sim \mathcal{N}(\boldsymbol{\mu}, \sigma^2) \\
 \boldsymbol{\mu} &= \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\phi}\mathbf{z} \\
 \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\
 \phi &\sim \mathcal{U}(-1, 1) \\
 \sigma &\sim \mathcal{HC}(25)
 \end{aligned}$$

### 86.2. Data

```

N <- 100
latitude <- runif(N,0,100); longitude <- runif(N,0,100)
J <- 3 #Number of predictors, including the intercept

```

```

X <- matrix(runif(N*J,0,3), N, J); X[,1] <- 1
beta.orig <- runif(J,0,3); phi <- runif(1,0,1)
D <- as.matrix(dist(cbind(longitude, latitude), diag=TRUE, upper=TRUE))
W <- exp(-D) #Inverse distance as weights
W <- ifelse(D == 0, 0, W)
epsilon <- rnorm(N,0,1)
y <- tcrossprod(X, t(beta.orig)) + sqrt(latitude) + sqrt(longitude) +
  epsilon
Z <- W / matrix(rowSums(W), N, N) * matrix(y, N, N, byrow=TRUE)
z <- as.vector(apply(Z, 1, sum))
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J), phi=0, sigma=0))
pos.beta <- grep("beta", parm.names)
pos.phi <- grep("phi", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,1000), runif(1,-1,1),
  rhalfcauchy(1,5)))
MyData <- list(J=J, PGF=PGF, X=X, latitude=latitude, longitude=longitude,
  mon.names=mon.names, parm.names=parm.names, pos.beta=pos.beta,
  pos.phi=pos.phi, pos.sigma=pos.sigma, y=y, z=z)

```

### 86.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  parm[Data$pos.phi] <- phi <- interval(parm[Data$pos.phi], -1, 1)
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  phi.prior <- dunif(phi, -1, 1, log=TRUE)
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, t(beta)) + phi*Data$z
  LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + phi.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rnorm(length(mu), mu, sigma), parm=parm)
  return(Modelout)
}

```

### 86.4. Initial Values

```
Initial.Values <- c(rep(0,J), 0.5, 1)
```

## 87. STARMA(1,1)

The data in this example of a space-time autoregressive moving average (STARMA) are coordinate-based, and the adjacency matrix  $\mathbf{A}$  is created from  $K$  nearest neighbors. Otherwise, an adjacency matrix may be specified as usual for areal data. Spatial coordinates are given in longitude and latitude for  $s = 1, \dots, S$  points in space and measurements are taken across time-periods  $t = 1, \dots, T$  for  $\mathbf{Y}_{s,t}$ .

### 87.1. Form

$$\begin{aligned}\mathbf{Y} &\sim \mathcal{N}(\mu, \sigma^2) \\ \mu_{s,t} &= \mathbf{X}_{s,t}\beta + \phi\mathbf{W}\mathbf{1}_{s,t-1} + \theta\mathbf{W}\mathbf{2}_{s,t-1}, \quad s = 1, \dots, S, \quad t = 2, \dots, T \\ \mathbf{W}\mathbf{1} &= \mathbf{V}\mathbf{Y} \\ \mathbf{W}\mathbf{2} &= \mathbf{V}\epsilon \\ \epsilon &= \mathbf{Y} - \mu \\ \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\ \phi &\sim \mathcal{U}(-1, 1) \\ \sigma &\sim \mathcal{HC}(25) \\ \theta &\sim \mathcal{N}(0, 1000)\end{aligned}$$

where  $\mathbf{V}$  is an adjacency matrix that is scaled so that each row sums to one,  $\beta$  is a vector of regression effects,  $\phi$  is the autoregressive space-time parameter,  $\sigma$  is the residual variance, and  $\theta$  is the moving average space-time parameter.

### 87.2. Data

```
S <- 100
T <- 10
K <- 5 #Number of nearest neighbors
latitude <- runif(S,0,100)
longitude <- runif(S,0,100)
X1 <- matrix(runif(S*T,-2,2), S, T)
X2 <- matrix(runif(S*T,-2,2), S, T)
for (t in 2:T) {
  X1[,t] <- X1[,t-1] + runif(S,-0.1,0.1)
  X2[,t] <- X2[,t-1] + runif(S,-0.1,0.1)}
beta.orig <- runif(3,-2,2); phi.orig <- 0.8; theta.orig <- 1
epsilon <- matrix(rnorm(S*T,0,0.1), S, T)
Z <- matrix(rnorm(S*T,0,0.1), S, T)
```

```

D <- as.matrix(dist(cbind(longitude, latitude), diag=TRUE, upper=TRUE))
A <- exp(-D)
A <- ifelse(D == 0, max(D), A)
A <- apply(A, 1, rank)
A <- ifelse(A <= K, 1, 0)
V <- A / rowSums(A) #Scaled matrix
V <- ifelse(is.nan(V), 1/ncol(V), V)
Y <- beta.orig[1] + beta.orig[2]*X1 + beta.orig*X2
W1 <- tcrossprod(V, t(Y))
Y <- Y + phi.orig*cbind(rep(0,S), W1[, -T])
W2 <- tcrossprod(V, t(epsilon))
Y <- Y + theta.orig*cbind(rep(0,S), W2[, -T])
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,3), phi=0, sigma=0, theta=0))
pos.beta <- grep("beta", parm.names)
pos.phi <- grep("phi", parm.names)
pos.sigma <- grep("sigma", parm.names)
pos.theta <- grep("theta", parm.names)
PGF <- function(Data) return(c(rnormv(3,0,1000), runif(1,-1,1),
  rhalfcauchy(1,5), rnormv(1,0,1000)))
MyData <- list(K=K, PGF=PGF, S=S, T=T, V=V, X1=X1, X2=X2, Y=Y,
  latitude=latitude, longitude=longitude, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.phi=pos.phi,
  pos.sigma=pos.sigma, pos.theta=pos.theta)

```

### 87.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  parm[Data$pos.phi] <- phi <- interval(parm[Data$pos.phi], -1, 1)
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  theta <- parm[Data$pos.theta]
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  phi.prior <- dunif(phi, -1, 1, log=TRUE)
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
  theta.prior <- dnormv(theta, 0, 1000, log=TRUE)
  ### Log-Likelihood
  W1 <- tcrossprod(Data$V, t(Data$Y))
  mu <- beta[1] + beta[2]*Data$X1 + beta[3]*Data$X2 +
    phi*cbind(rep(0, Data$S), W1[, -Data$T])
  epsilon <- Data$Y - mu
  W2 <- tcrossprod(Data$V, t(epsilon))

```

```

mu <- mu + theta*cbind(rep(0, Data$S), W2[, -Data$T])
LL <- sum(dnorm(Data$Y[, -1], mu[, -1], sigma, log=TRUE))
### Log-Posterior
LP <- LL + beta.prior + phi.prior + sigma.prior + theta.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
  yhat=rnorm(prod(dim(mu)), mu, sigma), parm=parm)
return(Modelout)
}

```

## 87.4. Initial Values

```
Initial.Values <- c(rep(0,3), 0, 1, 0)
```

## 88. State Space Model (SSM), Linear Regression

The data is presented so that the time-series is subdivided into three sections: modeled ( $t = 1, \dots, T_m$ ), one-step ahead forecast ( $t = T_m + 1$ ), and future forecast [ $t = (T_m + 2), \dots, T$ ]. Note that Dyn must also be specified for the SAMWG and SMWG MCMC algorithms.

### 88.1. Form

$$\begin{aligned}
 \mathbf{y}_t &\sim \mathcal{N}(\mu_t, \sigma_1^2), \quad t = 1, \dots, T_m \\
 \mathbf{y}_t^{new} &\sim \mathcal{N}(\mu_t, \sigma_1^2), \quad t = (T_m + 1), \dots, T \\
 \mu_t &= \alpha + \mathbf{x}_t \beta_t, \quad t = 1, \dots, T \\
 \alpha &\sim \mathcal{N}(0, 1000) \\
 \beta_1 &\sim \mathcal{N}(0, 1000) \\
 \beta_t &\sim \mathcal{N}(\beta_{t-1}, \sigma_2^2), \quad t = 2, \dots, T \\
 \sigma_j &\sim \mathcal{HC}(25), \quad j = 1, \dots, 2
 \end{aligned}$$

### 88.2. Data

```

T <- 20
T.m <- 14
beta.orig <- x <- rep(0, T)
for (t in 2:T) {
  beta.orig[t] <- beta.orig[t-1] + rnorm(1, 0, 0.1)
  x[t] <- x[t-1] + rnorm(1, 0, 0.1)}
y <- 10 + beta.orig*x + rnorm(T, 0, 0.01)
y[(T.m+2):T] <- NA
mon.names <- rep(NA, (T-T.m))
for (i in 1:(T-T.m)) mon.names[i] <- paste("mu[", (T.m+i), "]", sep="")
parm.names <- as.parm.names(list(alpha=0, beta=rep(0, T), sigma=rep(0, 2)))

```



```

pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnorm(1,0,1), rnorm(Data$T,0,1),
  rhalfcauchy(2,5)))
MyData <- list(PGF=PGF, T=T, T.m=T.m, mon.names=mon.names,
  parm.names=parm.names, pos.alpha=pos.alpha, pos.beta=pos.beta,
  pos.sigma=pos.sigma, x=x, y=y)
Dyn <- matrix(paste("beta[",1:T,"]",sep=""), T, 1)

```

### 88.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[Data$pos.alpha]
  beta <- parm[Data$pos.beta]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  alpha.prior <- dnorm(alpha, 0, 1000, log=TRUE)
  beta.prior <- sum(dnorm(beta[1], 0, 1000, log=TRUE),
    dnorm(beta[-1], beta[-Data$T], sigma[2], log=TRUE))
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
  ### Log-Likelihood
  mu <- alpha + beta*Data$x
  LL <- sum(dnorm(Data$y[1:Data$T.m], mu[1:Data$T.m], sigma[1],
    log=TRUE))
  yhat <- rnorm(length(mu), alpha + c(beta[1], rnorm(Data$T-1,
    beta[-Data$T], sigma[2]))) * Data$x, sigma[1]) #One-step ahead
  ### Log-Posterior
  LP <- LL + alpha.prior + beta.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=mu[(Data$T.m+1):Data$T],
    yhat=yhat, parm=parm)
  return(Modelout)
}

```

### 88.4. Initial Values

```
Initial.Values <- c(0, rep(0,T), rep(1,2))
```

## 89. State Space Model (SSM), Local Level

The local level model is the simplest, non-trivial example of a state space model (SSM). As such, this version of a local level SSM has static variance parameters.

### 89.1. Form

$$\begin{aligned} \mathbf{y}_t &\sim \mathcal{N}(\mu_t, \sigma_1^2), \quad t = 1, \dots, T \\ \mu_t &\sim \mathcal{N}(\mu_{t-1}, \sigma_2^2), \quad t = 2, \dots, T \\ \mu_1 &\sim \mathcal{N}(0, 1000) \\ \sigma_j &\sim \mathcal{HC}(25), \quad j = 1, \dots, 2 \end{aligned}$$

### 89.2. Data

```
T <- 20
T.m <- 14
mu.orig <- rep(0,T)
for (t in 2:T) {mu.orig[t] <- mu.orig[t-1] + rnorm(1,0,1)}
y <- mu.orig + rnorm(T,0,0.1)
y[(T.m+2):T] <- NA
mon.names <- rep(NA, (T-T.m))
for (i in 1:(T-T.m)) mon.names[i] <- paste("yhat[",(T.m+i),"]", sep="")
parm.names <- as.parm.names(list(mu=rep(0,T), sigma=rep(0,2)))
pos.mu <- grep("mu", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormmv(Data$T,0,1000), rhalfcauchy(2,5)))
MyData <- list(PGF=PGF, T=T, T.m=T.m, mon.names=mon.names,
  parm.names=parm.names, pos.mu=pos.mu, pos.sigma=pos.sigma, y=y)
Dyn <- matrix(paste("mu[",1:T,"]",sep=""), T, 1)
```

### 89.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  mu <- parm[Data$pos.mu]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  mu.prior <- sum(dnorm(mu[1], 0, 1000, log=TRUE),
    dnorm(mu[-1], mu[-Data$T], sigma[2], log=TRUE))
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
  ### Log-Likelihood
  LL <- sum(dnorm(Data$y[1:Data$T.m], mu[1:Data$T.m], sigma[1],
    log=TRUE))
  yhat <- rnorm(length(mu), c(mu[1], rnorm(Data$T-1, mu[-Data$T],
    sigma[2])), sigma[1]) #One-step ahead      ### Log-Posterior
  LP <- LL + mu.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=mu[(Data$T.m+1):Data$T],
```

```

      yhat=yhat, parm=parm)
return(Modelout)
}

```

#### 89.4. Initial Values

```
Initial.Values <- c(rep(0,T), rep(1,2))
```

## 90. State Space Model (SSM), Local Linear Trend

The local linear trend model is a state space model (SSM) that extends the local level model to include a dynamic slope parameter. For more information on the local level model, see section 89. This example has static variance parameters.

### 90.1. Form

$$\begin{aligned}
 \mathbf{y}_t &\sim \mathcal{N}(\mu_t, \sigma_1^2), \quad t = 1, \dots, T \\
 \mu_t &\sim \mathcal{N}(\mu_{t-1} + \delta_{t-1}, \sigma_2^2), \quad t = 2, \dots, T \\
 \mu_1 &\sim \mathcal{N}(0, 1000) \\
 \delta_t &\sim \mathcal{N}(\delta_{t-1}, \sigma_3^2), \quad t = 2, \dots, T \\
 \delta_1 &\sim \mathcal{N}(0, 1000) \\
 \sigma_j &\sim \mathcal{HC}(25), \quad j = 1, \dots, 3
 \end{aligned}$$

### 90.2. Data

```

T <- 20
T.m <- 14
mu.orig <- delta.orig <- rep(0,T)
for (t in 2:T) {
  delta.orig[t] <- delta.orig[t-1] + rnorm(1,0,0.1)
  mu.orig[t] <- mu.orig[t-1] + delta.orig[t-1] + rnorm(1,0,1)}
y <- mu.orig + rnorm(T,0,0.1)
y[(T.m+2):T] <- NA
mon.names <- rep(NA, (T-T.m))
for (i in 1:(T-T.m)) mon.names[i] <- paste("yhat[",(T.m+i),"]", sep="")
parm.names <- as.parm.names(list(mu=rep(0,T), delta=rep(0,T),
  sigma=rep(0,3)))
pos.mu <- grep("mu", parm.names)
pos.delta <- grep("delta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$T,0,10),
  rnormv(Data$T,0,10), rhalfcauchy(3,5)))

```

```
MyData <- list(PGF=PGF, T=T, T.m=T.m, mon.names=mon.names,
  parm.names=parm.names, pos.mu=pos.mu, pos.delta=pos.delta,
  pos.sigma=pos.sigma, y=y)
```

### 90.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  mu <- parm[Data$pos.mu]
  delta <- parm[Data$pos.delta]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  mu.prior <- sum(dnorm(mu[1], 0, 1000, log=TRUE),
    dnorm(mu[-1], mu[-Data$T]+delta[-Data$T], sigma[2],
    log=TRUE))
  delta.prior <- sum(dnorm(delta[1], 0, 1000, log=TRUE),
    dnorm(delta[-1], delta[-Data$T], sigma[3], log=TRUE))
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
  ### Log-Likelihood
  LL <- sum(dnorm(Data$y[1:Data$T.m], mu[1:Data$T.m], sigma[1],
    log=TRUE))
  yhat <- rnorm(length(mu), c(mu[1], rnorm(Data$T-1, mu[-Data$T],
    sigma[2])), sigma[1]) #One-step ahead
  ### Log-Posterior
  LP <- LL + mu.prior + delta.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=mu[(Data$T.m+1):Data$T],
    yhat=yhat, parm=parm)
  return(Modelout)
}
```

### 90.4. Initial Values

```
Initial.Values <- c(rep(0,T), rep(0,T), rep(1,3))
```

## 91. State Space Model (SSM), Stochastic Volatility (SV)

### 91.1. Form

$$\mathbf{y} \sim \mathcal{N}(0, \sigma^2)$$

$$\sigma^2 = \frac{1}{\exp(\theta)}$$

$$\begin{aligned}\beta &= \exp(\mu/2) \\ \theta_1 &\sim \mathcal{N}(\mu + \phi(\alpha - \mu), \tau) \\ \theta_t &\sim \mathcal{N}(\mu + \phi(\theta_{t-1} - \mu), \tau), \quad t = 2, \dots, T \\ \alpha &\sim \mathcal{N}(\mu, \tau) \\ \phi &\sim \mathcal{U}(-1, 1) \\ \mu &\sim \mathcal{N}(0, 10) \\ \tau &\sim \mathcal{HC}(25)\end{aligned}$$

## 91.2. Data

```
T <- 20
y <- rep(10,T); epsilon <- rnorm(T,0,1)
for (t in 2:T) {y[t] <- 0.8*y[t-1] + epsilon[t-1]}
mon.names <- c("LP",paste("sigma2[",1:T,"]",sep=""))
parm.names <- as.parm.names(list(theta=rep(0,T), alpha=0, phi=0, mu=0,
    tau=0))
pos.theta <- grep("theta", parm.names)
pos.alpha <- grep("alpha", parm.names)
pos.phi <- grep("phi", parm.names)
pos.mu <- grep("mu", parm.names)
pos.tau <- grep("tau", parm.names)
PGF <- function(Data) return(c(rnormmv(Data$T,0,10),
    rnormmv(1,rnorm(1,0,10),rhalfcauchy(1,5)),
    runif(1,-1,1), rnormmv(1,0,10), rhalfcauchy(1,5)))
MyData <- list(PGF=PGF, T=T, mon.names=mon.names, parm.names=parm.names, pos.theta=pos
pos.alpha=pos.alpha, pos.phi=pos.phi,
    pos.mu=pos.mu, pos.tau=pos.tau y=y)
Dyn <- matrix(paste("theta[",1:T,"]",sep=""), T, 1)
```

## 91.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  theta <- parm[Data$pos.theta]
  alpha <- parm[Data$pos.alpha]
  parm[Data$pos.phi] <- phi <- interval(parm[Data$pos.phi], -1, 1)
  mu <- parm[Data$pos.mu]
  parm[Data$pos.tau] <- tau <- interval(parm[Data$pos.tau], 1e-100, Inf)
  ### Log(Prior Densities)
  alpha.prior <- dnormmv(alpha, mu, tau, log=TRUE)
  theta.prior <- sum(dnormmv(theta[1], mu + phi*(alpha-mu), tau,
    log=TRUE), dnormmv(theta[-1], mu + phi*(theta[-Data$T]-mu), tau,
```

```

    log=TRUE))
  phi.prior <- dunif(phi, -1, 1, log=TRUE)
  mu.prior <- dnorm(mu, 0, 10, log=TRUE)
  tau.prior <- dhalfcauchy(tau, 25, log=TRUE)
  ### Log-Likelihood
  beta <- exp(mu / 2)
  sigma2 <- 1 / exp(theta)
  LL <- sum(dnorm(Data$y, 0, sigma2, log=TRUE))
  ### Log-Posterior
  LP <- LL + alpha.prior + theta.prior + phi.prior + mu.prior +
    tau.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP, sigma2),
    yhat=rnormv(length(Data$y), 0, sigma2), parm=parm)
  return(Modelout)
}

```

## 91.4. Initial Values

```
Initial.Values <- c(rep(0,T), 0, 0, 0, 1)
```

## 92. TARCH(1)

In this TARCH example, there are two regimes, one for positive residuals in the previous time-period, and the other for negative. The TARCH parameters are the  $\theta$  vector.

### 92.1. Form

$$\begin{aligned}
 \mathbf{y}_t &\sim \mathcal{N}(\mu_t, \sigma_t^2), \quad t = 2, \dots, T \\
 \sigma_t^2 &= \omega + \theta_1 \delta_{t-1} \epsilon_{t-1}^2 + \theta_2 (1 - \delta_{t-1}) \epsilon_{t-1}^2, \quad t = 2, \dots, T \\
 \delta_t &= \begin{cases} 1 & \text{if } \epsilon_t > 0 \\ 0 & \text{if } \epsilon_t \leq 0 \end{cases} \\
 \epsilon &= \mathbf{y} - \mu \\
 \mu_t &= \alpha + \phi \mathbf{y}_{t-1}, \quad t = 2, \dots, T \\
 \alpha &\sim \mathcal{N}(0, 1000) \\
 \phi &\sim \mathcal{U}(-1, 1) \\
 \omega &\sim \mathcal{HC}(25) \\
 \theta_j &\sim \mathcal{U}(0, 1), \quad j = 1, \dots, 2
 \end{aligned}$$

### 92.2. Data

```

T <- 20
phi <- 0.8

```

```

epsilon <- rnorm(T)
epsilon <- ifelse(epsilon < 0, epsilon * 2, epsilon)
y <- rep(0,T)
for (t in 2:T) {y[t] <- phi*y[t-1] + epsilon[t]}
mon.names <- c("LP","ynew","sigma2.new")
parm.names <- as.parm.names(list(alpha=0, phi=0, omega=0, theta=rep(0,2)))
pos.alpha <- grep("alpha", parm.names)
pos.phi <- grep("phi", parm.names)
pos.omega <- grep("omega", parm.names)
pos.theta <- grep("theta", parm.names)
PGF <- function(Data) return(c(rnormv(1,0,1000), runif(1,-1,1),
  rhalfcauchy(1,5), runif(2)))
MyData <- list(PGF=PGF, T=T, mon.names=mon.names, parm.names=parm.names,
  pos.alpha=pos.alpha, pos.phi=pos.phi, pos.omega=pos.omega,
  pos.theta=pos.theta, y=y)

```

### 92.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[Data$pos.alpha]
  parm[Data$pos.phi] <- phi <- interval(parm[Data$pos.phi], -1, 1)
  omega <- interval(parm[Data$pos.omega], 1e-100, Inf)
  parm[Data$pos.omega] <- omega
  theta <- interval(parm[Data$pos.theta], 0.001, 0.999)
  parm[Data$pos.theta] <- theta
  ### Log(Prior Densities)
  alpha.prior <- dnormv(alpha, 0, 1000, log=TRUE)
  phi.prior <- dunif(phi, -1, 1, log=TRUE)
  omega.prior <- dhalfcauchy(omega, 25, log=TRUE)
  theta.prior <- sum(dunif(theta, 0, 1, log=TRUE))
  ### Log-Likelihood
  mu <- alpha + c(0, Data$y[-Data$T]) * phi
  epsilon <- Data$y - mu
  delta <- (epsilon > 0) * 1
  sigma2 <- omega + theta[1] * c(0,delta[-Data$T]) *
    c(0, epsilon[-Data$T]^2)
  sigma2[-1] <- sigma2[-1] + theta[2] * (1 - delta[-Data$T]) *
    epsilon[-Data$T]^2
  sigma2.new <- omega + theta[1] * delta[Data$T] * epsilon[Data$T]^2 +
    theta[2] * (1 - delta[Data$T]) * epsilon[Data$T]^2
  ynew <- rnormv(1, alpha + Data$y[Data$T] * phi, sigma2.new)
  LL <- sum(dnormv(Data$y[-1], mu[-1], sigma2[-1], log=TRUE))
  ### Log-Posterior
  LP <- LL + alpha.prior + phi.prior + omega.prior + theta.prior

```

```

Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP, ynew, sigma2.new),
  yhat=rnormv(length(mu), mu, sigma2), parm=parm)
return(Modelout)
}

```

## 92.4. Initial Values

```
Initial.Values <- c(0, 0, 1, 0.5, 0.5)
```

# 93. Threshold Autoregression (TAR)

## 93.1. Form

$$\begin{aligned}
 \mathbf{y}_t &\sim \mathcal{N}(\nu_t, \sigma^2), \quad t = 1, \dots, T \\
 \mathbf{y}^{new} &= \alpha_2 + \phi_2 \mathbf{y}_T \\
 \nu_t &= \begin{cases} \alpha_1 + \phi_1 \mathbf{y}_{t-1}, & t = 1, \dots, T \quad \text{if } t \geq \theta \\ \alpha_2 + \phi_2 \mathbf{y}_{t-1}, & t = 1, \dots, T \quad \text{if } t < \theta \end{cases} \\
 \alpha_j &\sim \mathcal{N}(0, 1000) \in [-1, 1], \quad j = 1, \dots, 2 \\
 \phi_j &\sim \mathcal{N}(0, 1000) \in [-1, 1], \quad j = 1, \dots, 2 \\
 \theta &\sim \mathcal{U}(2, T-1) \\
 \sigma &\sim \mathcal{HC}(25)
 \end{aligned}$$

## 93.2. Data

```

y <- c(0.02, -0.51, -0.30, 1.46, -1.26, -2.15, -0.91, -0.53, -1.91,
  2.64, 1.64, 0.15, 1.46, 1.61, 1.96, -2.67, -0.19, -3.28,
  1.89, 0.91, -0.71, 0.74, -0.10, 3.20, -0.80, -5.25, 1.03,
  -0.40, -1.62, -0.80, 0.77, 0.17, -1.39, -1.28, 0.48, -1.02,
  0.09, -1.09, 0.86, 0.36, 1.51, -0.02, 0.47, 0.62, -1.36,
  1.12, 0.42, -4.39, -0.87, 0.05, -5.41, -7.38, -1.01, -1.70,
  0.64, 1.16, 0.87, 0.28, -1.69, -0.29, 0.13, -0.65, 0.83,
  0.62, 0.05, -0.14, 0.01, -0.36, -0.32, -0.80, -0.06, 0.24,
  0.23, -0.37, 0.00, -0.33, 0.21, -0.10, -0.10, -0.01, -0.40,
  -0.35, 0.48, -0.28, 0.08, 0.28, 0.23, 0.27, -0.35, -0.19,
  0.24, 0.17, -0.02, -0.23, 0.03, 0.02, -0.17, 0.04, -0.39,
  -0.12, 0.16, 0.17, 0.00, 0.18, 0.06, -0.36, 0.22, 0.14,
  -0.17, 0.10, -0.01, 0.00, -0.18, -0.02, 0.07, -0.06, 0.06,
  -0.05, -0.08, -0.07, 0.01, -0.06, 0.01, 0.01, -0.02, 0.01,
  0.01, 0.12, -0.03, 0.08, -0.10, 0.01, -0.03, -0.08, 0.04,
  -0.09, -0.08, 0.01, -0.05, 0.08, -0.14, 0.06, -0.11, 0.09,
  0.06, -0.12, -0.01, -0.05, -0.15, -0.05, -0.03, 0.04, 0.00,

```



```

-0.12, 0.04, -0.06, -0.05, -0.07, -0.05, -0.14, -0.05, -0.01,
-0.12, 0.05, 0.06, -0.10, 0.00, 0.01, 0.00, -0.08, 0.00,
0.00, 0.07, -0.01, 0.00, 0.09, 0.33, 0.13, 0.42, 0.24,
-0.36, 0.22, -0.09, -0.19, -0.10, -0.08, -0.07, 0.05, 0.07,
0.07, 0.00, -0.04, -0.05, 0.03, 0.08, 0.26, 0.10, 0.08,
0.09, -0.07, -0.33, 0.17, -0.03, 0.07, -0.04, -0.06, -0.06,
0.07, -0.03, 0.00, 0.08, 0.27, 0.11, 0.11, 0.06, -0.11,
-0.09, -0.21, 0.24, -0.12, 0.11, -0.02, -0.03, 0.02, -0.10,
0.00, -0.04, 0.01, 0.02, -0.03, -0.10, -0.09, 0.17, 0.07,
-0.05, -0.01, -0.05, 0.01, 0.00, -0.08, -0.05, -0.08, 0.07,
0.06, -0.14, 0.02, 0.01, 0.04, 0.00, -0.13, -0.17)
T <- length(y)
mon.names <- c("LP", "ynew")
parm.names <- as.parm.names(list(alpha=rep(0,2), phi=rep(0,2), theta=0,
sigma=0))
pos.alpha <- grep("alpha", parm.names)
pos.phi <- grep("phi", parm.names)
pos.theta <- grep("theta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rtrunc(4, "norm", a=-1, b=1, mean=0,
sd=sqrt(1000)), runif(1,2,Data$T-1), rhalfcauchy(1,5)))
MyData <- list(PGF=PGF, T=T, mon.names=mon.names, parm.names=parm.names,
pos.alpha=pos.alpha, pos.phi=pos.phi, pos.theta=pos.theta,
pos.sigma=pos.sigma, y=y)

```

### 93.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  parm[Data$pos.alpha] <- alpha <- interval(parm[Data$pos.alpha], -1, 1)
  parm[Data$pos.phi] <- phi <- interval(parm[Data$pos.phi], -1, 1)
  theta <- interval(parm[Data$pos.theta], 2, Data$T-1)
  parm[Data$pos.theta] <- theta
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  alpha.prior <- sum(dtrunc(alpha, "norm", a=-1, b=1, mean=0,
sd=sqrt(1000), log=TRUE))
  phi.prior <- sum(dtrunc(phi, "norm", a=-1, b=1, mean=0,
sd=sqrt(1000), log=TRUE))
  alpha.prior <- sum(dnormv(alpha, 0, 1000, log=TRUE))
  phi.prior <- sum(dnormv(phi, 0, 1000, log=TRUE))
  theta.prior <- dunif(theta, 2, Data$T-1, log=TRUE)
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
  ### Log-Likelihood

```

```

mu <- matrix(0, Data$T, 2)
mu[,1] <- c(alpha[1], alpha[1] + phi[1]*Data$y[-Data$T])
mu[,2] <- c(alpha[2], alpha[2] + phi[2]*Data$y[-Data$T])
nu <- mu[,2]; temp <- which(1:Data$T < theta)
nu[temp] <- mu[temp,1]
ynew <- rnorm(1, alpha[2] + phi[2]*Data$y[Data$T], sigma)
LL <- sum(dnorm(Data$y, nu, sigma, log=TRUE))
### Log-Posterior
LP <- LL + alpha.prior + phi.prior + theta.prior + sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,ynew),
  yhat=rnorm(length(nu), nu, sigma), parm=parm)
return(Modelout)
}

```

### 93.4. Initial Values

```
Initial.Values <- c(rep(0,4), T/2, 1)
```

## 94. Time Varying AR(1) with Chebyshev Series

This example consists of a first-order autoregressive model, AR(1), with a time-varying parameter (TVP)  $\phi$ , that is a Chebyshev series constructed from a linear combination of orthonormal Chebyshev time polynomials (CTPs) and parameter vector  $\beta$ . The user creates basis matrix  $\mathbf{P}$ , specifying polynomial degree  $D$  and time  $T$ . Each column is a CTP of a different degree, and the first column is restricted to 1, the linear basis. CTPs are very flexible for TVPs, and estimate quickly because each is orthogonal, unlike simple polynomials and splines.

### 94.1. Form

$$\begin{aligned}
 \mathbf{y}_t &\sim \mathcal{N}(\mu_t, \sigma^2), \quad t = 1, \dots, T \\
 \mu_t &= \alpha + \phi_{t-1} \mathbf{y}_{t-1} \\
 \phi_t &= \mathbf{P} \beta \\
 \alpha &\sim \mathcal{N}(0, 1000) \\
 \beta_d &\sim \mathcal{N}(0, 1000), \quad d = 1, \dots, (D+1) \\
 \sigma &\sim \mathcal{HC}(25)
 \end{aligned}$$

### 94.2. Data

```

D <- 6 #Maximum degree of Chebyshev time polynomials
T <- 100
P <- matrix(1, T, D+1)
for (d in 1:D) {P[,d+1] <- sqrt(2)*cos(d*pi*(c(1:T)-0.5)/T)}

```

```

alpha.orig <- 0; alpha.orig
beta.orig <- runif(D+1,-0.3,0.3); beta.orig
phi.orig <- tcrossprod(P, t(beta.orig))
e <- rnorm(T,0,1)
y <- rep(0,T)
for (t in 2:T) {y[t] <- alpha.orig + phi.orig[t-1]*y[t-1] + e[t]}
mon.names <- c("LP", "ynew", as.parm.names(list(phi=rep(0,T-1))))
parm.names <- as.parm.names(list(alpha=0, beta=rep(0,D+1), sigma=0))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(D+2,0,10), rhalfcauchy(1,5)))
MyData <- list(D=D, P=P, PGF=PGF, T=T, mon.names=mon.names,
  parm.names=parm.names, pos.alpha=pos.alpha, pos.beta=pos.beta,
  pos.sigma=pos.sigma, y=y)

```

### 94.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[Data$pos.alpha]
  beta <- parm[Data$pos.beta]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  alpha.prior <- dnormv(alpha, 0, 1000, log=TRUE)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
  ### Log-Likelihood
  phi <- tcrossprod(Data$P[-Data$T,], t(beta))
  mu <- c(alpha, alpha + phi*Data$y[-Data$T])
  ynew <- rnorm(1, alpha + tcrossprod(Data$P[Data$T,], t(beta))*
    Data$y[Data$T], sigma)
  LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + alpha.prior + beta.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,ynew,phi),
    yhat=rnorm(length(mu), mu, sigma), parm=parm)
  return(Modelout)
}

```

### 94.4. Initial Values

```
Initial.Values <- c(rep(0,D+2), 1)
```

## 95. Variable Selection, BAL

This approach to variable selection is one of several forms of the Bayesian Adaptive Lasso (BAL). The lasso applies shrinkage to exchangeable scale parameters,  $\gamma$ , for the regression effects,  $\beta$ .

### 95.1. Form

$$\begin{aligned} \mathbf{y} &\sim \mathcal{N}(\mu, \sigma^2) \\ \mu &= \mathbf{X}\beta \\ \beta_1 &\sim \mathcal{L}(0, 1000) \\ \beta_j &\sim \mathcal{L}(0, \gamma_j), \quad j = 2, \dots, J \\ \gamma_j &\sim \mathcal{G}^{-1}(\delta, \tau), \quad \in [0, \infty] \\ \delta &\sim \mathcal{HC}(25) \\ \tau &\sim \mathcal{HC}(25) \\ \sigma &\sim \mathcal{HC}(25) \end{aligned}$$

### 95.2. Data

```
data(demonsnacks)
J <- ncol(demonsnacks)
y <- log(demonsnacks$Calories)
X <- cbind(1, as.matrix(demonsnacks[,c(1,3:10)]))
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J), gamma=rep(0,J-1),
  log.delta=0, log.tau=0, log.sigma=0))
pos.beta <- grep("beta", parm.names)
pos.gamma <- grep("gamma", parm.names)
pos.delta <- grep("delta", parm.names)
pos.tau <- grep("tau", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,10),
  rgamma(Data$J-1,rhalfcauchy(1,5),rhalfcauchy(1,5)), rhalfcauchy(3,5)))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.gamma=pos.gamma,
  pos.delta=pos.delta, pos.tau=pos.tau, pos.sigma=pos.sigma, y=y)
```

### 95.3. Model

```
Model <- function(parm, Data)
{
```

```

### Hyperhyperparameters
delta <- interval(parm[Data$pos.delta], 1e-100, Inf)
parm[Data$pos.delta] <- delta
parm[Data$pos.tau] <- tau <- interval(parm[Data$pos.tau], 1e-100, Inf)
### Hyperparameters
gamma <- interval(parm[Data$pos.gamma], 1e-100, Inf)
parm[Data$pos.gamma] <- gamma
### Parameters
beta <- parm[Data$pos.beta]
sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
parm[Data$pos.sigma] <- sigma
### Log(Hyperhyperprior Densities)
delta.prior <- dhalfcauchy(delta, 25, log=TRUE)
tau.prior <- dhalfcauchy(tau, 25, log=TRUE)
### Log(Hyperprior Densities)
gamma.prior <- sum(dinvgamma(gamma, delta, tau, log=TRUE))
### Log(Prior Densities)
beta.prior <- sum(dlaplace(beta, 0, c(1000, gamma), log=TRUE))
sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
### Log-Likelihood
mu <- tcrossprod(Data$X, t(beta))
LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
### Log-Posterior
LP <- LL + beta.prior + gamma.prior + delta.prior + tau.prior +
      sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
                 yhat=rnorm(length(mu), mu, sigma), parm=parm)
return(Modelout)
}

```

## 95.4. Initial Values

```
Initial.Values <- c(rep(0,J), rep(0,J-1), rep(1,3))
```

# 96. Variable Selection, Horseshoe

## 96.1. Form

$$\mathbf{y} \sim \mathcal{N}(\mu, \sigma^2)$$

$$\mu = \mathbf{X}\beta$$

$$\beta_1 \sim \mathcal{N}(0, 1000)$$

$$\beta_j \sim \mathcal{HS}(0, \lambda_j, \tau, \sigma), \quad j = 2, \dots, J$$

$$\sigma \sim \mathcal{HC}(25)$$

## 96.2. Data

```
data(demonsnacks)
J <- ncol(demonsnacks)
y <- log(demonsnacks$Calories)
X <- cbind(1, as.matrix(demonsnacks[,c(1,3:10)]))
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J), lambda=rep(0,J-1),
  tau=0, sigma=0))
pos.beta <- grep("beta", parm.names)
pos.lambda <- grep("lambda", parm.names)
pos.tau <- grep("tau", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,10), runif(Data$J,0,100),
  runif(2,0,100)))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.lambda=pos.lambda,
  pos.tau=pos.tau, pos.sigma=pos.sigma, y=y)
```

## 96.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  lambda <- interval(parm[Data$pos.lambda], 1e-100, 1000)
  parm[Data$pos.lambda] <- lambda
  parm[Data$pos.tau] <- tau <- interval(parm[Data$pos.tau], 1e-100, 1000)
  sigma <- interval(parm[Data$pos.sigma], 1e-100, 1000)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta[1], 0, 1000, log=TRUE),
    dhs(beta[-1], lambda, tau, sigma, log=TRUE))
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, t(beta))
  LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rnorm(length(mu), mu, sigma), parm=parm)
  return(Modelout)
}
```

## 96.4. Initial Values

```
Initial.Values <- c(rep(0,J), rep(1,J), rep(1,2))
```

## 97. Variable Selection, RJ

This example uses the RJ (Reversible-Jump) algorithm of the `LaplacesDemon` function for variable selection and Bayesian Model Averaging (BMA). Other MCMC algorithms will not perform variable selection with this example, as presented. This is an example of variable selection in a linear regression. The only difference between the following example, and the example of linear regression (46), is that RJ specifications are also included for the RJ algorithm, and that the RJ algorithm must be used.

### 97.1. Form

$$\mathbf{y} \sim \mathcal{N}(\mu, \sigma^2)$$

$$\mu = \mathbf{X}\beta$$

$$\beta_j \sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J$$

$$\sigma \sim \mathcal{HC}(25)$$

### 97.2. Data

```
N <- 1000
J <- 100 #Number of predictors, including the intercept
X <- matrix(1,N,J)
for (j in 2:J) {X[,j] <- rnorm(N,runif(1,-3,3),runif(1,0.1,1))}
beta.orig <- runif(J,-3,3)
zero <- sample(2:J, round(J*0.9)) #Assign most parameters to be zero
beta.orig[zero] <- 0
e <- rnorm(N,0,0.1)
y <- as.vector(tcrossprod(beta.orig, X) + e)
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J), sigma=0))
pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,1), rhalfcauchy(1,5)))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.sigma=pos.sigma, y=y)
### Reversible-Jump Specifications bin.n <- J-1 #Maximum allowable model size
bin.p <- 0.4 #Most probable size: bin.p x bin.n is binomial mean and median
parm.p <- rep(1/J,J+1)
```

```
selectable=c(0, rep(1,J-1), 0)
```

### 97.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, t(beta))
  LL <- sum(dnorm(Data$y, mu, sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rnorm(length(mu), mu, sigma), parm=parm)
  return(Modelout)
}
```

### 97.4. Initial Values

```
Initial.Values <- GIV(Model, MyData, PGF=TRUE)
```

## 98. Variable Selection, SSVS

This example uses a modified form of the random-effects (or global adaptation) Stochastic Search Variable Selection (SSVS) algorithm presented in [O'Hara and Sillanpaa \(2009\)](#), which selects variables according to practical significance rather than statistical significance. Here, SSVS is applied to linear regression, though this method is widely applicable. For  $J$  variables, each regression effect  $\beta_j$  is conditional on  $\gamma_j$ , a binary inclusion variable. Each  $\beta_j$  is a discrete mixture distribution with respect to  $\gamma_j = 0$  or  $\gamma_j = 1$ , with precision 100 or  $\beta_\sigma = 0.1$ , respectively. As with other representations of SSVS, these precisions may require tuning.

The binary inclusion variables are discrete parameters, and discrete parameters are not supported in all algorithms. The example below is updated with the Griddy-Gibbs sampler.

When the goal is to select the best model, each  $\mathbf{X}_{1:N,j}$  is retained for a future run when the posterior mean of  $\gamma_j \geq 0.5$ . When the goal is model-averaging, the results of this model may be used directly, which would please L. J. Savage, who said that “models should be as big as an elephant” ([Draper 1995](#)).



### 98.1. Form

$$\begin{aligned}
 \mathbf{y} &\sim \mathcal{N}(\boldsymbol{\mu}, \sigma^2) \\
 \boldsymbol{\mu} &= \mathbf{X}\boldsymbol{\beta} \\
 (\beta_j | \gamma_j) &\sim (1 - \gamma_j)\mathcal{N}(0, 0.01) + \gamma_j\mathcal{N}(0, \beta_\sigma^2) \quad j = 1, \dots, J \\
 \beta_\sigma &\sim \mathcal{HC}(25) \\
 \gamma_j &\sim \mathcal{BERN}(1/(J-1)), \quad j = 1, \dots, (J-1) \\
 \sigma &\sim \mathcal{HC}(25)
 \end{aligned}$$

### 98.2. Data

```

data(demonsnacks)
J <- ncol(demonsnacks)
y <- log(demonsnacks$Calories)
X <- cbind(1, as.matrix(demonsnacks[,c(1,3:10)]))
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
mon.names <- c("LP", "min.beta.sigma")
parm.names <- as.parm.names(list(beta=rep(0,J), gamma=rep(0,J-1),
  b.sd=0, sigma=0))
pos.beta <- grep("beta", parm.names)
pos.gamma <- grep("gamma", parm.names)
pos.b.sd <- grep("b.sd", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnorm(Data$J,0,1), rcat(Data$J-1, p=rep(0.5,2)),
  rhalfcauchy(2,5)))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
  parm.names=parm.names, pos.beta=pos.beta, pos.gamma=pos.gamma,
  pos.b.sd=pos.b.sd, pos.sigma=pos.sigma, y=y)

```

### 98.3. Model

```

Model <- function(parm, Data)
{
  ### Hyperparameters
  beta.sigma <- interval(parm[Data$pos.b.sd], 1e-100, Inf)
  parm[Data$pos.b.sd] <- beta.sigma
  ### Parameters
  beta <- parm[Data$pos.beta]
  gamma <- parm[Data$pos.gamma]
  beta.sigma <- rep(beta.sigma, Data$J-1)
  beta.sigma[gamma == 0] <- 0.1
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  ### Log(Hyperprior Densities)

```

```

beta.sigma.prior <- sum(dhalfcauchy(beta.sigma, 25, log=TRUE))
### Log(Prior Densities)
beta.prior <- sum(dnorm(beta, 0, beta.sigma, log=TRUE))
gamma.prior <- sum(dbern(gamma, 1/(Data$J-1), log=TRUE))
sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
### Log-Likelihood
mu <- tcrossprod(Data$X, t(beta*c(1,gamma)))
LL <- sum(dnorm(y, mu, sigma, log=TRUE))
### Log-Posterior
LP <- LL + beta.prior + beta.sigma.prior + gamma.prior + sigma.prior
Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP, min(beta.sigma)),
  yhat=rnorm(length(mu), mu, sigma), parm=parm)
return(Modelout)
}

```

#### 98.4. Initial Values

```
Initial.Values <- c(rep(0,J), rep(1,J-1), rep(1,2))
```

### 99. Vector Autoregression, VAR(1)

#### 99.1. Form

$$\begin{aligned}
\mathbf{Y}_{t,j} &\sim \mathcal{N}(\mu_{t,j}, \sigma_j^2), \quad t = 1, \dots, T, \quad j = 1, \dots, J \\
\mu_{t,j} &= \alpha_j + \Phi_{1:J,j} \mathbf{Y}_{t-1,j} \\
\mathbf{y}_j^{new} &= \alpha_j + \Phi_{1:J,j} \mathbf{Y}_{T,j} \\
\alpha_j &\sim \mathcal{N}(0, 1000) \\
\sigma_j &\sim \mathcal{HC}(25) \\
\Phi_{i,k} &\sim \mathcal{N}(0, 1000), \quad i = 1, \dots, J, \quad k = 1, \dots, J
\end{aligned}$$

#### 99.2. Data

```

T <- 100
J <- 3
Y <- matrix(0,T,J)
for (j in 1:J) {for (t in 2:T) {
  Y[t,j] <- Y[t-1,j] + rnorm(1,0,0.1)}}
mon.names <- c("LP", as.parm.names(list(ynew=rep(0,J))))
parm.names <- as.parm.names(list(alpha=rep(0,J), Phi=matrix(0,J,J),
  sigma=rep(0,J)))
pos.alpha <- grep("alpha", parm.names)

```

```

pos.Phi <- grep("Phi", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormmv(Data$J,0,1),
  rnormmv(Data$J*Data$J,0,1), rhalfcauchy(Data$J,5)))
MyData <- list(J=J, PGF=PGF, T=T, Y=Y, mon.names=mon.names,
  parm.names=parm.names, pos.alpha=pos.alpha, pos.Phi=pos.Phi,
  pos.sigma=pos.sigma)

```

### 99.3. Model

```

Model <- function(parm, Data)
{
  ### Parameters
  alpha <- parm[Data$pos.alpha]
  Phi <- matrix(parm[Data$pos.phi], Data$J, Data$J)
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  alpha.prior <- sum(dnormmv(alpha, 0, 1000, log=TRUE))
  Phi.prior <- sum(dnormmv(Phi, 0, 1000, log=TRUE))
  sigma.prior <- sum(dhalfcauchy(sigma, 25, log=TRUE))
  ### Log-Likelihood
  mu <- matrix(alpha,Data$T,Data$J,byrow=TRUE)
  mu[-1,] <- mu[-1,] + tcrossprod(Data$Y[-Data$T,], Phi)
  Sigma <- matrix(sigma, Data$T, Data$J, byrow=TRUE)
  ynew <- rnorm(Data$J, alpha + as.vector(crossprod(Phi, Data$Y[Data$T,])),
    sigma)
  LL <- sum(dnorm(Data$Y, mu, Sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + alpha.prior + Phi.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=c(LP,ynew),
    yhat=rnorm(prod(dim(mu)), mu, Sigma), parm=parm)
  return(Modelout)
}

```

### 99.4. Initial Values

```
Initial.Values <- c(colMeans(Y), rep(0,J*J), rep(1,J))
```

## 100. Weighted Regression

It is easy enough to apply record-level weights to the likelihood. Here, weights are applied to the linear regression example in [section 46](#).

### 100.1. Form

$$\begin{aligned} \mathbf{y} &\sim \mathcal{N}(\boldsymbol{\mu}, \sigma^2) \\ \boldsymbol{\mu} &= \mathbf{X}\boldsymbol{\beta} \\ \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J \\ \sigma &\sim \mathcal{HC}(25) \end{aligned}$$

### 100.2. Data

```
data(demonsnacks)
N <- nrow(demonsnacks)
J <- ncol(demonsnacks)
y <- log(demonsnacks$Calories)
X <- cbind(1, as.matrix(demonsnacks[,c(1,3:10)]))
for (j in 2:J) {X[,j] <- CenterScale(X[,j])}
w <- c(rep(1,5), 0.2, 1, 0.01, rep(1,31))
w <- w * (sum(w) / N)
mon.names <- "LP"
parm.names <- as.parm.names(list(beta=rep(0,J), sigma=0))
pos.beta <- grep("beta", parm.names)
pos.sigma <- grep("sigma", parm.names)
PGF <- function(Data) return(c(rnormv(Data$J,0,1000), rhalfcauchy(1,5)))
MyData <- list(J=J, PGF=PGF, X=X, mon.names=mon.names,
               parm.names=parm.names, pos.beta=pos.beta, pos.sigma=pos.sigma, w=w,
               y=y)
```

### 100.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  beta <- parm[Data$pos.beta]
  sigma <- interval(parm[Data$pos.sigma], 1e-100, Inf)
  parm[Data$pos.sigma] <- sigma
  ### Log(Prior Densities)
  beta.prior <- sum(dnormv(beta, 0, 1000, log=TRUE))
  sigma.prior <- dhalfcauchy(sigma, 25, log=TRUE)
  ### Log-Likelihood
  mu <- tcrossprod(Data$X, t(beta))
  LL <- sum(w * dnorm(Data$y, mu, sigma, log=TRUE))
  ### Log-Posterior
  LP <- LL + beta.prior + sigma.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
                  yhat=rnorm(length(mu), mu, sigma), parm=parm)
```

```
return(Modelout)
}
```

#### 100.4. Initial Values

```
Initial.Values <- c(rep(0,J), 1)
```

### 101. Zero-Inflated Poisson (ZIP)

#### 101.1. Form

$$\begin{aligned}
 \mathbf{y} &\sim \mathcal{P}(\Lambda_{1:N,2}) \\
 \mathbf{z} &\sim \mathcal{BERN}(\Lambda_{1:N,1}) \\
 \mathbf{z}_i &= \begin{cases} 1 & \text{if } \mathbf{y}_i = 0 \\ 0 & \text{otherwise} \end{cases} \\
 \Lambda_{i,2} &= \begin{cases} 0 & \text{if } \Lambda_{i,1} \geq 0.5 \\ \Lambda_{i,2} & \text{otherwise} \end{cases} \\
 \Lambda_{1:N,1} &= \frac{1}{1 + \exp(-\mathbf{X}_1 \alpha)} \\
 \Lambda_{1:N,2} &= \exp(\mathbf{X}_2 \beta) \\
 \alpha_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J_1 \\
 \beta_j &\sim \mathcal{N}(0, 1000), \quad j = 1, \dots, J_2
 \end{aligned}$$

#### 101.2. Data

```
N <- 1000
J1 <- 4
J2 <- 3
X1 <- matrix(runif(N*J1,-2,2),N,J1); X1[,1] <- 1
X2 <- matrix(runif(N*J2,-2,2),N,J2); X2[,1] <- 1
alpha <- runif(J1,-1,1)
beta <- runif(J2,-1,1)
p <- invlogit(tcrossprod(X1, t(alpha)) + rnorm(N,0,0.1))
mu <- round(exp(tcrossprod(X2, t(beta)) + rnorm(N,0,0.1)))
y <- ifelse(p > 0.5, 0, mu)
z <- ifelse(y == 0, 1, 0)
mon.names <- "LP"
parm.names <- as.parm.names(list(alpha=rep(0,J1), beta=rep(0,J2)))
pos.alpha <- grep("alpha", parm.names)
pos.beta <- grep("beta", parm.names)
```

```
PGF <- function(Data) return(c(rnormv(Data$J1,0,5), rnormv(Data$J2,0,5)))
MyData <- list(J1=J1, J2=J2, N=N, PGF=PGF, X1=X1, X2=X2,
  mon.names=mon.names, parm.names=parm.names, pos.alpha=pos.alpha,
  pos.beta=pos.beta, y=y, z=z)
```

### 101.3. Model

```
Model <- function(parm, Data)
{
  ### Parameters
  parm[Data$pos.alpha] <- alpha <- interval(parm[Data$pos.alpha], -5, 5)
  parm[Data$pos.beta] <- beta <- interval(parm[Data$pos.beta], -5, 5)
  ### Log(Prior Densities)
  alpha.prior <- sum(dnormv(alpha, 0, 5, log=TRUE))
  beta.prior <- sum(dnormv(beta, 0, 5, log=TRUE))
  ### Log-Likelihood
  Lambda <- matrix(NA, Data$N, 2)
  Lambda[,1] <- invlogit(tcrossprod(Data$X1, t(alpha)))
  Lambda[,2] <- exp(tcrossprod(Data$X2, t(beta))) + 1e-100
  Lambda[which(Lambda[,1] >= 0.5),2] <- 0
  LL <- sum(dbern(Data$z, Lambda[,1], log=TRUE),
    dpois(Data$y, Lambda[,2], log=TRUE))
  ### Log-Posterior
  LP <- LL + alpha.prior + beta.prior
  Modelout <- list(LP=LP, Dev=-2*LL, Monitor=LP,
    yhat=rpois(nrow(Lambda), Lambda[,2]), parm=parm)
  return(Modelout)
}
```

### 101.4. Initial Values

```
Initial.Values <- GIV(Model, MyData, n=10000)
```

## References

- Barnard J, McCulloch R, Meng X (2000). “Modeling Covariance Matrices in Terms of Standard Deviations and Correlations, with Application to Shrinkage.” *Statistica Sinica*, **10**, 1281–1311.
- Congdon P (2003). *Applied Bayesian Modelling*. John Wiley & Sons, West Sussex, England.
- Draper D (1995). “Assessment and Propagation of Model Uncertainty.” *Journal of the Royal Statistical Society*, **B 57**(1), 45–97.

- Gelman A (2013). *R2WinBUGS: Running WinBUGS and OpenBUGS from R / S-PLUS*. R package version 2.1-19, URL <http://cran.r-project.org/web/packages/R2WinBUGS/index.html>.
- Gelman A, Carlin J, Stern H, Rubin D (2004). *Bayesian Data Analysis*. 2nd edition. Chapman & Hall, Boca Raton, FL.
- Ibrahim J, Chen M (2000). “Power Prior Distributions for Regression Models.” *Statistical Science*, **15**, 46–60.
- Kim J, Allenby G, Rossi P (2002). “Modeling Consumer Demand for Variety.” *Marketing Science*, **21**(3), 229–250.
- Kleine L (1950). *Economic Fluctuations in the United States 1921-1940*. John Wiley & Sons, New York, New York.
- Kotz S, Kozubowski T, Podgorski K (2001). *The Laplace Distribution and Generalizations: A Revisit with Applications to Communications, Economics, Engineering, and Finance*. Birkhauser, Boston.
- O’Hara R, Sillanpaa M (2009). “A Review of Bayesian Variable Selection Methods: What, How and Which.” *Journal of Bayesian Analysis*, **4**(1), 85–118.
- Spiegelhalter D, Thomas A, Best N, Lunn D (2003). *WinBUGS User Manual, Version 1.4*. MRC Biostatistics Unit, Institute of Public Health and Department of Epidemiology and Public Health, Imperial College School of Medicine, UK.
- Statisticat LLC (2014). *LaplacesDemon: Complete Environment for Bayesian Inference*. R package version 14.05.07, URL <http://www.bayesian-inference.com/software>.
- Zellner A (1962). “An Efficient Method of Estimating Seemingly Unrelated Regression Equations and Tests for Aggregation Bias.” *Journal of the American Statistical Association*, **57**, 348–368.

### Affiliation:

Statisticat, LLC

Farmington, CT

E-mail: [software@bayesian-inference.com](mailto:software@bayesian-inference.com)

URL: <http://www.bayesian-inference.com/software>