

# Program Reference

January 18, 2013

## Contents

<b>Interface</b>	<b>1</b>
C routine . . . . .	1
Fortran routine . . . . .	2
Supported angular momentum . . . . .	3
Data ordering . . . . .	3
Tensor . . . . .	4
<b>function list</b>	<b>4</b>

## Interface

### C routine

```
f(buf, shls, atm, natm, bas, nbas, env);
```

- buf: column-major double precision array.
  - for 1e integrals of shells (i,j), data are stored as [i1j1 i2j1 ... ]
  - for 2e integrals of shells (i,j|k,l), data are stored as  
[i1j1k1l1 i2j1k1l1 ... i1j2k1l1 ... i1j1k2l1 ... ]
  - complex data are stored as two double elements, first is real, followed by imaginary, e.g. [Re Im Re Im ...]
- shls: 0-based basis/shell indices.
  - int[2] for 1e integrals

- int[4] for 2e integrals
- atm: int[natm\*6], list of atoms. For ith atom, the 6 slots of atm[i] are
  - atm[i\*6+0] nuclear charge of atom i
  - atm[i\*6+1] env offset to save coordinates (env[atm[i\*6+1]], env[atm[i\*6+1]+1], env[atm[i\*6+1]+2]) are (x,y,z)
  - atm[i\*6+2] nuclear model of atom i, = 2 indicates gaussian nuclear model
  - atm[i\*6+3] unused
  - atm[i\*6+4] unused
  - atm[i\*6+5] unused
- natm: int, number of atoms, natm has no effect **except nuclear attraction** integrals
- bas: int[nbas\*8], list of basis. For ith basis, the 8 slots of bas[i] are
  - bas[i\*8+0] 0-based index of corresponding atom
  - bas[i\*8+1] angular momentum
  - bas[i\*8+2] number of primitive GTO in basis i
  - bas[i\*8+3] number of contracted GTO in basis i
  - bas[i\*8+4] kappa for spinor GTO.
    - < 0 the basis  $\sim j = l + 1/2$ .
    - > 0 the basis  $\sim j = l - 1/2$ .
    - = 0 the basis includes both  $j = l + 1/2$  and  $j = l - 1/2$
  - bas[i\*8+5] env offset to save exponents of primitive GTOs. e.g. 10 exponents env[bas[i\*8+5]] ... env[bas[i\*8+5]+9]
  - bas[i\*8+6] env offset to save column-major contraction coefficients. e.g. 10 primitive -> 5 contraction needs a  $10 \times 5$  array

env[bas[i*8+6]]	env[bas[i*8+6]]+10	env[bas[i*8+6]]+40
env[bas[i*8+6]]+1	env[bas[i*8+6]]+11	env[bas[i*8+6]]+41
.	.	...
.	.	.
env[bas[i*8+6]]+9	env[bas[i*8+6]]+19	env[bas[i*8+6]]+49

– ‘bas[i\*8+7]’ unused

- nbas: int, number of bases, nbas has no effect
- env: double[], save the value of coordinates, exponents, contraction coefficients

## Fortran routine

call f(buf, shls, atm, natm, bas, nbas, env)

- atm and bas are 2D integer array
  - atm(1:6,i) is the (charge, offset\_coord, nuclear\_model, unused, unused, unused) of the ith atom
  - bas(1:8,i) is the (atom\_index, angular, num\_primitive\_GTO, num\_contract\_GTO, kappa, offset\_exponent, offset\_coeff, unused) of the ith basis
- parameters are the same to the C function. Note those offsets atm(2,i) bas(6,i) bas(7,i) are 0-based.
- buf is 2D/4D double precision/double complex array

## Supported angular momentum

- 1e integrals  $l_{max} = 6$
- 2e integrals  $l_{max} = 4$

## Data ordering

- for Cartesian GTO, the output data in buf are sorted as

s shell	p shell	d shell	...
...	...	...	
s	p $x$	d $xx$	
s	p $y$	d $xy$	
...	p $z$	d $xz$	
	p $x$	d $yy$	
	p $y$	d $yz$	
	p $z$	d $zz$	
	...	...	

- for real spheric GTO, the output data in buf are sorted as

s shell	p shell	d shell	f shell	...
...	...	...	...	
s	p $y$	d $xy$	f $y(3x^2 - y^2)$	
s	p $z$	d $yz$	f $xyz$	
...	p $x$	d $z^2$	f $yz^2$	
	p $y$	d $xz$	f $z^3$	
	p $z$	d $x^2 - y^2$	f $xz^2$	
	p $x$	...	f $z(x^2 - y^2)$	
	...		f $x(x^2 - 3y^2)$	
			...	

- for spinor GTO, the output data in buf correspond to

...	kappa=0,p shell	kappa=1,p shell	kappa=0,d shell	...
...	...	...	...	
$p_{1/2}(-1/2)$	$p_{1/2}(-1/2)$	$d_{3/2}(-3/2)$		
$p_{1/2}(1/2)$	$p_{1/2}(1/2)$	$d_{3/2}(-1/2)$		
$p_{3/2}(-3/2)$	$p_{1/2}(-1/2)$	$d_{3/2}(1/2)$		
$p_{3/2}(-1/2)$	$p_{1/2}(1/2)$	$d_{3/2}(3/2)$		
$p_{3/2}(1/2)$	$p_{1/2}(-1/2)$	$d_{5/2}(-5/2)$		
$p_{3/2}(3/2)$	$p_{1/2}(1/2)$	$d_{5/2}(-3/2)$		
$p_{1/2}(-1/2)$	...	$d_{5/2}(-1/2)$		
$p_{1/2}(1/2)$		$d_{3/2}(-3/2)$		
$p_{3/2}(-3/2)$		$d_{3/2}(-1/2)$		
$p_{3/2}(-1/2)$		...		
...				

## Tensor

integrals like Gradients have tensor components. the output data is

- 3-component tensor
  - X buf(:,0)
  - Y buf(:,1)
  - Z buf(:,2)
- 9-component tensor
  - XX buf(:,0)
  - XY buf(:,1)
  - XZ buf(:,2)
  - YX buf(:,3)
  - YY buf(:,4)
  - YZ buf(:,5)
  - ZX buf(:,6)
  - ZY buf(:,7)
  - ZZ buf(:,8)

## function list

- Cartesian GTO integrals

• cgto_cart	
• cint1e_ovlp_cart	$\langle i j\rangle$
• cint1e_nuc_cart	$\langle i V_{nuc} j\rangle$
• cint1e_kin_cart	$.5\langle i \vec{p}\cdot\vec{p} j\rangle$
• cint1e_ia01p_cart	$\langle i \frac{\vec{r}}{r^3} \times\vec{\nabla} j\rangle$
• cint1e_irxp_cart	$\langle i \vec{r}_c\times\vec{\nabla} j\rangle$
• cint1e_iking_cart	$0.5i\langle\vec{p}\cdot\vec{p} U_g j\rangle$
• cint1e_iovlp_cart	$i\langle i U_g j\rangle$
• cint1e_inucg_cart	$i\langle i V_{nuc} U_g j\rangle$
• cint1e_ipovlp_cart	$\langle\vec{\nabla}i j\rangle$
• cint1e_ipkin_cart	$0.5\langle\vec{\nabla}i \vec{p}\cdot\vec{p} j\rangle$
• cint1e_ipnuc_cart	$\langle\vec{\nabla}i V_{nuc} j\rangle$
• cint1e_iprinv_cart	$\langle\vec{\nabla}i r^{-1} j\rangle$
• cint1e_rinv_cart	$\langle i r^{-1} j\rangle$
• cint2e_cart	$(ij kl)$
• cint2e_ig1_cart	$i(U_gj kl)$

• cint2e_ip1_cart	$(\vec{\nabla} i j   k l)$
• Spheric GTO integrals	
• cgto_spheric	
• cint1e_ovlp_sph	$\langle i   j \rangle$
• cint1e_nuc_sph	$\langle i   V_{nuc}   j \rangle$
• cint1e_kin_sph	$0.5 \langle i   \vec{p} \cdot p j \rangle$
• cint1e_ia01p_sph	$\langle i   \frac{\vec{r}}{r^3}   \times \vec{\nabla} j \rangle$
• cint1e_irxp_sph	$\langle i   \vec{r}_c \times \vec{\nabla} j \rangle$
• cint1e_iking_sph	$0.5 i \langle \vec{p} \cdot \vec{p} i   U_g j \rangle$
• cint1e_iovlp_sph	$i \langle i   U_g j \rangle$
• cint1e_inucg_sph	$i \langle i   V_{nuc}   U_g j \rangle$
• cint1e_ipovlp_sph	$\langle \vec{\nabla} i   j \rangle$
• cint1e_ipkin_sph	$0.5 \langle \vec{\nabla} i   \vec{p} \cdot p j \rangle$
• cint1e_ipnuc_sph	$\langle \vec{\nabla} i   V_{nuc}   j \rangle$
• cint1e_iprinv_sph	$\langle \vec{\nabla} i   r^{-1}   j \rangle$
• cint1e_rinv_sph	$\langle i   r^{-1}   j \rangle$
• cint2e_sph	$(i j   k l)$

• cint2e_ig1_sph	$i(iU_g j kl)$
• cint2e_ip1_sph	$(\vec{\nabla} i j kl)$
• Spinor GTO integrals	
• cgto_spinor	
• cint1e_ovlp	$\langle i j\rangle$
• cint1e_nuc	$\langle i V_{nuc} j\rangle$
• cint1e_nucg	$\langle i V_{nuc} U_g j\rangle$
• cint1e_srsr	$\langle \vec{\sigma} \cdot \vec{r} i   \vec{\sigma} \cdot \vec{r} j \rangle$
• cint1e_sr	$\langle \vec{\sigma} \cdot \vec{r} i   j \rangle$
• cint1e_srsp	$\langle \vec{\sigma} \cdot \vec{r} i   \vec{\sigma} \cdot \vec{p} j \rangle$
• cint1e_spsp	$\langle \vec{\sigma} \cdot \vec{p} i   \vec{\sigma} \cdot \vec{p} j \rangle$
• cint1e_sp	$\langle \vec{\sigma} \cdot \vec{p} i   j \rangle$
• cint1e_spspsp	$\langle \vec{\sigma} \cdot \vec{p} i   \vec{\sigma} \cdot \vec{p} \vec{\sigma} \cdot \vec{p} j \rangle$
• cint1e_spnuc	$\langle \vec{\sigma} \cdot \vec{p} i   V_{nuc}   j \rangle$
• cint1e_spnucsp	$\langle \vec{\sigma} \cdot \vec{p} i   V_{nuc}   \vec{\sigma} \cdot \vec{p} j \rangle$
• cint1e_srnucsr	$\langle \vec{\sigma} \cdot \vec{r} i   V_{nuc}   \vec{\sigma} \cdot \vec{r} j \rangle$
• cint1e_sa10sa01	$0.5 \langle \vec{\sigma} \times \vec{r}_e i   \vec{\sigma} \times \frac{\vec{r}}{r^3}   j \rangle$

• cint1e_ovlpg	$\langle i U_g j\rangle$
• cint1e_sa10sp	$0.5\langle \vec{r}_c \times \vec{\sigma}_i   \vec{\sigma} \cdot \vec{p} j\rangle$
• cint1e_sa10nucsp	$0.5\langle \vec{r}_c \times \vec{\sigma}_i   V_{nuc}   \vec{\sigma} \cdot \vec{p} j\rangle$
• cint1e_sa01sp	$\langle i   \frac{\vec{r}}{r^3} \times \vec{\sigma}   \vec{\sigma} \cdot \vec{p} j\rangle$
• cint1e_spgsp	$\langle U_g \vec{\sigma} \cdot \vec{p} i   \vec{\sigma} \cdot \vec{p} j\rangle$
• cint1e_spgnucsp	$\langle U_g \vec{\sigma} \cdot \vec{p} i   V_{nuc}   \vec{\sigma} \cdot \vec{p} j\rangle$
• cint1e_spgsa01	$\langle U_g \vec{\sigma} \cdot \vec{p} i   \frac{\vec{r}}{r^3} \times \vec{\sigma}   j\rangle$
• cint1e_ipovlp	$\langle \vec{\nabla} i   j\rangle$
• cint1e_ipkin	$0.5\langle \vec{\nabla} i   p \cdot p j\rangle$
• cint1e_ipnuc	$\langle \vec{\nabla} i   V_{nuc}   j\rangle$
• cint1e_iprinv	$\langle \vec{\nabla} i   r^{-1}   j\rangle$
• cint1e_ipspnucsp	$\langle \vec{\nabla} \vec{\sigma} \cdot \vec{p} i   V_{nuc}   \vec{\sigma} \cdot \vec{p} j\rangle$
• cint1e_ipsprinvsp	$\langle \vec{\nabla} \vec{\sigma} \cdot \vec{p} i   r^{-1}   \vec{\sigma} \cdot \vec{p} j\rangle$
• cint2e	$(ij kl)$
• cint2e_spsp1	$(\vec{\sigma} \cdot \vec{p} i \vec{\sigma} \cdot \vec{p} j   kl)$
• cint2e_spsp1spsp2	$(\vec{\sigma} \cdot \vec{p} i \vec{\sigma} \cdot \vec{p} j   \vec{\sigma} \cdot \vec{p} k \vec{\sigma} \cdot \vec{p} l)$



- cint2e\_srsr1  

$$(\vec{\sigma} \cdot \vec{r}_i \vec{\sigma} \cdot \vec{r}_j | kl)$$
- cint2e\_srsr1srsr2  

$$(\vec{\sigma} \cdot \vec{r}_i \vec{\sigma} \cdot \vec{r}_j | \vec{\sigma} \cdot \vec{r}_k \vec{\sigma} \cdot \vec{r}_l)$$
- cint2e\_sa10sp1  

$$0.5(\vec{r}_c \times \vec{\sigma}_i \vec{\sigma} \cdot \vec{p}_j | kl)$$
- cint2e\_sa10sp1spsp2  

$$0.5(\vec{r}_c \times \vec{\sigma}_i \vec{\sigma} \cdot \vec{p}_j | \vec{\sigma} \cdot \vec{p}_k \vec{\sigma} \cdot \vec{p}_l)$$
- cint2e\_g1  

$$(iU_{gj} | kl)$$
- cint2e\_spgsp1  

$$(\vec{\sigma} \cdot \vec{p}_i U_g \vec{\sigma} \cdot \vec{p}_j | kl)$$
- cint2e\_g1spsp2  

$$(iU_{gj} | \vec{\sigma} \cdot \vec{p}_k \vec{\sigma} \cdot \vec{p}_l)$$
- cint2e\_spgsp1spsp2  

$$(\vec{\sigma} \cdot \vec{p}_i U_g \vec{\sigma} \cdot \vec{p}_j | \vec{\sigma} \cdot \vec{p}_k \vec{\sigma} \cdot \vec{p}_l)$$
- cint2e\_ip1  

$$(\vec{\nabla}_{ij} | kl)$$
- cint2e\_ipspsp1  

$$(\vec{\nabla} \vec{\sigma} \cdot \vec{p}_i \vec{\sigma} \cdot \vec{p}_j | kl)$$
- cint2e\_ip1spsp2  

$$(\vec{\nabla}_{ij} | \vec{\sigma} \cdot \vec{p}_k \vec{\sigma} \cdot \vec{p}_l)$$
- cint2e\_ipspsp1spsp2  

$$(\vec{\nabla} \vec{\sigma} \cdot \vec{p}_i \vec{\sigma} \cdot \vec{p}_j | \vec{\sigma} \cdot \vec{p}_k \vec{\sigma} \cdot \vec{p}_l)$$
- cint2e\_ssp1ssp2  

$$(i\vec{\sigma} \vec{\sigma} \cdot \vec{p}_j | k\vec{\sigma} \vec{\sigma} \cdot \vec{p}_l)$$