

# Investigating Microbiome’s Role in Neuropsychiatric Disorders in Quest of Novel Therapeutics Using Computational Methods.

Shakil Ahmed Rafi

July 2, 2024

## Abstract

The gut microbiome has been extensively and fruitfully studied using recently using machine learning tools yielding novel connections between neuropsychiatric disorders and the presence and abundance of various microbiota. However despite this recent successes the oral microbiota has received little attention in the literature let alone the machine learning treatment that the gut microbiome receives. This presents a promising research gap. The author proposes that since an emerging consensus for an oral-microbiome-brain exists, since several robust datasets of the oral microbiome already exist, and since novel attentive transformer based models for regression, classification, and clustering of the microbiome do not exist yet, a growing and rather promising open gap exists in the literature for this to be fulfilled, and proposes this as their post-doctoral research goal.

## 1 Introduction

The role of human host microbiomes in neuropsychiatric diseases has been studied extensively in the literature, e.g. for reviews see [1], [2], and [3]. Furthermore they have been implicated in a number of neuropsychiatric disorders, such as ADHD, in [4], in increasing severity of autism spectrum disorders, ASD, in children in [5], in Alzheimer’s Disease in the elderly, in [6] and [7]

### 1.1 The gut microbiome in contrast to the oral microbiome

Of the different microbiomes in the human body, (e.g. gut, oral, skin, vaginal), the gut microbiome is the most extensively studied in relation to neuropsychiatric disorders, see [8]. It is not only the most studied but also the microbiome where modern machine learning techniques have been most frequently and fruitfully applied.

This leaves a research gap for other microbiomes of the human body. For instance both [1] and [9] both note the dearth of literature with respect to the oral microbiome.

**We propose, therefore, to take some of the tools, especially machine learning tools, used in gut microbiome analysis and apply them for the analysis of the oral microbiome.** We propose this for three reasons:

- i. The oral microbiome is yet under-studied with the tools applied to other microbiomes, as noted earlier [1],[9].
- ii. An emerging argument exists for an oral-microbiome-brain axis, OMBA, similar to the gut-brain axis, [10], [11], [12]. The consensus seems to be that this area still needs to be studied.
- iii. Large, robust, and mature datasets exist for the oral microbiomes, such as the Human Oral Microbiome Dataset, and extended Human Oral Microbiome Dataset[13], Cultivated Oral Bacterial Genome Reference [14], and even some smaller datasets such as the U.A.E. Healthy Future Study participants of 330 Emirati citizens [15], and less specialized datasets such as FinnGen, [16].

## 2 Machine Learning Methods to be Applied

Our main source of inspiration will be methods already applied to the gut microbiome, we will use the framework given in [17], and explore some of the hypotheses in [9].

## 2.1 Regression and Classification via TabNet

Regression in its various forms have a long history of use with predictions from microbiome, e.g. LASSO regression in using the blood microbiome to predict gut  $\alpha$ -diversity in [18].

Indeed logistic and linear regression models have also had some use in predicting autism spectrum disorders, ASD from the oral microbiome, [19].

What the literature seems to lack is more sophisticated neural network regression and classification techniques like TabNet (introduced in [20]). Because genomic data is well-known to be large and yet quite sparse simpler regression methods may not be the best at this task.

TabNet was introduced to tackle just this kind of problem. It uses an attention-based mechanism using a transformer-based [21] architecture where-in the model learns from a sparse but wide tabular data, extracting the salient features of the dataset as it reads the data along. It updates feature importances based on new data that it reads and assigns attention to these features updating the model.

Attention based models have shown incredible promise in other areas of AI research such as Large Language Models (ChatGPT) and in computer vision with vision transformers and indeed continues to do so with tabular data. Indeed benchmarking with the Tabzilla Benchmarking Suite [22] shows that TabNet may out-perform traditional gradient-boosted decision trees in contexts where the dataset is extremely large with high dimensionality and where there is large sparseness in the tabular structure.

These features make advanced attention based neural networks particularly appealing for genomic analysis. Combined with the fact that the literature is already thin in the case of the oral microbiome this presents a fertile area of research.

As an added bonus TabNet has a robust implementation in PyTorch [23] not only as a regressor but also as a classifier, making it easy to laterally transfer it from regression to classification.

We must note however the valid concern that neural network methods are thought to be less “explainable” compared to standard statistical methods. This is largely mitigated with TabNet, for instance [24] notes TabNet to be interpretable and provides “better or comparable” results to XGBoost & GLM with predicting insurance claims. On top of that modern tools like SHAPS [25] and LIME [26], largely bridge the interpretability gap.

The author therefore proposes using TabNet, and the related TabPFN [27] for smaller datasets as a

novel way of exploring the impact of the oral microbiome in neuropsychiatric disorders.

## 2.2 Clustering via Variational Autoencoders

In a similar vein to the previous section on regression and classification we note that microbiome research, especially gut microbiome research has had a long and fruitful use of clustering algorithms for research, e.g. [28] looks at four such clustering techniques, k-Means, hierarchical, partition around medoids, and Dirichlet multinomial model.

Because genomic data often has very high dimensions there has been some promising applications for variational autoencoders, VAEs, to reduce the dimensionality down and then feed the data to clustering algorithms. This technique has seen use especially in the detection of cancer e.g. [29] and [30], and in PubMed only one example that the author can find for metagenome binning, [31].

VAEs also have the added benefit of reducing noise, making the data smaller by discarding redundant features similar to principal component analysis (PCA) and related principal co-ordinate analysis (PoCA), making other, more traditional, analyses easier on the dataset.

Regardless, a research gap still exists in genomic analysis of the oral microbiome using variational autoencoders. This gap is even more salient considering that software tools like DeepMicro [32], a suite of tools for the use of variational autoencoders for genomic analysis, already exist, dramatically reducing the software overhead for such an analysis.

Finally, it must be noted that the exact mechanism by which the oral microbiome affects the brain is still up for debate. Several possible mechanisms have been suggested in [9]. What is clear is that this mechanism could possibly be more “direct” in the sense that for the gut microbiome, any toxins released into the bloodstream must first be filtered by the liver before it reaches the brain whereas this is not the case for the oral microbiome [9].

This, thus means that

## 3 Objective of Research

asdfsdf

## 4 Tentative Timeline and Expected Contributions and Impact

The author proposed the following timeline for their post-doc assuming a two year period. Again, this is subject to change but mostly in the direction of needing less time.

- i. Months 1-6: Explore the proposed datasets and run small scale pre-processing tasks including feature selection. This may include dimensionality reduction,
- ii. Months 7-12: Construct large scale models from the datasets, taking inspiration from existing literature. Steps include parameter tuning, benchmarking, and cross-validation.
- iii. Months 13-18:
- iv. Months 19-24:

## References

- [1] Goswami, A. *et al.* Role of Microbes in the Pathogenesis of Neuropsychiatric Disorders. *Front Neuroendocrinol* **62**, 100917 (2021). URL <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8364482/>.
- [2] Hashimoto, K. Emerging role of the host microbiome in neuropsychiatric disorders: overview and future directions. *Mol Psychiatry* **28**, 3625–3637 (2023).
- [3] Bonnechre, B., Amin, N. & van Duijn, C. The role of gut microbiota in neuropsychiatric diseases creation of an atlas-based on quantified evidence. *Frontiers in Cellular and Infection Microbiology* **12** (2022). URL <https://www.frontiersin.org/journals/cellular-and-infection-microbiology/articles/10.3389/fcimb.2022.831666>.
- [4] Bull-Larsen, S. & Mohajeri, M. H. The Potential Influence of the Bacterial Microbiome on the Development and Progression of ADHD. *Nutrients* **11** (2019). URL <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6893446/>. Publisher: Multidisciplinary Digital Publishing Institute (MDPI).
- [5] Tomova, A. *et al.* Gastrointestinal microbiota in children with autism in slovakia. *Physiology & Behavior* **138**, 179–187 (2015). URL <https://www.sciencedirect.com/science/article/pii/S0031938414005101>.
- [6] Yk, K. & C, S. The Microbiota-Gut-Brain Axis in Neuropsychiatric Disorders: Pathophysiological Mechanisms and Novel Treatments. *Current neuropharmacology* **16** (2018). URL <https://pubmed.ncbi.nlm.nih.gov/28925886/>. Publisher: Curr Neuropharmacol.
- [7] Escobar, Y.-N. H., O’Piela, D., Wold, L. E. & Mackos, A. R. Influence of the Microbiota-Gut-Brain Axis on Cognition in Alzheimer’s Disease. *J Alzheimers Dis* **87**, 17–31 (2022).
- [8] Sorboni, S. G., Moghaddam, H. S., Jafarzadeh-Esfehani, R. & Soleimanpour, S. A Comprehensive Review on the Role of the Gut Microbiome in Human Neurological Disorders. *Clin Microbiol Rev* **35**, e0033820 (2022).
- [9] Tao, K., Yuan, Y., Xie, Q. & Dong, Z. Relationship between human oral microbiome dysbiosis and neuropsychiatric diseases: An updated overview. *Behav Brain Res* **471**, 115111 (2024).
- [10] Bowland, G. B. & Weyrich, L. S. The Oral-Microbiome-Brain Axis and Neuropsychiatric Disorders: An Anthropological Perspective. *Front Psychiatry* **13**, 810008 (2022).
- [11] Xi, Y., Yu, M., Li, X., Zeng, X. & Li, J. The coming future: The role of the oral-microbiotabrain axis in aroma release and perception. *Comprehensive Reviews in Food Science and Food Safety* **23**, e13303 (2024). URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/1541-4337.13303>. eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/1541-4337.13303>.
- [12] Y, M. *et al.* Did the Brain and Oral Microbiota Talk to Each Other? A Review of the Literature. *Journal of clinical medicine* **9** (2020). URL <https://pubmed.ncbi.nlm.nih.gov/33260581/>. Publisher: J Clin Med.
- [13] Chen, T. *et al.* The Human Oral Microbiome Database: a web accessible resource for investigating oral microbe taxonomic and genomic information. *Database* **2010**, baq013 (2010). URL <https://doi.org/10.1093/database/baq013>. <https://academic.oup.com/database/article-pdf/doi/10.1093/database/baq013/1132285/baq013.pdf>.




Figure 1: Proposed mechanisms by which the oral microbiome may affect in neuropsychiatric disorders. Figure copied from [9], pg. 3.

- [14] Li, W. *et al.* A catalog of bacterial reference genomes from cultivated human oral bacteria. *npj Biofilms Microbiomes* **9**, 1–13 (2023). URL <https://www.nature.com/articles/s41522-023-00414-3>. Publisher: Nature Publishing Group.
- [15] Human Oral Microbiota Composition Data from UAE Healthy Future Study (UAEHFS) Pilot Participants. URL <https://datacatalog.med.nyu.edu/dataset/10404>.
- [16] FinnGen: an expedition into genomics and medicine | FinnGen. URL <https://www.finnngen.fi/en>.
- [17] Li, P., Luo, H., Ji, B. & Nielsen, J. Machine learning for data integration in human gut microbiome. *Microb Cell Fact* **21**, 241 (2022).
- [18] Wilmanski, T. *et al.* Blood metabolome predicts gut microbiome -diversity in humans. *Nat Biotechnol* **37**, 1217–1228 (2019). URL <https://www.nature.com/articles/s41587-019-0233-9>. Publisher: Nature Publishing Group.
- [19] Li, C. *et al.* A genetic association study reveals the relationship between the oral microbiome and anxiety and depression symptoms. *Front Psychiatry* **13**, 960756 (2022). URL <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9685528/>.
- [20] Arik, S. . & Pfister, T. TabNet: Attentive Interpretable Tabular Learning. *AAAI* **35**, 6679–6687 (2021). URL <https://ojs.aaai.org/index.php/AAAI/article/view/16826>.
- [21] Vaswani, A. *et al.* Attention is All you Need. In *Advances in Neural Information Processing Systems*, vol. 30 (Curran Associates, Inc., 2017). URL [https://proceedings.neurips.cc/paper\\_files/paper/2017/hash/3f5ee243547dee91fbd053c1c4a845aa-Abstract.html](https://proceedings.neurips.cc/paper_files/paper/2017/hash/3f5ee243547dee91fbd053c1c4a845aa-Abstract.html).
- [22] McElfresh, D. *et al.* When do neural nets outperform boosted trees on tabular data? (2023). 2305.02997.
- [23] pytorch-tabnet: PyTorch implementation of TabNet. URL <https://github.com/dreamquark-ai/tabnet>.
- [24] McDonnell, K., Murphy, F., Sheehan, B., Masello, L. & Castignani, G. Deep learning

- in insurance: Accuracy and model interpretability using tabnet. *Expert Systems with Applications* **217**, 119543 (2023). URL <https://www.sciencedirect.com/science/article/pii/S0957417423000441>.
- [25] Lundberg, S. M. & Lee, S.-I. A Unified Approach to Interpreting Model Predictions. In *Advances in Neural Information Processing Systems*, vol. 30 (Curran Associates, Inc., 2017). URL [https://proceedings.neurips.cc/paper\\_files/paper/2017/hash/8a20a8621978632d76c43dfd28b67767-Abstract.html](https://proceedings.neurips.cc/paper_files/paper/2017/hash/8a20a8621978632d76c43dfd28b67767-Abstract.html).
- [26] Ribeiro, M. T., Singh, S. & Guestrin, C. "why should i trust you?": Explaining the predictions of any classifier. In *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining, KDD '16*, 1135–1144 (Association for Computing Machinery, New York, NY, USA, 2016). URL <https://doi.org/10.1145/2939672.2939778>.
- [27] Hollmann, N., Mller, S., Eggenberger, K. & Hutter, F. TabPFN: A transformer that solves small tabular classification problems in a second (2022). 2207.01848.
- [28] Shi, Y., Zhang, L., Peterson, C. B., Do, K.-A. & Jenq, R. R. Performance determinants of unsupervised clustering methods for microbiome data. *Microbiome* **10**, 25 (2022). URL <https://doi.org/10.1186/s40168-021-01199-3>.
- [29] Hira, M. T. *et al.* Integrated multi-omics analysis of ovarian cancer using variational autoencoders. *Sci Rep* **11**, 6265 (2021). URL <https://www.nature.com/articles/s41598-021-85285-4>. Publisher: Nature Publishing Group.
- [30] Zhang, X. *et al.* Integrated multi-omics analysis using variational autoencoders: Application to pan-cancer classification. In *2019 IEEE International Conference on Bioinformatics and Biomedicine (BIBM)*, 765–769 (2019).
- [31] Nissen, J. N. *et al.* Improved metagenome binning and assembly using deep variational autoencoders. *Nat Biotechnol* **39**, 555–560 (2021).
- [32] Oh, M. & Zhang, L. DeepMicro: deep representation learning for disease prediction based on microbiome data. *Sci Rep* **10**, 6026 (2020). URL <https://www.nature.com/articles/s41598-020-63159-5>. Publisher: Nature Publishing Group.