

Annals of DIAGNOSTIC PATHOLOGY

Annals of Diagnostic Pathology 13 (2009) 140-145

Review Article

Molecular biology: an early detector of oral cancers Siddiq M. Ahmed, MD, DNB^a, Mubeen, MDS^b, V.R. Jigna^{b,*}

^aDepartment of General Pathology, Bangalore Medical College and Research Institute, Bangalore, India ^bDepartment of Oral Medicine and Radiology, Government Dental College and Research Institute, Bangalore 560002, India

Abstract

Oral cancers have been one of the leading causes of deaths particularly in the developing countries. Prime reason for this high mortality and morbidity is attributed to the delay in diagnosis and prompt treatment. Relentless research in the field of oncology has led to advent of novel procedures for the early detection of oral cancers. Molecular biology is highly promising in this regard. It is a procedure that detects alterations at a molecular level much before they are seen under a microscope and much before clinical changes occur. Molecular studies serve as basis by which we will eventually be able not only to augment clinical assessment and classification of oral lesions but also predict malignant potential of oral lesions, thus reducing incidence and increasing the scope for early diagnosis and treatment of oral cancers. However, making such sophisticated tools available for the common man in developing countries is one of the most important challenges faced today.

© 2009 Elsevier Inc. All rights reserved.

© 2009 Elsevier Inc. All rights reserved

Keywords:

Oral cancer; Early detection; Molecular biology; Developing countries

1. Introduction

Oral cancer is the sixth most common cancer of both sexes in the general population and the third most common cancer in developing nations. About half of patients afflicted die within 5 years of diagnosis, whereas surviving patients may be left with severe esthetic and functional compromises [1-3]. India has one of the highest incidence of oral cancer in the world [4]. Oral cancer ranks number 1 among men and number 3 among women in India [5].

Such high impact diseases are challenging to diagnose with clinical evaluation and existing laboratory testing. Even with the advent of newer technologies and sophisticated laboratory tools, definitive diagnosis has always been elusive. More than two thirds of oral cancers documented are diagnosed only at an advanced stage. To address this fatal problem, an accurate, qualitative, convenient, and noninvasive diagnostic tool needs to be devised. Molecular biological analysis is one such promising platform that can be used for early detection of oral cancers, their manage-

E-mail address: jigna_ad@yahoo.com (V.R. Jigna).

ment, and thus improve prognosis for the patients. When considering these tools for diagnosis, cost is also a very important factor to be considered, especially in developing nations like India where estimated per capita income is \$275.

Molecular biology is defined as the study of biology at a molecular level. It chiefly concerns itself with understanding the interactions between the various systems of a cell including interactions between DNA, RNA, and protein biosynthesis and learning how these interactions are regulated. Extensive research has been carried out in the field of oncology, and molecular tools have assumed great importance. Because our area of concern is the oral and perioral structures, it is of importance to highlight the molecular analytical tools that have been used and that could be used in future in diagnosis and treatment of oral cancers.

Molecular tools most commonly used are the following:

- Chromosome in situ hybridization
- Cytomorphometry
- Immunohistochemistry
- Polymerase chain reaction
- DNA image cytometry
- DNA miroarrays
- Proteomics
- Gene therapy.

^{*} Corresponding author. Tel.: +91 9731333083 (Mobile), +91 080 26932803 (Residence).

2. Chromosome in situ hybridization

Chromosome in situ hybridization (CISH) is a cytogenic technique that is used to detect and localize the presence or absence of specific DNA sequence or chromosome. It uses fluorescent probes that bind to only those parts of the chromosomes with which they show a high degree of sequence similarity. Fluorescence microscopy is used to find out where the fluorescent probe is bound to the chromosome, thus providing an ability to directly visualize the genetic change in tissue sections or exfoliated cells.

In oral cancers, this technique is used to form the diagnosis and evaluate prognosis and remission of the disease. Analyses of normal and premalignant lesions adjacent to tumors have demonstrated that chromosome instability can be detected in the field of the tumor (ie, in normal and premalignant cells in a tissue at 100% risk of tumor development), and the degree of chromosome instability increases with the degree of histologic progression toward cancer. Analyses of premalignant lesions, for example, oral leukoplakia and erythroplakia from individuals at risk for cancers by CISH have uncovered varying degrees of chromosome instability [6]. Studies indicate that most leukoplakia lesions contain an abnormal number of chromosomes 7 and 17, and lesions with greater than 3% proportion of cells with trisomy 9 have a significant higher likelihood of progression to cancer [7,8]. Studies have resulted in detection of new molecular markers like HER-2/neu at the protein level in salivary duct carcinomas (SDCs). Tissue sections from 12 previously diagnosed SDCs were evaluated by immunohistochemistry (IHC) and CISH for HER-2/neu status. A total of 4 SDCs were positive by IHC; all 4 cases showed amplification with CISH. The remaining 8 cases were negative by IHC and showed no gene amplification with CISH. Salivary duct carcinomas in this study show HER-2/neu overexpression on both the protein and gene levels in approximately 30% of cases [9]. Use of CISH technique can prove to be useful as a molecular diagnostic tool for oral cancers, and its relatively lower cost when compared to techniques like microarrays makes it more affordable for the low socioeconomic groups of the society.

3. Cytomorphometry

It is a quantitative technique that evaluates parameters such as nuclear area (NA), cytoplasmic area (CA), and nuclear/cytoplasmic ratio (NA/CA). Papanicolaou smears are prepared, and the NA, CA, and NA/CA are recorded.

In oral cancers, malignant changes are detected through estimation of NA/CA. They typically show a reduction in CA before the reduction in NA. Cytomorphometric techniques are used to assess nuclear diameter and cytoplasmic diameter in dysplastic lesions and oral squamous cell carcinomas. Cytoplasmic diameter was highest in normal mucosa, lower in dysplastic lesions, and lowest in squamous cell carcinomas. By contrast, nuclear diameter was lowest in

normal mucosa, higher in dysplastic lesions, and highest in squamous cell carcinoma. Thus, it suggested that reduced nuclear size and increased cytoplasmic size are useful indications of malignant transformation [10-12]. Cytomorphometric techniques have also been used to assess the effects of smoking crack cocaine on the oral squamous epithelial cells. The study revealed increase in the NA and decrease in the CA and consequent increase in NA/CA, thus concluding that crack cocaine was able to induce significant changes on the oral epithelial cells. Because this illicit drug is normally used in association with other risk factors for oral cancer (tobacco and alcohol), crack cocaine abusers should have frequent preventive oral exams [13].

4. Immunohistochemistry

Immunohistochemistry is a popular technique that has been used as a part of histopathological examination. It can also be used in assessing molecular alterations in premalignant and malignant oral lesions.

p53 mutations are one of the most common events in tumorogenesis of oral cancers [7]. The gene is located at 17p 13.1. It functions as a 'guardian of cell' by providing the molecular brake and maintaining the genomic stability. When DNA damage occurs, cell produces p53, which stops cell division by arresting cells at G,-S boundary, induces DNA repair and triggers programed cell death or apoptosis. Tumor suppression of p53 is negated by point mutation, deletion of alleles, and binding to viral protein (E6 or E7 of HPV 16/18). Mutant p53 demonstrates a longer half-life than wild type, and its mutant form is often detectable by molecular biology techniques [14]. p53 protein expression has been detected by IHC in 90% of oral leukoplakia, whereas it is absent in normal mucosa [15]. Several studies have demonstrated that p53 detection by IHC alone or with other markers appears to be associated with greater risk for malignant progression [16-22]. Parabasal detection of expression of p53 by IHC has been shown to have a stronger correlation with progression to cancer in several studies [15,23-25], lending credence to the potential application of p53 detection by IHC for stratification of risk of malignant transformation in oral leukoplakia [7,26-29].

The immunohistochemical investigations have also been employed in detection of human papillomavirus (HPV) infection in oral precancerous lesions and oral cancers. The study suggest that HPV type 16/18 has a close association with the development of oral squamous cell carcinoma and that the infection is maintained even after neoplastic change of the infected cells [30].

Immunohistochemistry is widely accessible and easy to perform at a reasonable cost. However, this semiquantitative procedure is beset by technical artifacts, sensitivity differences between different antibodies, and subjective interpretation, resulting in interobserver variability between pathologists.

5. Polymerase chain reaction

Polymerase chain reaction (PCR) is an extremely versatile technique used for copying of DNA. It allows a single DNA sequence to be copied millions of times or altered in a predetermined way. PCR technique is used for quantitative measurements of DNA or RNA molecules. It has also been used to introduce restriction enzymes sites or to mutate particular bases of DNA.

Polymerase chain reaction technique has been used to amplify DNA in samples from oral carcinomas and has been analyzed with restriction fragment length polymorphism. It detects the commonly implicated molecular alterations in oral cancers such as loss of heterozygosity, microsatellite instability, and changes in methylation pattern [12].

The causal link between HPV and a subset of head and neck squamous cell carcinoma (HNSCC) cases has also been established in recent years. Human papillomavirus DNA has been identified in approximately 15% to 26% of HNSCC cases. The most commonly detected HPV in HNSCC is HPV-16, which has been demonstrated in 90% to 95% of all HPV positive HNSCC cases, followed by HPV-18 and HPV-33. To identify this, PCR technique has been used [31-35].

Another interesting trend currently being investigated is the use of the PCR to determine if surgical margins obtained at the time of surgery that are histopathologically free of tumor contain a small amount of histologically undetectable tumor cells. Specifically, the use of PCR to detect specific mutations identified in the primary tumor in the histopathologically negative surgical margin could be very useful because these mutations would indicate the presence of residual (histopathologically undetected) tumor cells [36]. It will be very important to establish whether the presence of submicroscopic tumor cells contributes to prognosis and clinical outcome.

Use of versatile techniques like PCR must be made readily available and more affordable to the general population by getting third parties to cover them. These techniques enable the clinicians to identify the pathology at the earliest, thus contributing to the future therapeutic improvements.

6. DNA Image cytometry

DNA image cytometry is a technique that permits quantification of nuclear DNA content, thereby assessing the DNA ploidy status.

Molecular and genetic changes arise during carcinogenesis and could lend themselves to be markers of transformation. Mutations in p53, loss of heterozygosity, and chromosomal polysomy are all associated with progression to carcinoma and may be predictive when used and analyzed in combination [37-40]. Routine use of these techniques is, however, hampered by the complexity of the tests, the lack of facilities in many routine laboratories, and the high cost involved [41,42]. As a surrogate for such individual

molecular markers, measurement of gross genomic damage, in the form of aberrant DNA content, could be a valuable method for prognostication of malignant and premalignant lesions [43-49]. Relativity recently, there has been a major advancement in this area with the use of automated image cytometry to measure ploidy in nuclei extracted from routinely processed paraffin sections. This system may be more sensitive in the evaluation of oral potentially malignant lesions into 'low' and 'high' risk [50,51].

Evaluation of ploidy status in oral leukoplakia and oral lichen planus allows the identification of gross genomic alterations and has been shown to be a useful tool in identifying lesions with high risk of malignant transformation to oral cancers. Studies show that anueploid lesions have the maximum incidence of malignant transformation [34,52,53]. DNA image cytometry has also been used to detect cancer cells with abnormal DNA content at the invasive tumor front, which is associated with poor prognosis of the patients with oral carcinomas [54,55]. Thus, it could help to find the appropriate treatment option for the patients. The sensitivity of cytological diagnosis combined with DNA cytometry is 98% and specificity 100% when compared with gold standards of histology [12].

7. DNA microarrays

DNA microarray is a high throughput technology used in molecular biology and helps in the study of sequence of genes. It consists of an arrayed series of thousands of microscopic spots of DNA oligonucleotides, each containing a specific DNA sequence. This is a short section of gene or other DNA element that are used as probes to hybridize DNA or RNA samples. Because an array can contain tens of thousands of probes, a microarray experiment can accomplish many genetic tests in parallel [56,57].

Gene profiling method has been used for comparing the normal and cancerous oral mucosa, and a distinct pattern of gene expression was observed when normal and cancerous cells were compared in genome-wide analysis of oral cancer [58]. Gene profiling by DNA microarrays is capable of identifying up-regulated or down-regulated genes correlated with oral tumor recurrences and lymph node metastasis, thus helping clinician to plan a treatment that prevents recurrence and reduce chances of metastasis [34].

Microarray platforms have to be developed to accomplish the procedures. A number of issues must be addressed before establishing a microarray platform and beginning expression profiling studies, in particular, the overall cost. For a cDNA microarray platform, one must purchase a clone set, robot, printing pins, and reagents needed for DNA amplification and purification. The cost of these materials can vary significantly, but one can expect to need at least \$100 000 to establish such a platform. However, once the process of printing and hybridizing microarrays has been optimized, the cost per experiment will fall dramatically. Thus, one must

decide if the number of planned experiments is enough to warrant the time and cost of establishing a microarray platform. Another challenge is the efficient management and analysis of the large volume of data generated by microarray approaches. However, increasingly sophisticated computational methods continue to be developed that are amenable to large data sets generated from microarray experiments. Certainly, as key disease pathways are identified, custom arrays containing relevant subsets of genes may eventually be integrated into clinical settings for more widespread use [59].

8. Proteomics

The term "proteomics" indicates PROTEins expressed by a genOME and is the systematic analysis of protein profiles of tissues, paralleling the related field of genomics [57]. The major workhorse of proteomics-based expression profiling is still the combination of high resolution 2-Dgel electrophoresis and mass spectrometry. Laser Capture Microdissection, protein biochips, and isotope-coded affinity tag peptide labeling are among the technologies that are currently having an important impact on cancer research [60]. A new protein analysis system, based on the surface enhanced laser desorption/ionization has been recently applied for the separation, detection, and analysis of multiple proteins in very small amounts (10 ng) of microdissected cancer tissue. This system facilitates protein capture, purification, analysis, and processing from complex biological mixtures directly onto protein chip array surfaces, and the detection of the purified proteins is performed by time-of-flight mass spectrometry [56]. For a better understanding of how patterns of protein expression shape the tissue microenvironment, protein expression in tissue derived by Laser Capture Microdissection from squamous cell carcinoma of the oral cavity was analyzed using a high-throughput antibody microarray approach [61]. A reproducible correlation was found between the expression patterns of multiple proteins within epithelial cells and the progression of oral cavity tumor. A comparison of the protein maps of normal and malignant prostate were used to identify 20 proteins lost in malignant transformation, including PSA, a-1 antichymotrypsin, haptoglobin, and lactylglutathione lyase [62].

Hence, proteomic technologies have the potential to greatly aid the development of molecular diagnostics and serve as markers for the early detection of cancer. These technologies will also accelerate the anticancer drug target discovery and validation. Furthermore, proteomic technologies will be used to design rational drugs according to the molecular profile of the cancer cell and thus facilitate the development of personalized cancer therapy [57].

9. Gene therapy

Gene therapy is a promising new approach for the management of oral cancers. It is the insertion of gene into an

individual's cells and tissues to treat a disease in which a definitive mutant allele is replaced with a functional one. Gene therapy uses an adenovirus vector, which is used to introduce modified DNA into a human cell. If the treatment is successful, the new gene will make a functional protein.

Gene therapy approaches to oral cancers and precancers include the following:

- Addition gene therapy: Aim of approach is to regulate the tumor growth by introducing tumor suppressor gene that inactivates the carcinogenic cells.
- Gene therapy using oncolytic viruses: This approach uses viruses that replicate only in the tumor cells and thus kills them.
- Suicide gene therapy: In this therapy, enzyme encoding gene is introduced into the tumor cell that stimulate the generation of products that are toxic for the cells.
- Immunotherapy: The aim of immunotherapy is to increase the patient's immune response to the tumor.
- Introduction of genes to inhibit tumor angiogenesis: This technique uses microencapsulated cells for the release of therapeutic proteins to encapsulate recombinant cells [63-70].

Use of molecular biology for management of oral cancers is still at its infancy. In the future, it may be the treatment of choice that will overcome all the complications associated with the present trend of management of oral cancers, thus proving to be a blessing to the patients.

10. Conclusion

Molecular analysis is a fast emerging and fascinating technique that can replace the present available diagnostic procedures in the field of oral oncology for early detection of oral cancers, thus saving the patients of physical and mental trauma they go through when oral cancers are detected at an advanced stage. However, practical problems such as affordability of these tests have to be considered when such tools are used for common man living in developing and underdeveloped countries. Governments of such countries in association with organizations like the World Health Organization and the United Nations Educational, Scientific, and Cultural Organization should conduct nationalized health program schemes for regular screening of high-risk patients with molecular analytical tools at subsidized rates so that oral cancers can be detected at the earliest and hence reduce morbidity and mortality in countries like India, where the incidence of oral cancers is high.

References

- [1] Siverman SI. Early diagnosis of oral cancer. Cancer 1988;62:1796-9.
- [2] Vokes EE, Weichselbaum RR, Lippman SM, et al. Head and neck cancer (review). New Eng J Med 1993;328:184-94.
- [3] Sidransksy D. Molecular genetics of head and neck cancer. Curr Opin Oncol 1995;7:229-33.

- [4] Hamada GS, Bos AJ, Kasuga H, et al. Comparative epidemiology of oral cancer in Brazil & India. Tokai J Exp Clin Med 1991:16(1):63-72.
- [5] Sankaranarayanan R. Oral caner in India: an epidemiologic and clinical review. Oral Surg Oral Med Oral Path 1990;69:325-30.
- [6] Walter NH, Hyo JK, Jin SL, et al. Detection of chromosome instability of tissue fields at risk: in situ hybridization. J Cell Biochem 1997;25S: 57-62.
- [7] Mithani SK, Mydlarz WK, Grumbine FL, et al. Molecular genetics of premalignant oral lesions. Oral Diseases 2007;13:126-33.
- [8] Kim J, Yook JI, Lee EH, et al. Evaluation of premalignant potential in oral lichen planus using interphase cytogenetics. J Oral Pathol Med 2001;30(2):65-70.
- [9] Johnson CJ, Barry MB, Vasef MA, et al. HER-2/neu expression in salivary duct carcinoma: an immunohistochemical and chromogenic in situ hybridization study. Appl Immunohistochem Mol Morphol 2008; 16(1):54-8.
- [10] Ogden GR, Cowpe JG, Chisholm DM, et al. Detection of oral cancer: role of oral exfoliative cytology. Presented at the International Symposium on Cofactor Interactions and Cancer Prevention; Nice, France; March 17-19; in the section on Early Cancer Detection; 1993.
- [11] Mollaoulu N, Cowpe JG, Walker R. Cytomorphologic analysis of Papanicolaou stained smears collected from floor of the mouth mucosa in patients with or without oral malignancy. Turk J Med Sci 2001;31: 225-8.
- [12] Mehrotra R, Gupta A, Singh M, et al. Application of cytology and molecular biology in diagnosing premalignant or malignant oral lesions. Mol Cancer 2006;5:11.
- [13] Woyceichoski IE, de Arruda EP, Resende LG, et al. Cytomorphometric analysis of crack cocaine effects on the oral mucosa. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2008;105(6):745-9.
- [14] Reibel J. Prognosis of oral pre-malignant lesions: significance of clinical, histopathological, and molecular biological characteristics. Crit Rev Oral Biol Med 2003;14:47-62.
- [15] Lippman SM, Shin DM, Lee JJ, et al. p53 and retinoid chemoprevention of oral carcinogenesis. Cancer Res 1995;55:16-9.
- [16] Nishioka H, Hiasa Y, Hayashi I, et al. Immunohistochemical detection of p53 oncoprotein in human oral squamous cell carcinomas and leukoplakias: comparison with proliferating cell nuclear antigen staining and correlation with clinicopathological findings. Oncology 1993;50:426-9.
- [17] Kaur J, Srivastava A, Ralhan R. Overexpression of p53 protein in betel- and tobacco-related human oral dysplasia and squamous-cell carcinoma in India. Int J Cancer 1994;58:340-5.
- [18] Girod SC, Pape HD, Krueger GR. p53 and PCNA expression in carcinogenesis of the oropharyngeal mucosa. Eur J Cancer B Oral Oncol 1994;30B:419-23.
- [19] Huang XF, Zhang WG, Song LJ, et al. Immunohistochemical detection of PCNA and p53 protein in the premalignant lesions and squamous cell carcinomas of the oral mucosa. Shanghai Kou Qiang Yi Xue 1997; 6:73-4.
- [20] Ries JC, Schreiner D, Steininger H, et al. p53 mutation and detection of p53 protein expression in oral leukoplakia and oral squamous cell carcinoma. Anticancer Res 1998;18:2031-6.
- [21] Scheifele C, Schlechte H, Bethke G, et al. Detection of TP53-mutations in brush biopsies from oral leukoplakias. Mund Kiefer Gesichtschir 2002;6:410-4.
- [22] Kovesi G, Szende B. Changes in apoptosis and mitotic index, p53 and Ki67 expression in various types of oral leukoplakia. Oncology 2003; 65:331-6
- [23] Cruz IB, Snijders PJ, Meijer CJ, et al. p53 expression above the basal cell layer in oral mucosa is an early event of malignant transformation and has predictive value for developing oral squamous cell carcinoma. J Pathol 1998;184:360-8.
- [24] Cruz I, Napier SS, van der Waal I, et al. Suprabasal p53 immunoexpression is strongly associated with high grade dysplasia and risk for malignant transformation in potentially malignant oral lesions from Northern Ireland. J Clin Pathol 2002;55:98-104.

- [25] Cruz I, Snijders PJ, Van Houten V, et al. Specific p53 immunostaining patterns are associated with smoking habits in patients with oral squamous cell carcinomas. J Clin Pathol 2002;55:834-40.
- [26] Wong DTW, Todd R, Tsuji T, et al. Molecular biology of human oral cancers. Crit Rev Oral Biol Med 1996;7(4):319-28.
- [27] Ibrahim SO, Johannessen AC, Idris AM, et al. immunohistochemical detection of p53 in non-malignant and malignant oral lesions associated with snuff dipping in the Sudan and Sweden. Int J Cancer 1996;68:749-53.
- [28] Williams HK. Molecular pathogenesis of oral squamous carcinoma. Mol Pathol 2000;53:165-72.
- [29] Fregonesi PAG, Teresa DB, Duarte RA, et al. p16INK4A immunohistochemical overexpression in premalignant and malignant oral lesions infected with human papillomavirus. J Histochem Cytochem 2003;51(10):1291-7.
- [30] Miki I, Kazunari S, Masahiro U. Immunohistochemical investigation of the human papilloma virus (HPV) infection type 16/18 in oral precancerous lesions and oral cancer. Jpn J Cancer Clin 2002;48(4): 229-35
- [31] Herrero R, Castellsagué X, Pawlita M, et al. Human papillomavirus and oral cancer: the International Agency for Research on Cancer multicenter study. J Natl Cancer Inst 2003;95(23):1772-83.
- [32] Kreimer AR, Clifford GM, Boyle P, et al. Human papillomavirus types in head and neck squamous cell carcinomas worldwide: a systematic review. Cancer Epidemiol Biomark Prev 2005;14(2):467-75.
- [33] Syrjanen S. Human papillomavirus (HPV) in head and neck cancer. J Clin Virol 2005;32S:S59-66.
- [34] Perez-Ordonez B, Beauchemin M, Jordan RCK. Molecular biology of squamous cell carcinoma of the head and neck. J Clin Pathol 2006;59: 445-53.
- [35] Ribeiro da Silva CEX, Guerreiro da Silva IDC, Cerri A. Prevalence of human papillomavirus in squamous cell carcinoma of the tongue. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2007;104:497-500.
- [36] Breman JA, Mao L, Hruban RH, et al. Molecular assessment of histopathological staging in squamous cell carcinoma of head and neck. N Eng J Med 1995;332:429-35.
- [37] Garcia-Pola Vallejo MJ, Anitua Roldan MJ, Fernández Álvarez BE, et al. Study comparative of Ki-67 expression in oral lichen planus and oral leukoplakia. Quant Anal Med Oral 2001;6:364-70.
- [38] Saiz-Rodriguez A. Molecular basis of oral cancer. Med Oral 2001;6: 342-9.
- [39] Burgos JS. Absence of p53 alterations in nasopharyngeal carcinoma Spanish patients with Epstein-Barr virus infection. Virus Genes 2003; 27:263-8.
- [40] Sheard MA, Uldrijan S, Vojtesek B. Role of p53 in regulating constitutive and X-radiation-inducible CD95 expression and function in carcinoma cells. Cancer Res 2003;63:7176-84.
- [41] Zhang C, Zhang P, Sung CJ, et al. Overexpression of p53 is correlated with stromal invasion in extramammary Paget's disease of the vulva. Hum Pathol 2003;34:880-5.
- [42] Wang J, Taylor CR. Apoptosis and cell cycle-related genes and proteins in classical Hodgkin lymphoma: application of tissue microarray technique. Appl Immunohistochem Mol Morphol 2003; 11:206-13.
- [43] Abdel-Salam M, Mayall BH, Hansen LS, et al. Nuclear DNA analysis of oral hyperplasia and dysplasia using image cytometry. J Oral Pathol Med 1987;16(9):431-5.
- [44] Sarker SK, Ghufoor K, Patel KS, et al. Nuclear DNA content using computerized image cytometry of squamous cell carcinomas of the head and neck. J Laryngol Otol 1997;111:43-7.
- [45] Rihakova P, Brychtova S, Kotrsova L, et al. DNA ploidy correlates with grade, proliferation and clinical outcome but not with presence of human oncogenic HPVs or expression of Bcl-2 in preneoplastic and neoplastic lesions of the uterine cervix. Neoplasma 2001;48:274-7.
- [46] Sudbo J, Reith A. Which putatively pre-malignant oral lesions become oral cancers? Clinical relevance of early targeting of high-risk individuals. J Oral Pathol Med 2003;32:63-70.

- [47] Remmerbach TW, Weidenbach H, Hemprich A, et al. Earliest detection of oral cancer using non-invasive brush biopsy including DNA-imagecytometry; report on four cases. Anal Cell Pathol 2003;25(4):159-66.
- [48] Maraki D, Becker J, Boecking A. Cytologic and DNA-cytometric very early diagnosis of oral cancer. J Oral Pathol Med 2004;33:398-404.
- [49] Zafer Özgür P, Ahmet K, Ömer G, et al. Evaluation of nuclear morphometry and DNA ploidy status for detection of malignant and premalignant oral lesions: quantitative cytologic assessment and review of methods for cytomorphometric measurements. J Oral Maxillofac Surg 2006;64:628-35.
- [50] VanDiest PJ, Smeulders AW, Thunnissen FB, et al. Cytomorphometry. A methodologic study of preparation techniques, selection methods and sample sizes. Anal Quant Cytol Histol 1989;11:225-31.
- [51] Shirata NK, Zerbini MC, Longatto Filho A, et al. DNA ploidy in cervical lesions assessed by computed image analysis: relation to histopathology. Pathologica 2003;95:88-91.
- [52] Femiano F, Scully C. DNA cytometry of oral leukoplakia and oral lichen planus. Med Oral Patol Oral Cir Bucal 2005;10:E9-E14.
- [53] Dimitra M, Sebnem Y, Natalia P. Cytologic and DNA-cytometric examination of oral lesions in lichen planus. J Oral Pathol Med 2006; 35:227-32.
- [54] Noguchi M, Kinjyo H, Kohama G, et al. Invasive front in oral squamous cell carcinoma: image and flow cytometric analysis with clinicopathologic correlation. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2002;93:682-7.
- [55] Handschel J, Oz D, Pomjanski N, et al. Additional use of DNA-image cytometry improves the assessment of resection margins. J Oral Pathol Med 2007;36(8):472-5.
- [56] Patel V, Leethanakul C, Gutkind JS. New approaches to the understanding of the molecular basis of oral cancer. Crit Rev Oral Biol Med 2001;12(1):55-63.
- [57] Nagpala JK, Das BR. Oral cancer: reviewing the present understanding of its molecular mechanism and exploring the future directions for its effective management. Oral Oncology 2003;39:213-21.

- [58] Shillitoe EJ, May M, Patel V, et al. Genome-wide analysis of oral cancer—early results from the Cancer Genome Anatomy Project. Eur J Cancer B Oral Oncol 2000;90:340-7.
- [59] Nazmul-Hossain ANM, Patel KJ, Rhodus NL, et al. Microarrays: applications in dental research. Oral Diseases 2008;14:25-9.
- [60] Jain KK. Recent advances in oncoproteomics. Curr Opin Mol Ther 2002;4(3):203-9.
- [61] Hanash SM. Global profiling of gene expression in cancer using genomics and proteomics. Curr Opin Mol Ther 2001;3: 538-45.
- [62] Meehan KL, Holland JW, Dawkins HJ. Proteomic analysis of normal and malignant prostate tissue to identify novel proteins lost in cancer. Prostate 2002;50:54-63.
- [63] O'Malley BW, Cope KA, Chen S, et al. Combination gene therapy for oral cancer in a murine model. Cancer Res 1996;56:1737-41.
- [64] Cusack Jr JC, Tanabe KK. Cancer gene therapy. Surg Oncol Clin N Am 1998:7:421-69.
- [65] Gleich LL. Gene therapy for head and neck cancer. Laryngoscope 2000;110(5 Pt 1):708-26.
- [66] Fukui T, Hayashi Y, Kagami H, et al. Suicide gene therapy for human oral squamous cell carcinoma cell lines with adeno-associated virus vector. Oral Oncol 2001;37:211-5.
- [67] Xi S, Grandis JR. Gene therapy for the treatment of oral squamous cell carcinoma. J Dent Res 2003;82(1):11-6.
- [68] Ladeinde AL, Ogunlewe MO, Adeyemo WL, et al. Gene therapy in the management of oral cancer: a review of recent developments. Niger Postgrad Med J 2005;12(1):18-22.
- [69] Young M, Overlid N, Konopka K, et al. Gene therapy for oral cancer: efficient delivery of a 'suicide gene' to murine oral cancer cells in physiological milieu. Med Oral Patol Oral Cir Bucal 2008;13(1): E15-21.
- [70] Barbellido SA, Trapero JC, Sánchez JC, et al. Gene therapy in the management of oral cancer: review of the literature. Med Oral Patol Oral Cir Bucal 2008;13(1):E15-21.