

Prove of Constant Radius Model

Hanchai Nonprasart and Sakepisit Maysamat

Mahidol Wittayanusorn School

SCI 30196: Science Project 1

Amornsri Amornvatcharapong, Teerapong Suksumran

February 22, 2022

Abstract

This is the first generation of the model. It's a draft. Who would even care to read the abstract when it's not done anyways?

Keywords: None 1, None 2, None 3

DRAFT

Prove of Constant Radius Model

Contents

Rewrite note

This proof will be organized using the following guideline

1. reviewing existing definitions and theorem
2. proving certain property of the existing mathematical objects
3. defining the conditions of the model
4. formulating of the model
5. parameterize the model
6. asserting defined property of the model

It turned out that figures will crash overleaf, if you want to view such file, please compile them separately. To merge the file, it is needed to either pay the subscription or download and run it on your computer.

Tensor

Preliminary

(1)

$$(A + B)^{i_1 \dots i_n}_{j_1 \dots j_m} = A^{i_1 \dots i_n}_{j_1 \dots j_m} + B^{i_1 \dots i_n}_{j_1 \dots j_m}$$

(2)

$$(\alpha A)^{i_1 \dots i_n}_{j_1 \dots j_m} = \alpha A^{i_1 \dots i_n}_{j_1 \dots j_m}$$

(3)

$$(A \otimes B)^{i_1 \dots i_l i_{l+1} \dots i_{l+n}}_{j_1 \dots j_k j_{k+1} \dots j_{k+m}} = A^{i_1 \dots i_l}_{j_1 \dots j_k} B^{i_{l+1} \dots i_{l+n}}_{j_{k+1} \dots j_{k+m}}$$

(4)

$$(\text{contr } T)^{i_1 \dots i_n}_{j_1 \dots j_m} = \sum_a T^{i_1 \dots i_n a}_{a j_1 \dots j_m}$$

(5)

$$(AB)^i_j = \text{contr}(A \otimes B) = \sum_k A^i_k B^k_j$$

(6)

$$\left(\begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix} \right)_{ij} = \sum_k A_{ik} B_{kj}$$

$$(7) \quad IA = A = AI$$

$$(8) \quad I_n = \text{diag}(1, 1, \dots, 1)$$

$$(9) \quad I_{a+b} = \begin{bmatrix} I_a & 0_{a \times b} \\ 0_{b \times a} & I_b \end{bmatrix}$$

$$(10) \quad T_{a,b}T_{a,b} = I$$

Differential

$$(11) \quad J_x F = \left(\frac{\partial F_i}{\partial x_j} \right)_{i,j}$$

$$(12) \quad Df|_x : T_x \mathbb{R}^k \rightarrow T_{f(x)} \mathbb{R}^n = (x, v) \mapsto (f(x), J_x f(v))$$

Group

Definition 1 (Lie group (lee_2013)). A *Lie group* is a smooth manifold G (without boundary) that is also a group in the algebraic sense, with the property that the multiplication map $m : G \times G \rightarrow G$ and the inversion map $i : G \rightarrow G$, given by

$$m(g, h) = gh, \quad i(g) = g^{-1}$$

are both smooth.

Preposition 1 (Lie group (lee_2013)). If G is a smooth manifold with a group structure such that the map $G \times G \rightarrow G$ given by $(g, h) \mapsto gh^{-1}$ is smooth, then G is a Lie group.

Definition 2 (Semidirect product). Suppose H and N are groups, and $\theta : H \times N \rightarrow N$ is a smooth left action of H on N . It is said to be an *action by automorphisms* if for each $h \in H$, the map $\theta_h : N \rightarrow N$ is a group automorphism of N (i.e., an isomorphism from N to itself). Given such action, we define a new group $N \rtimes_\theta H$, called a *semidirect product* of H and N , as follows. $N \rtimes_\theta H$ is just the Cartesian product $N \times H$; but the group multiplication is defined by

$$(n, h)(n', h') = (n\theta_h(n'), hh').$$

Definition 3. An orthogonal group $O(n)$ is a group of orthogonal matrix M where $M^{-1} = M^T$ with matrix multiplication.

Definition 4. An special orthogonal group $SO(n)$ is a subgroup of the orthogonal group $O(n)$ whose element M have the property that $\det M = 1$.

Definition 5. An indefinite orthogonal group $O(m, n)$ is a group of indefinite orthogonal matrix M where $M^{-1} = gM^Tg$ for $g = \text{diag}(-1, \dots, -1, 1, \dots, 1)$ with matrix multiplication.

Definition 6. An orthochronous indefinite orthogonal group $O^+(m, n)$ is a subgroup of the indefinite orthogonal group $O(m, n)$ whose element M have the property that $M^1_1 > 0$.

Definition 7. An special orthochronous indefinite orthogonal group $SO^+(m, n)$ is a subgroup of the orthochronous indefinite orthogonal group $O^+(m, n)$ whose element M have the property that $\det M = 1$.

Definition 8. An translation group $T(n)$ is a group of vectors in \mathbb{R}^n with vector addition.

Definition 9. An euclidean group $E(n)$ is a semidirect product of orthogonal group $O(n)$ extended by translation group $T(n)$.

Definition 10. A *Klein geometry* is a pair (G, H) where G is a Lie group and H is a closed Lie subgroup of G such that the (left) coset space

$$X := G/H$$

is connected.

Example 10.1. KleinGeometryExamples (I)

Manifold

Definition 11 (Abstract differentiable manifold ([kuhnelwolfgang_2006](#))). A *k-dimensional differentiable manifold* (briefly: a *k-manifold*) is a set M together with a family $(M_i)_{i \in I}$ of subsets such that

1. $M = \bigcup_{i \in I} M_i$ (union),
2. for every $i \in I$ there is an injective map $\varphi_i : M_i \rightarrow \mathbb{R}^k$ so that $\varphi_i(M_i)$ is open in \mathbb{R}^k , and
3. for $M_i \cap M_j \neq \emptyset$, $\varphi_i(M_i \cap M_j)$ is open in \mathbb{R}^k .

Definition 12 (Structures on a manifold ([kuhnelwolfgang_2006](#))). Given a k -dimensional differentiable manifold, one gets additional structure by replacing additional requirements on the transformation functions $\varphi_j \circ \varphi_i^{-1}$, which belong to the atlas of the manifold; if all $\varphi_j \circ \varphi_i^{-1}$ are (left-hand side), then one speaks of (right-hand side) as follows:

continuous	\leftrightarrow	topological manifold
differentiable	\leftrightarrow	differentiable manifold
C^1 -differentiable	\leftrightarrow	C^1 -manifold
C^r -differentiable	\leftrightarrow	C^r -manifold
C^∞ -differentiable	\leftrightarrow	C^∞ -manifold
real analytic	\leftrightarrow	real analytic manifold
complex analytic	\leftrightarrow	complex analytic manifold of dimension $\frac{k}{2}$
affine	\leftrightarrow	affine manifold
projective	\leftrightarrow	projective manifold
conformal	\leftrightarrow	manifold with a conformal structure
orientation-preserving	\leftrightarrow	orientable manifold

Definition 13 (Tangent vector ([kuhnelwolfgang_2006](#))). A *tangent vector* X at p is a derivation (derivative operator) defined on the set of *germs of functions*

$$\mathcal{F}_p(M) := \{ f : M \rightarrow \mathbb{R} \mid f \text{ differentiable} \} / \sim,$$

where the equivalence relation \sim is defined by declaring $f \sim f^*$ if and only if f and f^* coincide in a neighborhood of p . The value $X(f)$ is also referred to as the *directional derivative* of f in the direction X .

This definition means more precisely the following. X is a map $X : \mathcal{F}_p(M) \rightarrow \mathbb{R}$ with the two following properties:

1. $X(\alpha f + \beta g) = \alpha X(f) + \beta(g)$, $f, g \in \mathcal{F}_p(M)$ (\mathbb{R} -linearity);
2. $X(f \cdot g) = X(f) \cdot g(p) + f(p) \cdot X(g)$ for $f, g \in \mathcal{F}_p(M)$ (product rule).

(For this to make sense, both f and g have to be defined in a neighborhood of p .)

Briefly: *tangent vectors are derivations acting on scalar functions.*

Corollary 1 (Tangent space ([kuhnelwolfgang_2006](#))). *The tangent space $T_p M$ of M at p is defined in all cases as the set of all tangent vectors at the point p . By definition $T_p M$ and $T_q M$ are disjoint if $p \neq q$.*

Definition 14 (Derivative ([kuhnelwolfgang_2006](#))). Let $F : M \rightarrow N$ be a differentiable map, and let p, q be two fixed points with $F(p) = q$. Then the *derivative* or the *differential* of F at p is defined as the map

$$DF|_p : T_p M \rightarrow T_q N$$

whose value at $X \in T_p M$ is given by $(DF|_p(X))(f) = X(f \circ F)$ for every $f \in \mathcal{F}_q(N)$ (which automatically implies the relation $f \circ F \in \mathcal{F}_p(M)$).

Lemma 1 (Chain rule ([kuhnelwolfgang_2006](#))). *For the derivative as defined in this manner, one has the chain rule in the form*

$$D(G \circ F)|_p = DG|_{F(p)} \circ DF|_p$$

for every composition $M \xrightarrow{F} N \xrightarrow{G} Q$ of maps, or, more briefly, $D(G \circ F) = DG \circ DF$.

Definition 15 (Riemannian metric ([kuhnelwolfgang_2006](#))). A *Riemannian metric* g on M is an association $p \mapsto g_p \in L^2(T_p M; \mathbb{R})$ such that the following conditions are satisfied:

1. $g_p(X, Y) = g_p(Y, X)$ for all X, Y , (symmetry)
2. $g_p(X, X) > 0$ for all $X \neq 0$, (positive definiteness)
3. The coefficient g_{ij} in every local representation (i.e., in every chart)

$$g_p = \sum_{i,j} g_{ij}(p) \cdot dx^i|_p \otimes dx^j|_p$$

are differentiable functions. (differentiability)

Remark 1. The pair (M, g) is then called *Riemannian manifold*. One also refers to the Riemannian metric as the *metric tensor*. In local coordinates the metric tensor is given by the matrix (g_{ij}) of functions. In Ricci calculus this is simply written as g_{ij} .

(kuhnelwolfgang_2006)

Remark 2. A Riemannian metric g defines at every point p an *inner product* g_p on the tangent space $T_p M$, and therefore the notation $\langle X, Y \rangle$ instead of $g_p(X, Y)$ is also used. The notions of angles and lengths are determined by this inner product, just as these notions are determined by the first fundamental form on surface elements. The length or norm of vector X is given by $\|X\| := \sqrt{g(X, X)}$, and the angle β between two tangent vectors X and Y can be defined by the validity of the equation $\cos \beta \cdot \|X\| \cdot \|Y\| = g(X, Y)$. (kuhnelwolfgang_2006)

Curvature

Definition 16 (Christoffel symbols (kuhnelwolfgang_2006)).

1. The quantities $\partial_k \Gamma_{ij}$ defined by the expressions

$$\partial_k \Gamma_{ij} := \left\langle \nabla_{\frac{\partial f}{\partial u^i}} \frac{\partial f}{\partial u^j}, \frac{\partial f}{\partial u^k} \right\rangle$$

are called the *Christoffel symbols of the first kind*.

2. The quantities Γ^k_{ij} defined by the expressions

$$\nabla_{\frac{\partial f}{\partial u^i}} \frac{\partial f}{\partial u^j} = \sum_k \Gamma^k_{ij} \frac{\partial f}{\partial u^k}$$

are called the *Christoffel symbols of the second kind*.

3. By definition one has $\partial_k \Gamma_{ij} = \partial_k \Gamma_{ji}$, $\Gamma^k_{ij} = \Gamma^k_{ji}$ as well as $\partial_k \Gamma_{ij} = \sum_m \Gamma^m_{ij} g_{mk}$.

Definition 17 (Gauss map (kuhnelwolfgang_2006)). For a surface element

$f : U \rightarrow \mathbb{R}^3$, the *Gauss map*

$$\nu : U \rightarrow S^2$$

is defined by the formula

$$\nu(u_1, u_2) := \frac{\frac{\partial f}{\partial u_1} \times \frac{\partial f}{\partial u_2}}{\left\| \frac{\partial f}{\partial u_1} \times \frac{\partial f}{\partial u_2} \right\|}.$$

Definition 18 (Weingarten map, shape operator (kuhnelwolfgang_2006)). Let

$f : U \rightarrow \mathbb{R}^3$ be a surface element with Gauss map $\nu : U \rightarrow S^2 \subset \mathbb{R}^3$.

1. For every $u \in U$ the image plane of the bilinear map

$$D\nu|_u : T_u U \rightarrow T_{\nu(u)} \mathbb{R}^3$$

is parallel to the tangent plane $T_u f$. By canonically identifying $T_{\nu(u)} \mathbb{R}^3 \cong \mathbb{R}^3 \cong T_{f(u)} \mathbb{R}^3$ we may therefore view $D\nu$ at every point as the map

$$D\nu|_u : T_u U \rightarrow T_u f.$$

Moreover, by restricting to the image, we may view the map $Df|_u$ as a linear isomorphism

$$Df|_u : T_u U \rightarrow T_u f.$$

In this sense the inverse mapping $(Df|_u)^{-1}$ is well-defined and is also an isomorphism

2. The map $L := -D\nu \circ (Df)^{-1}$ defined pointwise by

$$L_u := -(D\nu|_u) \circ (Df|_u)^{-1} : T_u f \rightarrow T_u f$$

is called the *Weingarten map* or the *shape operator* of f . Obviously, for every parameter u this is a linear endomorphism of the tangent plane at the corresponding point $f(u)$.

3. L is independent of the parameterization f (up to the choice of the sign of the unit normal vector ν), and it is self-adjoint with respect to the first fundamental form I .

Definition 19 (Hypersurface element ([kuhnelwolfgang_2006](#))). $f : U \rightarrow \mathbb{R}^{n+1}$ is called a *regular hypersurface element*, if $U \subset \mathbb{R}^n$ is open and f is a (C^2-) immersion. The parameter $u = (u_1, \dots, u_n)$ is associated with the point $f(u)$ with $n+1$ coordinates $f(u) = (f_1(u), \dots, f_{n+1}(u))$. The *tangent hyperplane* $T_u f$ is defined to be the image of $T_u U$ under the map $Df|_u$. Similarly, one defines

- the *Gauss map* $\nu : U \rightarrow S^n$ by the unit normal vector $\nu(u)$, which is

perpendicular to $T_u f$ (but note: in \mathbb{R}^{n+1} for $n \geq 3$ there is no bilinear vector product of tangent vectors; still one can formally define ν as an n -linear vector product),

- the *Weingarten map* $L = -D\nu \circ (Df)^{-1}$,

- the first, second, and third fundamental forms

$$I = (g_{ij})_{i,j=1,\dots,n} = \left(\left\langle \frac{\partial f}{\partial u_i}, \frac{\partial f}{\partial u_j} \right\rangle \right),$$

$$\begin{aligned} II &= (h_{ij})_{i,j=1,\dots,n} & = \left(\left\langle \frac{\partial^2 f}{\partial u_i \partial u_j}, \nu \right\rangle \right), \\ III &= (e_{ij})_{i,j=1,\dots,n} & = \left(\left\langle \frac{\partial \nu}{\partial u_i}, \frac{\partial \nu}{\partial u_j} \right\rangle \right). \end{aligned}$$

Definition 20 (Curvature tensor ([kuhnelwolfgang_2006](#))).

$$R(X, Y)Z := \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z$$

is a tensor field, which is called the *curvature tensor* of the surface.

Remark 3 (Curvature tensor ([kuhnelwolfgang_2006](#))). The left-hand side of the Gauss equation is called the *curvature tensor* and is in general expressed in the form

$$R^s{}_{ikj} := \frac{\partial}{\partial u^k} \Gamma^s{}_{ij} - \frac{\partial}{\partial u^j} \Gamma^s{}_{ik} + \sum_r (\Gamma^r{}_{ij} \Gamma^s{}_{rk} - \Gamma^r{}_{ik} \Gamma^s{}_{rj}).$$

Example 20.1.

$$S(v) = \pm \nabla_v n$$

Example 20.2. Eigenvalue of second fundamental form (or shape operator)

Objective

Objective. The objective is that given a natural number n and a real number κ , one can construct

1. a set M
2. a Lie group M with operation \otimes_M ,
3. an n -dimensional C^∞ -manifold M with chart $\varphi_i \subset M \rightarrow \mathbb{R}^n$,
4. a Riemannian manifold M with inner product $g_p \in T_p M \times T_p M \rightarrow \mathbb{R}$

(to be determined) such that

- group action is distance preserved.
- model is continuous with respect to κ (and smooth with respect to each basis).
- for all two-dimensional linear subspaces of the manifold, the sectional curvature is κ .

(cannot figure formal definition out yet.)

Conjecture 1. If the parameters (κ, n) is associated with Klein geometry (G, H) then $(M, \otimes_M) \cong G/H$.

That is, from ??,

- For $\kappa > 0$, $G \cong O(n+1)$ and $H \cong O(n)$.
- For $\kappa = 0$, $G \cong E(n)$ and $H \cong O(n)$.
- For $\kappa < 0$, $G \cong O^+(1, n)$ and $H \cong O(n)$.

Model Foundation

Trigonometry

Definition 21. Generalized trigonometric functions $f_k : \mathbb{R} \rightarrow \mathbb{R}$ and $f_k^* : \mathbb{R} \rightarrow \mathbb{R}$ are defined as

$$f_k(\theta) := \begin{cases} g(k\theta) & \text{if } k \geq 0, \\ h(k\theta) & \text{otherwise,} \end{cases}$$

$$f_k^*(\theta) := \begin{cases} g(k\theta) & \text{if } k \geq 0, \\ h(-k\theta) & \text{otherwise,} \end{cases}$$

where g (resp. h) are the associated trigonometric (resp. hyperbolic) function.

Example 21.1 (Generalized sine functions).

$$\sin_k \theta := \begin{cases} \sin(k\theta) & \text{if } k \geq 0, \\ \sinh(k\theta) & \text{otherwise.} \end{cases}$$

$$\sin_k^* \theta := \begin{cases} \sin(k\theta) & \text{if } k \geq 0, \\ \sinh(-k\theta) & \text{otherwise.} \end{cases}$$

$$= \begin{cases} \sin(k\theta) & \text{if } k \geq 0, \\ -\sinh(k\theta) & \text{otherwise.} \end{cases}$$

$$= \begin{cases} \sin(|k|\theta) & \text{if } k \geq 0, \\ \sinh(|k|\theta) & \text{otherwise.} \end{cases}$$

(see ??)

Example 21.2 (Generalized cosine functions).

$$\cos_k \theta := \begin{cases} \cos(k\theta) & \text{if } k \geq 0, \\ \cosh(k\theta) & \text{otherwise.} \end{cases}$$

$$\cos_k^* \theta := \begin{cases} \cos(k\theta) & \text{if } k \geq 0, \\ \cosh(-k\theta) & \text{otherwise.} \end{cases}$$

$$= \begin{cases} \cos(k\theta) & \text{if } k \geq 0, \\ \cosh(k\theta) & \text{otherwise.} \end{cases}$$

$$= \cos_k \theta$$

(see ??)

Theorem 1 (Pythagorean's identity equivalence).

$$\cos_k^2 \theta + \operatorname{sgn} k \sin_k^2 \theta = 1$$

Proof of ??. Proof by exhaustion. □

Preposition 2 (Generalized trigonometric functions of sum of arguments).

$$\begin{aligned} \sin_k(\theta + \phi) &= \sin_k \theta \cos_k \phi + \cos_k \theta \sin_k \phi \\ \sin_k^*(\theta + \phi) &= \sin_k^* \theta \cos_k \phi + \cos_k \theta \sin_k^* \phi \\ \cos_k(\theta + \phi) &= \cos_k \theta \cos_k \phi - \operatorname{sgn} k \sin_k \theta \sin_k \phi \\ &= \cos_k \theta \cos_k \phi - \sin_k^* \theta \sin_k \phi \\ &= \cos_k \theta \cos_k \phi - \sin_k \theta \sin_k^* \phi \end{aligned}$$

Proof of ??. Proof by exhaustion. □

Preposition 3 (Derivative of generalized trigonometric functions).

$$\sin'_k \theta = k \cos_k \theta$$

$$\sin_k'^* \theta = |k| \cos_k \theta$$

$$\cos'_k \theta = -k \sin_k^* \theta$$

Proof of ??. Proof by exhaustion. □

Matrices

Definition 22. *Generalized rotation matrix* is defined as

$$R_k(\theta) := \begin{bmatrix} \cos_k \theta & -\sin_k^* \theta \\ \sin_k \theta & \cos_k \theta \end{bmatrix},$$

where $\theta \in \mathbb{R}$.

Corollary 2 (Generalized rotation matrix at zero).

$$R_k(0) = I_2$$

Proof of ??. Obvious □

Corollary 3 (Generalized rotation matrix of sum of arguments).

$$R_k(\theta) R_k(\phi) = R_k(\theta + \phi)$$

Proof of ??.

$$R_k(\theta) R_k(\phi) = \begin{bmatrix} \cos_k \theta & -\sin_k^* \theta \\ \sin_k \theta & \cos_k \theta \end{bmatrix} \begin{bmatrix} \cos_k \phi & -\sin_k^* \phi \\ \sin_k \phi & \cos_k \phi \end{bmatrix} \quad (??)$$

$$= \begin{bmatrix} \cos_k \theta \cos_k \phi + (-\sin_k^* \theta) \sin_k \phi & \cos_k \theta (-\sin_k^* \phi) + (-\sin_k^* \theta) \cos_k \phi \\ \sin_k \theta \cos_k \phi + \cos_k \theta \sin_k \phi & \sin_k \theta (-\sin_k^* \phi) + \cos_k \theta \cos_k \phi \end{bmatrix} \quad (??)$$

$$= \begin{bmatrix} \cos_k \theta \cos_k \phi - \sin_k^* \theta \sin_k \phi & -(\sin_k^* \theta \cos_k \phi + \cos_k \theta \sin_k^* \phi) \\ \sin_k \theta \cos_k \phi + \cos_k \theta \sin_k \phi & \cos_k \theta \cos_k \phi - \sin_k \theta \sin_k^* \phi \end{bmatrix} \quad (\text{simplify})$$

$$= \begin{bmatrix} \cos_k \theta + \phi & -\sin_k^* \theta + \phi \\ \sin_k \theta + \phi & \cos_k \theta + \phi \end{bmatrix} \quad (??)$$

$$= R_k(\theta + \phi) \quad (??)$$

$$R_k(\theta) R_k(\phi) = R_k(\theta + \phi)$$

□

Corollary 4 (Inverse of generalized rotation matrix).

$$R_k(\theta)^{-1} = R_k(-\theta)$$

Proof of ??.

$$R_k(\theta) R_k(-\theta) = R_k(0) \quad (\text{??})$$

$$= I_2 \quad (\text{??})$$

$$R_k(-\theta) R_k(\theta) = R_k(0) \quad (\text{??})$$

$$= I_2 \quad (\text{??})$$

$$R_k(\theta) R_k(-\theta) = R_k(-\theta) R_k(\theta) = I_2$$

$$R_k(\theta)^{-1} = R_k(-\theta)$$

□

Definition 23. *Position matrix* is defined recursively as

$$P_{k,n}(\{\theta^1, \dots, \theta^n\}) := \begin{bmatrix} P_{k,n-1}(\{\theta^1, \dots, \theta^{n-1}\}) & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} T_{2,n+1} \begin{bmatrix} R_k(\theta^n) & 0_{2 \times n-1} \\ 0_{n-1 \times 2} & I_{n-1} \end{bmatrix} T_{2,n+1},$$

$$P_{k,0} := I_1,$$

where $\theta = \{\theta^i\} \in \mathbb{R}^n$ for $i \in \{1..n\}$.

Definition 24. Let $P(n, k)$ be set of position matrices.

Corollary 5 (Position matrix at zero).

$$P_{k,n}(0_n) = I_{n+1}$$

Proof of ??. Prove by mathematical induction on n , Let

$$(13) \quad P_{k,n-1}(0_{n-1}) = I_n$$

$$P_{k,n}(0_n) = \begin{bmatrix} P_{k,n-1}(0_{n-1}) & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} T_{2,n+1} \begin{bmatrix} R_k(0) & 0_{2 \times n-1} \\ 0_{n-1 \times 2} & I_{n-1} \end{bmatrix} T_{2,n+1} \quad (\text{??})$$

$$= \begin{bmatrix} I_n & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} T_{2,n+1} \begin{bmatrix} R_k(0) & 0_{2 \times n-1} \\ 0_{n-1 \times 2} & I_{n-1} \end{bmatrix} T_{2,n+1} \quad (\text{??})$$

$$= \begin{bmatrix} I_n & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} T_{2,n+1} \begin{bmatrix} I_2 & 0_{2 \times n-1} \\ 0_{n-1 \times 2} & I_{n-1} \end{bmatrix} T_{2,n+1} \quad (\text{??})$$

$$= I_{n+1} T_{2,n+1} I_{n+1} T_{2,n+1} \quad (\text{??})$$

$$= T_{2,n+1} T_{2,n+1} \quad (\text{??})$$

$$= I_{n+1} \quad (\text{??})$$

$$P_{k,n} (0_n) = I_{n+1}$$

□

Definition 25. *Orientation matrix* is defined as

$$Q_n^\pm (\phi_{n-1}, \phi_{n-2}, \dots, \phi_1) := \begin{bmatrix} 1 & 0_{1 \times n} \\ 0_{n \times 1} & X_{+1,n-1}^\pm (\phi_{n-1}, \phi_{n-2}, \dots, \phi_1) \end{bmatrix},$$

$$Q_0^\pm := \pm I_1,$$

where $\phi_m \in \mathbb{R}^m$ for $m \in \{1..n-1\}$.

Definition 26. Let $Q(n)$ be set of orientation matrices.

Corollary 6 (Orientation matrix at zero).

$$Q_n^+ (0_{n-1}, 0_{n-2}, \dots) = I_{n+1}$$

Proof of ??. Prove by mathematical induction on n , Let

$$(14) \quad Q_{n-1}^+ (0_{n-2}, 0_{n-3}, \dots) = I_n$$

$$Q_n^+ (0_{n-1}, 0_{n-2}, \dots) = \begin{bmatrix} 1 & 0_{1 \times n} \\ 0_{n \times 1} & X_{+1,n-1}^\pm (0_{n-1}, 0_{n-2}, \dots) \end{bmatrix} \quad (\text{??})$$

$$= \begin{bmatrix} 1 & 0_{1 \times n} \\ 0_{n \times 1} & P_{+1,n-1} (0_{n-1}) Q_{n-1}^+ (0_{n-2}, 0_{n-3}, \dots) \end{bmatrix} \quad (\text{??})$$

$$= \begin{bmatrix} 1 & 0_{1 \times n} \\ 0_{n \times 1} & P_{+1,n-1} (0_{n-1}) I_n \end{bmatrix} \quad (\text{??})$$

$$= \begin{bmatrix} 1 & 0_{1 \times n} \\ 0_{n \times 1} & P_{+1,n-1} (0_{n-1}) \end{bmatrix} \quad (\text{??})$$

$$= \begin{bmatrix} 1 & 0_{1 \times n} \\ 0_{n \times 1} & I_n \end{bmatrix} \quad (\text{??})$$

$$= I_{n+1} \quad (??)$$

$$Q_n^+(0_{n-1}, 0_{n-2}, \dots) = I_{n+1}$$

□

Definition 27. *Point matrix* is defined as

$$X_{k,n}^\pm(\theta, \phi_{n-1}, \phi_{n-2}, \dots, \phi_1) := P_{k,n}(\theta) Q_n^\pm(\phi_{n-1}, \phi_{n-2}, \dots, \phi_1),$$

where $\theta \in \mathbb{R}^n$ and $\phi_m \in \mathbb{R}^m$ for $m \in \{1..n-1\}$.

Definition 28. Let $X(n, k)$ be set of point matrices.

Corollary 7 (Position matrix as subset of point matrix).

$$P(k, n) \subset X(k, n)$$

Proof of ??.

$$\forall P \in P(k, n) \forall Q \in Q(n), PQ \in X(k, n) \quad (??)$$

$$\implies PI \in X(k, n) \quad (??)$$

$$\implies P \in X(k, n) \quad (??)$$

$$P(k, n) \subset X(k, n)$$

□

Corollary 8 (Point matrix at zero).

$$X_{k,n}^+(0_n, 0_{n-1}, \dots) = I_{n+1}$$

Proof of ??. It can be implied from ????.

□

Group structure

Preposition 4. *Group of position matrix with multiplication is a subgroup of an orthogonal group for $k > 0$.*

$$(P(n, k), \cdot) \cong O(n+1)$$

Proof.

$$P = P_{k,n}(\{\theta^1, \dots, \theta^n\})$$

$$:= \begin{bmatrix} P_{k,n-1}(\{\theta_1^n, \dots, \theta^{n-1}\}) & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} T_{2,n+1} \begin{bmatrix} R_k(\theta^n) & 0_{2 \times n-1} \\ 0_{n-1 \times 2} & I_{n-1} \end{bmatrix} T_{2,n+1} \quad (??)$$

$$P^T = T_{2,n+1}^T \begin{bmatrix} R_k(\theta^n) & 0_{2 \times n-1} \\ 0_{n-1 \times 2} & I_{n-1} \end{bmatrix}^T T_{2,n+1}^T \begin{bmatrix} P_{k,n-1}(\{\theta_1^n, \dots, \theta^{n-1}\}) & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix}^T \quad (\text{Transpose of product})$$

$$= T_{2,n+1}^T \begin{bmatrix} R_k(\theta^n)^T & 0_{2 \times n-1} \\ 0_{n-1 \times 2} & I_{n-1} \end{bmatrix} T_{2,n+1}^T \begin{bmatrix} P_{k,n-1}(\{\theta_1^n, \dots, \theta^{n-1}\})^T & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} \quad (\text{Transpose of block})$$

$$= T_{2,n+1} \begin{bmatrix} R_k(\theta^n)^T & 0_{2 \times n-1} \\ 0_{n-1 \times 2} & I_{n-1} \end{bmatrix} T_{2,n+1} \begin{bmatrix} P_{k,n-1}(\{\theta_1^n, \dots, \theta^{n-1}\})^T & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} \quad (\text{Transpose of permutation})$$

$$PP^T = \left(\begin{bmatrix} P_{k,n-1}(\{\theta_1^n, \dots, \theta^{n-1}\}) & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} T_{2,n+1} \begin{bmatrix} R_k(\theta^n) & 0_{2 \times n-1} \\ 0_{n-1 \times 2} & I_{n-1} \end{bmatrix} T_{2,n+1} \right) \left(\begin{bmatrix} P_{k,n-1}(\{\theta_1^n, \dots, \theta^{n-1}\})^T & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} \right) \quad (\text{Equation above})$$

$$= \begin{bmatrix} P_{k,n-1}(\{\theta_1^n, \dots, \theta^{n-1}\}) & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} T_{2,n+1} \begin{bmatrix} R_k(\theta^n) & 0_{2 \times n-1} \\ 0_{n-1 \times 2} & I_{n-1} \end{bmatrix} \begin{bmatrix} P_{k,n-1}(\{\theta_1^n, \dots, \theta^{n-1}\})^T & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} \quad ()$$

$$= \begin{bmatrix} P_{k,n-1}(\{\theta_1^n, \dots, \theta^{n-1}\}) & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} T_{2,n+1} \begin{bmatrix} R_k(\theta^n) R_k(\theta^n)^T & 0_{2 \times n-1} \\ 0_{n-1 \times 2} & I_{n-1} \end{bmatrix}$$

$$T_{2,n+1} \begin{bmatrix} P_{k,n-1}(\{\theta_1^n, \dots, \theta^{n-1}\})^T & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} \quad (??)$$

$$= \begin{bmatrix} P_{k,n-1}(\{\theta_1^n, \dots, \theta^{n-1}\}) & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} T_{2,n+1}$$

$$T_{2,n+1} \begin{bmatrix} P_{k,n-1}(\{\theta_1^n, \dots, \theta^{n-1}\})^T & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} \quad ()$$

$$= \begin{bmatrix} P_{k,n-1}(\{\theta_1^n, \dots, \theta^{n-1}\}) & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} \begin{bmatrix} P_{k,n-1}(\{\theta_1^n, \dots, \theta^{n-1}\})^T & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} \quad ()$$

$$\begin{aligned}
&= \begin{bmatrix} P_{k,n-1}(\{\theta^1_n, \dots, \theta^{n-1}\}) P_{k,n-1}(\{\theta^1_n, \dots, \theta^{n-1}\})^T & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} \quad (??) \\
&= I_{n+1} \quad (\text{mathematical induction})
\end{aligned}$$

□

Preposition 5. Group of position matrix with multiplication is a subgroup of an $(1, n)$ -orthochronous indefinite orthogonal group for $k < 0$.

$$(P(n, k), \cdot) \cong O^+(1, n)$$

Proof.

$$\begin{aligned}
P &= P_{k,n}(\{\theta^1, \dots, \theta^n\}) \\
&:= \begin{bmatrix} P_{k,n-1}(\{\theta^1_n, \dots, \theta^{n-1}\}) & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} T_{2,n+1} \begin{bmatrix} R_k(\theta^n) & 0_{2 \times n-1} \\ 0_{n-1 \times 2} & I_{n-1} \end{bmatrix} T_{2,n+1} \quad (??) \\
P^T &= T_{2,n+1}^T \begin{bmatrix} R_k(\theta^n) & 0_{2 \times n-1} \\ 0_{n-1 \times 2} & I_{n-1} \end{bmatrix}^T T_{2,n+1}^T \begin{bmatrix} P_{k,n-1}(\{\theta^1_n, \dots, \theta^{n-1}\}) & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix}^T \quad (\text{Transpose of product}) \\
&= T_{2,n+1}^T \begin{bmatrix} R_k(\theta^n)^T & 0_{2 \times n-1} \\ 0_{n-1 \times 2} & I_{n-1} \end{bmatrix} T_{2,n+1}^T \begin{bmatrix} P_{k,n-1}(\{\theta^1_n, \dots, \theta^{n-1}\})^T & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} \quad (\text{Transpose of block}) \\
&= T_{2,n+1}^T \begin{bmatrix} R_k(\theta^n)^T & 0_{2 \times n-1} \\ 0_{n-1 \times 2} & I_{n-1} \end{bmatrix} T_{2,n+1} \begin{bmatrix} P_{k,n-1}(\{\theta^1_n, \dots, \theta^{n-1}\})^T & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} \quad (\text{Transpose of permutation}) \\
g &= \text{diag } -1, 1, \dots, 1 \\
g P g P^T &= I_{n+1} \quad (\text{mathematical induction})
\end{aligned}$$

$$P_1^1 > 0$$

□

Preposition 6. Group of position matrix with multiplication is isomorphic to translation group for $k \rightarrow 0$.

$$(P(n, k), \cdot) \cong E(n)$$

Proof.

$$\begin{aligned}
 R_k(\theta) &:= \begin{bmatrix} \cos_k(\theta) & -\sin_k^*(\theta) \\ \sin_k(\theta) & \cos_k(\theta) \end{bmatrix} & (\text{??}) \\
 &\rightarrow \begin{bmatrix} 1 & 0 \\ k\theta & 1 \end{bmatrix} & (\text{Limits of the first row}) \\
 P_{k,n}(\{\theta^1, \dots, \theta^n\}) &:= \begin{bmatrix} P_{k,n-1}(\{\theta^1, \dots, \theta^{n-1}\}) & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} T_{2,n+1} \begin{bmatrix} R_k(\theta^n) & 0_{2 \times n-1} \\ 0_{n-1 \times 2} & I_{n-1} \end{bmatrix} T_{2,n+1} & (\text{??}) \\
 &\rightarrow \begin{bmatrix} P_{k,n-1}(\{\theta^1, \dots, \theta^{n-1}\}) & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} T_{2,n+1} \begin{bmatrix} 1 & 0 \\ k\theta^n & 0 \\ 0_{n-1 \times 2} & I_{n-1} \end{bmatrix} T_{2,n+1} & (\text{Equation above}) \\
 &= \begin{bmatrix} P_{k,n-1}(\{\theta^1, \dots, \theta^{n-1}\}) & 0_{n \times 1} \\ 0_{1 \times n} & 1 \end{bmatrix} \begin{bmatrix} I_n & 0_{n \times 1} \\ k\theta^n & 0_{1 \times n-1} \\ 0_{n-1 \times 2} & 1 \end{bmatrix} & (\text{property of permuting}) \\
 &= \begin{bmatrix} P_{k,n-1}(\{\theta^1, \dots, \theta^{n-1}\}) & 0_{n \times 1} \\ k\theta^n & 0_{1 \times n-1} \end{bmatrix} & (\text{??}) \\
 P_{k,n}(\theta) &\rightarrow \begin{bmatrix} 1 & 0_{n \times 1} \\ k\theta & I_n \end{bmatrix} & (\text{mathematical induction}) \\
 P_{k,n}(\theta) P_{k,n}(\phi) &\rightarrow \begin{bmatrix} 1 & 0_{n \times 1} \\ k\theta & I_n \end{bmatrix} \begin{bmatrix} 1 & 0_{n \times 1} \\ k\phi & I_n \end{bmatrix} & (\text{Equation above}) \\
 &= \begin{bmatrix} 1 & 0_{n \times 1} \\ k\theta + k\phi & I_n \end{bmatrix} & (\text{??})
 \end{aligned}$$

□

Preposition 7. Group of orientation with multiplication is isomorphic to orthogonal group.

$$(Q(n), \cdot) \cong O(n)$$

Preposition 8. Group of point with multiplication is isomorphic to orthogonal group for $k > 0$.

$$(X(n), \cdot) \cong O(n)$$

Preposition 9. Group of point with multiplication is isomorphic to $(1, n)$ -orthochronous indefinite orthogonal group for $k < 0$.

$$(X(n), \cdot) \cong O^+(1, n)$$

Preposition 10. Group of point with multiplication is isomorphic to Euclidean group for $k \rightarrow 0$.

$$(X(n), \cdot) \cong E(n)$$

Model Parametrization

Definition 29. For any point matrix $X_{k,n}^\pm(\theta, \phi_1, \phi_2, \dots, \phi_n)$, n -dimensional vector θ is defined as *position parameter*.

Definition 30. For point matrix X , $(n+1)$ -dimensional column vector

$p := \frac{1}{k} X \cdot e^1 = \frac{1}{k} X_1$ is defined as *position vector*.

Definition 31. $P^*(k, n)$ is a set of position vectors.

Lemma 2. For point matrix $X = PO$ where P and O are position and orientation matrix respectively, $p = \frac{1}{k} X_1 = \frac{1}{k} P_1$.

Proof of ??.

$$\begin{aligned} p^i &= \frac{1}{k} X_1^i && ?? \\ &= \frac{1}{k} \sum_j P_j^i O_j^1 && ?? \\ &= \frac{1}{k} P_1^i && ?? \\ p &= \frac{1}{k} P_1 \end{aligned}$$

□

Lemma 3. Given position parameter θ , position vector can be evaluated as the following.

$$\psi_0^{-1} : \theta \mapsto p = \frac{1}{k} \begin{pmatrix} \prod_{j \in \{1..n\}} \cos_k \theta^j \\ \sin_k \theta^{i-1} \prod_{j \in \{i..n\}} \cos_k \theta^j \\ \sin_k \theta^n \end{pmatrix}.$$

Proof of ??. Simplify ????. □

Lemma 4. Given position vector p , position parameter can be calculated as the following.

$$\psi_0 : p \mapsto \theta = \begin{cases} \arcsin_k^{\operatorname{sgn} p^1} \frac{kp^2}{\prod_{j \in \{2..n\}} \cos_k \theta^j} \\ \arcsin_k \frac{kp^{i+1}}{\prod_{j \in \{i+1..n\}} \cos_k \theta^j} \\ \arcsin_k kp^{n+1} \end{cases}$$

$$\in \begin{cases} P \rightarrow \left(-\frac{\pi}{k}, \frac{\pi}{k}\right] \times \left[-\frac{1}{2}\frac{\pi}{k}, \frac{1}{2}\frac{\pi}{k}\right]^{n-1} & \text{if } k > 0 \\ P \rightarrow \mathbb{R}^n & \text{if } k \leq 0 \end{cases}$$

where $\cos_k(\arcsin_k^\pm(x)) = \pm \cos_k(\arcsin_k(x))$.

Proof of ??. From ??,

$$kp = \begin{pmatrix} \prod_{j \in \{1..n\}} \cos_k \theta^j \\ \sin_k \theta^{i-1} \prod_{j \in \{i..n\}} \cos_k \theta^j \\ \sin_k \theta^n \end{pmatrix}$$

$$\sin_k \theta^n = k\theta^{n+1}$$

$$\theta^n = \arcsin_k k\theta^{n+1}$$

$$\sin_k \theta^{i-1} \prod_{j \in \{i..n\}} \cos_k \theta^j = k\theta^i$$

$$\sin_k \theta^{i-1} = \frac{k\theta^i}{\prod_{j \in \{i..n\}} \cos_k \theta^j}$$

$$\theta^{i-1} = \arcsin_k \frac{k\theta^i}{\prod_{j \in \{i..n\}} \cos_k \theta^j}$$

$$\theta^i = \arcsin_k \frac{k\theta^{i+1}}{\prod_{j \in \{i+1..n\}} \cos_k \theta^j}$$

$$p^1 = \prod_{j \in \{1..n\}} \cos_k \theta^j$$

$$p^1 = \cos_k \theta^1 \prod_{j \in \{2..n\}} \cos_k \theta^j$$

$$\operatorname{sgn} p^1 = \operatorname{sgn} \cos_k \theta^1 \prod_{j \in \{2..n\}} \operatorname{sgn} \cos_k \theta^j$$

$$\operatorname{sgn} p^1 = \operatorname{sgn} \cos_k \theta^1 \prod_{j \in \{2..n\}} 1$$

$$\operatorname{sgn} p^1 = \operatorname{sgn} \cos_k \theta^1$$

$$\theta = \begin{pmatrix} \arcsin_k^{\operatorname{sgn} p^1} \frac{kp^2}{\prod_{j \in \{2..n\}} \cos_k \theta^j} \\ \arcsin_k \frac{kp^{i+1}}{\prod_{j \in \{i+1..n\}} \cos_k \theta^j} \\ \arcsin_k kp^{n+1} \end{pmatrix}$$

□

Lemma 5.

$$\Psi = \{ \psi \mid \psi^{-1} \in S^n \rightarrow P : \theta \mapsto P_{k,n}(\theta + x) \text{ for } x \in \mathbb{R}^n \}$$

is a coordinate chart of a C^∞ differential structure on P for

$$S = \begin{cases} \left(-\frac{1}{2}\frac{\pi}{k}, +\frac{1}{2}\frac{\pi}{k}\right) & k > 0 \\ \mathbb{R} & k \leq 0 \end{cases}$$

Proof of ??. From ??, It is sufficient to shows that

1. R_ψ is an open subset of real vector space (defined),
2. $\bigcup_{\psi \in \Psi} D_\psi = P$ (obvious),
3. transition map is in differentiability class C^∞ .

To show that $\bigcup_{\psi \in \Psi} D_\psi = P$.

$$\begin{aligned} M \in P &\implies \exists \theta_0, M = P_{k,n}(\theta_0) \\ &\implies \exists \theta_0, M = P_{k,n}(0 + \theta_0) \\ &\implies M \in R_{\psi^{-1}} \\ &\implies M \in D_\psi \\ &\implies M \in \bigcup_{\psi \in \Psi} D_\psi \end{aligned}$$

$$P \subset \bigcup_{\psi \in \Psi} D_\psi$$

$$M \in \bigcup_{\psi \in \Psi} D_\psi \implies \exists \psi \in \Psi, M \in D_\psi$$

$$\implies \exists \psi \in \Psi, M \in R_{\psi^{-1}}$$

$$\begin{aligned}
&\implies \exists x_0 \exists \theta \in S^n, M = P_{k,n}(\theta + x_0) \\
&\implies \exists x_0, M = P_{k,n}(0 + x_0) \\
&\implies \exists x_0, M = P_{k,n}(x_0) \\
&\implies M \in P
\end{aligned}$$

$$\bigcup_{\psi \in \Psi} D_\psi \subset P$$

$$\bigcup_{\psi \in \Psi} D_\psi = P$$

To show that every transition map is in differentiability class C^∞ .

Consider $\psi_1, \psi_2 \in \Psi$ and $x_1, x_2 \in \mathbb{R}^n$ where

$$\psi_i^{-1} \in S^n \rightarrow R : \theta \mapsto P(\theta + x_i).$$

If $\psi_1^{-1}(\theta_1) = \psi_2^{-1}(\theta_2)$, let $\phi_i = \theta_i + x_i$.

$$\begin{aligned}
\frac{1}{k} \begin{pmatrix} \prod_{j \in \{1..n\}} \cos_k \phi_1^j \\ \sin_k \phi_1^{i-1} \prod_{j \in \{i..n\}} \cos_k \phi_1^j \\ \sin_k \phi_1^n \end{pmatrix} &= \frac{1}{k} \begin{pmatrix} \prod_{j \in \{1..n\}} \cos_k \phi_2^j \\ \sin_k \phi_2^{i-1} \prod_{j \in \{i..n\}} \cos_k \phi_2^j \\ \sin_k \phi_2^n \end{pmatrix} \quad ?? \\
\begin{pmatrix} \prod_{j \in \{1..n\}} \cos_k \phi_1^j \\ \sin_k \phi_1^{i-1} \prod_{j \in \{i..n\}} \cos_k \phi_1^j \\ \sin_k \phi_1^n \end{pmatrix} &= \begin{pmatrix} \prod_{j \in \{1..n\}} \cos_k \phi_2^j \\ \sin_k \phi_2^{i-1} \prod_{j \in \{i..n\}} \cos_k \phi_2^j \\ \sin_k \phi_2^n \end{pmatrix}
\end{aligned}$$

If $k \geq 0$, by mathematical induction, $\phi_1 = m \frac{\pi}{k} \pm \phi_2$.

Otherwise, by mathematical induction, $\phi_1 = \phi_2$.

Hence, the transition map $\tau_{1,2} = \psi_2 \circ \psi_1^{-1}$ is in the form of $\theta \mapsto c \pm \theta$ and is in differentiability class C^∞ .

□

Locus of position vector

Lemma 6. For $k > 0$, P^* is a $(n+1)$ -sphere of radius k^{-1} .

Proof of ??. Simplify ?? using ??.

□

Lemma 7. For $k < 0$, P^* is a forward sheet of a two-sheeted $(n + 1)$ -hyperboloid of radius k^{-1} .

Proof of ??. Simplify ?? using ??.

□

Lemma 8. For $k \rightarrow 0$, P^* is a n -Euclidean manifold at infinity.

Proof of ??. Using limits.

□

Geometric properties

Embedding

Definition 32. Let $N = (Q, g)$ be a n -dimension Riemannian manifold on position parameter space with such inner product g that the map $\cdot \mapsto \frac{1}{k} \cdot e^1 \in P \rightarrow P^*$ is an isometric embedding to $(n + 1)$ -Euclidean manifold.

Remark 4. It have been proved that there is no isometric embedding from H^n to E^{n+1} . Hence, this section is based on a faulty assumption and would not provide a useful information.

Lemma 9. Position parameter θ^i is associated to the following vector in position vector space.

$$\frac{\partial}{\partial \theta^i} = \begin{pmatrix} -k \tan_k^* \theta^i p^j \\ k \frac{1}{\tan_k \theta^i} p^{i+1} \\ 0 \end{pmatrix} \frac{\partial}{\partial p^j}$$

Proof of ??.

$$\begin{aligned} \frac{\partial}{\partial \theta^i} &= \frac{\partial p^j}{\partial \theta^i} \frac{\partial}{\partial p^j} \\ &= \frac{\partial}{\partial \theta^i} \left[\frac{1}{k} \begin{pmatrix} \prod_{l \in \{1..n\}} \cos_k \theta^l \\ \sin_k \theta^{j-1} \prod_{l \in \{j..n\}} \cos_k \theta^l \\ \sin_k \theta^n \end{pmatrix} \right] \frac{\partial}{\partial p^j} \quad ?? \\ &= \frac{1}{k} \frac{\partial}{\partial \theta^i} \left[\begin{pmatrix} \prod_{l \in \{1..n\}} \cos_k \theta^l \\ \sin_k \theta^{j-1} \prod_{l \in \{j..n\}} \cos_k \theta^l \\ \sin_k \theta^n \end{pmatrix} \right] \frac{\partial}{\partial p^j} \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{k} \begin{bmatrix} -k \sin_k^* \theta^i \prod_{l \in \{1..n\}/\{i\}} \cos_k \theta^l \\ -k \sin_k^* \theta^i \sin_k \theta^{j-1} \prod_{l \in \{j..n\}/\{i\}} \cos_k \theta^l \\ k \cos_k \theta^i \prod_{l \in \{i+1..n\}} \cos_k \theta^l \\ 0 \end{bmatrix} \frac{\partial}{\partial p^j} \quad ?? \\
&= \begin{bmatrix} -\tan_k^* \theta^i \prod_{l \in \{1..n\}} \cos_k \theta^l \\ -\tan_k^* \theta^i \sin_k \theta^{j-1} \prod_{l \in \{j..n\}} \cos_k \theta^l \\ \frac{1}{\tan_k \theta^i} \prod_{l \in \{i+1..n\}} \cos_k \theta^l \\ 0 \end{bmatrix} \frac{\partial}{\partial p^j} \\
&= \begin{bmatrix} -k \tan_k^* \theta^i p^j \\ k \frac{1}{\tan_k \theta^i} p^{i+1} \\ 0 \end{bmatrix} \frac{\partial}{\partial p^j}
\end{aligned}$$

□

Lemma 10. The metric tensor of N is

$$g_{ij} = \begin{cases} (1 - \operatorname{sgn} k) \tan_k^2(\theta^a) \prod_{1 \leq j} \cos_k^2 \theta^j + \prod_{a < j} \cos_k^2 \theta^j & \text{if } i = j, \\ 0 & \text{otherwise.} \end{cases}$$

Proof of ??.

$$\begin{aligned}
g_{ab} \left[\frac{\partial}{\partial \theta^i} \right] &= \sum_{l,m=1}^{n+1} \frac{\partial p^l}{\partial \theta^a} g_{lm} \left[\frac{\partial}{\partial p^i} \right] \frac{\partial p^m}{\partial \theta^b} \\
&= \sum_{l=1}^{n+1} \frac{\partial p^l}{\partial \theta^a} \frac{\partial p^l}{\partial \theta^b}
\end{aligned}$$

If $a < b$,

$$\begin{aligned}
g_{ab} &= \begin{bmatrix} -k \tan_k^*(\theta^a) p^j \\ k \frac{1}{\tan_k(\theta^a)} p^{a+1} \\ 0 \end{bmatrix} \cdot \begin{bmatrix} -k \tan_k^*(\theta^b) p^j \\ k \frac{1}{\tan_k(\theta^b)} p^{b+1} \\ 0 \end{bmatrix} \\
&= \sum k^2 \tan_k^*(\theta^a) \tan_k^*(\theta^b) p^{j^2} - k^2 \frac{\tan_k^*(\theta^b)}{\tan_k(\theta^a)} p^{a+1^2}
\end{aligned}$$

$$\begin{aligned}
&= \tan_k(\theta^b) \tan_k(\theta^a) \prod_{a \leq j \leq n+1} \cos_k^2 \theta^j \left(\prod_{1 \leq j < a} \cos_k^2 \theta^j + \sum \sin_k^2 \theta^i \prod_{i < j < a} \cos_k^2 \theta^j - \operatorname{sgn} k \right) \\
&= \tan_k(\theta^b) \tan_k(\theta^a) \prod_{a \leq j \leq n+1} \cos_k^2 \theta^j \left(\prod_{1 \leq j < a} \cos_k^2 \theta^j + \operatorname{sgn} k \left(1 - \prod_{1 \leq j < a} \cos_k^2 \theta^j \right) - \operatorname{sgn} k \right)
\end{aligned}$$

$$g_{ab} = 0.$$

If $a > b$, $g_{ab} = g_{ba} = 0$.

If $a = b$,

$$\begin{aligned}
g_{aa} &= \begin{pmatrix} -k \tan_k^* \theta^a p^j \\ k \frac{1}{\tan_k \theta^a} p^{a+1} \\ 0 \end{pmatrix} \cdot \begin{pmatrix} -k \tan_k^* \theta^a p^j \\ k \frac{1}{\tan_k \theta^a} p^{a+1} \\ 0 \end{pmatrix} \\
&= \sum_{j=1}^a \left(kp^j \tan_k^* \theta^a \right)^2 + \left(kp^{a+1} \cot_k \theta^a \right)^2 \\
&= \sum_{j=1}^a \left(kp^j \tan_k \theta^a \right)^2 + \left(kp^{a+1} \cot_k \theta^a \right)^2 \\
&= \sin_k^2(\theta^a) \prod_{a < j} \cos_k^2 \theta^j \left(\prod_{1 \leq j < a} \cos_k^2 \theta^j + \sum_{1 \leq i < a} \sin_k^2 \theta^i \prod_{i < j < a} \cos_k^2 \theta^j + \cot_k^2(\theta^a) \right) \\
&= \sin_k^2(\theta^a) \prod_{a < j} \cos_k^2 \theta^j \left(\prod_{1 \leq j < a} \cos_k^2 \theta^j + \sum_{1 \leq i < a} \sin_k^2 \theta^i \prod_{i < j < a} \cos_k^2 \theta^j + \cot_k^2(\theta^a) \right) \\
&= \sin_k^2(\theta^a) \prod_{a < j} \cos_k^2 \theta^j \left(\prod_{1 \leq j < a} \cos_k^2 \theta^j - \operatorname{sgn} k \prod_{1 \leq j < a} \cos_k^2 \theta^j + \operatorname{sgn} k \prod_{1 \leq j < a} \cos_k^2 \theta^j + \sum_{1 \leq i < a} \sin_k^2 \theta^i \prod_{i < j < a} \cos_k^2 \theta^j \right) \\
&= \sin_k^2(\theta^a) \prod_{a < j} \cos_k^2 \theta^j \left(\prod_{1 \leq j < a} \cos_k^2 \theta^j - \operatorname{sgn} k \prod_{1 \leq j < a} \cos_k^2 \theta^j + \operatorname{sgn} k + \cot_k^2(\theta^a) \right) \\
&= \sin_k^2(\theta^a) \prod_{a < j} \cos_k^2 \theta^j \left(\prod_{1 \leq j < a} \cos_k^2 \theta^j - \operatorname{sgn} k \prod_{1 \leq j < a} \cos_k^2 \theta^j + \operatorname{sgn} k + \csc_k^2(\theta^a - \operatorname{sgn} k) \right) \\
&= \sin_k^2(\theta^a) \prod_{a < j} \cos_k^2 \theta^j \left((1 - \operatorname{sgn} k) \prod_{1 \leq j < a} \cos_k^2 \theta^j + \csc_k^2(\theta^a) \right) \\
&= (1 - \operatorname{sgn} k) \tan_k^2(\theta^a) \prod_j \cos_k^2 \theta^j + \prod_{a < j} \cos_k^2 \theta^j
\end{aligned}$$

□

Curvature

Curvature (Method I)

This method may be easier to generalize to Model II where each direction can have partially independent curvature (or even Model III where extrinsic curvature become a thing). But it may be challenging to define Gauss map properly.

Lemma 11. *Given a position parameter θ , the tangent vector in position vector space can be calculated as follows*

$$\nu(p) = \begin{cases} \begin{bmatrix} kp^1 \\ +kp^i \\ 1 \\ 0 \end{bmatrix} & k > 0 \\ \frac{1}{\sqrt{-1+2(kp^1)^2}} \begin{bmatrix} kp^1 \\ -kp^i \end{bmatrix} & k = 0 \\ \begin{bmatrix} kp^1 \\ -kp^i \end{bmatrix} & k < 0 \end{cases}$$

Proof of ??. From ?????,

$$p \in P \iff \begin{cases} \sum_i p^{i2} = k^{-2} & k > 0, \\ p^1 = k^{-1} & k \rightarrow 0, \\ p^{12} - \sum_{1 < i} p^{i2} = k^{-2} \wedge p^1 > 0 & k < 0. \end{cases}$$

Let

$$F(p) = \begin{cases} \sum_i p^{i2} - k^{-2} & k > 0, \\ p^1 - k^{-1} & k \rightarrow 0, \\ p^{12} - \sum_{1 < i} p^{i2} - k^{-2} & k < 0. \end{cases}$$

$$n = l \nabla F(p)$$

$$= l \begin{cases} \nabla \sum_i p^{i2} - k^{-2} & k > 0, \\ \nabla p^1 - k^{-1} & k \rightarrow 0, \\ \nabla p^{12} - \sum_{1 < i} p^{i2} - k^{-2} & k < 0. \end{cases}$$

$$\begin{aligned}
&= l \begin{cases} \begin{pmatrix} 2p^i \\ 1 \\ 0 \end{pmatrix} & k > 0, \\ \begin{pmatrix} 1 \\ 0 \end{pmatrix} & k \rightarrow 0, \\ \begin{pmatrix} -2p^1 \\ 2p^i \end{pmatrix} & k < 0. \end{cases} \\
\hat{n} &= \pm \begin{cases} \begin{pmatrix} \frac{1}{||p||} (p^i) \\ 1 \\ 0 \end{pmatrix} & k > 0, \\ \begin{pmatrix} 1 \\ 0 \end{pmatrix} & k \rightarrow 0, \\ \begin{pmatrix} \frac{1}{||p||} (-p^1) \\ p^i \end{pmatrix} & k < 0. \end{cases} \\
&= \pm \begin{cases} \begin{pmatrix} \frac{1}{k^{-1}} (p^i) \\ 1 \\ 0 \end{pmatrix} & k > 0, \\ \begin{pmatrix} 1 \\ 0 \end{pmatrix} & k \rightarrow 0, \\ \begin{pmatrix} \frac{1}{\sqrt{-1+2(kp^1)^2}} (-p^1) \\ p^i \end{pmatrix} & k < 0. \end{cases} \\
\nu = \hat{n} &:= \begin{cases} \begin{pmatrix} kp^i \\ 1 \\ 0 \end{pmatrix} & k > 0, \\ \begin{pmatrix} 1 \\ 0 \end{pmatrix} & k \rightarrow 0, \\ \begin{pmatrix} \frac{1}{\sqrt{-1+2(kp^1)^2}} (p^1) \\ -p^i \end{pmatrix} & k < 0. \end{cases} \quad \square
\end{aligned}$$

Lemma 12.

$$L(p) = \begin{cases} -kI & k > 0, \\ 0 & k \rightarrow 0, \\ \frac{1}{\|p\|} \left(\frac{1}{\|p\|^2} pp^T \text{diag}(-1, 1 \dots, 1) - I \right) & k < 0. \end{cases}$$

Proof of ??.

$$\begin{aligned} \frac{\partial}{\partial p^j} \frac{1}{\|p\|} &= \frac{\partial}{\partial \|p\|} \frac{1}{\|p\|} \frac{\partial}{\partial p^j} \|p\| \\ &= -\frac{1}{\|p\|^2} \frac{\partial}{\partial p^j} \sqrt{\sum_i p^{i2}} \\ &= -\frac{1}{\|p\|^2} \frac{\partial}{\partial \sum_i p^{i2}} \sqrt{\sum_i p^{i2}} \frac{\partial}{\partial p^j} \sum_i p^{i2} \\ &= -\frac{1}{\|p\|^2} \frac{1}{2\sqrt{\sum_i p^{i2}}} 2p^j \\ &= -\frac{1}{\|p\|^2} \frac{1}{2\|p\|} 2p^j \\ &= -\frac{p^j}{\|p\|^3} \end{aligned}$$

From ??,

$$\begin{aligned} L(p) &= -(D\nu \circ (Df)^{-1})(p) \\ &= -(D\nu)(p) \\ &= - \left(\frac{\partial}{\partial p^j} \nu^i \Big|_p \right)_{i,j} \\ (L(p))_{i,j} &= \begin{cases} -\frac{\partial}{\partial p^j} kp^i & k > 0, \\ -\frac{\partial}{\partial p^j} 1 & k \rightarrow 0 \text{ and } i = 1, \\ -\frac{\partial}{\partial p^j} 0 & k \rightarrow 0 \text{ and } i \neq 1, \\ -\frac{\partial}{\partial p^j} \left(\frac{1}{\|p\|} p^1 \right) & k < 0 \text{ and } i = 1, \\ -\frac{\partial}{\partial p^j} \left(-\frac{1}{\|p\|} p^i \right) & k < 0 \text{ and } i \neq 1, \end{cases} \end{aligned}$$

$$\begin{aligned}
&= \begin{cases} -k & k > 0 \text{ and } i = j, \\ 0 & k > 0 \text{ and } i \neq j, \\ 0 & k \rightarrow 0, \\ -\left(\frac{1}{\|p\|} \frac{\partial}{\partial p^j} p^1 + p^1 \frac{\partial}{\partial p^j} \frac{1}{\|p\|}\right) & k < 0 \text{ and } i = 1, \\ \left(\frac{1}{\|p\|} \frac{\partial}{\partial p^j} p^i + p^i \frac{\partial}{\partial p^j} \frac{1}{\|p\|}\right) & k < 0 \text{ and } i \neq 1, \end{cases} \\
&= \begin{cases} -k & k > 0 \text{ and } i = j, \\ 0 & k > 0 \text{ and } i \neq j, \\ 0 & k \rightarrow 0, \\ -\frac{1}{\|p\|} + \frac{p^1{}^2}{\|p\|^3} & k < 0 \text{ and } i = j = 1, \\ \frac{p^1 p^j}{\|p\|^3} & k < 0 \text{ and } j \neq i = 1, \\ \frac{1}{\|p\|} - \frac{p^i{}^2}{\|p\|^3} & k < 0 \text{ and } i = j \neq 1, \\ -\frac{p^i p^j}{\|p\|^3} & k < 0, i \neq j \text{ and } i \neq 1, \end{cases}
\end{aligned}$$

For $k > 0$, $L(p) = -kI$.

For $k \rightarrow 0$, $L(p) = 0$.

For $k < 0$, $L(p) = \frac{1}{\|p\|} \left(\frac{1}{\|p\|^2} pp^T \operatorname{diag}(-1, 1 \dots, 1) - I \right)$. □

Lemma 13.

Proof of ??. The proof is left as an exercise to the other author.

For $k > 0$,

$$\begin{aligned}
0 &= \det L(p) - \lambda I \\
&= \det -kI - \lambda I \\
&= \det (-k - \lambda)I \\
&= (-k - \lambda)^{n+1} \det I \\
&= (-k - \lambda)^{n+1} \\
\lambda &\in \{-k, \dots, -k\}
\end{aligned}$$

For $k < 0$,

$$\begin{aligned}
0 &= \det L(p) - \lambda I \\
&= \det \frac{1}{\|p\|} \left(\frac{1}{\|p\|^2} pp^T \operatorname{diag}(-1, 1 \dots, 1) - I \right) - \lambda I \\
&= \det \frac{1}{\|p\|} \left(\frac{1}{\|p\|^2} pp^T \operatorname{diag}(-1, 1 \dots, 1) - I \right) - \left(\lambda' - \frac{1}{\|p\|} \right) I \quad \lambda' := \lambda + \frac{1}{\|p\|} \\
&= \det \frac{1}{\|p\|^3} pp^T \operatorname{diag}(-1, 1 \dots, 1) - \frac{1}{\|p\|} I - \left(\lambda' - \frac{1}{\|p\|} \right) I \\
&= \det \frac{1}{\|p\|^3} pp^T \operatorname{diag}(-1, 1 \dots, 1) - \lambda' I \\
&= \det \frac{1}{\|p\|^3} pp^T \operatorname{diag}(-1, 1 \dots, 1) - \frac{1}{\|p\|^3} \lambda'' I \\
&= \frac{1}{\|p\|^{3(n+1)}} \det pp^T \operatorname{diag}(-1, 1 \dots, 1) - \lambda'' I \quad \lambda'' := \|p\|^3 \lambda' \\
&= \det pp^T \operatorname{diag}(-1, 1 \dots, 1) - \lambda'' I \\
&= \lambda''' \left(\lambda'' + p^{12} - \sum_{i \neq 1} p^{i2} \right) \\
0 &= \lambda''' \left(\lambda'' + \frac{1}{k} \right) \\
\lambda'' &\in \left\{ -\frac{1}{k}, 0, \dots, 0 \right\} \\
\lambda' &\in \left\{ -\frac{1}{k\|p\|^3}, 0, \dots, 0 \right\} \\
\lambda &\in \left\{ -\frac{1}{k\|p\|^3} - \frac{1}{\|p\|}, -\frac{1}{\|p\|}, \dots, -\frac{1}{\|p\|} \right\} \\
&= \left\{ -\frac{1 + \|p\|^2}{k\|p\|^3}, -\frac{1}{\|p\|}, \dots, -\frac{1}{\|p\|} \right\} \\
&= \left\{ -\frac{2k^2 p^{12}}{k\|p\|^3}, -\frac{1}{\|p\|}, \dots, -\frac{1}{\|p\|} \right\} \\
&= \left\{ -\frac{2kp^{12}}{\|p\|^3}, -\frac{1}{\|p\|}, \dots, -\frac{1}{\|p\|} \right\} \\
&= \left\{ k^2 \|p\| \frac{-2p^{12}}{k\|p\|^4}, -\frac{1}{\|p\|}, \dots, -\frac{1}{\|p\|} \right\} \\
&= \left\{ k^2 \|p\| \frac{-2p^{12}}{k\|p\|^4}, -\frac{1}{\|p\|}, \dots, -\frac{1}{\|p\|} \right\}
\end{aligned}$$

□

Curvature (Method II)

This method may be easier to be done (despite the fact that it never finished). But it raises problems when trying to generalize e.g. dealing with extrinsic curvature (which may be introduced in Model III if not to mess with other basis geometries).

Lemma 14.

Lemma 15.

Curvature (Conclusion)

Lemma 16.

CurvatureParameter. It can be seen that $\sec(p) = \kappa = \operatorname{sgn}(k)k^2$. Hence, when provided κ, k can be determined and used to evaluate the model.

The Model

Definition 1.1. For any parameter $\kappa, n,$

$$\begin{aligned} \mathbb{M} &:= M \\ \otimes_{\mathbb{M}} &:= \cdot \\ \varphi_{\mathbb{M}} &:= X \mapsto \theta \\ g_{\mathbb{M}} &:= g \\ \mathbb{P}_{\mathbb{M}} &\equiv R \\ \mathbb{T}_{\mathbb{M}} &\equiv M \end{aligned}$$

for injective smooth function $K : \kappa \mapsto k = \operatorname{sgn} \kappa \sqrt{|\kappa|} \in \mathbb{R} \rightarrow \mathbb{R}$.

Assertion 1.1.

Assertion 1.2.

Future plan

Model II

It is very simple to be able to model composite geometries e.g. $S^2 \times E$ by tensor product of the existing model. But to be able to merge them as smooth model may be

challenging since not all combination of basis curvature have their own intrinsic geometry. So it may be to find independent variable for each basis or to introduce extrinsic curvature (Model A).

Model B

It is known that E emerged at $n \geq 1$ while S and H emerged at $n \geq 2$ and there's more complex pure geometries than these that emerged in higher dimension. It is interesting and challenging to explore such geometries and prove whether the curvature still works as indicator in such geometries or are there any patterns for their symmetries.

Model A

This model is based on curvature and mostly just 3 basis geometries and extrinsic curvature which seems to be interesting despite some critical result in some combination e.g. $S^1 \times S^1$ vs S^2 . It can be even more challenging to have variable curvature with respect to other intrinsic position.




Figure 1

Generalized trigonometric functions as function of k

Note. This graph shows the value of generalized trigonometric functions as solid line and trigonometric and hyperbolic functions in the unused domain as dashed line.




Figure 2

Generalized sine function




Figure 3

Generalized sine function variant




Figure 4

Generalized cosine function