

# The Galactic Plane

Lukas Finkbeiner

May 11, 2020

## Abstract

We study

1. velocity patterns of various H clouds using Doppler shifts
  2. black hole mass
  3. accuracy of a spiral arm fit
- results, uncertainties.

## 1 Introduction and Background

Given certain patterns, how well does a spiral fit? We need to control for the total number of data. We want the normalization of any two compared sets to be the same (so, we scale accordingly). What numpy functions I used.

$$R_{\text{arm}} = R_0 e^{\kappa(\phi - \phi_0)} \quad (1)$$

To explore these questions, we want to be able to examine the Doppler velocities and brightness temperatures associated with different parts of the galactic plane of the Milky Way. We want to examine power spectra collected for values of the galactic longitude  $\ell$  on the range  $-10^\circ$  to  $250^\circ$ . We will provide a more comprehensive analysis of our particular setup in the methods section. For now, we consider, in the abstract, two equations for calibrating the spectra.

$$T_{\text{line}} = G \frac{s_{\text{on}}}{s_{\text{off}}} \quad (2)$$

$$G = \frac{T_{\text{sys, cal}} - T_{\text{sys, cold}}}{\sum (s_{\text{cal}} - s_{\text{cold}})} \sum s_{\text{cold}} \quad (3)$$

The first equation will be used to demonstrate a consistent HI signal by combining spectra for two different frequencies in the local oscillator. The second equation will be used to scale the power spectrum so that the y-axis carries information about the brightness temperatures associated with different frequencies on our spectra.

The following equation is a conclusion of the tangent-point method ( $0^\circ < \ell \leq 90^\circ$ ) for converting galactic rotation to Doppler velocity.  $R_\odot \approx 220 \text{ km / s}$  and  $R_\odot \approx 8.5 \text{ kpc} \equiv 2.6 \times 10^{17} \text{ km}$ .




Figure 1: This is a simple illustration demonstrating how the value of  $\ell$  can place constraints on valid values for  $R$  (specifically,  $R_2$  here represents the right-triangle case, for which the radius is minimized). However, since our universal value for galactic latitude  $b = 0^\circ$  has no relevance to radial distances, no pair of galactic coordinates will specify the radius at which we are working. Consequently, the tangent-point method must explicitly establish that we work with a specific radius  $R = R_{\min}$ .

$$V_{\text{Dopp}} = \left[ \frac{V(R)}{R} - \frac{V(R_\odot)}{R_\odot} \right] R_\odot \sin(\ell) \quad (4)$$

The following representation will be used in establishing the velocity curve inside the solar circle:

$$V(R) = \frac{R}{R_\odot} \left[ \frac{V_{\text{Dopp}}}{\sin \ell} - V(R_\odot) \right] \quad (5)$$

In preparation for our mass estimations, we start with the conventional

$$v^2 = \frac{GM_{\text{gravitational}}}{R} \quad (6)$$

Then, we re-arrange to get the desired form:

$$M_{\text{gravitational}} = \frac{Rv^2}{G} \quad (7)$$

As we collect Doppler velocities for each galactic longitude  $\ell$ , it is important to distinguish between  $\ell$  and the radial distance  $R$  of a hydrogen cloud to the galactic center. To estimate mass for the part of the galaxy inside the solar circle, we use the tangent point method by applying equation 5 with  $R = R_{\min} = R_\odot \sin \ell$ . In figure 1, we illustrate the range of radii associated with an arbitrary  $\ell$ . The tangent-point method necessarily excludes these other radii. Later on, in the observations section, we will visualize some consequences of this.




Figure 2: This is a visibility demonstration to clarify our observation process: we hold the local sidereal time fixed and plot the topocentric coordinate conversions for all values of the galactic longitude  $\ell$  on our range. We began this project with the nominal range of  $\ell \in [-10^\circ, 250^\circ]$ . By calculating similar visibility curves for a full sidereal day, we are able to classify some of our observations with higher risk for obstruction (particularly when the target altitude is close to the dish’s minimum attainable altitude). We also calculated that  $\ell = -10^\circ$  is not visible for any sufficiently wide segment of the day. This visibility plot was generated at 19:40 on the tenth of May.

## 2 Methods

To conduct all of our measurements of the galactic plane of the Milky Way, we run a tracking script on a remote computer linked to the Leuschner dish. Our full observation consists of 260 galactic coordinate pairs where the galactic latitude is a constant  $0^\circ$  and the galactic longitude begins at  $-9^\circ$  and increases in  $1^\circ$  increments. We convert from galactic to topocentric coordinates using the rotation matrices [as described in a previous report](#). We schedule our measurements by entering a value for the local sidereal time and considering which galactic coordinate pairs correspond to topocentric coordinate pairs which are currently accessible by the Leuschner dish (see figure 2 for an example case).

As a demonstration of the Leuschner dish signal chain (illustrated in figure 3), we show how we map a set of input signals to a final frequency axis for our plots (frequency versus brightness temperature). Since we are considering only frequency, this allows us to skip over the amplifiers and attenuators.

The initial band-pass filter ranges from 1.4 to 1.63 GHz. There is a down-converter that mixes this filtered signal with the signal from the local oscillator LO. This leads to a new




Figure 3: This is a simplification of the Leuschner signal chain, where the horizontal polarization has been ignored (we do not use those data in this report). Furthermore, we abstract-away the clock rate of the ROACH and the various stages of digitization. For our ‘on’ LO frequency of 1270 MHz, the output spectra cover a frequency range 1415 to 1425 MHz.

frequency band (intermediate frequency IF) of 130 MHz to 360 MHz. Since we are using a real-valued mixer, we apply a band-pass filter to eliminate the sum frequencies. This filter is centered at 150 MHz and has a bandwidth of 10 MHz. Consequently, we expect the following frequency range in our outputs (which take the form of spectra in .fits files): a 10 MHz wide range beginning at 145 MHz IF (which maps to 1.415 GHz RF) and ending at 155 MHz IF (which maps to 1.425 GHz RF).

Before we analyze these results, we need to combine each pair of four averaged spectra into a single calibrated spectrum which comprehensively describes our observations for a single galactic coordinate pair. We want to change our local oscillator frequency (as indicated in equation 2) between two values. This allows us to, via the mixer, adjust where the HI line appears in our spectra. We thereby create ‘on’ (LO at 1270 MHz) and ‘off’ (LO at 1268 MHz) spectra. Our ‘off’ frequency, 1268 MHz, leads to an output range of 1.417 to 1.419 GHz.

The gain,  $G$  (indicated in equation 3), is calculated with an additional spectrum (*cal*) that we deliberately contaminate with thermal noise. We do this by activating a noise diode wired close to the Leuschner dish receiver. For the vertical-polarization receiver (with which we are exclusively concerned here), the noise diode was independently estimated to contribute about 80 Kelvins. For our ‘cold’ spectrum, we simply use the spectrum measured for the same galactic coordinates but without the noise diode enabled. We control for any time-dependent environmental inconsistencies by taking these two spectra consecutively.

We sample at 24 MHz (we keep 1/32 of the samples from the 768 MHz clock). We have a 12MHz Nyquist bandwidth, and we are operating in the twelfth Nyquist window. To protect our results from noise, we record 10 spectra for each case (on spectrum with noise diode, on spectrum without, off spectrum with noise diode, off spectrum without) and independently average them. Figure 4, in our observations section, is an example of a fully calibrated




Figure 4: This is an example of a fully calibrated spectrum ( $\ell = 80^\circ$ ), for which we have created 260. There is a black vertical line indicating the location of the 21 cm HI line at rest (1420.405751786 MHz at rest). Despite the forty spectra used to produce a single spectrum of this kind, there remain clear limitations to the utility. There is a clear sinusoidal trend (with a sloped baseline) as well as systematics spikes (here visible at 1.424 GHz) which persist across spectra. Our intensity maximizer function is generally resistant to these annoyances due to the comparatively high signal to noise ratio of our setup. Nevertheless, there do exist some exceptions, as our later plots potentially feature.

spectrum which shows a sort of HI mirroring effect (a direct consequence of dividing the on and off spectra) as well as a y-axis that is physically meaningful due to this procedure.

### 3 Observations

By maximizing the brightness temperature for the fully calibrated spectrum (consider how the shape of figure 4 would be friendly to a primitive maximization algorithm) of every galactic longitude on our range, we are able to produce a single plot which represents an overview of our data for this project. Figure 5 is close to what we would consider our image  $T_A(\ell, V_{LSR})$  except that the graph is not colored to reflect varying brightness temperatures at the different angles. However, on its own, the figure does reflect the peculiarity of the various gas clouds in that the plot does not always appear to follow consistent patterns.

Additionally, we may more quantitatively explore the idea pursued in the illustration for figure 1. Figure 6 represents a dispersion of lines relating linear velocity to the distance from the galactic center. We create this visualization by treating radial distance and galactic longitude as parameters essentially independent but for the constraint that  $R_{\min} = R_\odot \sin \ell$  and  $R_{\max} = R_\odot$ . This particular distribution appears highly symmetric, which may at first counter worries as to the rigor of a tangent-point approach.

However, since the tangent-point method is applied exclusively to the range  $\ell \in (0^\circ, 90^\circ]$ , a more relevant comparison is offered in figure 7. Selecting the minimum value for radial distance seems more natural as we approach ninety degrees (the arc length vanishes and we




Figure 5: We take the peak intensity of each spectrum to represent the shifted HI line for that galactic longitude. Observe that, for our tangent-point considerations ( $\ell \in [0, 90]$ ), we have an initial region featuring wild fluctuations followed by a relatively flat region for the second half of the angles. While this could represent physical realities, this seeming discontinuity of the derivative could also be a consequence of the primitive intensity-maximizing function, or it could be a consequence of pointing the telescope at angles close to mechanical boundaries. This is *almost* the image  $T_A(\ell, V_{\text{LSR}})$ ; we need the brightness temperature associated with each Doppler velocity, then we can color this.




Figure 6: This is an overplot of all 90 possible velocity curves. Specifically, when we treat the solar circle of the galaxy, we know that for each  $\ell$ ,  $R$  can take on a maximum value of  $R_{\odot}$  (the radius of the solar circle) and a minimum value of  $R_{\odot} \sin \ell$ . The tangent-point method mandates the usage of  $R_{\min}$ , but we here demonstrate how that method necessitates the collapse of line segments into points.






Figure 7: Here we are essentially representing similar information to that of figure 6. We use a semilog vertical axis to verify that there is one consistently-varying multiplicative offset ( $\frac{1}{\sin \ell}$ ) as we examine different galactic longitudes. This plot also agrees with our expectation that our two boundary cases are meaningless:  $\ell = 0^\circ$  approaches an infinite linear velocity and  $\ell = 90^\circ$  corresponds to a velocity curve for which the minimum and maximum radii are the same (an infinitesimal segment).

are treating one value for radius anyway). However, as we approach the other end, the arc lengths begin to explode.

Ten degrees is still far from our lower bound of zero, but we can see here that the single galactic longitude spans a linear velocity of around an order of magnitude. Furthermore, we need radius anyway for our classical mass considerations (refer to equation 7), and the problem is more concerning for that (we nearly span the full range of the solar circle). With a square term on velocity and a first-order radius term, we can expect a range of values which span nearly three orders of magnitude on mass. This author lacks sufficient understanding of the tangent-point method to make conclusions about its accuracy, which means that this situation is neither a problem to be ignored nor, necessarily, a problem to which to draw academic attention.

## 4 Analysis

If we take the  $V(R) \approx 220$  km/s for  $R > R_\odot \approx 8.5$  kpc, we may apply equation 7 out to radius of the Milky Way, estimated by [people](#) to be about 18.5 kpc. This corresponds to the ideal cumulative mass plot, figure 8. The final mass, for the range exclusively defined as the outer region of the Milky Way, rests at  $2.23808 \times 10^{32}$  kg (about one hundred solar masses).

Finally, we can estimate the mass of the super-massive black hole at the center of the milky way if we use the Doppler velocity for our closest angle ( $\ell = 1^\circ$ ) and combine equations 5 and 7 and integrate over radii from the galactic center out to a single light-year. According to [blah](#), the active galactic nucleus is about  $4.4 \times 10^7$  km in radius. By comparison, a light year is six orders of magnitude larger than the estimated radius of the black hole. **I have no**




Figure 8: This is a cumulative mass distribution for radii larger than that of the solar circle. The plot begins with a mass of zero because we subtract-off whatever the function calculates for radii smaller than that of the solar circle; we know the calculations to be erroneous for the smaller radii because we are applying different calculation methods. Observe that the rate of increase of mass drops off rather steeply. We hope for this to be the case as the galaxy thins out toward its edges and is most dense at smaller radii. However, the transition is rather jarring to be occurring as far out as the Sun. It may point to a deficiency in the mass approximation near  $R_{\odot}$ .




Figure 9: This is a plot of galactic rotation velocities by radius, calculated using the tangent-point method. If we approximated the galactic plane as a rigid disk, we would expect the velocity to increase with greater radius (as the same angular distance must be covered in the same amount of time at a greater radius). While this line does not disqualify a spiral, we might initially expect a spiral shape to come from *similar* linear velocities manifesting as curved arms at greater radii. It is not clear if the fall-off of gravitational influence would agree with this model.






Figure 10: This is a semilog plot of interstellar cloud mass versus radius. The calculation rests conceptually on a cumulative distribution function, so we may immediately spot imperfections in the data. Specifically, after about 3 kpc, the data do not leveling off consistently. Instead, the mass response fluctuates and then slowly declines. However, the initial rise does fit with our expectation of a general decrease in density as we move outward from the center.

idea what is going on.

The result?  $\text{Mass} \approx 7.6405 \times 10^{13} \text{ kg}$ . This is seventeen orders of magnitude below the mass of the sun. Something has gone terribly wrong. Even if I were to expand to expand my range of radii to 10000 times the radius of the black hole, I would still be at three-quarters the mass of our sun.

## 5 Conclusions

We seem to have at least a couple of results! How do I estimate the uncertainty on the Leuschner dish?

## 6 Acknowledgments

Who did what?

Theory and background provided by Aaron Parsons. “LAB 4: Mapping the Galactic HI Line.” Updated April 2020.

## References

- [1] Michel Goossens, Frank Mittelbach, and Alexander Samarin. *The L<sup>A</sup>T<sub>E</sub>X Companion*. Addison-Wesley, Reading, Massachusetts, 1993.

- [2] Albert Einstein. *Zur Elektrodynamik bewegter Körper*. (German) [*On the electrodynamics of moving bodies*]. Annalen der Physik, 322(10):891–921, 1905.
- [3] Knuth: Computers and Typesetting,  
<http://www-cs-faculty.stanford.edu/~uno/abcde.html>