

# The Galactic Plane

Lukas Finkbeiner

May 10, 2020

## Abstract

We study

1. velocity patterns of various H clouds using Doppler shifts
  2. black hole mass
  3. accuracy of a spiral arm fit
- results, uncertainties.

## 1 Introduction and Background

Given certain patterns, how well does a spiral fit? We need to control for the total number of data. We want the normalization of any two compared sets to be the same (so, we scale accordingly). What numpy functions I used.

21 cm line transition: 1420.405751786 MHz at rest.

$$R_{\text{arm}} = R_0 e^{\kappa(\phi - \phi_0)} \quad (1)$$

The following equation is a conclusion of the tangent-point method for converting galactic rotation to Doppler velocity.  $R_\odot \approx 220 \text{ km / s}$  and  $R_\odot \approx 8.5 \text{ kpc} \equiv 2.6 \times 10^{17} \text{ km}$ .

$$V_{\text{Dopp}} = \left[ \frac{V(R)}{R} - \frac{V(R_\odot)}{R_\odot} \right] R_\odot \sin(\ell) \quad (2)$$

The following representation will be used in establishing the velocity curve inside the solar circle:

$$V(R) = \frac{R}{R_\odot} \left[ \frac{V_{\text{Dopp}}}{\sin \ell} - V(R_\odot) \right] \quad (3)$$

You have to cite sources and find a better common unit.

Before we analyze the output of our spectrometer, we need to calibrate the spectra. The concepts are the same as in lab 2 (this should be a citation?).

We want to change our local oscillator frequency between two values. This allows us to, via the mixer, adjust where the HI line appears in our spectra. We thereby create ‘on’ (LO at 1270 MHz) and ‘off’ (LO at 1268 MHz) spectra.

$$T_{\text{line}} = G \frac{s_{\text{on}}}{s_{\text{off}}} \quad (4)$$




Figure 1: This is a simplification of the Leuschner signal chain, where the horizontal polarization has been ignored (we do not use those data in this report). Furthermore, we abstract-away the clock rate of the ROACH and the various stages of digitization. For our ‘on’ LO frequency of 1270 MHz, the output spectra cover a frequency range 1415 to 1425 MHz.

The gain,  $G$ , is calculated with an additional spectrum (*cal*) that we deliberately contaminate with thermal noise. We do this by activating a noise diode which, for the vertical-polarization receiver (with which we are exclusively concerned here) is independently estimated to contribute about 80 Kelvins. Finally, we compare the magnitudes with a spectrum without this contamination and we compute:

$$G = \frac{T_{\text{sys, cal}} - T_{\text{sys, cold}}}{\sum (s_{\text{cal}} - s_{\text{cold}})} \sum s_{\text{cold}} \quad (5)$$

As a demonstration of the Leuschner dish signal chain (illustrated in figure 1), we show how we map a set of input signals to a final frequency axis for our plots (frequency versus brightness temperature). Since we are considering only frequency, this allows us to skip over the amplifiers and attenuators.

The initial band-pass filter ranges from 1.4 to 1.63 GHz. There is a down-converter that mixes this filtered signal with the signal from the local oscillator LO. This leads to a new frequency band (intermediate frequency IF) of 130 MHz to 360 MHz. Since we are using a real-valued mixer, we apply a band-pass filter to eliminate the sum frequencies. This filter is centered at 150 MHz and has a bandwidth of 10 MHz. Consequently, we expect the following frequency range in our outputs (which take the form of spectra in .fits files): a 10 MHz wide range beginning at 145 MHz IF (which maps to 1.415 GHz RF) and ending at 155 MHz IF (which maps to 1.425 GHz RF).

We used 1268 MHz for our second LO frequency, the ‘off’ spectrum. This leads to an output range of 1.417 to 1.419 GHz.

We sample at 24 MHz (we keep 1/32 of the samples from the 768 MHz clock), -12 MHz Nyquist bandwidth. so we are operating in the twelfth Nyquist window.

## 2 Methods

To observe the galactic plane, we have our trusty tracking script. After three labs' worth of development, it is almost correct!

How we calibrate: for each value  $\ell$ , we take four sets of ten spectra: ...

We convert from galactic to topocentric coordinates using rotation matrices, as described in a previous report. How much should I include? If I describe the matrices again, that will certainly have to go in the introduction.

We probably want a basic run-down of a .fits file, but perhaps some of that can go in the introduction.

## 3 Observations

What would be a good *introduction* to the data? We have hundreds of raw spectra. I am not sure how to *succinctly* describe observations before we get into the analysis.

## 4 Analysis

I should explain the process of taking a spectrum to a point on our galactic plane map (Doppler shift calculations, then orthographic projection).

We can estimate the mass of the super-massive black hole at the center of the milky way if we use Doppler velocities for our three closest angles ( $\ell \in \{-1, 0, 1\}$  degrees) and integrate over radii from 0 to 44 million kilometers.

## 5 Conclusions

We seem to have at least a couple of results! How do I estimate the uncertainty on the Leuschner dish?

## 6 Acknowledgments

Who did what?

Theory and background provided by Aaron Parsons. "LAB 4: Mapping the Galactic HI Line." Updated April 2020.