

# COL 352 Introduction to Automata and Theory of Computation

Nikhil Balaji

Bharti 420  
Indian Institute of Technology, Delhi  
[nbalaji@cse.iitd.ac.in](mailto:nbalaji@cse.iitd.ac.in)

February 1, 2023

Lecture 10: Pumping Lemma: Examples

# Pumping Lemma for regular languages

## Pumping Lemma

$L \subseteq \Sigma^*$  is a regular language  $\implies$

there exists  $n \geq 1$  such that

for all strings  $w \in L$  with  $|w| \geq n$  we have that

there exists  $x, y, z \in \Sigma^*$  with  $w = xyz$ ,  $|y| > 0$ ,  $|xy| \leq n$  such that

for all  $i \geq 0$  we have that

$xy^iz \in L$ .

# Pumping Lemma for regular languages

## Pumping Lemma

$L \subseteq \Sigma^*$  is a regular language  $\implies$

there exists  $n \geq 1$  such that

for all strings  $w \in L$  with  $|w| \geq n$  we have that

there exists  $x, y, z \in \Sigma^*$  with  $w = xyz$ ,  $|y| > 0$ ,  $|xy| \leq n$  such that

for all  $i \geq 0$  we have that

$xy^iz \in L$ .

## Contrapositive

If for all  $n \geq 1$  such that

there exists a string  $w \in L$  with  $|w| \geq n$  such that

for all breakups  $x, y, z \in \Sigma^*$  with  $w = xyz$ ,  $|y| > 0$ ,  $|xy| \leq n$  we have that

there exists  $i \geq 0$  such that

$xy^iz \in L$ .

$\implies L \subseteq \Sigma^*$  is not a regular language

# Applying Pumping Lemma: Example 1

Consider

$$L_{a,b} = \{a^n b^n \mid n \geq 0\}$$

# Applying Pumping Lemma: Example 1

Consider

$$L_{a,b} = \{a^n b^n \mid n \geq 0\}$$

- For each  $n$ , we need a word. Let  $w = a^n b^n$ .

# Applying Pumping Lemma: Example 1

Consider

$$L_{a,b} = \{a^n b^n \mid n \geq 0\}$$

- ▶ For each  $n$ , we need a word. Let  $w = a^n b^n$ .
- ▶ The first  $n$  characters of  $w$  are  $a^n$ . The breaks  $x$  and  $y$  are to be from within  $a^n$ .

# Applying Pumping Lemma: Example 1

Consider

$$L_{a,b} = \{a^n b^n \mid n \geq 0\}$$

- ▶ For each  $n$ , we need a word. Let  $w = a^n b^n$ .
- ▶ The first  $n$  characters of  $w$  are  $a^n$ . The breaks  $x$  and  $y$  are to be from within  $a^n$ .
- ▶ Let  $x = a^i, y = a^j$  where  $i + j \leq n$  and  $j \neq 0$ .

$$w = a^i a^j a^{n-i-j} b^n$$

# Applying Pumping Lemma: Example 1

Consider

$$L_{a,b} = \{a^n b^n \mid n \geq 0\}$$

- ▶ For each  $n$ , we need a word. Let  $w = a^n b^n$ .
- ▶ The first  $n$  characters of  $w$  are  $a^n$ . The breaks  $x$  and  $y$  are to be from within  $a^n$ .
- ▶ Let  $x = a^i, y = a^j$  where  $i + j \leq n$  and  $j \neq 0$ .

$$w = a^i a^j a^{n-i-j} b^n$$

- ▶ Choose  $k = 0$  for each  $i, j$ . The corresponding word is

$$a^i (a^j)^0 a^{n-i-j} b^n = a^{n-j} b^n \notin L_{a,b}$$



## Example 2

Consider

$$L = \{w \in \{a, b\}^* \mid w \text{ has equal number of a's and b's}\}$$

## Example 2

Consider

$$L = \{w \in \{a, b\}^* \mid w \text{ has equal number of a's and b's}\}$$

Exercise: What is  $L \cap a^*b^*$ ?

## Example 3

Consider

$$PAL = \{w \cdot w^R \mid w \in \{a, b\}^*\}$$

## Example 3

Consider

$$PAL = \{w \cdot w^R \mid w \in \{a, b\}^*\}$$

- ▶ For each  $n$ , let  $w = a^n b b a^n$ .

## Example 3

Consider

$$PAL = \{w \cdot w^R \mid w \in \{a, b\}^*\}$$

- ▶ For each  $n$ , let  $w = a^n b b a^n$ .
- ▶ The first  $n$  characters of  $w$  are  $a^n$

## Example 3

Consider

$$PAL = \{w \cdot w^R \mid w \in \{a, b\}^*\}$$

- ▶ For each  $n$ , let  $w = a^n b b a^n$ .
- ▶ The first  $n$  characters of  $w$  are  $a^n$
- ▶ Let  $x = a^i, y = a^j$  where  $i + j \leq n$  and  $j \neq 0$ .

$$w = a^i a^j a^{n-i-j} b b a^n$$

## Example 3

Consider

$$PAL = \{w \cdot w^R \mid w \in \{a, b\}^*\}$$

- ▶ For each  $n$ , let  $w = a^n b b a^n$ .
- ▶ The first  $n$  characters of  $w$  are  $a^n$
- ▶ Let  $x = a^i, y = a^j$  where  $i + j \leq n$  and  $j \neq 0$ .

$$w = a^i a^j a^{n-i-j} b b a^n$$

- ▶ Choose  $k = 2$  for each  $i, j$ . The corresponding word is

$$a^i (a^j)^2 a^{n-i-j} b b a^n = a^{n+j} b b a^n \notin PAL$$

# Food for thought

- ▶ If  $L_1 \cup L_2$  is regular, are  $L_1$  and  $L_2$  regular?
- ▶ Is this regular:  $\{a^n b^m \mid n \neq m\}$ ?
- ▶ Is this regular:  $\{a^n b^m \mid n \geq m\}$ ?
- ▶ Is this regular:  $\{a^n b^{n+1} \mid n \geq 0\}$ ?



# Need for infinite memory

Feels like all non-regular languages needed to remember infinite memory.

In  $\{a^n b^n \mid n \geq 0\}$  we need to remember the number of seen  $a$ 's and count the  $b$ 's to match.

Finite number of states cannot count unboundedly increasing number.

## Example 4: Primes

$$L_{prime} = \{1^p \in |p \text{ prime}\}$$

- Pick  $w = 1^p$  ( $p$  larger than pumping length).

## Example 4: Primes

$$L_{prime} = \{1^p \in |p \text{ prime}\}$$

- ▶ Pick  $w = 1^p$  ( $p$  larger than pumping length).
- ▶  $w = xyz$ ,  $y \neq \epsilon$ .
- ▶ For all  $k \geq 0$  Let  $x = 1^r, y = 1^s, z = 1^t$  where  $r + s + t = p$

## Example 4: Primes

$$L_{prime} = \{1^p \in |p \text{ prime}\}$$

- ▶ Pick  $w = 1^p$  ( $p$  larger than pumping length).
- ▶  $w = xyz$ ,  $y \neq \epsilon$ .
- ▶ For all  $k \geq 0$  Let  $x = 1^r, y = 1^s, z = 1^t$  where  $r + s + t = p$
- ▶ Consider  $xy^kz = 1^r 1^{ks} 1^t = 1^{p+(k-1)s}$
- ▶ We need  $p + (k-1)s$  to be prime for all  $k \geq 0$  to satisfy pumping condition.

## Example 4: Primes

$$L_{\text{prime}} = \{1^p \in |p \text{ prime}\}$$

- ▶ Pick  $w = 1^p$  ( $p$  larger than pumping length).
- ▶  $w = xyz$ ,  $y \neq \epsilon$ .
- ▶ For all  $k \geq 0$  Let  $x = 1^r, y = 1^s, z = 1^t$  where  $r + s + t = p$
- ▶ Consider  $xy^kz = 1^r 1^{ks} 1^t = 1^{p+(k-1)s}$
- ▶ We need  $p + (k-1)s$  to be prime for all  $k \geq 0$  to satisfy pumping condition.
- ▶ Choose  $k = p + 1$ , we have  $p + (k-1)s = p + ps = (s+1)p$ .

# Not a sufficient condition

Pumping lemma gives only a necessary condition for regularity, not a sufficient condition!

# Not a sufficient condition

Pumping lemma gives only a necessary condition for regularity, not a sufficient condition!

**Exercise:** Find a language that is not regular, but which satisfies the pumping conditions.

## Example 5

$$L = \{a^n b^m \mid n > m\}$$



## Example 5

$$L = \{a^n b^m \mid n > m\}$$

- For each  $n$ , let  $w = a^n b^n$

## Example 5

$$L = \{a^n b^m \mid n > m\}$$

- ▶ For each  $n$ , let  $w = a^n b^n$
- ▶ Any break-up of  $w$  has  $x = a^i$  and  $y = a^j$ , where  $i + j \leq n$  and  $j \neq 0$

$$\overbrace{a^i}^x \overbrace{a^j}^y \overbrace{a^{n-j-i} b^n}^z$$

## Example 5

$$L = \{a^n b^m \mid n > m\}$$

- ▶ For each  $n$ , let  $w = a^n b^n$
- ▶ Any break-up of  $w$  has  $x = a^i$  and  $y = a^j$ , where  $i + j \leq n$  and  $j \neq 0$

$$\overbrace{a^i}^x \overbrace{a^j}^y \overbrace{a^{n-j-i} b^n}^z$$

- ▶ if  $k > 2$ , we get  $xy^k z = a^i a^{2j} a^{n-j-i} b^n \in L$ , so NOT enough!

## Example 5

$$L = \{a^n b^m \mid n > m\}$$

- ▶ For each  $n$ , let  $w = a^n b^n$
- ▶ Any break-up of  $w$  has  $x = a^i$  and  $y = a^j$ , where  $i + j \leq n$  and  $j \neq 0$

$$\overbrace{a^i}^x \overbrace{a^j}^y \overbrace{a^{n-j-i} b^n}^z$$

- ▶ if  $k > 2$ , we get  $xy^k z = a^i a^{2j} a^{n-j-i} b^n \in L$ , so NOT enough!
- ▶ However, let  $k = 0$  (pumping down).

## Example 5

$$L = \{a^n b^m \mid n > m\}$$

- ▶ For each  $n$ , let  $w = a^n b^n$
- ▶ Any break-up of  $w$  has  $x = a^i$  and  $y = a^j$ , where  $i + j \leq n$  and  $j \neq 0$

$$\underbrace{a^i}_x \underbrace{a^j}_y \underbrace{a^{n-j-i} b^n}_z$$

- ▶ if  $k > 2$ , we get  $xy^k z = a^i a^{2j} a^{n-j-i} b^n \in L$ , so NOT enough!
- ▶ However, let  $k = 0$  (pumping down). Then,  $xy^k z = a^i a^{n-j-i} b^n \notin L$ .

# Homomorphisms: A tool to show regularity

# Homomorphisms: A tool to show regularity

## Definition

A homomorphism is a map  $h : \Sigma^* \rightarrow \Gamma^*$  such that for all  $x, y \in \Sigma^*$

# Homomorphisms: A tool to show regularity

## Definition

A homomorphism is a map  $h : \Sigma^* \rightarrow \Gamma^*$  such that for all  $x, y \in \Sigma^*$

$$h(xy) = h(x)h(y)$$



# Homomorphisms: A tool to show regularity

## Definition

A homomorphism is a map  $h : \Sigma^* \rightarrow \Gamma^*$  such that for all  $x, y \in \Sigma^*$

$$h(xy) = h(x)h(y)$$

$$h(\varepsilon) = \varepsilon$$

# Homomorphisms: A tool to show regularity

## Definition

A homomorphism is a map  $h : \Sigma^* \rightarrow \Gamma^*$  such that for all  $x, y \in \Sigma^*$

$$h(xy) = h(x)h(y)$$

$$h(\varepsilon) = \varepsilon$$

We can derive the second condition from the first!

# Homomorphisms: A tool to show regularity

## Definition

A homomorphism is a map  $h : \Sigma^* \rightarrow \Gamma^*$  such that for all  $x, y \in \Sigma^*$

$$h(xy) = h(x)h(y)$$

$$h(\varepsilon) = \varepsilon$$

We can derive the second condition from the first!

$$|h(\varepsilon)| = |h(\varepsilon\varepsilon)|$$

# Homomorphisms: A tool to show regularity

## Definition

A homomorphism is a map  $h : \Sigma^* \rightarrow \Gamma^*$  such that for all  $x, y \in \Sigma^*$

$$h(xy) = h(x)h(y)$$

$$h(\varepsilon) = \varepsilon$$

We can derive the second condition from the first!

$$\begin{aligned} |h(\varepsilon)| &= |h(\varepsilon\varepsilon)| \\ &= |h(\varepsilon)h(\varepsilon)| \end{aligned}$$

# Homomorphisms: A tool to show regularity

## Definition

A homomorphism is a map  $h : \Sigma^* \rightarrow \Gamma^*$  such that for all  $x, y \in \Sigma^*$

$$h(xy) = h(x)h(y)$$

$$h(\varepsilon) = \varepsilon$$

We can derive the second condition from the first!

$$\begin{aligned} |h(\varepsilon)| &= |h(\varepsilon\varepsilon)| \\ &= |h(\varepsilon)h(\varepsilon)| \\ &= |h(\varepsilon)| + |h(\varepsilon)| \end{aligned}$$

# Homomorphisms: A tool to show regularity

## Definition

A homomorphism is a map  $h : \Sigma^* \rightarrow \Gamma^*$  such that for all  $x, y \in \Sigma^*$

$$h(xy) = h(x)h(y)$$

$$h(\varepsilon) = \varepsilon$$

We can derive the second condition from the first!

$$\begin{aligned} |h(\varepsilon)| &= |h(\varepsilon\varepsilon)| \\ &= |h(\varepsilon)h(\varepsilon)| \\ &= |h(\varepsilon)| + |h(\varepsilon)| \\ \implies |h(\varepsilon)| &= 0 \end{aligned}$$

# Homomorphisms: A tool to show regularity

## Definition

A homomorphism is a map  $h : \Sigma^* \rightarrow \Gamma^*$  such that for all  $x, y \in \Sigma^*$

$$h(xy) = h(x)h(y)$$

$$h(\varepsilon) = \varepsilon$$

We can derive the second condition from the first!

$$\begin{aligned} |h(\varepsilon)| &= |h(\varepsilon\varepsilon)| \\ &= |h(\varepsilon)h(\varepsilon)| \\ &= |h(\varepsilon)| + |h(\varepsilon)| \\ \implies |h(\varepsilon)| &= 0 \\ \implies h(\varepsilon) &= \varepsilon \end{aligned}$$

# Homomorphisms : Examples

- ▶ Given  $\forall a \in \Sigma, h(a)$ , we can uniquely determine  $h(w)$  for every  $w \in \Sigma^*$
- ▶ Any map  $h : \Sigma \rightarrow \Gamma^*$  extends by induction uniquely to a map  $h' : \Sigma^* \rightarrow \Gamma^*$



# Homomorphisms : Examples

- ▶ Given  $\forall a \in \Sigma, h(a)$ , we can uniquely determine  $h(w)$  for every  $w \in \Sigma^*$
- ▶ Any map  $h : \Sigma \rightarrow \Gamma^*$  extends by induction uniquely to a map  $h' : \Sigma^* \rightarrow \Gamma^*$
- ▶ For  $A \subseteq \Sigma^*$

$$h(A) = \{h(x) \mid x \in A\} \subseteq \Gamma^*$$

# Homomorphisms : Examples

- ▶ Given  $\forall a \in \Sigma, h(a)$ , we can uniquely determine  $h(w)$  for every  $w \in \Sigma^*$
- ▶ Any map  $h : \Sigma \rightarrow \Gamma^*$  extends by induction uniquely to a map  $h' : \Sigma^* \rightarrow \Gamma^*$

- ▶ For  $A \subseteq \Sigma^*$

$$h(A) = \{h(x) \mid x \in A\} \subseteq \Gamma^*$$

- ▶ Problem set 1: If  $A$  is regular then so is  $h(A)$  (image of  $A$ ).

# Homomorphisms : Examples

- ▶ Given  $\forall a \in \Sigma, h(a)$ , we can uniquely determine  $h(w)$  for every  $w \in \Sigma^*$
- ▶ Any map  $h : \Sigma \rightarrow \Gamma^*$  extends by induction uniquely to a map  $h' : \Sigma^* \rightarrow \Gamma^*$

- ▶ For  $A \subseteq \Sigma^*$

$$h(A) = \{h(x) \mid x \in A\} \subseteq \Gamma^*$$

- ▶ Problem set 1: If  $A$  is regular then so is  $h(A)$  (image of  $A$ ).
- ▶ For  $B \subseteq \Gamma^*$ ,

$$h^{-1}(B) := \{x \in \Sigma^* \mid h(x) \in B\}$$

- ▶ Prove: If  $B$  is regular then so is  $h^{-1}(B)$ .

# Homomorphisms : Examples

- ▶ Given  $\forall a \in \Sigma, h(a)$ , we can uniquely determine  $h(w)$  for every  $w \in \Sigma^*$
- ▶ Any map  $h : \Sigma \rightarrow \Gamma^*$  extends by induction uniquely to a map  $h' : \Sigma^* \rightarrow \Gamma^*$

- ▶ For  $A \subseteq \Sigma^*$

$$h(A) = \{h(x) \mid x \in A\} \subseteq \Gamma^*$$

- ▶ Problem set 1: If  $A$  is regular then so is  $h(A)$  (image of  $A$ ).
- ▶ For  $B \subseteq \Gamma^*$ ,

$$h^{-1}(B) := \{x \in \Sigma^* \mid h(x) \in B\}$$

- ▶ Prove: If  $B$  is regular then so is  $h^{-1}(B)$ .
- ▶ **Caveat:** This **does not** say that  $A$  is regular if  $h(A)$  is. Example:

# Homomorphisms : Examples

- ▶ Given  $\forall a \in \Sigma, h(a)$ , we can uniquely determine  $h(w)$  for every  $w \in \Sigma^*$
- ▶ Any map  $h : \Sigma \rightarrow \Gamma^*$  extends by induction uniquely to a map  $h' : \Sigma^* \rightarrow \Gamma^*$

- ▶ For  $A \subseteq \Sigma^*$

$$h(A) = \{h(x) \mid x \in A\} \subseteq \Gamma^*$$

- ▶ Problem set 1: If  $A$  is regular then so is  $h(A)$  (image of  $A$ ).
- ▶ For  $B \subseteq \Gamma^*$ ,

$$h^{-1}(B) := \{x \in \Sigma^* \mid h(x) \in B\}$$

- ▶ Prove: If  $B$  is regular then so is  $h^{-1}(B)$ .
- ▶ **Caveat:** This **does not** say that  $A$  is regular if  $h(A)$  is. Example:

$$L_{a,b} = \{a^n b^n \mid n \in \mathbb{N}\}$$

# Homomorphisms : Examples

- ▶ Given  $\forall a \in \Sigma, h(a)$ , we can uniquely determine  $h(w)$  for every  $w \in \Sigma^*$
- ▶ Any map  $h : \Sigma \rightarrow \Gamma^*$  extends by induction uniquely to a map  $h' : \Sigma^* \rightarrow \Gamma^*$

- ▶ For  $A \subseteq \Sigma^*$

$$h(A) = \{h(x) \mid x \in A\} \subseteq \Gamma^*$$

- ▶ Problem set 1: If  $A$  is regular then so is  $h(A)$  (image of  $A$ ).
- ▶ For  $B \subseteq \Gamma^*$ ,

$$h^{-1}(B) := \{x \in \Sigma^* \mid h(x) \in B\}$$

- ▶ Prove: If  $B$  is regular then so is  $h^{-1}(B)$ .
- ▶ **Caveat:** This **does not** say that  $A$  is regular if  $h(A)$  is. Example:

$$L_{a,b} = \{a^n b^n \mid n \in \mathbb{N}\}$$

- ▶ What is  $h(L_{a,b})$  under  $h : \Sigma \rightarrow \Sigma$  given by  $h(a) = h(b) = a$ ?

# More non-regularity tricks

$$L = \{a^n b^m \mid n \geq m\}$$

## More non-regularity tricks

$$L = \{a^n b^m \mid n \geq m\}$$

Consider

$$L^R = \{b^m a^n \mid n \geq m\}$$

- ▶ If  $L$  was regular so is  $L^R$  (reverse of  $L$ ) - Prove!



# More non-regularity tricks

$$L = \{a^n b^m \mid n \geq m\}$$

Consider

$$L^R = \{b^m a^n \mid n \geq m\}$$

- ▶ If  $L$  was regular so is  $L^R$  (reverse of  $L$ ) - Prove!
- ▶ Apply  $h : \Sigma \rightarrow \Sigma$  ( $h(a) = b, h(b) = a$ ).

# More non-regularity tricks

$$L = \{a^n b^m \mid n \geq m\}$$

Consider

$$L^R = \{b^m a^n \mid n \geq m\}$$

- ▶ If  $L$  was regular so is  $L^R$  (reverse of  $L$ ) - Prove!
- ▶ Apply  $h : \Sigma \rightarrow \Sigma$  ( $h(a) = b, h(b) = a$ ).

$$L' = h(L^R) = \{a^m b^n \mid n \geq m\}$$

# More non-regularity tricks

$$L = \{a^n b^m \mid n \geq m\}$$

Consider

$$L^R = \{b^m a^n \mid n \geq m\}$$

- ▶ If  $L$  was regular so is  $L^R$  (reverse of  $L$ ) - Prove!
- ▶ Apply  $h : \Sigma \rightarrow \Sigma$  ( $h(a) = b, h(b) = a$ ).

$$L' = h(L^R) = \{a^m b^n \mid n \geq m\}$$

- ▶ What is  $L \cap L'$ ?

# More non-regularity tricks

$$L = \{a^n b^m \mid n \geq m\}$$

Consider

$$L^R = \{b^m a^n \mid n \geq m\}$$

- ▶ If  $L$  was regular so is  $L^R$  (reverse of  $L$ ) - Prove!
- ▶ Apply  $h : \Sigma \rightarrow \Sigma$  ( $h(a) = b, h(b) = a$ ).

$$L' = h(L^R) = \{a^m b^n \mid n \geq m\}$$

- ▶ What is  $L \cap L'$ ?

$$L \cap L' = \{a^n b^n \mid n \in \mathbb{N}\}$$

# Converse of Pumping Lemma

# Converse of Pumping Lemma

## Example

Consider

$$L = \overbrace{\{ca^n b^n\}}^{L_1} \cup \overbrace{\{c^k w \mid k \neq 1, w \text{ starts with } a \text{ or } b\}}^{L_2}$$

# Converse of Pumping Lemma

## Example

Consider

$$L = \overbrace{\{ca^n b^n\}}^{L_1} \cup \overbrace{\{c^k w \mid k \neq 1, w \text{ starts with } a \text{ or } b\}}^{L_2}$$

- ▶  $L_2$  is regular (why?)

# Converse of Pumping Lemma

## Example

Consider

$$L = \overbrace{\{ca^n b^n\}}^{L_1} \cup \overbrace{\{c^k w \mid k \neq 1, w \text{ starts with } a \text{ or } b\}}^{L_2}$$

- ▶  $L_2$  is regular (why?)
- ▶  $L_1$  is not regular (why??)



# Converse of Pumping Lemma

## Example

Consider

$$L = \overbrace{\{ca^n b^n\}}^{L_1} \cup \overbrace{\{c^k w \mid k \neq 1, w \text{ starts with } a \text{ or } b\}}^{L_2}$$

- ▶  $L_2$  is regular (why?)
- ▶  $L_1$  is not regular (why??)