

# COL 352 Introduction to Automata and Theory of Computation

Nikhil Balaji

Bharti 420  
Indian Institute of Technology, Delhi  
[nbalaji@cse.iitd.ac.in](mailto:nbalaji@cse.iitd.ac.in)

January 19, 2023

## Lecture 7: Pattern Matching and Regular Expressions

# Recap

- ▶  $\text{DFA} = \text{NFA} = \varepsilon\text{-NFA}$ .
- ▶ All of them recognize (compute/decide) exactly regular languages
- ▶ Regular languages are closed under union, intersection, complement, concatenation, Kleene star, ...

A number of widely different and equiexpressive formalisms precisely capture the same class of languages:

# Why regular?

- ▶ DFA = NFA =  $\varepsilon$ -NFA.
- ▶ All of them recognize (compute/decide) exactly regular languages
- ▶ Regular languages are closed under union, intersection, complement, concatenation, Kleene star, ...

A number of widely different and equiexpressive formalisms precisely capture the same class of languages:

- ▶ Type-3 languages in Chomsky's hierarchy.

# Why regular?

- ▶ DFA = NFA =  $\varepsilon$ -NFA.
- ▶ All of them recognize (compute/decide) exactly regular languages
- ▶ Regular languages are closed under union, intersection, complement, concatenation, Kleene star, ...

A number of widely different and equiexpressive formalisms precisely capture the same class of languages:

- ▶ Type-3 languages in Chomsky's hierarchy.
- ▶ Monadic second-order logic definable languages (Buchi '60, Elgot '61, Trackhtenbrot'62)

# Why regular?

- ▶ DFA = NFA =  $\varepsilon$ -NFA.
- ▶ All of them recognize (compute/decide) exactly regular languages
- ▶ Regular languages are closed under union, intersection, complement, concatenation, Kleene star, ...

A number of widely different and equiexpressive formalisms precisely capture the same class of languages:

- ▶ Type-3 languages in Chomsky's hierarchy.
- ▶ Monadic second-order logic definable languages (Buchi '60, Elgot '61, Trackhtenbrot'62)
- ▶ Certain Algebraic connection (acceptability via finite semi-group): Eilenberg'76

# Why regular?

- ▶ DFA = NFA =  $\varepsilon$ -NFA.
- ▶ All of them recognize (compute/decide) exactly regular languages
- ▶ Regular languages are closed under union, intersection, complement, concatenation, Kleene star, ...

A number of widely different and equiexpressive formalisms precisely capture the same class of languages:

- ▶ Type-3 languages in Chomsky's hierarchy.
- ▶ Monadic second-order logic definable languages (Buchi '60, Elgot '61, Trackhtenbrot'62)
- ▶ Certain Algebraic connection (acceptability via finite semi-group): Eilenberg'76
- ▶ Regular expressions (Kleene'50s).
- ▶ Rational languages.

# Algebraic view of regular languages

Consider any regular language

# Algebraic view of regular languages

Consider any regular language

- ▶ It can be letters 1 or a or  $\epsilon$  etc.
- ▶ It can be emptyset.
- ▶ It can be a word, e.g., 110: got as (finite) concatenation of letters.
- ▶ It can be several words: got as (finite) union of sets of words.



# Algebraic view of regular languages

Consider any regular language

- ▶ It can be letters 1 or a or  $\epsilon$  etc.
- ▶ It can be emptyset.
- ▶ It can be a word, e.g., 110: got as (finite) concatenation of letters.
- ▶ It can be several words: got as (finite) union of sets of words. Can these be used to generate all regular languages?
- ▶ It can be something that "repeats"!

# Algebraic view of regular languages

Consider any regular language

- ▶ It can be letters 1 or a or  $\epsilon$  etc.
- ▶ It can be emptyset.
- ▶ It can be a word, e.g., 110: got as (finite) concatenation of letters.
- ▶ It can be several words: got as (finite) union of sets of words. Can these be used to generate all regular languages?
- ▶ It can be something that "repeats"!

## Kleene star

For a language  $L$ , its Kleene closure, denoted  $L^*$  is the set of all strings obtained by taking any number of strings from  $L$  with possible repetitions and concatenating all of them.

$$L^* = \cup_{i \geq 0} L^i$$

$$L^0 = \{\epsilon\}, L^i = L \circ L^{i-1}$$

# Application: Pattern Matching

# Application: Pattern Matching

- ▶ `rm *.pdf`
- ▶ `ls`, `grep`, `awk` etc.

# Application: Pattern Matching

- ▶ `rm *.pdf`
- ▶ `ls`, `grep`, `awk` etc.
- ▶ Textual (declarative) way to represent regular languages (compared to automata).

# Application: Pattern Matching

- ▶ `rm *.pdf`
- ▶ `ls`, `grep`, `awk` etc.
- ▶ Textual (declarative) way to represent regular languages (compared to automata).

# Application: Pattern Matching

- ▶ `rm *.pdf`
- ▶ `ls`, `grep`, `awk` etc.
- ▶ Textual (declarative) way to represent regular languages (compared to automata).

## Definition (Pattern)

A pattern  $\alpha$  is a string of symbols of a certain form representing a (possibly infinite) set of strings in  $\Sigma^*$ .

# Application: Pattern Matching

- ▶ `rm *.pdf`
- ▶ `ls`, `grep`, `awk` etc.
- ▶ Textual (declarative) way to represent regular languages (compared to automata).

## Definition (Pattern)

A pattern  $\alpha$  is a string of symbols of a certain form representing a (possibly infinite) set of strings in  $\Sigma^*$ .

$$L(\alpha) = \{x \in \Sigma^* \mid x \text{ matches } \alpha\}$$



# Atomic Patterns

1  $a \in \Sigma, L(a) = \{a\}$

# Atomic Patterns

- 1  $a \in \Sigma, L(a) = \{a\}$
- 2  $\varepsilon, L(\varepsilon) = \{\varepsilon\}$
- 3  $\emptyset, L(\emptyset) = \emptyset$
- 4  $\Sigma$ , matching any alphabet
- 5  $\Sigma^*$ , matching any finite string

# Compound Patterns

- 1  $a \in \Sigma, L(a) = \{a\}$
- 2  $\varepsilon, L(\varepsilon) = \{\varepsilon\}$
- 3  $\emptyset, L(\emptyset) = \emptyset$
- 4  $\Sigma$ , matching any alphabet
- 5  $\Sigma^*$ , matching any finite string
- 6  $x$  matches  $\alpha + \beta$  if  $L(\alpha + \beta) = L(\alpha) \cup L(\beta)$
- 7  $x$  matches  $\alpha \cap \beta$  if  $L(\alpha \cap \beta) = L(\alpha) \cap L(\beta)$
- 8  $x$  matches  $\alpha\beta$  if  $x = yz$  where  $L(\alpha\beta) = L(\alpha)L(\beta)$
- 9  $x$  matches  $\overline{\alpha}$  if  $L(\overline{\alpha}) = \overline{L(\alpha)} = \Sigma^* \setminus L(\alpha)$

# Compound Patterns

- ❶  $a \in \Sigma, L(a) = \{a\}$
- ❷  $\varepsilon, L(\varepsilon) = \{\varepsilon\}$
- ❸  $\emptyset, L(\emptyset) = \emptyset$
- ❹  $\Sigma$ , matching any alphabet
- ❺  $\Sigma^*$ , matching any finite string
- ❻  $x$  matches  $\alpha + \beta$  if  $L(\alpha + \beta) = L(\alpha) \cup L(\beta)$
- ❼  $x$  matches  $\alpha \cap \beta$  if  $L(\alpha \cap \beta) = L(\alpha) \cap L(\beta)$
- ❽  $x$  matches  $\alpha\beta$  if  $x = yz$  where  $L(\alpha\beta) = L(\alpha)L(\beta)$
- ❾  $x$  matches  $\overline{\alpha}$  if  $L(\overline{\alpha}) = \overline{L(\alpha)} = \Sigma^* \setminus L(\alpha)$
- ❿  $x$  matches  $\alpha^*$  if  $x$  can be expressed as zero or more of strings that match  $\alpha$ , i.e.,  $L(\alpha^*) = L(\alpha)^*$
- ⓫  $x$  matches  $\alpha^+$  if  $x$  can be expressed as one or more of strings that match  $\alpha$ , i.e.,  $L(\alpha^+) = L(\alpha)^+$

# Examples of Patterns

Given patterns for the following languages

# Examples of Patterns

Given patterns for the following languages

- ▶ All words containing at least one  $a$ .

# Examples of Patterns

Given patterns for the following languages

- ▶ All words containing at least one  $a$ .  $\Sigma^* a \Sigma^*$

# Examples of Patterns

Given patterns for the following languages

- ▶ All words containing at least one  $a$ .  $\Sigma^* a \Sigma^*$
- ▶ Words containing at least one  $a$  or one  $b$ .
- ▶ Words containing at least one  $a$  and one  $b$ .



# Examples of Patterns

Given patterns for the following languages

- ▶ All words containing at least one  $a$ .  $\Sigma^* a \Sigma^*$
- ▶ Words containing at least one  $a$  or one  $b$ .
- ▶ Words containing at least one  $a$  and one  $b$ .
- ▶ Words of even length.

# Examples of Patterns

Given patterns for the following languages

- ▶ All words containing at least one  $a$ .  $\Sigma^*a\Sigma^*$
- ▶ Words containing at least one  $a$  or one  $b$ .
- ▶ Words containing at least one  $a$  and one  $b$ .
- ▶ Words of even length.  $(\Sigma\Sigma)^*$
- ▶ Strings containing the substring  $abaab$ ?
- ▶ Strings containing no occurrence of  $a$

# Examples of Patterns

Given patterns for the following languages

- ▶ All words containing at least one  $a$ .  $\Sigma^* a \Sigma^*$
- ▶ Words containing at least one  $a$  or one  $b$ .
- ▶ Words containing at least one  $a$  and one  $b$ .
- ▶ Words of even length.  $(\Sigma\Sigma)^*$
- ▶ Strings containing the substring  $abaab$ ?
- ▶ Strings containing no occurrence of  $a$   $(\Sigma \cap \bar{a})^*$
- ▶ Strings where every occurrence of  $a$  is followed by an occurrence of  $b$ .

# Examples of Patterns

Given patterns for the following languages

- ▶ All words containing at least one  $a$ .  $\Sigma^* a \Sigma^*$
- ▶ Words containing at least one  $a$  or one  $b$ .
- ▶ Words containing at least one  $a$  and one  $b$ .
- ▶ Words of even length.  $(\Sigma\Sigma)^*$
- ▶ Strings containing the substring  $abaab$ ?
- ▶ Strings containing no occurrence of  $a$   $(\Sigma \cap \bar{a})^*$
- ▶ Strings where every occurrence of  $a$  is followed by an occurrence of  $b$ .

Express each of these languages as a pattern.

# Questions

- ▶ How hard is to determine whether a given string  $x$  matches a pattern?
- ▶ Is every set of strings represented by some pattern?

# Questions

- ▶ How hard is to determine whether a given string  $x$  matches a pattern?
- ▶ Is every set of strings represented by some pattern?
- ▶ Patterns  $\alpha \equiv \beta$  if  $L(\alpha) = L(\beta)$ . Can you tell whether two given patterns are equivalent?

# Questions

- ▶ How hard is to determine whether a given string  $x$  matches a pattern?
- ▶ Is every set of strings represented by some pattern?
- ▶ Patterns  $\alpha \equiv \beta$  if  $L(\alpha) = L(\beta)$ . Can you tell whether two given patterns are equivalent?
- ▶ Which of the operators are redundant?

# Questions

- ▶ How hard is to determine whether a given string  $x$  matches a pattern?
- ▶ Is every set of strings represented by some pattern?
- ▶ Patterns  $\alpha \equiv \beta$  if  $L(\alpha) = L(\beta)$ . Can you tell whether two given patterns are equivalent?
- ▶ Which of the operators are redundant?
  - ▶  $\varepsilon$



# Questions

- ▶ How hard is to determine whether a given string  $x$  matches a pattern?
- ▶ Is every set of strings represented by some pattern?
- ▶ Patterns  $\alpha \equiv \beta$  if  $L(\alpha) = L(\beta)$ . Can you tell whether two given patterns are equivalent?
- ▶ Which of the operators are redundant?
  - ▶  $\varepsilon = \overline{\Sigma\Sigma^*}$

# Questions

- ▶ How hard is to determine whether a given string  $x$  matches a pattern?
- ▶ Is every set of strings represented by some pattern?
- ▶ Patterns  $\alpha \equiv \beta$  if  $L(\alpha) = L(\beta)$ . Can you tell whether two given patterns are equivalent?
- ▶ Which of the operators are redundant?
  - ▶  $\varepsilon = \overline{\Sigma\Sigma^*}$
  - ▶  $\Sigma^*$

# Questions

- ▶ How hard is to determine whether a given string  $x$  matches a pattern?
- ▶ Is every set of strings represented by some pattern?
- ▶ Patterns  $\alpha \equiv \beta$  if  $L(\alpha) = L(\beta)$ . Can you tell whether two given patterns are equivalent?
- ▶ Which of the operators are redundant?
  - ▶  $\varepsilon = \overline{\Sigma\Sigma^*}$
  - ▶  $\Sigma^* = (\Sigma)^*$

# Questions

- ▶ How hard is to determine whether a given string  $x$  matches a pattern?
- ▶ Is every set of strings represented by some pattern?
- ▶ Patterns  $\alpha \equiv \beta$  if  $L(\alpha) = L(\beta)$ . Can you tell whether two given patterns are equivalent?
- ▶ Which of the operators are redundant?
  - ▶  $\varepsilon = \overline{\Sigma\Sigma^*}$
  - ▶  $\Sigma^* = (\Sigma)^*$
  - ▶  $\Sigma$

# Questions

- ▶ How hard is to determine whether a given string  $x$  matches a pattern?
- ▶ Is every set of strings represented by some pattern?
- ▶ Patterns  $\alpha \equiv \beta$  if  $L(\alpha) = L(\beta)$ . Can you tell whether two given patterns are equivalent?
- ▶ Which of the operators are redundant?
  - ▶  $\varepsilon = \overline{\Sigma\Sigma^*}$
  - ▶  $\Sigma^* = (\Sigma)^*$
  - ▶  $\Sigma = a_1 + \cdots + a_k$

# Questions

- ▶ How hard is to determine whether a given string  $x$  matches a pattern?
- ▶ Is every set of strings represented by some pattern?
- ▶ Patterns  $\alpha \equiv \beta$  if  $L(\alpha) = L(\beta)$ . Can you tell whether two given patterns are equivalent?
- ▶ Which of the operators are redundant?
  - ▶  $\varepsilon = \overline{\Sigma\Sigma^*}$
  - ▶  $\Sigma^* = (\Sigma)^*$
  - ▶  $\Sigma = a_1 + \cdots + a_k$
  - ▶  $\alpha \cap \beta$

# Questions

- ▶ How hard is to determine whether a given string  $x$  matches a pattern?
- ▶ Is every set of strings represented by some pattern?
- ▶ Patterns  $\alpha \equiv \beta$  if  $L(\alpha) = L(\beta)$ . Can you tell whether two given patterns are equivalent?
- ▶ Which of the operators are redundant?
  - ▶  $\varepsilon = \overline{\Sigma \Sigma^*}$
  - ▶  $\Sigma^* = (\Sigma)^*$
  - ▶  $\Sigma = a_1 + \dots + a_k$
  - ▶  $\alpha \cap \beta = \overline{\overline{\alpha} + \overline{\beta}}$

# Questions

- ▶ How hard is to determine whether a given string  $x$  matches a pattern?
- ▶ Is every set of strings represented by some pattern?
- ▶ Patterns  $\alpha \equiv \beta$  if  $L(\alpha) = L(\beta)$ . Can you tell whether two given patterns are equivalent?
- ▶ Which of the operators are redundant?
  - ▶  $\varepsilon = \overline{\Sigma \Sigma^*}$
  - ▶  $\Sigma^* = (\Sigma)^*$
  - ▶  $\Sigma = a_1 + \cdots + a_k$
  - ▶  $\alpha \cap \beta = \overline{\overline{\alpha} + \overline{\beta}}$
- ▶ Removing redundancy: Useful theoretically at the expense of losing succinctness.



# Questions

- ▶ How hard is to determine whether a given string  $x$  matches a pattern?
- ▶ Is every set of strings represented by some pattern?
- ▶ Patterns  $\alpha \equiv \beta$  if  $L(\alpha) = L(\beta)$ . Can you tell whether two given patterns are equivalent?
- ▶ Which of the operators are redundant?
  - ▶  $\varepsilon = \overline{\Sigma\Sigma^*}$
  - ▶  $\Sigma^* = (\Sigma)^*$
  - ▶  $\Sigma = a_1 + \cdots + a_k$
  - ▶  $\alpha \cap \beta = \overline{\overline{\alpha} + \overline{\beta}}$
- ▶ Removing redundancy: Useful theoretically at the expense of losing succinctness.
- ▶ Can you get rid of complementation?

# Regular expressions

For a regular expression  $E$  we write  $L(E)$  for its language. The set of valid regular expressions  $RegEx$  can be defined recursively as the following:

	Syntax	Semantics
Empty String	$\epsilon$	$L(\epsilon) = \{\epsilon\}$
Empty Set	$\emptyset$	$L(\emptyset) = \emptyset$
Single Letter	$a$	$L(a) = \{a\}$
Union	$E + F$	$L(E + F) = L(E) \cup L(F)$
Concatenation	$E.F$	$L(E.F) = L(E) \circ L(F)$
Kleene Star	$E^*$	$L(E)^*$

# Regular expressions

For a regular expression  $E$  we write  $L(E)$  for its language. The set of valid regular expressions  $RegEx$  can be defined recursively as the following:

	Syntax	Semantics
Empty String	$\epsilon$	$L(\epsilon) = \{\epsilon\}$
Empty Set	$\emptyset$	$L(\emptyset) = \emptyset$
Single Letter	$a$	$L(a) = \{a\}$
Union	$E + F$	$L(E + F) = L(E) \cup L(F)$
Concatenation	$E.F$	$L(E.F) = L(E) \circ L(F)$
Kleene Star	$E^*$	$L(E)^*$

## Associativity of $+$ and $\circ$ :

- ▶  $L(\alpha + (\beta + \gamma)) = L((\alpha + \beta) + \gamma)$
- ▶  $L(\alpha(\beta\gamma)) = L((\alpha\beta)\gamma)$

# Regular expressions

For a regular expression  $E$  we write  $L(E)$  for its language. The set of valid regular expressions  $RegEx$  can be defined recursively as the following:

	Syntax	Semantics
Empty String	$\epsilon$	$L(\epsilon) = \{\epsilon\}$
Empty Set	$\emptyset$	$L(\emptyset) = \emptyset$
Single Letter	$a$	$L(a) = \{a\}$
Union	$E + F$	$L(E + F) = L(E) \cup L(F)$
Concatenation	$E.F$	$L(E.F) = L(E) \circ L(F)$
Kleene Star	$E^*$	$L(E)^*$

## Associativity of $+$ and $\circ$ :

- ▶  $L(\alpha + (\beta + \gamma)) = L((\alpha + \beta) + \gamma)$
- ▶  $L(\alpha(\beta\gamma)) = L((\alpha\beta)\gamma)$
- ▶  $\alpha + (\beta\gamma) \neq (\alpha + \beta)\gamma,$

# Regular expressions

For a regular expression  $E$  we write  $L(E)$  for its language. The set of valid regular expressions  $RegEx$  can be defined recursively as the following:

	Syntax	Semantics
Empty String	$\epsilon$	$L(\epsilon) = \{\epsilon\}$
Empty Set	$\emptyset$	$L(\emptyset) = \emptyset$
Single Letter	$a$	$L(a) = \{a\}$
Union	$E + F$	$L(E + F) = L(E) \cup L(F)$
Concatenation	$E.F$	$L(E.F) = L(E) \circ L(F)$
Kleene Star	$E^*$	$L(E)^*$

## Associativity of $+$ and $\circ$ :

- ▶  $L(\alpha + (\beta + \gamma)) = L((\alpha + \beta) + \gamma)$
- ▶  $L(\alpha(\beta\gamma)) = L((\alpha\beta)\gamma)$
- ▶  $\alpha + (\beta\gamma) \neq (\alpha + \beta)\gamma, \alpha + \beta^* \neq (\alpha + \beta)^*$

# Regular expressions

For a regular expression  $E$  we write  $L(E)$  for its language. The set of valid regular expressions *RegEx* can be defined recursively as the following:

	Syntax	Semantics
Empty String	$\epsilon$	$L(\epsilon) = \{\epsilon\}$
Empty Set	$\emptyset$	$L(\emptyset) = \emptyset$
Single Letter	$a$	$L(a) = \{a\}$
Union	$E + F$	$L(E + F) = L(E) \cup L(F)$
Concatenation	$E.F$	$L(E.F) = L(E) \circ L(F)$
Kleene Star	$E^*$	$L(E)^*$

## Associativity of $+$ and $\circ$ :

- ▶  $L(\alpha + (\beta + \gamma)) = L((\alpha + \beta) + \gamma)$
- ▶  $L(\alpha(\beta\gamma)) = L((\alpha\beta)\gamma)$
- ▶  $\alpha + (\beta\gamma) \neq (\alpha + \beta)\gamma, \alpha + \beta^* \neq (\alpha + \beta)^*$

## Precedence rules: $*$ $>$ $\circ$ $>$ $+$

# Language defined by regular expression

## Lemma

The language defined by any regular expression is regular.

## Example

$$(a + b)^*$$





# Language defined by regular expression

## Lemma

The language defined by any regular expression is regular.

## Example

$$(a + b)^*$$








# Language defined by regular expression

## Lemma

The language defined by any regular expression is regular.

## Example

$$(a + b)^*$$







# Language defined by regular expression

## Lemma

The language defined by any regular expression is regular.

## Example

$$(a + b)^*$$



# Language defined by regular expression

## Lemma

The language defined by any regular expression is regular.

# Language defined by regular expression

## Lemma

The language defined by any regular expression is regular.

Proof idea

# Language defined by regular expression

## Lemma

The language defined by any regular expression is regular.

## Proof idea

It is easy to construct NFAs for 1.,2.,3.

# Language defined by regular expression

## Lemma

The language defined by any regular expression is regular.

## Proof idea

It is easy to construct NFAs for 1.,2.,3.

If we inductively have NFAs for  $L(R_1)$ ,  $L(R_2)$  then we can create an NFA for  $L(R_1 + R_2)$  and  $L(R_1 \circ R_2)$ .

# Language defined by regular expression

## Lemma

The language defined by any regular expression is regular.

## Proof idea

It is easy to construct NFAs for 1.,2.,3.

If we inductively have NFAs for  $L(R_1)$ ,  $L(R_2)$  then we can create an NFA for  $L(R_1 + R_2)$  and  $L(R_1 \circ R_2)$ .

Similarly, if we inductively have NFAs for  $L(R_1)$  then we can create an NFA for  $(L(R_1))^*$

# Language defined by regular expression

## Lemma

The language defined by any regular expression is regular.

## Proof idea

It is easy to construct NFAs for 1.,2.,3.

If we inductively have NFAs for  $L(R_1)$ ,  $L(R_2)$  then we can create an NFA for  $L(R_1 + R_2)$  and  $L(R_1 \circ R_2)$ .

Similarly, if we inductively have NFAs for  $L(R_1)$  then we can create an NFA for  $(L(R_1))^*$

What about the converse?



# Example

- ▶  $(aaa)^* + (aaaaa)^*$
- ▶  $(11 + 0)^* (00 + 1)^*$

# A few axioms we can use for simplification

## Associativity

$$\alpha + (\beta + \gamma) \equiv (\alpha + \beta) + \gamma$$

$$\alpha(\beta\gamma) \equiv (\alpha\beta)\gamma$$

# A few axioms we can use for simplification

Associativity

$$\alpha + (\beta + \gamma) \equiv (\alpha + \beta) + \gamma$$

$$\alpha(\beta\gamma) \equiv (\alpha\beta)\gamma$$

Commutativity

$$\alpha + \beta \equiv \beta + \alpha$$

Identity

$$\alpha + \emptyset \equiv \alpha$$

Idempotent

$$\alpha + \alpha \equiv \alpha$$

# A few axioms we can use for simplification

Associativity

$$\alpha + (\beta + \gamma) \equiv (\alpha + \beta) + \gamma$$

$$\alpha(\beta\gamma) \equiv (\alpha\beta)\gamma$$

Commutativity

$$\alpha + \beta \equiv \beta + \alpha$$

Identity

$$\alpha + \emptyset \equiv \alpha$$

Idempotent

$$\alpha + \alpha \equiv \alpha$$

Left Distributivity

$$\alpha(\beta + \gamma) \equiv \alpha\beta + \alpha\gamma$$

Right Distributivity

$$(\alpha + \beta)\gamma \equiv \alpha\gamma + \beta\gamma$$

# A few axioms we can use for simplification

Associativity

$$\alpha + (\beta + \gamma) \equiv (\alpha + \beta) + \gamma$$

$$\alpha(\beta\gamma) \equiv (\alpha\beta)\gamma$$

Commutativity

$$\alpha + \beta \equiv \beta + \alpha$$

Identity

$$\alpha + \emptyset \equiv \alpha$$

Idempotent

$$\alpha + \alpha \equiv \alpha$$

Left Distributivity

$$\alpha(\beta + \gamma) \equiv \alpha\beta + \alpha\gamma$$

Right Distributivity

$$(\alpha + \beta)\gamma \equiv \alpha\gamma + \beta\gamma$$

Closure

$$\epsilon + \alpha\alpha^* \equiv \alpha^*; \epsilon + \alpha^*\alpha \equiv \alpha^*$$

# A few axioms we can use for simplification

Associativity

$$\alpha + (\beta + \gamma) \equiv (\alpha + \beta) + \gamma$$

$$\alpha(\beta\gamma) \equiv (\alpha\beta)\gamma$$

Commutativity

$$\alpha + \beta \equiv \beta + \alpha$$

Identity

$$\alpha + \emptyset \equiv \alpha$$

Idempotent

$$\alpha + \alpha \equiv \alpha$$

Left Distributivity

$$\alpha(\beta + \gamma) \equiv \alpha\beta + \alpha\gamma$$

Right Distributivity

$$(\alpha + \beta)\gamma \equiv \alpha\gamma + \beta\gamma$$

Closure

$$\epsilon + \alpha\alpha^* \equiv \alpha^*; \epsilon + \alpha^*\alpha \equiv \alpha^*$$

DeMorgan-type laws

$$(\alpha + \beta)^* = (\alpha^*\beta^*)^*$$

# A few axioms we can use for simplification

Associativity

$$\alpha + (\beta + \gamma) \equiv (\alpha + \beta) + \gamma$$

$$\alpha(\beta\gamma) \equiv (\alpha\beta)\gamma$$

Commutativity

$$\alpha + \beta \equiv \beta + \alpha$$

Identity

$$\alpha + \emptyset \equiv \alpha$$

Idempotent

$$\alpha + \alpha \equiv \alpha$$

Left Distributivity

$$\alpha(\beta + \gamma) \equiv \alpha\beta + \alpha\gamma$$

Right Distributivity

$$(\alpha + \beta)\gamma \equiv \alpha\gamma + \beta\gamma$$

Closure

$$\epsilon + \alpha\alpha^* \equiv \alpha^*; \epsilon + \alpha^*\alpha \equiv \alpha^*$$

DeMorgan-type laws

$$(\alpha + \beta)^* = (\alpha^*\beta^*)^*$$

Subset order

$$\beta + \alpha\gamma \leq \gamma \implies \alpha^*\beta \leq \gamma$$

$$\beta + \gamma\alpha \leq \gamma \implies \beta\alpha^* \leq \gamma$$

# A few axioms we can use for simplification

Associativity

$$\alpha + (\beta + \gamma) \equiv (\alpha + \beta) + \gamma$$

$$\alpha(\beta\gamma) \equiv (\alpha\beta)\gamma$$

Commutativity

$$\alpha + \beta \equiv \beta + \alpha$$

Identity

$$\alpha + \emptyset \equiv \alpha$$

Idempotent

$$\alpha + \alpha \equiv \alpha$$

Left Distributivity

$$\alpha(\beta + \gamma) \equiv \alpha\beta + \alpha\gamma$$

Right Distributivity

$$(\alpha + \beta)\gamma \equiv \alpha\gamma + \beta\gamma$$

Closure

$$\epsilon + \alpha\alpha^* \equiv \alpha^*; \epsilon + \alpha^*\alpha \equiv \alpha^*$$

DeMorgan-type laws

$$(\alpha + \beta)^* = (\alpha^*\beta^*)^*$$

Subset order

$$\beta + \alpha\gamma \leq \gamma \implies \alpha^*\beta \leq \gamma$$

$$\beta + \gamma\alpha \leq \gamma \implies \beta\alpha^* \leq \gamma$$

where

$$\alpha \leq \beta \iff L(\alpha) \subseteq L(\beta)$$

$$\iff L(\alpha + \beta) = L(\beta)$$

$$\iff \alpha + \beta = \beta$$



# A few consequences that follow

## Exercise!

$$(\alpha\beta)^*\alpha \equiv \alpha(\beta\alpha)^*$$

$$(\alpha^*\beta)^*\alpha^* \equiv (\alpha + \beta)^*$$

$$\alpha^*(\beta\alpha^*)^* \equiv (\alpha + \beta)^*$$

$$(\epsilon + \alpha)^* \equiv \alpha^*$$

$$\alpha\alpha^* \equiv \alpha^*\alpha$$

# Example

- ▶  $(aaa)^* + (aaaaa)^*$
- ▶  $(11 + 0)^*(00 + 1)^*$
- ▶  $(1 + 01 + 001)^*(\varepsilon + 0 + 00)$

# Example

- ▶  $(aaa)^* + (aaaaa)^*$
- ▶  $(11 + 0)^*(00 + 1)^*$
- ▶  $(1 + 01 + 001)^*(\varepsilon + 0 + 00)$

$$(1 + 01 + 001)^*(\varepsilon + 0 + 00) \equiv ((\varepsilon + 0 + 00)1)^*(\varepsilon + 0 + 00)$$

## Example

- ▶  $(aaa)^* + (aaaaa)^*$
- ▶  $(11 + 0)^*(00 + 1)^*$
- ▶  $(1 + 01 + 001)^*(\varepsilon + 0 + 00)$

$$\begin{aligned}(1 + 01 + 001)^*(\varepsilon + 0 + 00) &\equiv ((\varepsilon + 0 + 00)1)^*(\varepsilon + 0 + 00) \\ &\equiv ((\varepsilon + 0)(\varepsilon + 0)1)^*(\varepsilon + 0)(\varepsilon + 0)\end{aligned}$$

## Example

- ▶  $(aaa)^* + (aaaaa)^*$
- ▶  $(11 + 0)^*(00 + 1)^*$
- ▶  $(1 + 01 + 001)^*(\varepsilon + 0 + 00)$

$$\begin{aligned}(1 + 01 + 001)^*(\varepsilon + 0 + 00) &\equiv ((\varepsilon + 0 + 00)1)^*(\varepsilon + 0 + 00) \\ &\equiv ((\varepsilon + 0)(\varepsilon + 0)1)^*(\varepsilon + 0)(\varepsilon + 0)\end{aligned}$$

$(1 + 01 + 001)^*(\varepsilon + 0 + 00)$  = all strings over  $\{0, 1\}$  with no substring of more than two adjacent 0's.

# DFA to regular expression

## Lemma

Any regular language can be specified by a regular expression

# DFA to regular expression

## Lemma

Any regular language can be specified by a regular expression

**Want:** Given any DFA, convert it into a regular expression.

## Lemma

Given any DFA  $A$ , we can obtain a regular expression, say  $R_A$ , such that  $L(A) = L(R_A)$ .

# Computing with labelled graphs

## Lemma

Any regular language can be specified by a regular expression

**Want:** Given any DFA, convert it into a regular expression.

## Lemma

Given any DFA  $A$ , we can obtain a regular expression, say  $R_A$ , such that  $L(A) = L(R_A)$ .