

Available Sample Cores:




Figure 2: Map of coring locations of previous (dark blue) and anticipated (pink) Antarctic expeditions from which sediment samples will be utilised (see also Table 1).

Table 1: Details of *sedaDNA* samples available to this research. Regular font indicates completed, italic font indicates upcoming expeditions. Kyr = 1,000 years, myr = million years, *sequence of field campaigns (1 complete).

Expedition	Number of samples	Age range	Location	Participation
IN2017_V01	316	25 kyr	IMAS	CI Armbrecht, PI Post
Exp. 382	196	1 myr	IMAS	CI Armbrecht
COLLAPS	449	240 kyr	IMAS	Collaborator De Santis
IN2023_V01	360	2.7 myr	IMAS	PI's Post & Suter, CI Armbrecht
CROTALE	2x6m piston core	1.2 & 3 myr	Univ. Bordeaux	Collaborator Etourneau
PS118	84	9 kyr	Univ. Bordeaux	Collaborator Crosta
SWAIS-2C*	15	recent	Univ. Waikato	CI Armbrecht
Denman*	3x1m gravity core	11 kyr	IMAS	PI Post, collab. Noble & Chase
Ross Sea Polynya	20	recent	Univ. Otago	Collaborator Riesselman
<i>IN2026_V01 Cook Ice Shelf</i>	<i>400</i>	<i>1 myr</i>	<i>IMAS</i>	CI Armbrecht, PI's Post & Suter
<i>Ross Sea Marine Protected Area</i>	<i>100</i>	<i>tbd</i>	<i>Victoria Univ. Wellington</i>	Collaborator Bertler
*1 completed field campaign out of a sequence		kyr=1,000 years	myr=million years	