

अतिसार (Atisāra)

Introduction

During *Satyayuga*, the sacrificial animals were actually released soon after the recitation of the sacrificial quotations, and these animals were not actually killed during the course of sacrifice. However in the age following the performance of the sacrifice by *Dakṣa Prajāpati*, the sons of *Manu*, like *Nariṣyan*, *Nābhāga*, *Ikṣāku*, *Nṛga* & *Śaryāti* started actually assassinating these sacrificial animals during the performance of *Yajña*, thinking that it was so ordained in the scriptures or being pressed by the animals themselves. Subsequently, it became quite impossible to get other animals in the required number, and *Prṣadhra* who was performing a sacrifice for longer duration and started sacrificing cows & bulls when other animals were not available. On seeing this the creatures became shocked and when in this afflicted mental state & consequent loss of *agni* they ingested the killed cows & bulls, it caused *Atisāra* due to heaviness, hotness, unsuitability & use of inauspicious substance. It can be concluded that *Atisāra* originated initially from the sacrifice performed by *Prṣadhra*.

अतिसार हेतु (Atisāra Hetu)

गुर्वतिस्निग्धरूक्षोष्णद्रवस्थूलातिशीतलैः ।
 विरुद्धाध्यशनाजीर्णरसात्म्यैश्चापि भोजनैः ॥
 स्नेहाद्यैरतियुक्तैश्च मिथ्यायुक्तैर्विषाद्भयात् ।
 शोकाद् दुष्टाम्युपमद्यातिपानात् सात्त्व्यर्तुपर्ययात् ॥
 जलातिरमणैर्वेगविघातैः कृमिदोषतः ।
 नृणां भवत्यतीसारो लक्षणं तस्य वक्ष्यते ॥

S.S.Utt. 40/3-5

The following etiological factors are responsible for the causation of *Atisāra* : Excessive consumption of foods which are heavy for digestion, very fatty, rough, hot, liquid, hard & cold; incompatible food consumption; excessive

quantity; uncooked & unaccustomed; improper or excessive administration of oleation and other therapies; consumption of poison; fear; sorrow; drinking contaminated water & wine in excess; adaptation of unwholesome substances contrary to seasonal regimen; excessive indulgence in water sports; suppression of natural urges and suffering from *Kṛmidoṣa*.

अतिसार संप्राप्ति (Atisāra Samprāpti)

संशय्यापां धातुरन्तःकृशानुं वर्चोमिश्रो मारुतेन प्रणुन्नः ।

वृद्धोऽतीवाधः सरत्येष यस्माद् व्याधिं घोरं तं त्वतीसारमाहुः ॥

S.S.Utt. 40/6

Due to consumption of above etiological factors *Āpa Dhātus* (body fluids) are enhanced in excess and brings diminution of strength of *Agni*, gets mixed with stool, causes downward movement in *Annavaha srotas* by *Vāta* leading to development of excess watery stool is known as *Atisāra*.

Nidāna Sevana (Consumption of Etiological factors)

↓
 Aggravation of *Pittādi doṣās* and *Āpa Dhātus*

↓
 Diminution of *Agni*

↓
 Development of *Āma* inside the gastro intestinal tract

↓
 Agitated *doṣās* mixed with stool

↓
 causes downward movement in *Annavaha srotas* by *Vāta*

↓
Atisāra (Excess watery stool)

अतिसार संप्राप्ति घटक (Samprāpti Ghaṭaka of Atisāra)

Doṣa : *Vāta pradhāna tridoṣaja*
Vāta → *Samāna*, *Apāna*

	<i>Pitta</i> → <i>Pācaka</i>
	<i>Kapha</i> → <i>Kledaka</i>
Dūṣya	: <i>Dhātu</i> → <i>Rasa, Rakta, Māmsa, Meda</i>
	<i>Upadhātu</i> →
	<i>Śārīrika Mala</i> → <i>Mūtra & Puriṣa</i>
	<i>Dhātumala</i> → <i>Kapha, Sveda, Pitta</i>
Agni	: <i>Jaṭharāgni, Dhātvaṅni</i>
Agniduṣṭi	: <i>Mandāgni</i>
Srotas	: <i>Annavaḥ Puriṣavaha, Srotas, Udaḥvaha</i>
Srotoduṣṭi	: <i>Atipravṛtti, Vimārgagamana</i>
Adhiṣṭhāna	: <i>Mahāsrotas</i>
Udbhavasthāna	: <i>Āmāśayottha & Pakvāśa-yottha</i>
Sancārasthāna	: <i>Annavaḥ Srotas, Puriṣavaha</i>
Vyaktasthāna	: <i>Guda</i>
Svabhāva	: <i>Āśukāri, Cirakāri</i>
Prabhāva	: With complications → Incurable Manifested newly without complications → Curable
Roga bheda	: 6 types
Roga Mārga	: <i>Abhyantara</i>

पूर्वरूप (Pūrvarupa)

हृन्नाभिपायूदरकुक्षितोदगात्रावसादानिलसन्निरोधाः ।
विट्सङ्ग आध्मानमथाविपाको भविष्यतस्तस्य पुरःसराणि ॥
S.S.Utt. 40/8-9

Premonitory signs & symptoms of Atisāra
are as follows : Pricking type of pain in heart, umbilicus, rectum, lower abdomen and flanks; general malaise; non elimination of flatus, constipation, distension of abdomen and indigestion.

रोग भेद (Roga bheda)

एकैकशः सर्वशश्चापि दोषैः शोकेनान्यः षष्ठ आमेन चोक्तः ।
केचित् प्राहुर्नैकरूपप्रकारं नैवेत्येवं काशिराजस्त्वोचत् ॥
दोषावस्थास्तस्य नैकप्रकाराः काले काले व्याधितस्योद्भवन्ति ॥
S.S.Utt. 40/7-8

Atisāra is classified into six types i.e. 1. *Vātaja*, 2. *Pittaja*, 3. *Kaphaja*, 4. *Sannipātaja*, 5. *Śokaja* & 6. *Āmaja*; Some people says it is classified into various types but *Kāśirāja* says it is not right, because it has many different stages of the *Doṣās* and appear at various times during the course of the disease.

According to Caraka

1. *Vātaja*, 2. *Pittaja*, 3. *Kaphaja*, 4. *Sannipātaja* & 5. *Āgantuja* (*Bhayaja* & *Śokaja*).

Classification of Atisāra

सामं चेत् कुणपं शूलैरुपविशेदरोचकी ।
स घनातिविषाकुष्ठनतदारुवचाः पिबेत् ॥
शकृद्वातमसृक् पित्तं कफं वा योजितिसार्यते ।
पक्वं, तत्र स्ववर्गीयैर्वस्तिः श्रेष्ठं भिषग्जितम् ॥

C.S.Si. 8/19-20

1. *Āmaja*, 2. *Puriṣaja* 3. *Vātaja*, 4. *Raktaja*, 5. *Pittaja*, 6. *Kaphaja*.

If patient passes stool which is *Āmayukta* associated with cadaveric smell, colic pain, anorexia, then he should be given to drink the decoction of *Ghana*, *Ativiṣa*, *Nata*, *Devadāru* & *Vacā*.

If the patient passes excessively mature stool, flatus, blood, *Pitta* or *Kapha*, this condition should be treated with enema prepared with appropriate drugs.

षण्णामेषां द्विसंस्पर्गात् त्रिंशद्भेदा भवन्ति तु ।
केवलैः सह षट्त्रिंशद्विद्यात् सोपद्रवानपि ॥

C.S.Si. 8/21

These six varieties of *Atisāra* are subdivided into thirty more varieties depending on the predominance of one or the other *Doṣa*. If six varieties are combined with them, their number becomes thirty six and are manifested along with their complications.

अतीसारः समासेन द्विधा-सामो निरामकः ॥
सासृङ्गनिरस्रः तत्राद्ये गौरवादप्सु मज्जति ।
शकृद्दुर्गन्धमाटोपविष्टम्भातिप्रसेकिनः ॥
विपरीतो निरामस्तु कफात्पक्वोऽपि मज्जति ।

A.H.Ni. 8/13-14

It is classified briefly into two types i.e. *Sāma*

& *Nirāma* and *Sāṣṛk* mixed with blood & *Nirāṣṛk* not mixed with blood.

Sāma variety is characterised by heavy stools and which sinks in water associated with foul smell, intestinal gurgling, stasis of undigested food in the stomach, pain in abdomen and excessive salivation.

Nirāma variety is characterised by opposite to *Āma* variety symptoms.

Note : Faeces containing more of *Kapha* also sinks but it is mature.

वातज अतिसार (Vātaja Atisāra)

अथावरकालं वातलस्य वातातपव्यायामातिमात्रनिषेविणो रुक्षात्प्रेमिताशिनस्तीक्ष्णमद्यव्यवायनित्यस्योदावर्तयतश्च वेगान् वायुः प्रकोपमापद्यते । C.S.Ci. 19/5

वातज अतिसार हेतु (Vātaja Atisāra Hetu)

Etiological factors of Vātaja Atisāra are as follows : If *Vāta Prakṛti* person adopts the following etiology leads to development of *Vātika Atisāra* by aggravating *Vāta*. Excessive exposure to strong wind, hot sun and physical exercise; consumption of rough food; less intake of food; irregular meals; alcohol; excessive sexual intercourse & suppression of natural urges.

वातज अतिसार संग्राप्ति (Vātaja Atisāra Samprāpti)

पक्ता चोपहन्यन्ते; स वायुः कुपितोऽग्नावुपहते मूत्रस्वेदौ पुरीषाशयमुपहत्य, ताभ्यां पुरीषं द्रवीकृत्य, अतीसाराय प्रकल्पते । C.S.Ci. 19/5

Due to above etiological factors *Vāta* gets exacerbated and *Agni* is diminished. At this stage agitated *Vāta* forcefully brings down the *Mūtra* & *Sveda* to the *Purīṣāśaya* and liquifies the stool to cause *Ātisāra*.

वातज अतिसार लक्षण/आमातिसार (Vātaja Atisāra Lakṣaṇa/Āmātisāra)

तस्य रुपाणि विज्जलमामं विप्लुतमवसादि रुक्षं द्रवं सशूलमामगन्धमीषच्छब्दमशब्दं वा विबद्धमूत्रवातमतिसार्यते पुरीषं, वायुश्चान्तःकोष्ठे सशब्दशूलस्तिर्यक् चरति विबद्ध इत्यामातिसारो वातात् । C.S.Ci. 19/5

The Signs & symptoms of Vātaja Atisāra during Āma stage are as follows : Stool is slimy

mixed with mucus; floats on water or spreads on water; precipitating, rough and liquid nature of stool; diarrhoea associated with colic pain, foul smell of the stool; patient passes stool associated with less sound or no sound; retention of urine and flatus; obstructed *Vāta* moves inside the *Koṣṭha* in haphazard direction and causes pain & gurgling sound.

वातज अतिसार लक्षण/पक्वातिसार (Vātaja Atisāra Lakṣaṇa/ Pakvatīsaara)

पक्वं वा विबद्धमल्पाल्पं सशब्दं सशूलफेनपिच्छापरिकर्तिकं हृष्टरोमा विनिःश्वसज् शुष्कमुखः कट्यूरुत्रिकजानुपृष्ठपार्श्वशूली भ्रष्टगुदो मुहुर्मुहुर्विग्रथितमुपवेश्यते पुरीषं वातात्; तमाहुरनुग्रथितमित्येके, वातनुग्रथितवर्चस्त्वात् ॥ C.S.Ci. 19/5

The signs & symptoms of Vātaja Atisāra during Pakva stage are as follows : Constipation; passes hard stool in little quantity associated with sound & colic pain; stool is frothy & slimy; cutting pain in the anus; horripilation; increased respiration; dryness of mouth; pain in waist, thigh, sacral region, knee, back and sides of the chest; prolapse of rectum; patient frequently passes scabulous stool (*Grathita*).

It is called *Anugrathita Atisāra* because of stool being scabulous by *Vāta*.

पित्तज अतिसार हेतु (Pittaja Atisāra Hetu)

पित्तलस्य पुनरम्ललवणकटुकक्षारोष्णतीक्ष्णातिमात्रनिषेविणः प्रतताग्निसूर्यसंतापोष्ण-मारुतोपहतगात्रस्य क्रोधेर्ष्याबहुलस्य पित्तं प्रकोपमापद्यते । C.S.Ci. 19/6

The following etiological factors manifest Pittaja Atisāra due to aggravation of Pitta : *Pittaja prakṛti* individuals if indulges the following etiological factors i.e regular consumption of sour, saline substances in excess; excessive exposure to strong fire, scorching sun & hot winds; psychologically stressed with anger & jealous.

पित्तज अतिसार संग्राप्ति (Pittaja Atisāra Samprāpti)

तत् प्रकुपितं द्रवत्वादूष्माणमुपहत्य पुरीषाशयविसृतमौष्ण्याद् द्रवत्वात् सरत्वाच्च भित्त्वा पुरीषमतिसाराय प्रकल्पते ।

C.S.Ci. 19/6

Due to above etiological factors *Pitta* gets aggravated and suppresses the functions of *Agni*

due to liquidity of *Pitta* and reaches the colon and disintegrates the stool because of heat, liquidity & mobility and causes *Pittaja Atisāra*.

पित्तज अतिसार लक्षण (Pittaja Atisāra Lakṣaṇa)

तस्य रूपाणि- हरिद्रं हरितं नीलं कृष्णं रक्तपित्तोपहित-
मतिदुर्गन्धमतिसार्यते पुरीषं, तृष्णादाहस्वेदमूच्छाशूलबद्ध-
संतापपाकपरीत इति पित्तातिसारः ॥ C.S.Ci. 19/6

Signs & Symptoms of Pittaja Atisāra are as follows : Patient passes frequent stool which are either yellow, green, blue or black in colour; stool is associated with *Rakta* and *Pitta* and emits foul smell; thirst; burning sensation; excessive sweating; fainting; colic pain; inflammation & suppuration in anus.

कफज अतिसार हेतु (Kaphaja Atisāra Hetu)

श्लेष्मलस्य तु गुरुमधुरशीतस्निग्धोपसेविनः संपूरकस्या-
चिन्तयतो दिवास्वप्नपरस्यालसस्य श्लेष्मा प्रकोपमापद्यते ।

C.S.Ci. 19/7

If *Kaphaja Prakṛti* individual indulges following etiological factors leading to aggravation of *Kapha* as a result *Kaphaja Atisāra* manifest.

Consumption of heavy, sweet, cold & unctuous ingredients in excessive quantity; over saturation; free from mental exercises; day sleep & sitting idle.

कफज अतिसार संप्राप्ति (Kaphaja Atisāra Samprāpti)

स स्वभावाद् गुरुमधुरशीतस्निग्धः स्रस्तोऽग्निमुपहृत्य
सौम्यस्वभावात् पुरीषाशयमुपहत्योपक्लेद्य पुरीषमतिसाराय कल्पते ।

C.S.Ci. 19/7

Due to above etiological factors *Kapha* gets aggravated. Exacerbated *Kapha* diminishes the power of *Agni* by its heavy, sweet, cold & unctuous nature and makes downward movement and reaches the colon and further due to watery nature affects the colon with excessive fluid and manifest *Kaphaja Atisāra*.

कफज अतिसार लक्षण (Kaphaja Atisāra Lakṣaṇa)

तस्य रूपाणि स्निग्धं श्वेतं पिच्छिलं तन्तुमदामं गुरु दुर्गन्धं
श्लेष्मोपहितमनुबद्धशूलमल्पाल्पमभीक्ष्णमतिसार्यते सप्रवाहिकं,
गुरुदरगुदबस्तिवक्ष्णदेशः कृतेऽप्यकृतसंज्ञ- सलोमहर्षः सोत्क्लेशो
निद्रालस्यपरीतः सदनोऽन्तद्वेषी चेति श्लेष्मातिसारः ॥

C.S.Ci. 19/7

Signs and symptoms of Kaphaja Atisāra are as follows- Patient passes stool which is unctuous, white, slimy, thready, *Āma*, heavy, foetid, mixed with mucus associated with frequent colic pain; patient passes little quantity of stool frequently associated with tenesmus; patient suffers from heaviness in abdomen, anus, pelvis, & groin; has motions without knowledge; horripilation; nausea; excessive sleep; idleness; lassitude and aversion towards food.

सन्निपातज अतिसार हेतु (Sannipātaja Atisāra Hetu)

अतिशीतस्निग्धरुक्षोष्णागुरुखरकङ्गिनविषमविरुद्धासात्य-
भोजनात् कालातीतभोजनाद् यत्किंचिदभ्यवहरणात् प्रदुष्टमद्य-
पानीयपानादतिमद्यपानादसंशोधनात् प्रतिकर्मणां विषमग-
मनादनुपचाराज्ज्वलनादित्यपवनसलिलातिसेवनादस्वप्नादति -
स्वप्नाद्वेगविधारणादुतुविपर्ययादयथाबलमारम्भाद्भयशोकचित्तो-
द्वेगातियोगात् कुमिशोषज्वराशौविकारातिकर्षणाद्वा व्यापन्नाग्नेस्त्रयो
दोषाः प्रकुपिताः C.S.Ci. 19/8

The following etiological factors are said to be responsible for the manifestation of *Sannipātaja Atisāra* by simultaneous aggravation of all the three *Doṣās* : Excessive consumption of cold, unctuous, hot, heavy, coarse and hard ingredients; irregular meals; consumption of mutually contradictory food substances; unwholesome food; avoiding intake of food; delayed meals, taking food without considering dietetic rules & regulations; drinking polluted water & wine; drinking of alcohol in excess; not undergoing elimination therapies as per seasons. Inappropriate or non administration of therapeutics; excessive exposure to fire, hot season sun, strong wind and bath etc. in strong force of water; no sleep or excessive sleep; suppression of natural urges; seasonal perversations; excessive physical exercise beyond one's own physical strength; excessive exposure to fear, grief and anxiety; excessive emaciation due to *Kṛmi*, *Śoṣa*, *Jvara*, *Arśa* and derangement of *Agni*.

सन्निपातज अतिसार संप्राप्ति एवं लक्षण (Sannipātaja Atisāra Samprāpti evaṁ Lakṣaṇa)

भूय एवाग्निमुपहृत्य पक्वाशयमनुप्रविश्यातीसारं सर्वदोषलिङ्गं
जनयन्ति ॥ C.S.Ci. 19/8

Due to above etiological factors all the three *Doṣās* gets aggravated and brings affliction in *Agni* and reaches *Pakvāśaya* and cause *Sannipātaja Atisāra* with signs & symptoms of all the three *Doṣās*.

सन्निपातज अतिसार लक्षण (Sannipātaja Atisāra Lakṣaṇa)

अपि च शोणितादीन् धातून्तिप्रकृष्टं दूषयन्तो धातुदोषस्व-
भावकृतानतीसारवर्णानुपदर्शयन्ति । तत्र शोणितादिषु धातुष्वति-
प्रदुष्टेषु हारिद्रहरितनीलमाज्जिष्ठमांसधावनसन्निकाशं रक्तं कृष्णं श्वेतं
वराहमेदःसदृशमनुबद्धवेदअनमवेदनं वा समासव्यत्यादुपवेश्यते
शकृद् ग्रथितमामं सकृत्, सकृदपि पक्वमनतिक्षीणमांसशोणितबलो
मन्दाग्निर्विहतमुखरसश्च, तादृशमातुरं कृच्छ्रसाध्यं विद्यात् ।

एभिर्वर्णैरतिसार्यमाणं सोपद्रवमातुरमसाध्योऽयमिति प्रत्या-
चक्षीत, तद्यथा पक्वशोणिताभं यकृतखण्डोपमं मेदोमांसोद-
कसन्निकाशं दधिघृतमज्जतैलवसाक्षीरवेसवाराभमतिनील-
मतिरक्तमतिवृष्णमुदकमिवाच्छं पुनर्मैचकाभमतिस्निग्धं
हरितनीलकषायवर्णं कर्बुरमाविलं पिच्छिलं तन्तुमदामं चन्द्रकोप-
तमतिकृष्णपूपितपूयगन्ध्यामाममत्स्यगन्धि मक्षिकाक्रान्तं
कुथितबहुधातुस्त्रावमल्पपुरीषमपुरीषं वाऽतिसार्यमाणं तृष्णादाह-
ज्वरभ्रमतमकहिकवाश्वासानुबन्धमतिवेदनमवेदनं वा स्वस्तपक्वगुदं
पतितगुदविलं मुक्तनालमतिक्षीणबलमांसशोणितं सर्वपर्वस्थि-
शूलिनमरोचकारतिप्रलापसंमोहपरीतं सहसोपरतविकारम-
तिसारिणमचिकित्स्यं विद्यात्, इति सन्निपातातिसारः ॥

C.S.Ci. 19/9

Excessively aggravated *Vātādi* three *Doṣās* brings abnormality in *Raktādidhātus* as a result it manifest different coloration of stool; nature of vitiation of *Doṣa* & *Dhātu* exhibit various colours in stool. If not taken care, further vitiation of *Doṣās* & *Dhātus* manifest following signs & symptoms- Patient passes stool which is either yellow, green, blue, red like *Mañjiṣṭhā*, pink (water in which meat is washed), red, black, white and yellowish (like the colour of pig-fat) painful abdomen or painless.

The above mentioned signs & symptoms manifest in their entirety or only few may be observed. Patient passes stool which is scabulous, *Āma*, or may be free from mucus; diminution of *Māmsa*, *Rakta* and *Bala*; suppression of the power of *Agni*; distaste in mouth. Above symptomatology indicates that the disease is curable with difficulty.

The following signs and symptoms indicates the incurable nature of the disease- Patient passes stool appears like cooked blood, liver pieces, fat, meat water, curd, ghee, marrow, oil, muscle fat, milk and *Vesavāra* (a spicy preparations); patient passes stool which is blue, red, black, transparent like water or tar coloured, greasy; colour of the stool appears like green, blue and brown colours; stool appears like variegated in colour (*Karbura*), dirty, slimy, fibrous, mixed with mucus and spotted with *Candraka* (moony lustre), cadaverous smell, foetid, putrid, smells like undigested products or like raw fish, attracts flies, stool contain profuse discharge of increased tissues, passing little or no stool associated with thirst, burning sensation, fever, giddiness, fainting, hiccup, asthma excessive pain or no pain, prolapse & suppuration of anus, sphincters come out of their sites or whole of the rectum comes out; excessive loss of strength, *Māmsa* & *Rakta*; pain in all the joints & bones; patient suffers from anorexia, dislike for everything, delirium & unconsciousness and sudden cessation of the signs & symptoms of the disease.

सन्निपातज अतिसार (Sannipātaja Atisāra)

तन्त्रायुक्तो मोहसादाम्यशोषी वर्चः कुर्यान्नैकवर्णं तृषार्तः ।

सर्वोद्भूते सर्वलिङ्गोपपत्तिः कृच्छ्रश्चायं बालवृद्धेष्वसाध्यः ॥

S.S.Utt. 40/13

Sannipātaja Atisāra is characterised by stupor, delusion, debility, dryness of mouth, eliminates faeces of many colours, thirst and has the appearance of specific symptoms of all the *Doṣās*.

It is very difficult to treat and incurable in young & old persons.

आगन्तुज अतिसार (Āgantuja Atisāra)

आगन्तू द्वावतीसारौ मानसौ भयशोकजौ ।

तत्तयोर्लक्षणं वायोर्यदतीसारलक्षणम् ॥

C.S.Ci. 19/11

The exogenous (*Āgantuja Atisāra*) manifest due to mental factors, and it is of two types i.e. *Bhayaja* & *Śokaja*. Their signs & symptoms are similar to those of *Vātaja Atisāra*.

शोकज अतिसार (Śokaja Atisāra)

तैस्तैर्भावैः शोचतोऽल्पाशनस्य बाष्पावेगः पक्तिमाविध्य जन्तोः ।
कोष्ठं गत्वा क्षोभयत्यस्य रक्तं तच्चाधस्तात् काकणन्तीप्रकाशम् ॥
वर्चोमिश्रं निःपुरीषं सगन्धं निर्गन्धं वा सार्यते तेन कृच्छ्रात् ।
शोकोत्पन्नो दुश्चिकित्सोऽतिमात्रं रोगो वैद्यैः कष्ट एष प्रदिष्टः ॥
S.S.Utt. 40/14-15

Person who is suffering from various kinds of grief eat less quantity of food, the force of the tears (grief) vitiates the digestive power of man and reaches the gastrointestinal tract, vitiates his *Rakta*; begins to move in the downward direction and passes stool mixed with *Rakta*; the blood resembles *Kākaṇanti* in colour either mixed with stool or without stool comes out from *anus* associated with either bad smell or without smell & passes with difficulty. It is very difficult to treat says the physicians.

आमज अतिसार (Āmaja Atisāra)

आमाजीर्णोपद्रुताः क्षोभयन्तः कोष्ठं दोषा सम्प्रदुष्टाः सभक्तम् ।
नानावर्णं नैकशः सारयन्ति कृच्छ्राज्जन्तोः षष्ठमेनं वदन्ति ॥
S.S.Utt. 40/15-16

Due to *Amājirṇa*, *Doṣās* excacerbates in excess in abdomen gives rise to elimination of faeces of various colours, passes stool with difficulty and it is difficult to treat.

रक्तातिसार (Raktātisāra)

पित्तातिसारी यस्त्वेतां क्रियां मुक्त्वा निषेवते ।
पित्तलान्यन्नपानानि तस्य पित्तं महाबलम् ॥
कुर्याद्रक्तातिसारं तु रक्तमाशु प्रदूषयेत् ।
तृष्णां शूलं विदाहं च गुदपाकं च दारुणम् ॥
C.S.Ci. 19/69-70

पित्तातिसारी यो मर्त्यः पित्तलान्यतिषेवते ।
पित्तं प्रदुष्टं तस्याशु रक्तातीसारमावहेत् ॥
ज्वरं शूलं तृष्णां दाहं गुदपाकञ्च दारुणम् ॥
S.S.Utt. 40/116-117

One who is suffering from *Pittaja Atisāra* and does not follow the treatment of *Atisāra* and indulges foods & drinks which cause further aggravation of *Pitta* upto great extent as a result *Rakta* gets vitiates and manifest *Raktātisāra*. It is very difficult for management because it manifest

serious condition and patient suffers from thirst, colic pain, burning sensation & inflammation of the ano-rectal region.

Raktātisāra may manifest as a complication of *Pittātisāra* and it may also manifest independently without *Pittātisāra* if person adopts *Pitta* provocative etiological factors in excess.

उपद्रव (Upadrava)

All the *Lakṣaṇa* mentioned under *Sannipātaja Atisāra*.

शूलप्रवाहिकाध्मानपरिकर्त्यरुचिज्वरान् ।
तृष्णोष्णादाहमूर्च्छादींश्चैषां विद्यादुपद्रवान् ॥
C.S.Si. 8/22

Complications of *Atisāra* are colic pain, gripping pain, flatulence, thirst, feeling of hotness, burning sensation, fainting etc.

साध्यासाध्यता एवं अरिष्ठ (Sādhyāsādhyatā evam Ariṣṭa)

सर्पिर्मंदोवेसवाराम्बुतैलमज्जाक्षीरक्षौद्ररूपं स्रवेद् यत् ।
मज्जिष्ठाऽऽभं मस्तुलुङ्गोपमं वा विस्त्रं शीतं प्रेतगन्ध्यञ्जनाभम् ॥
राजीमद् वा चन्द्रकैः सन्ततं वा पूयप्रख्यं कर्दमाभं तथोष्णम् ।
हन्यादेतद् यत् प्रतीपं भवेच्च क्षीणं हन्युश्चोपसर्गाः प्रभूताः ॥
असंवृतगुदं क्षीणं दुराध्मातमुपद्रुतम् ।
गुदे पक्वे गतोष्माणमतीसारकिणं त्यजेत् ॥
S.S.Utt. 40/19-21

Stool resembling ghee, fat, water of meat soup, oil, bone marrow, milk, honey, *Mañjiṣṭā* or *Mastuluṅga*, putrid smell, cold, cadaveric smell, appears like *Añjana* (*Sauvirāñjana*), has separating lines, glistening particles, resembling like pus or slush and hot, these features greatly enhanced and acting as complications and kill the weak persons. The patient whose anal orifice does not close at all, who is emaciated, having severe flatulence, inflammation in the anus, and coldish touch should be rejected.

अतिसार निवृत्ति लक्षण (Atisāra Nivṛtti Lakṣaṇa)

यस्योच्चारं विना मूत्रं सम्यग्वायुश्च गच्छति ।
दीप्ताग्नेर्लघुकोष्ठस्य स्थितस्तस्योदरामयः ॥
S.S.Utt. 40/162

Proper elimination of urine, flatus and stool,

enhancement of digestive fire and feeling of lightness in gastro-intestinal tract indicates that patient is recovered from *Atisāra*.

उपशय/पथ्य (Upaśaya/Pathya)

Auśadha

Kālīṅgakādi yoga, Nāgarādi Yoga, Mustādi Yoga, Pippalyādi Yoga, Cāṅgeri ghṛta, Cavyādi ghṛta, Dāryādi ghṛta, Kapitthāṣṭaka Cūrṇa, Dādimāṣṭaka Cūrṇa.

Anna

In appropriate time, when patient feels hungry, he should be given light food to eat. This helps the patient to regain appetite, his *Agni* get stimulated and strength is promoted immediately.

In the initial stage, depending upon the wholesomeness, patient should take light food along with butter milk, *Avanti-soma* or *Kāñji* (sour drink), *Yavāgu* (thick gruel), *Tarpaṇa* (roasted flour of cereals mixed with water), wine, honey, then gradually, the patient should gradually be given *Yavāgu* (thick gruel), *Vilepī* (a sticky gruel), *Khaḍa* (a sour appetiser) *Yūṣa* (vegetable soup) and boiled rice mixed with meat soup, which are prepared by adding digestive stimulants and astringent drugs.

Vihāra

Vāta gets aggravated soon, even by fear and grief, that's why *Vāta* mitigating factors are beneficial and it should be done by providing pleasure & courage to the patient as per patient need.

If the patient of *Atisāra* is hungry, he should be given light food to eat. *Peya* (thin gruel without containing any grains in it) prepared by boiling with appropriate drugs, or fried paddy or roasted-corn flour mixed along with water is also useful for him. The *Maṇḍa* (that does not contain *Siktaka*) can also be taken by the patient. (B.R. 7/11-12)

Pathya (Meals to be taken) for a patient of *Atisāra* and *Pravāhikā*: Vomiting, *Laṅgana*, sleeping are good for the patient of *Atisāra*. The following edibles and meals are helpful for the patient: old *Śālī* variety of paddy, *Ṣaṣṭhī* variety of rice, *Vilepī* (watery rice), *Maṇḍa* of poppy, soups of lentil and pigeon pea, meatsoup of rabbit, *Aiṇ*

(a kind of black deer), *Lāva* bird, deer, *Kapīñjala* (partridge), all kinds of small fishes, the *Śṛīṅgī* fish, *Madhurālikā* (a kind of fish), medicinal oil, *Ghṛta* and milk obtained from goats and cow, buttermilk, butter, fresh banana fruits, honey (obtained from smaller bees), *Jambūphalā*, dried ginger, common ginger, *Kamalakanda*, *Kapittha*, *Vikañṭaka*, *Bakula*, *Bilva*, *Tenduka*, *Dāḍima* (both ripe and unripe ones), palm fruits, *Kaṅcaṭa* leaves, *Cāṅgerī*, *Vijayā*, *Mañjiṣṭhā*, *Jātiphala*, *Hṛbera*, cumin seeds, *Girimallikā*, *Kustumburu*, *Mahānimba*, all kinds of stomachiac foods, light meals and drinks. The point about two fingers below the naval as well as the point around bottom of backbone, both should be heated in a crescent shape. This measure also helps one to get cure of *Atisāra*. (B.R. 7/183-189)

प्रवाहिका (Pravāhikā)

वायुः प्रवृद्धो निचितं बलासं नुदत्यधस्तादहिताशनस्य ।

प्रवाहमाणस्य मुहुर्मलाक्तं प्रवाहिकां तां प्रवदन्ति तज्ज्ञाः ॥

S.S.Utt. 40/138

One who consumes unwholesome foods & drinks leading to aggravation of *Vāta* and pushes the *Kapha* downwards again & again as a result *Kapha* comes out along with stool is known as *Pravāhikā*.

प्रवाहिका भेद एवं लक्षण (Pravāhikā Bheda Evam Lakṣaṇa)

प्रवाहिका वातकृता सशूला पित्तात् सदाहा सकफा कफाच्च ।

सशोणिता शोणितसम्भवा तु ताः स्नेहलक्षप्रभवा मतास्तु ॥

तासामतीसारवदादिशेच लिङ्गं क्रमं चामविषक्वताञ्च ॥

S.S.Utt. 40/140

1. *Vātaja*, 2. *Pittaja*, 3. *Kaphaja* and 4. *Raktaja* *Pravāhikā* are described along with their symptomatology & these may be considered as classification of *Pravāhikā*.

Pravāhika caused by exacerbated *Vāta* produces pain, that caused by *Pitta* produces burning sensation, that caused by *Kapha* produces elimination of stool mixed with *Kapha* and that caused by *Rakta* produces stool mixed with *Rakta*. These kind of *Pravāhika* are produced by more of unctuousness & roughness in *Pakvāsaya*. Stages of *Āma* & *Pakva* and the management principles are similar to that of *Atisāra*.

Differentiate between Atisāra and Pravāhikā

S.No.	Atisāra	Pravāhikā
1	Frequent passing of excessive watery stool	Evacuation of stool mixed with Kapha
2	No strain while evacuating stool	Strain while evacuating stool

Differentiate doṣaja Atisāra

S.No	Vātaja	Pittaja	Kaphaja	Sannipātaja
Etiology	If <i>Vāta prakṛti</i> person adopts the following etiology leads to development of <i>Vātika Atisāra</i> by aggravating <i>Vāta</i> . Excessive exposure to strong wind, hot sun and physical exercise; consumption of rough food; less intake of food; irregular meals; alcohol; excessive sexual intercourse & suppression of natural urges.	<i>Pittaja prakṛti</i> individuals if indulges the following etiological factors i.e regular consumption of sour, saline substances in excess; excessive exposure to strong fire, scorching sun & hot winds; psychologically stressed with anger & jealous.	If <i>Kaphaja Prakṛti</i> individual indulges following etiological factors leading to aggravation of <i>Kapha</i> as a result <i>Kaphaja Atisāra</i> manifest. Consumption of heavy, sweet, cold & unctuous ingredients in excessive quantity; over saturation; free from mental exercises; day sleep & sitting idle.	The following etiological factors are said to be responsible for the manifestation of <i>Sannipātaja Atisāra</i> by simultaneous aggravation of all the three <i>Doṣās</i> : Excessive consumption of cold, unctuous, hot, heavy, coarse and hard ingredients; irregular meals; consumption of mutually contradictory food substances; unwholesome food; avoiding intake of food; delayed meals, taking food without considering dietetic rules & regulations; drinking polluted water & wine; drinking of alcohol in excess; not undergoing elimination therapies as per seasons. Inappropriate or non administration of therapeutics; excessive exposure to fire, hot season sun, strong wind and bath etc. in strong force of water; no sleep or excessive sleep; suppression of natural urges; seasonal perversations; excessive physical exercise beyond once own physical strength; excessive exposure to

				fear, grief and anxiety; excessive emaciation due to <i>Kṛmi</i> , <i>Śoṣa</i> , <i>Jvara</i> , <i>Arśa</i> and derangement of <i>Agni</i> .
Clinical Feature	Constipation; passes hard stool in little quantity associated with sound & colic pain; stool is frothy & slimy. Patient frequently passes scabulous stool (<i>Grathita</i>).	Patient passes frequent stool which are either yellow, green, blue or black in colour; stool is associated with <i>Rakta</i> and <i>Pitta</i> and emits foul smell	Patient passes stool which is unctuous, white, slimy, thready, <i>Āma</i> , heavy, foetid, mixed with mucus associated with frequent colic pain; patient passes little quantity of stool frequently associated with tenesmus;	Excessively aggravated <i>Vātādi</i> three <i>Doṣās</i> brings abnormality in <i>Raktādhātus</i> as a result it manifest different coloration of stool; nature of vitiation of <i>Doṣa</i> & <i>Dhātu</i> exhibit various colours in stool Patient passes stool which is either yellow, green, blue, red like <i>Mañjiṣṭhā</i> , pink (water in which meat is washed), red, black, white and yellowish (like the colour of pig-fat) painful abdomen or painless.
Clinical Feature	patient suffers from pain in waist, thigh, sacral region, knee, back and sides of the chest;	patient suffers from thirst; burning sensation; excessive sweating; fainting; colic pain	patient suffers from heaviness in abdomen, anus, pelvis, & groin;	patient suffers from excessive loss of strength, <i>Māmsa</i> & <i>Rakta</i> ; pain in all the joints & bones;
Complication	patients of <i>Vātika Atisāra</i> may not develop <i>Raktātisāra</i> as a complication	patients of <i>Pittātisāra</i> may develop <i>Raktātisāra</i> as a complication	patients of <i>Kaphaja Atisāra</i> may not develop <i>Raktātisāra</i> as a complication	

MCQs

- 'Visramsi' according to Acharya Bhoja is
 - Rakta-Pitta
 - Atisara
 - Pravahika
 - Grahani
- Type of *Atisara* not mentioned by *Susruta*
 - Bhayaja
 - Sokaja
 - Tridoshaja
 - Kaphaja
- Types of *Atisara* according to *Sharangadhara*
 - 6
 - 7
 - 5
 - 4
- Bhrastaguda muhurmuhur vibrathitha* features observed in
 - Vatika Atisara*
 - Paithika Atisara*
 - Tridoshaja Atisara*
 - Sokaja Atisara*
- Krite-apyakritasamjna* sa lomaharsha is the feature of
 - Pravahika*
 - Tridoshaja Atisara*
 - Kaphaja Atisara*
 - Bhayaja Atisara*
- Nana varna naikashah sarayanthi* features observed in
 - Pittaja Atisara*
 - Tridoshaja Atisara*
 - Shokaja Atisara*
 - Amaja Atisara*

7. Types of *Pravahika* as per *Madavanidana*

- a. 5 b. 4
c. 3 d. 2

8. विज्जलभामं विप्लुतमवसादि features observed in

- a. Rakta-Pitta b. Vataja Atisara
c. Pitttaja Atisara d. Grahani

9. According to *Ashtanga Hridaya* the types of *Atisara* include all except

- a. Bhayaja b. Shokaja
c. Amaja d. Both a & c

10. *Purvarupa* of *Atisara* include all except

- a. Gatrasada b. Malagraha
c. Adhmana d. Hrillasa

11. *Kshipramushnamdravamlavam* is the feature of which *atisara*

- a. Paittika b. Bhayaja
c. Shokaja d. Both b & c

Answer

1. c	2. a	3. b
4. a	5. c	6. d
7. b	8. b	9. c
10. d	11. d	