

Management of patients with Gastrointestinal diseases

Dr . Bardies shaban





الحديث الجانبي يشتت الانتباه



من فضلك اغلق الهاتف

2- Cholelithiasis & Cholecystitis


- ◉ Definition
- ◉ Causes
- ◉ Manifestations
- ◉ Diagnosis
- ◉ Complications
- ◉ Managements

Definition of cholecystitis


acute inflammation of the gallbladder, causes pain, tenderness, and rigidity of the upper right abdomen that may radiate to right shoulder.

Cholecystitis

Gallbladder inflammation



Causes




Calculus cholecystitis

is the cause of more than **90%** of cholecystitis. In which a gallbladder stone obstructs bile outflow.

Bacteria


Function of the gall bladder:

- ▶ The gallbladder functions as a **storage for bile**. The bile is a thick, brown liquid and composed of water and electrolytes (sodium, potassium, calcium, chloride, and bicarbonate).
- ▶ Bile: digestion of fat, neutralize of gastric acid



Definition of Cholelithiasis

Calculi or **gallstones**, usually form in the **gallbladder** from the **solid constituents of bile**, they vary in **size, shape and composition**.



- 1



Risk Factors for Gallbladder Stones



Obesity



Cholesterol-lowering Drugs



Diabetes



Rapid Weight Loss



Fasting



Oestrogen

Types of gallstones

Cholesterol gallstones



Pigment gallstones



Types of Gallstones



Cholesterol Stones



Pigment Stones



Mixed Stones

Symptoms of Gallbladder Stones



Pain in the Right
Abdomen and Back



Nausea and
Vomiting



Fever & Chills




Jaundice



Abdominal Bloating

Gall Stone

the real pain



How Gallbladder Problems Affect the Body



management



SOUTHLAKE
CENTRAL CALIFORNIA

TREATMENT FOR GALLSTONES

By Dr Valeria Simone MD

CONTACT US
+1 813-748-0200

www.southlakegeneralsurgery.com

The advertisement features two circular images on the right side. The top image shows a person's bare midsection with a hand pointing to the upper right abdomen. The bottom image is a medical illustration of a gallbladder filled with numerous small, green, oval-shaped gallstones. The background of the ad is white with a blue wavy shape at the bottom and a hexagonal pattern on the left.

Pharmacological management

- Ursodeoxycholic acid (**UDCA**) has been used to dissolve small gallstones composed primarily of cholesterol.
- Treatment with UDCA can **reduce the size** of existing stones, **dissolve small stones**, and prevent new stones from **forming**
- **Six to 12 months** of therapy are required in many patients to dissolve stones, and monitoring of the patient for recurrence of symptoms.



Medical management


▶ **Nutritional**

- ▶ Low-fat diet.
- ▶ Cooked fruits, rice or lean meats, mashed potatoes, non-gas-forming vegetables and bread.
- ▶ The patient should avoid eggs, cream, fried foods, cheese, and alcohol.
- ▶ Approximately 80% of the patients achieve remission with rest, IV fluids, nasogastric suction, analgesia, and antibiotic.

Non invasive / non surgical management

► Extracorporeal Shock Wave Lithotripsy


Repeated **shockwaves** directed at the **gallstones** in the gallbladder to fragment the stones.



Non surgical management

► Intracorporeal Lithotripsy

Using of laser pulse technology



Surgical management


▶ Cholecystectomy

Laparoscopic

- 4 small incision made through the abdominal wall.

Surgical /open

- large incision through abdominal.



Postoperative Nursing Interventions

1- Relieving Pain

- **Analgesic** as prescribed to control the pain.
- Use of a pillow over the incision may **reduce pain**.
- Complete frequent **assessment** of pain.
- Note and report **right upper quadrant** abdominal pain

Postoperative Nursing Interventions

2- Improving Respiratory Status

- Places the patient in the semi fowler's position.
- Nasogastric suction to relieve abdominal distention.
- Encourage deep breath and cough every hour to expand the lung fully and prevent atelectasis.
- Early ambulation

Postoperative Nursing Interventions

3- Maintaining Skin Integrity & Promoting Biliary Drainage

- Frequently change of **dressing** and protection of the skin from irritation because bile is corrosive to the skin.
- Every 24 h the nurse measures the bile collected and record the amount, color and character of the **drainage**.
- Within **7 to 14 days** the drainage tube is **removed**

Postoperative Nursing Interventions

4- Improving Nutritional Status

- Water and fluids are given in about 24 hours
- Maintain a careful record of fluid intake and output.
- soft diet is started when bowel sounds return.
- low fats diet and high in carbohydrate and proteins.

Postoperative Nursing Interventions

5– Monitoring and Managing Complications

- Closely monitor vital signs and inspect the surgical incision and drain.
- Instructs the patient and family to report any changes in color of stool.
- Assess the patient for loss of appetite, vomiting, pain, distention of the abdomen and increase temperature.


Postoperative Nursing Interventions

6– general health teaching

- Inform the patient about symptoms that should be reported to the physician which include :
- jaundice, dark urine, pale colored stool, signs of inflammation and infection as pain or fever.
- Take a shower or bath after 1 or 2 days.
- Drive a car after 3 or 4 days.
- Avoid lifting objects exceeding 5 pounds (2.5 kg) after surgery, usually for 1 week.

- Call surgeon if experience any signs or symptoms of infection at or around the puncture site: redness, tenderness, or swelling.
- Wash puncture site with mild soap and water.





Summary



- List **3 methods of management** for gallbladder stones
- List **4 nursing interventions** for patient post Cholecystectomy
- List 4 risk factors **for gallbladder stones**



Thank
you