

## **Ortho: 1- Fracture**

- 1- Which type of fracture occurs when the bone is broken into more than two fragments?  
A. **Comminuted fracture**  
B. Greenstick fracture  
C. Oblique fracture  
D. Transverse fracture
- 2- Which type of fracture occurs when bone breaks into two parts that are completely separated?  
A. Comminuted fracture  
B. .Greenstick fracture  
C. **Complete fracture**  
D. in complete fracture
- 3- Which type of fracture occurs with no break in the skin?  
A. Complete fracture  
B. in complete fracture  
C. **Closed fracture**  
D. Open fracture
4. Which type of fracture involves a break where the bone pierces the skin?  
A. Closed fracture  
B. Greenstick fracture  
C. **Open fracture**  
D. Impacted fracture
5. A fracture in which the bone bends and cracks but does not completely break is called:  
A. Comminuted fracture  
B. Spiral fracture  
C. **Greenstick fracture**  
D. Transverse fracture
6. What is the most common cause of fractures?  
A. Infection  
B. **Trauma or injury**  
C. Cancer  
D. Genetic mutation
7. Which fracture occurs due to repeated stress over time?  
A. Open fracture  
B. **Stress fracture**  
C. Impacted fracture  
D. Pathological fracture

8. **What is the first step in managing a suspected fracture?**
- A. Massage the area
  - B. Apply heat
  - C. Immobilize the limb
  - D. Give antibiotics
9. **Which of the following is NOT a sign of fracture?**
- A. Pain
  - B. Swelling
  - C. Deformity
  - D. Increased mobility without pain
10. **Which of the following is a complication of fractures?**
- A. Hypertension
  - B. Fat embolism
  - C. Diabetes
  - D. Asthma
11. **The term "closed fracture" means:**
- A. Bone is shattered
  - B. Skin over the fracture is intact
  - C. Bone is infected
  - D. Bone is bent
12. **Which of the following is a sign of an open fracture?**
- A. Swelling only
  - B. Bruising
  - C. Bone visible through the skin
  - D. Mild pain
13. **What does "reduction" mean in fracture treatment?**
- A. Breaking the bone further
  - B. Aligning the bone fragments
  - C. Removing the bone
  - D. Strengthening muscles
14. **What is the main goal of fracture management?**
- A. Eliminate pain only
  - B. Restore normal function
  - C. Increase bone size
  - D. Prevent movement permanently

**15-List signs and symptoms of fractures.**

- Pain , Swelling
- Tenderness
- Deformity
- Ecchymoses
- loss of function
- Crepitation with any movement

#### 16-List the care of patient with traction

- T: traction
- R: rope hang free
- A: alignment
- C: check circulation
- T: type and location of traction
- I: increase fluid intake
- O: overhead trapeze
- N: no weight on bed or floor

#### 17- List Complications of Fractures

- Hemorrhagic shock
- Damage to nerves from fragments of bone
- Fat emboli
- External pressure from dressings or casts that are too tight
- Infection

#### 18-List 6 P of neurovascular assessment of patient with fracture

- Pain
- Pulse
- Paresthesia
- Pallor
- Paralysis
- Pressure

#### 19- List nursing management for patient with fracture

- Put patient in a high Fowler's position
- Establish a peripheral IV fluids.
- Elevation of the affected part to prevent swelling , toes should be higher than the trunk
- If a cast is in place, be bivalved (split through all layers of the material). Dressings will be cut or replaced
- Wash hands
- Use aseptic technique when caring for wound
- Report excessive drainage or inflammation to physician
- Culture drainage
- Frequent monitoring of temperature

### **Ortho: 2- Hip replacement**

#### 1- The most common indication for hip replacement is:

- A. Fracture of arm
- B. Osteoarthritis
- C. Asthma
- D. Migraine

#### 2- Which part of the hip is replaced in total hip replacement?

- A. Only femur
- B. Only pelvis
- C. Both femoral head and acetabulum
- D. Only cartilage

**3- The artificial hip joint is usually made of:**

- A. Wood
- B. Plastic only
- C. Metal and plastic/ceramic
- D. Glass

**4- Which of the following is a goal of hip replacement surgery?**

- A. Increase bone length
- B. Relieve pain and improve mobility
- C. Stop blood flow
- D. Cure infection

**5- What is “total hip arthroplasty”?**

- A. Partial repair
- B. Complete replacement of hip joint
- C. Muscle surgery
- D. Bone cutting only

**6- Which condition may require hip replacement?**

- A. Rheumatoid arthritis
- B. Common cold
- C. Hypertension
- D. Gastritis

**7- What is a key part of recovery after surgery?**

- A. Bed rest only
- B. Physiotherapy
- C. Avoid movement completely
- D. No follow-up

**8- Which age group most commonly undergoes hip replacement?**

- A. Children
- B. Young adults only
- C. Older adults
- D. Infants

**9- Which of the following is an early complication?**

- A. Implant wear
- B. Deep vein thrombosis (DVT)
- C. Prosthesis loosening after years
- D. Bone remodeling

**10- What is the purpose of physiotherapy after surgery**

- A. Prevent infection
- B. Strengthen muscles and restore movement
- C. Replace the implant
- D. Stop blood circulation

**11- Which symptom may indicate hip prosthesis infection?**

- A. Better sleep
- B. Improved mobility
- C. Fever and persistent pain
- D. Increased appetite

**12- Which type of hip replacement only the femoral head is replaced?**

- A. Total hip replacement
- B. Hip Resurfacing
- C. Hip arthroplasty
- D. Partial hip replacement

**13- Avascular Necrosis is:**

- A. Complication of hip replacement
- B. Side effect of hip replacement
- C. Cause of hip replacement
- D. Type of hip replacement

**14- Ankle and calf pumping exercises are exercises done for patients post orthopedic surgery to:**

- A. Increase pain
- B. Increase swelling
- C. Enhance circulation
- D. Decrease tissue perfusion

**15- Homan's sign is:**

- A. Abdominal pain on deep palpation
- B. Swelling of the hand when the arm is elevated
- C. Chest pain during deep breathing
- D. Calf pain on dorsiflexion of the foot

**16- What is the purpose of early mobilization after surgery?**

- A. Increase pain
- B. Delay healing
- C. Prevent complications like DVT
- D. Increase dislocation

## **17-List nursing diagnosis for patient post orthopedic surgery**

- Pain related to the surgical procedure.
- Impaired physical mobility related to pain, swelling.
- Altered body image and self-esteem related to impact of musculoskeletal problems.
- High risk for altered peripheral tissue perfusion related to swelling and impaired circulation.
- High risk for fluid volume deficit related to excessive losses from surgery.
- High risk for infection related to traumatized tissues.

## **18- list common causes of hip replacement**

- Osteoarthritis
- Rheumatoid Arthritis
- Hip Fracture
- Avascular Necrosis

## **19- list nursing interventions to decrease pain for patients post orthopedic surgery**

- Analgesic as prescribed to control the pain (used as a preventive).
- Elevation of the operative extremity and application of cold compresses , if prescribed, help to control edema.
- Portable suction of the wound decreases fluid accumulation and hematoma formation.
- After 3 to 4 days, most patients require only oral analgesia.

## **Ortho: 3- Rheumatoid arthritis**

### **1. What type of disease is rheumatoid arthritis?**

- A. Degenerative disease
- B. Autoimmune disease
- C. Infectious disease
- D. Metabolic disease

### **2. What is a characteristic symptom of rheumatoid arthritis?**

- A. Morning stiffness lasting less than 10 minutes
- B. Pain only at night
- C. Morning stiffness lasting more than 1 hour
- D. Sudden paralysis

### **3. What is the main pathological process in RA?**

- A. Bone fracture
- B. Muscle rupture
- C. Synovial inflammation (synovitis)
- D. Nerve compression

**4. Which lab finding is commonly elevated in active RA?**

- A. ESR and CRP
- B. Platelets only
- C. Sodium
- D. Potassium

**5. What is the primary goal of RA treatment?**

- A. Cure infection
- B. Increase weight
- C. Improve vision
- D. Control inflammation and prevent joint damage

**6. Which joint involvement pattern is typical in RA?**

- A. Symmetrical small joints
- B. Asymmetrical joints
- C. Spine only
- D. Lower limb only

**7. Which imaging finding suggests early RA?**

- A. Bone fracture
- B. Complete joint destruction
- C. Soft tissue swelling
- D. Dislocation

**8. Which feature suggests RA become more severe ?**

- A. Decreased pain
- B. Normal ESR
- C. Increased appetite
- D. Increased joint swelling and stiffness

**9. A 35-year-old female presents with symmetrical joint pain in hands for 3 months, morning stiffness >1 hour, and fatigue. What is the MOST likely diagnosis?**

- A. Osteoarthritis
- B. Rheumatoid arthritis
- C. Gout
- D. Fibromyalgia

**10. A long-standing RA patient presents with deformity of fingers (ulnar deviation). What caused this?**

- A. Trauma
- B. Chronic synovial inflammation
- C. Infection
- D. Nerve injury

11. Which of the following is NOT cause of R.A?

- A. Obesity
- B. Advanced age
- C. Genetic factors
- D. **Helicobacter pylori infection**

12. Which of the following is manifestation not related to joint of R.A?

- A. Joint pain
- B. Swelling joint
- C. Warm joint
- D. **Dry eye and mouth**

13. list JOINT manifestation of R.A.

- Joint pain, tenderness
- Red joint
- Joint swelling
- Joint stiffness
- Numbness of hand and feet

14. list manifestation of R.A (not including joint)

- Disturbed sleep
- Low grade fever
- Fatigue
- Depression and mood change
- Dry eye and mouth
- Weight loss

15. List nursing managements of R.A.

- Pain relief measures
- Relief of fatigue
- Increasing mobility
- Monitoring for potential complications
- Increase knowledge of disease
- Promoting self-care

## **Ortho: 4- Osteomyelitis & Osteoporosis**

1. What is the most common cause of osteomyelitis?

- A. Viral infection
- B. Fungal infection only
- C. **Bacterial infection**
- D. Autoimmune reaction

2. The most common organism causing osteomyelitis is:

- A. E.coli
- B. **Staphylococcus aureus**
- C. Pseudomonas
- D. Streptococcus pneumoniae



**3. Which route of infection is most common in osteomyelitis?**

- A. Direct trauma only
- B. Hematogenous spread**
- C. Food contamination
- D. Airborne spread

**4. Which of the following is NOT manifestations of Osteomyelitis?**

- A. Drainage of pus from affected bone
- B. Bone pain
- C. Swelling of affected area
- D. Heart burn**

**5. A classic symptom of osteomyelitis is:**

- A. Local bone pain with fever**
- B. Headache
- C. Chest pain
- D. Abdominal pain

**6. Which complication may result from untreated osteomyelitis?**

- A. Stroke
- B. Cataract
- C. Gastritis
- D. Bone necrosis and chronic infection**

**7. Osteoporosis is characterized by:**

- A. Increased bone mass
- B. Decreased bone density and strength**
- C. Bone infection
- D. Joint inflammation

**8. The most common complication of osteoporosis is:**

- A. Infection
- B. Paralysis
- C. Fracture**
- D. Tumor

**9. Which population is most at risk of osteoporosis?**

- A. Young males
- B. Postmenopausal women**
- C. Children
- D. Athletes

**10. A major nursing intervention in osteoporosis is:**

- A. Bed rest only
- B. Fall prevention**
- C. Fluid restriction
- D. Oxygen therapy

**11. Which medication increases bone resorption and leads to osteoporosis?**

- A. Long-term corticosteroids**
- B. Antibiotics
- C. Antacids
- D. Insulin

**12. Which hormone helps protect bone density?**

- A. Estrogen**
- B. Adrenaline
- C. Insulin
- D. Prolactin

**13. What is the main goal of osteoporosis treatment?**

- A. Cure infection
- B. Prevent fractures**
- C. Increase fever
- D. Improve vision

**14. Which vitamin deficiency is strongly linked to osteoporosis?**

- A. Vitamin A
- B. Vitamin B12
- C. Vitamin D**
- D. Vitamin K only

**15. What is the main principle of hyperbaric oxygen therapy?**

- A. Delivering oxygen at normal atmospheric pressure only
- B. Providing pure oxygen at higher-than-normal atmospheric pressure in a closed chamber**
- C. Reducing oxygen supply to tissues
- D. Using carbon dioxide to improve breathing

**16. list clinical manifestations of osteomyelitis**

- **Bone pain**
- **Fever and chills**
- **Redness and swelling of affected part**
- **Lack of energy**
- **Irritability**
- **Drainage of pus from affected bone**
- **Inability to move affected limb**

**17. list nursing diagnosis of osteomyelitis**

- Acute pain related to infection
- Impaired physical mobility related to pain
- Risk for sepsis
- Risk for impaired skin integrity

**18. list nursing managements of osteoporosis.**

- Fall prevention
- Administer medications as prescribed
- Encourage calcium-rich diet and exercise
- Educate on medication adherence

**GIT: 1- Peptic ulcer**

**1. The most common cause of peptic ulcer disease is:**

- A. Stress only
- B. *Helicobacter pylori* infection
- C. High calcium diet
- D. Viral infection

**2. Another important cause of peptic ulcer disease is:**

- A. Antihypertensives
- B. NSAIDs use
- C. Antibiotics
- D. Insulin

**3. A common symptom of peptic ulcer is:**

- A. Burning epigastric pain
- B. Chest pain only
- C. Leg pain
- D. Headache

**4. pain of duodenal ulcer is typically:**

- A. Worse after eating
- B. Only at night always
- C. Relieved by eating
- D. Not related to food

**5. Which complication is most dangerous in peptic ulcer?**

- A. Hair loss
- B. Perforation
- C. Diarrhea
- D. Constipation

**6. The diagnostic test of choice for peptic ulcer is:**

- A. Endoscopy
- B. ECG
- C. MRI brain
- D. X-ray chest

**7. Which drug is commonly used to reduce gastric acid?**

- A. Antibiotics only
- B. Antiviral drugs
- C. Proton pump inhibitors (PPIs)
- D. Insulin

**8. A complication of long-term NSAID use is:**

- A. Peptic ulcer
- B. Asthma
- C. Hypertension
- D. Cataract

**9. A key nursing advice for peptic ulcer patients is:**

- A. Skip meal
- B. Avoid NSAIDs and spicy food
- C. Increase smoking
- D. Bed rest only

**10. Which type of ulcer is more likely to cause weight loss?**

- A. Duodenal ulcer
- B. Gastric ulcer
- C. Both equally
- D. Neither

**11. A patient with peptic ulcer has sudden severe abdominal pain and rigid abdomen. What is the most likely complication?**

- A. Gastritis
- B. Constipation
- C. Appendicitis
- D. Perforation

**12. Which sign may indicate GIT bleeding in peptic ulcer?**

- A. Hematuria
- B. Melena
- C. Jaundice
- D. Rash

13. Which of the following increases risk of peptic ulcer?

- A. Regular exercise
- B. High water intake
- C. Smoking
- D. Fruit diet

14. The main action of proton pump inhibitors is:

- A. Increase acid secretion
- B. Block gastric acid production
- C. Kill bacteria directly
- D. Increase digestion speed

15. Which symptom is more typical in gastric ulcer than duodenal ulcer?

- A. Pain worsened by food
- B. Pain relieved by food
- C. No pain at all
- D. Only night pain

16. A patient with peptic ulcer is vomiting blood. This condition is called:

- A. Hemoptysis
- B. Hematemesis
- C. Hematuria
- D. Melena

17. Which bacteria test confirms H. pylori infection?

- A. ECG
- B. EEG
- C. MRI
- D. Stool antigen test

18. Differentiate between gastric and duodenal ulcer in relation to pain

| Duodenal Ulcers                                                                                                                                                                                                                                                           | Gastric Ulcers                                                                                                                                                                                                                                             |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <ul style="list-style-type: none"><li>• Pain relieved by meal.</li><li>• Occurs 2-3 hrs after meal.</li><li>• Most common type.</li><li>• Dark, Tarry Stools (Melena) Occur</li></ul>  | <ul style="list-style-type: none"><li>• Pain increased by meal.</li><li>• Occurs 30m to 1hr after meal.</li><li>• Not as common.</li><li>• Vomiting occurs.</li></ul>  |

**19. list complications of peptic ulcer.**

- Bleeding
- Perforation
- Obstruction

**20. list nursing interventions for patient teaching about peptic ulcer**

- Instruct patient about importance of continuing taking medication even after signs and symptoms have decreased.
- Avoid smoking
- Rest and stress reduction.
- Awareness of complications
- Follow up care

## **GIT: 2- Cholelithiasis & Cholecystitis**

**1- Cholelithiasis means:**

- A. Infection of liver
- B. Gallstones in the gallbladder
- C. Kidney stones
- D. Pancreatic inflammation

**2. The most common type of gallstones is:**

- A. Calcium stones
- B. Protein stones
- C. Iron stones
- D. Cholesterol stones

**3. The classic symptom of gallstones is:**

- A. Right upper quadrant pain after fatty meals
- B. Left chest pain
- C. Headache
- D. Knee pain

**4. Cholecystitis is:**

- A. Liver infection
- B. Inflammation of the gallbladder
- C. Kidney infection
- D. Stomach ulcer

**5. The best diagnostic tool for gallstones is:**

- A. X-ray chest
- B. Abdominal ultrasound
- C. ECG
- D. MRI brain

**6. A diet that increases gallstone symptoms is:**

- A. High fat diet
- B. High fiber diet
- C. Low fat diet
- D. Water diet

**7. Treatment of choice for symptomatic gallstones is:**

- A. Antibiotics only
- B. Cholecystectomy
- C. Insulin
- D. Chemotherapy

**8. Which organ is affected in cholecystitis?**

- A. Liver
- B. Gallbladder
- C. Stomach
- D. Pancreas

**9. Which condition increases risk of gallstones?**

- A. Obesity
- B. Anemia
- C. Asthma
- D. Hypertension

**10. Which of the following is NOT a risk factor for gallstones?**

- A. Obesity
- B. Fasting
- C. Rapid weight loss
- D. High fiber diet

**11. Radiation of gallbladder pain is commonly to:**

- A. Left arm
- B. Right shoulder/scapula
- C. Left leg
- D. Neck only

**12. patient with cholecystitis may present with:**

- A. Fever and nausea
- B. Rash only
- C. Chest pain only
- D. Vision loss

**13. Which complication can occur if gallstone blocks common bile duct?**

- A. Jaundice
- B. Asthma
- C. Diabetes
- D. Migraine

**14. A 40-year-old obese female presents with RUQ pain after fatty meals. what is diagnosis?**

- A. Hepatitis
- B. Cholelithiasis
- C. Pancreatitis
- D. Gastritis

**15. List causes of gall bladder stones.**

- Obesity
- Cholesterol lowering drugs
- Diabetes
- Rapid weight loss
- Fasting
- Estrogen

**16. list post-operative nursing interventions for gall bladder stones**

- Relieving pain
- Improving respiratory status
- Maintaining skin integrity & promoting biliary drainage
- Improving nutritional status
- Monitoring and managing complications
- General health teaching

### **GIT: 3- liver cirrhosis & Esophageal varices**

**1. liver cirrhosis is defined as:**

- A. Acute liver inflammation
- B. Chronic liver damage with fibrosis and nodules
- C. Liver infection only
- D. Increased liver size only

**2. The most common cause of liver cirrhosis worldwide is:**

- A. Trauma
- B. Chronic viral hepatitis
- C. Asthma
- D. Kidney disease



**3. Another major cause of cirrhosis is:**

- A. Alcohol abuse
- B. Exercise
- C. Low salt diet
- D. Vitamin deficiency

**4. Common complication of cirrhosis is:**

- A. Asthma
- B. Migraine
- C. Diabetes
- D. Portal hypertension

**5. Portal hypertension can lead to:**

- A. Hypertension only
- B. Diabetes
- C. Esophageal varices
- D. Pneumonia

**6. Esophageal varices are:**

- A. Gastric ulcers
- B. Dilated veins in the esophagus
- C. Liver tumors
- D. Intestinal obstruction

**7. The most dangerous complication of esophageal varices is:**

- A. Hematemesis
- B. Constipation
- C. Fever
- D. Headache

**8. Patient with cirrhosis presents with abdominal distension due to fluid. This is:**

- A. Edema
- B. Ascites
- C. Tumor
- D. Obesity

**9. Late complication of cirrhosis affecting brain function is:**

- A. Stroke
- B. Migraine
- C. Hepatic encephalopathy
- D. Seizures only

**10. What causes esophageal varices in cirrhosis?**

- A. Increased portal pressure
- B. Infection
- C. Low sugar
- D. High calcium

**11. Which substance accumulates in hepatic encephalopathy?**

- A. Glucose
- B. Ammonia
- C. Calcium
- D. Oxygen

**12. Which sign indicates severe liver disease?**

- A. Mild fatigue
- B. Mild pain
- C. Headache
- D. Jaundice & ascites

**13. A patient with cirrhosis should avoid:**

- A. Alcohol
- B. Water
- C. Fruits
- D. Walking

**14. Which of the following is NOT complication of liver cirrhosis?**

- A. Hepatic encephalopathy
- B. Portal hypertension
- C. Melena
- D. Bone necrosis

**15. List clinical manifestation of liver cirrhosis.**

- Ascites
- Jaundice
- Weight loss
- Epistaxis
- Clubbing fingers
- Fever

**16. list complications of liver cirrhosis**

- Recurrent infection
- Hepatic encephalopathy
- Lower limb edema
- Ascites
- Portal hypertension
- Jaundice
- Liver cancer

**17. list nursing interventions for patient with liver cirrhosis**

- Promoting rest
- Improving Nutritional Status:
- Providing Skin Care:
- Reducing risk of injury:
- Teaching Patients Self-Care

**18. List nursing interventions for patient with esophageal varices**

- Monitor vital signs every 15–30 minutes
- Maintain NPO status until cleared
- Establish large-bore IV access for fluids/blood transfusion
- Administer blood products as ordered
- Position patient in high fowler or side-lying to reduce aspiration risk
- Administer medications :
  - vasopressin to reduce portal pressure
  - Lactulose if hepatic encephalopathy is present
- Prepare for emergency endoscopy
- Monitor for signs of hypovolemic shock: ↓ BP, ↑ HR, cold/clammy skin

**GIT: 4- Hepatitis**

**1. Which type of Hepatitis is transmitted through blood?**

- A. Hepatitis A
- B. **Hepatitis B**
- C. Hepatitis E
- D. Hepatitis A and E

**2. Which of the following is a common symptom of Hepatitis?**

- A. Headache only
- B. Chronic cough
- C. **Yellowing of the skin and eyes**
- D. Ear pain

**3. Which type of Hepatitis is commonly transmitted through contaminated food or water?**

- A. Hepatitis B
- B. Hepatitis C
- C. **Hepatitis A**
- D. Hepatitis D

**4. What is the primary mode of transmission of Hepatitis A?**

- A. **Fecal-oral route**
- B. Blood transfusion
- C. Sexual contact
- D. Airborne droplets

**5. Which of the following Hepatitis viruses has a vaccine available?**

- A. Hepatitis A only
- B. Hepatitis B only
- C. Hepatitis A and B
- D. Hepatitis C

**6. Which Hepatitis virus is most likely to cause acute, self-limited infection?**

- A. Hepatitis A
- B. Hepatitis B
- C. Hepatitis C
- D. Hepatitis D

**7. Which of the following is a common route of Hepatitis B transmission?**

- A. Contaminated water
- B. Blood and body fluids
- C. Airborne spread
- D. Insect bites

**8. Serious complication of chronic Hepatitis is:**

- A. Asthma
- B. Diabetes
- C. Liver cirrhosis
- D. Hypertension

**9- List 3 modes of transmission of hepatitis A**

- Fecal oral route
- Contaminated foods or water
- Hands or other objects contaminated with infected feces
- Sexual intercourse (oral –anal)
- Undercooked shellfish like oysters, mussels, and clams

**10- List 4 manifestations of hepatitis B.**

- Loss of appetite
- Jaundice
- Fever
- Dark urine
- Pale stool
- Nausea, vomiting
- Headache, fatigue
- Joint pain.

**11- List 4 modes of transmission of hepatitis B , C**

- Infected blood
- Unsafe infected syringe
- Unprotected sexual contact
- From mother to baby during childbirth

## Neuro : 1- Disc prolapse

- 1- Which part of the spine is most commonly affected?
  - A. Cervical spine
  - B. Thoracic spine
  - C. Lumbar spine
  - D. Sacral spine
- 2- What is the main cause of disc prolapse?
  - A. Viral infection
  - B. Trauma and degeneration
  - C. Poor diet
  - D. Allergy
- 3- Which symptom is most common?
  - A. Cough
  - B. Radiating limb pain
  - C. Skin rash
  - D. Fever
- 4- Sciatica is associated with:
  - A. Lumbar disc prolapse
  - B. Cervical disc prolapse
  - C. Thoracic disc prolapse
  - D. Sacral fracture
- 5- Which nerve is commonly affected in lumbar disc prolapse?
  - A. Facial nerve
  - B. Sciatic nerve
  - C. Vagus nerve
  - D. Optic nerve
- 6- What is a common risk factor?
  - A. Regular exercise
  - B. Physical trauma
  - C. Adequate sleep
  - D. Hydration
- 7- Which of the following is part of conservative treatment?
  - A. Bed rest and physiotherapy
  - B. Antibiotics
  - C. Insulin
  - D. Antivirals
- 8- When is surgery indicated?
  - A. Mild pain
  - B. No symptoms
  - C. Severe neurological deficit
  - D. Headache

**9- Which part of the intervertebral disc herniates in disc prolapse?**

- A. Annulus fibrosus
- B. **Nucleus pulposus**
- C. Vertebral body
- D. Spinal ligament

**10- Which level is most commonly affected in lumbar disc prolapse?**

- A. L1–L2
- B. C1–C2
- C. **L4–L5**
- D. T12–L1

**11- Straight Leg Raise (SLR) test is used to detect:**

- A. Cervical Prolapse
- B. **Lumbar nerve root irritation**
- C. Shoulder injury
- D. Hip fracture

**12- What is the role of physiotherapy?**

- A. Worsens condition
- B. **Strengthens muscles and reduces pain**
- C. Causes infection
- D. Replaces surgery completely

**13- Which drug class is commonly used for pain relief?**

- A. Antibiotics
- B. Antivirals
- C. Antifungals
- D. **NSAIDs**

**14- Which clinical sign is positive in lumbar disc herniation?**

- A. Murphy's sign
- B. Kernig's sign
- C. **Straight Leg Raise test**
- D. Tinel's sign

**15- Which investigation is most sensitive for soft diagnosis of disc prolapse?**

- A. CT scan
- B. X-ray
- C. **MRI**
- D. Ultrasound

**16- Which surgical procedure is commonly performed?**

- A. Appendectomy
- B. **Discectomy**
- C. Cholecystectomy
- D. Nephrectomy

**17- Which symptom indicates urgent surgical need?**

- A. Mild pain
- B. Intermittent discomfort
- C. **Progressive neurological deficit**
- D. Occasional stiffness

**18- Which preventive measure is most effective?**

- A. Proper posture and exercise
- B. Heavy lifting
- C. Smoking
- D. Prolonged sitting

**19- Which of the following surgical procedures involve removal of part of the vertebral bone?**

- A. Discectomy
- B. Removal of the herniated disc
- C. Endoscopic discectomy
- D. Laminectomy

**20- Which of the following surgical procedures involve removal of the herniated portion of the disc using endoscope?**

- A. Discectomy
- B. Endoscopic discectomy
- C. Laminectomy
- D. Open surgery to remove disc

**21- List 4 causes of disc prolapse**

- Degenerative disc disease ; age-related wear and tear
- Trauma or injury ;e.g., lifting heavy objects improperly
- Repetitive strain or twisting motions
- Obesity ; increased spinal load
- Poor posture
- Sedentary lifestyle
- Genetic predisposition

**22- List 3 manifestations of cervical disc prolapse**

- Neck pain
- Shoulder radiation
- Numbness and tingling in upper limb

**23- List 3 manifestations of thoracic disc prolapse**

- Pain in mid back and chest
- Abdominal pain
- Numbness and sensory loss around trunk

**24- List 4 manifestations of lumbar disc prolapse**

- Low back pain
- Numbness and tingling in lower limb
- loss of bladder control
- Sciatica leg pain

**25- List items that should be assessed during neurological examination for patient with lumbar disc prolapse.**

- Reflexes
- Muscle strength
- Sensory deficits
- Straight leg raise test

**26- List 5 non-surgical management for disc prolapse**

- Bed rest (short-term) – avoid prolonged immobility
- NSAIDs (e.g., ibuprofen)
- Muscle relaxants
- Oral steroids or epidural steroid injections
- Hot/cold compresses
- Activity modification (avoid lifting, bending)
- Physiotherapy :as core strengthening, stretching exercises and postural training

**27- Enumerate 5 nursing assessment for patient with disc prolapse.**

- Assess s&s ; complete assessment of pain
- Observe for numbness, tingling, muscle weakness
- Monitor gait and mobility
- Check for signs of bowel/bladder dysfunction
- assess impact on ADLs (activities of daily living)

**28- List 4 nursing diagnosis of disc prolapse**

- Acute Pain related to nerve compression
- Impaired physical mobility related to pain and discomfort
- Anxiety related to limited mobility
- Deficient knowledge related to condition and self-care
- Risk for injury related to muscle weakness or sensory deficits

**29- Enumerate 5 nursing management for patient with disc prolapse**

**For Pain Management**

- Apply hot/cold compresses
- Encourage use of firm mattress or back support
- Position patient with knees slightly bent (lumbar cases)
- Administer analgesics, NSAIDs, muscle relaxants as prescribed

**For Mobility support**

- Encourage mobilization as tolerated
- Assist with physical therapy
- Teach log-rolling technique when turning in bed
- Use of braces or lumbar support belts if prescribed

**To reduce anxiety**

- Allow patient to express feeling.
- Explanation about diagnosis, treatment and medication.
- Good interaction between patient and nurse.
- Encourage the family to participate in patient care and emotional support

**General education**

- Teach proper body mechanics: bend at knees, not waist
- Avoid heavy lifting, prolonged sitting, or twisting
- Encourage weight loss if overweight
- Emphasize importance of exercise, core strengthening



## • **Neuro : 2- Fibromyalgia**

**1- What is fibromyalgia characterized by?**

- A. Joint inflammation
- B. **Widespread musculoskeletal pain**
- C. Bone fractures
- D. Muscle rupture

**2. Fibromyalgia is more common in:**

- A. Men
- B. **Women**
- C. Children only
- D. Elderly men only

**3- Which of the following is a common symptom?**

- A. High fever
- B. Bleeding
- C. **Fatigue**
- D. Rash

**4- Fibromyalgia pain is typically:**

- A. **Widespread**
- B. Localized
- C. Only in joints
- D. Only in bones

**5- Which of the following is associated with fibromyalgia?**

- A. Improved sleep
- B. **Sleep disturbances**
- C. Increased appetite
- D. Hair loss

**6- Which condition commonly coexists with fibromyalgia?**

- A. Diabetes
- B. **Depression**
- C. Asthma
- D. Hypertension

**7- How is fibromyalgia diagnosed?**

- A. Blood test
- B. X-ray
- C. **Clinical criteria**
- D. Biopsy

**8- Which medication is commonly used?**

- A. Antibiotics
- B. NSAIDs only
- C. **Antidepressants**
- D. Antivirals

**9- Which lifestyle change helps management?**

- A. Complete rest
- B. **Regular exercise**
- C. Smoking
- D. High sugar diet

**10- Which symptom worsens fibromyalgia pain?**

- A. Regular exercise
- B. **Stress**
- C. Good sleep
- D. Hydration

**11- Which of the following is commonly normal in fibromyalgia?**

- A. MRI
- B. ESR
- C. CRP
- D. **All of the above**

**12- Which diagnostic tool is used along with widespread pain index (WPI)?**

- A. ESR
- B. **Symptom Severity Scale (SSS)**
- C. X-ray
- D. CT scan

**13- Which of the following is TRUE about lab findings?**

- A. Elevated CRP
- B. Positive rheumatoid factor
- C. **Normal lab results**
- D. High WBC count

**14- Which symptom cluster defines fibromyalgia besides pain?**

- A. Fever and rash
- B. **Fatigue, sleep disturbance, cognitive dysfunction**
- C. Bleeding and bruising
- D. Paralysis

**15- Which patient description fits fibromyalgia best?**

- A. Local joint swelling with redness
- B. **Chronic widespread pain with normal labs**
- C. Acute infection with fever
- D. Bone deformities

**16- Which management principle is essential?**

- A. Focus on cure only
- B. **Patient education and self-management**
- C. Immediate surgery
- D. Antibiotics

**17- Which environmental factor can trigger flares?**

- A. Warm weather
- B. **Emotional stress**
- C. Balanced diet
- D. Exercise

**18- List causes of fibromyalgia**

- **Sleep disturbance**
- **Hormonal imbalance**
- **Genetic**
- **Psychological factors**
- **Infection**

- Physical stress
- Emotional stress

#### 19- Enumerate diagnostic criteria for fibromyalgia

1. Generalized pain in 4 or 5 regions of body
2. Symptoms at similar level for at least 3 months
3. Widespread pain index more than 7 and severity of symptoms more than 5

#### 20- List non pharmacological management for fibromyalgia

- Physical therapy; exercise programs, Avoid overexertion – balance activity and rest
- Stress management techniques; deep breathing exercises, meditation
- Massage, warm bath, heat therapy
- Encourage regular bedtime routine
- Limit caffeine and electronics before bed
- Provide information on sleep hygiene
- Provide reassurance and active listening
- Refer to mental health services if needed
- Encourage support group participation
- Educate on chronic nature of disease and need for lifestyle changes
- Reinforce importance of stress reduction
- Discuss medication side effects and adherence

### Neuro : 3- Epilepsy

#### 1- What is a seizure?

- A. Sudden increase in heart rate
- B. **Abnormal electrical activity in the brain**
- C. Muscle growth
- D. Bone fracture

#### 2- Which of the following is a common symptom of a generalized seizure?

- A. **Loss of consciousness**
- B. Hair loss
- C. Increased appetite
- D. Sneezing

#### 3- Which type of seizure involves only one part of the brain?

- A. Generalized seizure
- B. **Focal seizure**
- C. Febrile seizure
- D. Tonic seizure

#### 4- What is an aura in epilepsy?

- A. A type of medication
- B. **A warning sign before a seizure**
- C. A cure for epilepsy
- D. A surgical procedure

**5- Which of the following can trigger seizures in some people with epilepsy?**

- A. Sleep deprivation
- B. Flashing lights
- C. Stress
- D. All of the above

**6- What is the most common treatment for epilepsy?**

- A. Antibiotics
- B. Antiepileptic drugs
- C. Painkillers
- D. Vitamins

**7- What should you do if someone is having a seizure?**

- A. Put something in their mouth
- B. Hold them down tightly
- C. Turn them on their side and keep them safe
- D. Give them water immediately

**8- Which imaging modality is most useful for identifying structural brain abnormalities in epilepsy?**

- A. X-ray
- B. CT scan
- C. MRI
- D. Ultrasound

**9- Which of the following is NOT a common cause of epilepsy?**

- A. Head trauma
- B. Genetic factors
- C. Brain tumors
- D. Hypertension alone

**10- Which seizure type involves sudden muscle stiffening?**

- A. Clonic seizure
- B. Atonic seizure
- C. Tonic seizure
- D. Absence seizure

**11- Which seizure type is associated with sudden loss of muscle tone and falls?**

- A. Tonic seizure
- B. Clonic seizure
- C. Atonic seizure
- D. Myoclonic seizure

**12- Which surgical option may be considered for drug-resistant epilepsy?**

- A. Appendectomy
- B. Lobectomy
- C. Nephrectomy
- D. Thyroidectomy

**13- Which of the following the nurse SHOULD NOT do it during seizure?**

- A. Place the patient in a side-lying position
- B. Clear the area of harmful objects
- C. Restrain the patient
- D. Monitor breathing and skin color

**14- Which of the following the nurse SHOULD NOT do it in seizure?**

- A. Maintain airway and keep the patient in the side-lying position
- B. Check vital signs
- C. Allow the patient to rest or sleep
- D. Give oral fluid to patient

**15- List 5 risk factors for epilepsy in adult**

- Trauma
- Brain tumors
- Stroke
- Infections
- Neurodegenerative diseases

**16- List three types of generalized epilepsy**

- Tonic
- A tonic
- Tonic clonic
- Myoclonic

**17- List nursing interventions for aura phase of seizure**

- Observe for early warning signs (e.g., dizziness, unusual smells, anxiety, visual changes)
- Help the patient sit or lie down in a safe place
- Remove nearby dangerous objects (sharp or hard items)
- Loosen tight clothing, especially around the neck
- Stay with the patient (do not leave them alone)
- Prepare emergency equipment (oxygen, suction)
- Provide reassurance and reduce anxiety

**18- List nursing interventions during seizure**

- Ensure patient safety first
- Place the patient in a side-lying position (recovery position) to prevent aspiration
- Clear the area of harmful objects
- Do NOT restrain the patient
- Do NOT put anything in the patient's mouth
- Time the duration of the seizure
- Monitor breathing and skin color (watch for cyanosis)
- Do not give food or fluids during the seizure

**19- List nursing interventions after seizure**

- Maintain airway and keep the patient in the side-lying position
- Check vital signs (pulse, respiration, blood pressure)
- Assess level of consciousness (confusion and drowsiness are common)

- Reassure and reorient the patient
- Check for injuries (e.g., tongue biting, head trauma)
- Allow the patient to rest or sleep
- Document: Duration of seizure ,Type of movements ,Presence of aura
- Administer prescribed medications if ordered, such as Diazepam
- Clear secretions if needed (suction if necessary)

## **20- List 5 signs of aura phase of seizure**

- Strange or unusual smells (e.g., burning rubber)
- Odd tastes in the mouth
- Tingling sensation
- Visual disturbances (flashing lights, blurred vision)
- Hearing unusual sounds (buzzing, ringing)
- Sudden fear or anxiety
- Sweating
- Palpitations
- Dizziness
- Sudden confusion
- Difficulty speaking or understanding

## **Oral questions (just examples)**

### **1- Give the rational of:**

- Presence of jaundice in patient with gallbladder stone
- Presence of dark urine in patient with gallbladder stone
- Presence of pale stool in patient with gallbladder stone
- Presence of dark stool in patient with peptic ulcer
- giving vit B12 and iron for patient with peptic ulcer

### **2- Discuss the complications of peptic ulcer**

### **3- Discuss hyperbaric oxygen therapy in patient with osteomyelitis**

### **4- Manifestation of rheumatic arthritis**

### **5- What is human's sign in patient with hip replacement?**

### **6- Nursing interventions for disc prolapse**

### **7- What are the neurological tests for patient with disc prolapse?**

### **8- Discuss the diagnostic criteria for fibromyalgia**