

Management of patients with Gastrointestinal diseases

Dr . Bardies shaban





الحديث الجانبي يشتت الانتباه




من فضلك اغلق الهاتف


3- Hepatitis

- ◉ Definition
- ◉ Types
- ◉ Manifestations
- ◉ Diagnosis
- ◉ Complications
- ◉ Managements


Hepatitis: inflammation and damage of the liver cells,
Caused by viral infection (**viral hepatitis**) , exposure to alcohol,
or toxic chemical substances or disorder in the immune system.

HEPATITIS






Hepatitis A



- Mild infection, resolve within 2 months without leaving long-term effects
- Also called (infectious jaundice).
- Once the person had the infection, cannot get it again.
- The mortality rate is approximately 0.5% for patients under 40 years and 1-2% for patients over 40 years.



Mode of transmission:

- Fecal oral route
- Contaminated foods or water
- Hands or other objects contaminated with infected feces
- Sexual intercourse (oral –anal)
- Undercooked shellfish like oysters, mussels, and clams

Methods of Transmission of Hepatitis A & E

Contaminated Food & Water



Fecal-Oral Route

Feces to Mouth




Poor Sanitation

Lack of Clean Water & Hygiene



Person-to-Person Contact

Close Personal Contact



Clinical Manifestations of Hepatitis A & E



Fever



Nausea & Vomiting



Jaundice



Joint Pain



Loss of Appetite



Dark Urine



Fatigue



Abdominal Pain



Hepatitis A Virus



Hepatitis E Virus



Abdominal Pain



Legs

PREVENTION OF HEPATITIS A & E



Get Vaccinated



Practice Good Hygiene



Drink Safe Water



**Eat Clean and
Cooked Food**



Avoid Street Food



Improve Sanitation

Hepatitis E

Prevention is the key





**Drinking
clean water**



**Clean practices like
washing hands**




**Eating
healthy food**

Nursing management




- ▶ Provide **gradual** activity and **rest** to prevent fatigue.
- ▶ Small **frequent meals**, high **protein** diet.
- ▶ Monitor **fluid** balance.
- ▶ If anorexia and vomiting persists, **enteral** feeding is recommended.
- ▶ Educate patient to **seek medical care** if symptoms persists or worsen.
- ▶ Drink **plenty of fluids** to avoid or prevent dehydration. **(I&O)**
- ▶ Avoid **alcoholic** beverages because it worsens the effect of the virus.

Hepatitis B




- ▶ Hepatitis B virus (HBV) is the major cause of **cirrhosis** and **hepatocellular carcinoma**
- ▶ It may be either **acute** (self- limiting) or **chronic** (long standing infection).
- ▶ People with self-limiting infection clear the infection **spontaneously** within weeks to months.

SYMPTOMS OF HEPATITIS B




SIGNS OF HEPATITIS B

- ✓ Fatigue
- ✓ Nausea
- ✓ Fever
- ✓ Vomiting
- ✓ Muscle aches
- ✓ Abdominal pain
- ✓ Loss of appetite
- ✓ A general feeling of being unwell



Prevention

- Blood test of donors or blood transfusions.
- Use of disposable syringes and needles
- Good personnel hygiene.
- Use protective barriers as gloves when dealing with body fluids.
- Immunization (HB vaccine)




Hepatitis C

- ▶ It was known as **none A, non B** hepatitis.
- ▶ Between age **40 to 59** years.
- ▶ The infection can now be **cured**
- ▶ **15 to 25** percent of people with HCV clear the infection **without** treatment



Clinical manifestations

How Hepatitis C Affects the Body



Prevention

- ❑ Cover cuts and blisters
 - ❑ Properly dispose of any used bandages, tissues, or anything containing blood
 - ❑ Wash hands or objects that have come in contact with blood
 - ❑ Clean spilled blood on surfaces
- 
- ❑ Don't share personal items that have blood on it
 - ❑ Do not breastfeed if your nipples become cracked and bleed
 - ❑ Do not donate blood, sperm, or organs

HAV

Hepatitis A virus
acute infection

HBV

Hepatitis B virus
acute or chronic infection

HCV

Hepatitis C virus
acute or chronic infection
(usually chronic)

Vaccine



YES



YES



NO

Transmission



Feces
(contaminated food,
water, objects)
**Direct person-
to-person contact**



**Blood, certain
body fluids,**
childbirth, sexual
contact, sharing
used needles
or syringes



Blood,
sharing used
needles or
syringes,
needlestick
injuries

Treatment



**Rest, fluids,
nutrition**



**Rest, fluids,
nutrition** (acute)
Medications
(chronic)



Medications
can cure over
95% of people
with this virus

Complications



Rare



**Cirrhosis,
liver failure,
liver cancer**




**Cirrhosis,
liver failure,
liver cancer**

Hepatitis D

- ▶ It is a small virus that requires concomitant **infection** with **HBV** to survive.
- ▶ HDV **cannot survive on its own** because it requires a protein that the HBV makes to enable it to infect liver cells.
- ▶ Individuals who **already** have chronic **HBV** infection can acquire HDV infection at the same time.

Mode of transmission

- ▶ Shared **needles** among drug abusers
- ▶ **Contaminated** blood
- ▶ **Sexual** contact
- ▶ **No vaccine** for hepatitis D is available. However, HBV vaccine can prevent an individual not infected with HBV from contracting hepatitis D because hepatitis D virus requires live HBV to replicate in the body.



Summary



- List 3 mode of transmission of hepatitis A
- List 4 manifestations of hepatitis B.
- Differentiate between hepatitis A and hepatitis C



Thank
you