



Lecture 4

Normal Flora & Opportunistic Infections




Dr Rania Hamed


Lecturer of Microbiology

2025 – 2026

GOOD BACTERIA



BAD BACTERIA



Lecture Learning Outcomes

01

Define Normal Flora

Understand the concept of normal flora and explain its role in human health and disease prevention.

02

Identify Key Sites

Recognize the main **anatomical locations** where normal flora colonizes the human body.

03

Differentiate Flora Types

Distinguish between **resident flora** that permanently colonizes the body and **transient flora** that temporarily inhabits body surfaces.

04

Explain Benefits and Risks

Describe both the **protective functions** and **pathogenic role** of normal flora under certain conditions.

05

Apply Clinical Knowledge

Understand the nurse's role in maintaining microbial balance and **preventing opportunistic infections** in patient care.

Introduction to Microorganisms

- ❑ **Microorganisms** are microscopic living organisms that exist all around us and within us. They include **bacteria, fungi, and protozoa.**
- ❑ These tiny life forms play important roles in **ecosystems, human health, and disease processes.**
- ❑ Understanding microbiology is **fundamental to nursing practice and infection control.**

Pathogenic vs Non-Pathogenic Microbes


Non-Pathogenic Microbes

- Do not cause disease in healthy hosts
- Often beneficial to human health
- Comprise most of normal flora
- Support vital body functions

Pathogenic Microbes

- Capable of causing disease
- overcome host defenses
- Produce toxins or damage tissues
- Require specific treatment

 **Think: Not all microbes are harmful! Most microorganisms in and on our bodies are actually helpful partners in health.**



Definition of Normal Flora

Normal flora is the mixture of microorganisms primarily bacteria and fungi—that are regularly found at specific anatomical sites of the human body.

- ❑ These microorganisms establish residence on residence on body surfaces exposed to the the external environment, including **skin, mucous membranes, and the gastrointestinal gastrointestinal tract.**

Normal flora represents a complex ecosystem that exists **in balance** with the human host.

We Are More Microbe Than Human

"We're all just petri dishes with shoes."

30- 40T

Human Cells


**Approximate number of cells that make
make up the human body**

38- 40T

Microbial Cells

**Microbes living in and on the human
body—outnumbering human cells**

Microbes are an essential part of our biological identity and contribute significantly to our health, immunity, and metabolism.



Resident vs Transient Flora

Resident Flora

- Permanently colonize the body
- Stable and fixed population
- Acquired during and shortly after birth
- Reestablish quickly if disturbed
- Provide continuous protective benefits

Transient Flora

- Temporarily inhabit the body
- Present for hours to months
- Acquired from the environment
- Do not permanently establish

Development of Normal Flora

Before Birth

Fetus is sterile in the uterus, protected by placental barrier and amniotic sac.

1

First Days

Skin contact, feeding, and environmental exposure introduce diverse microorganisms.

3

During Birth

Rapid colonization begins as infant passes through birth canal, acquiring maternal vaginal and fecal flora.

2

Establishment

By age 2-3, normal flora with stable resident populations.

4

Normal Flora OF THE HUMAN BODY

Throat:

Staphylococcus epidermidis
Streptococci viridans
Streptococci pyogenes
Streptococci pneumonia
Haemophilus influenzae

Nose:

Staphylococcus aureus
Staphylococcus
epidermidis
Streptococci

Mouth:

Candida
Streptococci
Trichomonas tenax
Bacteroides
actinomyces

Intestine:

Lactobacillus
Clostridium
Escherichia coli

Skin:

Staphylococcus aureus
Staphylococcus epidermidis
Candida

Vagina:

Candida
Lactobacilli

Lab+
Mate


Normal Flora of the Skin

These microbes form a **protective barrier** against pathogenic invaders.

The skin is **the body's largest organ** and home to diverse microbial communities.

Common Skin Residents:

- *Staphylococcus epidermidis* – most abundant





Normal Flora of the Mouth

The oral cavity is one of the most populated microbial habitats in the body, with over 700 different bacterial species identified.

Common Oral Microbes

- *Streptococcus* species
- *Lactobacillus* species
- *Actinomyces* species
- *Neisseria* species

Functions

- Prevent colonization by pathogens
- Initiate digestion of carbohydrates
- Stimulate local immune responses

Normal Flora of the Gastrointestinal Tract

The **GI tract** contains the largest and most diverse microbial community in the human body, with trillions of organisms playing critical roles in **digestion, immunity, and health**.




Stomach

Low pH limits microbial growth; mainly acid-tolerant *Helicobacter* and *Lactobacillus* species.



Small Intestine

Moderate populations including *Lactobacillus*, *Streptococcus*, and some *Bacteroides*.



Large Intestine

Highest microbial density with *E. coli*, *Clostridium*, and *Bifidobacterium*

? Question: Why does the large intestine have more bacteria than the small intestine?

Normal Flora of the Respiratory Tract

Upper Respiratory Tract


Nose, pharynx, and throat harbor resident flora including:

- *Staphylococcus aureus*
- *Streptococcus* species

Lower Respiratory Tract

Lungs and bronchi are typically **sterile** due to mucus clearance and immune defenses.


Human Respiratory System



Watch this **short video** to see Nasal Swab Culture: Demonstration of Normal Flora”



Normal Flora of the Genitourinary Tract



Female Genital Tract

Dominated by *Lactobacillus* species that produce **lactic acid**, maintaining **acidic pH** (3.8-4.5) that **inhibits pathogens**.




Male Genital Tract

Lower microbial density compared to females. Distal urethra contains *Staphylococcus epidermidis*.



Urinary Tract

Bladder and upper urinary tract are normally **sterile**..



Sterile Sites in the Healthy Body

In a healthy individual, certain internal tissues and fluids are completely sterile—free from all microorganisms.




Blood and Cardiovascular System

Circulating blood is sterile; presence of bacteria indicates bloodstream infection (**septicemia**).



Central Nervous System

Brain tissue and cerebrospinal fluid (CSF) are sterile; microbial presence causes **meningitis** or **encephalitis**.



Muscle and Deep Tissues

Internal muscles and organs are sterile unless compromised by **trauma or surgery**.


Functions and Benefits of Normal Flora

Normal flora provides numerous protective and metabolic functions essential for human health.




Competitive Exclusion

Occupy ecological niches, preventing colonization by pathogenic microorganisms through competition for **nutrients and attachment sites**.




Vitamin Synthesis

Intestinal bacteria produce essential vitamins including vitamin K (clotting) and B vitamins (metabolism).



Immune Development

Stimulate and train the immune system, enhancing ability to distinguish self from non-self.



Digestive Assistance

Break down complex carbohydrates and produce short-chain fatty acids that feed intestinal cells.

Competitive Exclusion of Pathogens

Microbial Antagonism: Normal flora actively inhibits pathogen growth through competition and antimicrobial production.



Site Competition

Occupy attachment sites on epithelial surfaces, blocking pathogen adherence.



Nutrient Competition

Consume available nutrients, limiting resources for potential pathogens.




Antimicrobial Production

Produce fatty acids, hydrogen peroxide, and bacteriocins that kill or inhibit competing microbes.


💡 **Clinical Application:** Probiotic supplements aim to restore competitive exclusion after antibiotic treatment disrupts normal flora.

DYSBIOSIS



NORMAL

VS



DYSBIOSIS

Disturbance of Normal Flora

When the delicate balance of normal flora is disrupted, harmful consequences can occur. **This imbalance is called dysbiosis.**

1 Antibiotic Therapy

Broad-spectrum antibiotics kill beneficial bacteria along with pathogens, allowing resistant organisms to overgrow.

2 Stress and Illness

Physical or psychological stress can alter microbial populations and weaken immune defenses.

3 Dietary Changes

Sudden shifts in diet composition can rapidly change gut microbiota, affecting digestive health.

4 Immunosuppression

Weakened immunity from disease or medication allows normally harmless flora to become pathogenic.


Opportunistic Infections

Opportunistic microorganisms are members of normal flora that cause disease only when **host defenses are compromised** or when they access **abnormal body sites**.

Escherichia coli

Normal location: Large intestine

Pathogenic when: Enters urinary tract, causing UTIs



Staphylococcus aureus

Normal location: Nose, skin

Pathogenic when: Enters wounds, causing serious infections

💡 **Consider: The same bacterium that helps you digest food can cause life-threatening threatening infection if it enters your bloodstream.**

Symbiotic Relationships

Symbiosis describes the close, long-term biological interaction between different organisms.
Normal flora exists in symbiotic relationships with humans.

Relationship Type	Organism 1	Organism 2	Example
Mutualism	Benefits	Benefits	<i>E. coli</i> in human colon—bacteria get nutrients, humans get vitamin K
Commensalism	Benefits	Not affected	<i>Staphylococcus epidermidis</i> on skin—bacteria benefit, human unaffected
Parasitism	Benefits	Harmed	<i>Mycobacterium tuberculosis</i> in lungs—bacteria benefit, human develops TB

Most normal flora exists in mutualistic or commensalistic relationships with humans.

The Nurse's Role in Maintaining Healthy Flora

Nurses play a critical role in preserving patients' normal flora and preventing opportunistic infections.

01

Patient Education

Teach patients about appropriate antibiotic use, probiotic benefits, and maintaining healthy microbial balance.

02

Antibiotic Management

Support for appropriate antibiotic selection and duration to minimize disruption of normal flora.

03

Infection Prevention

proper hand hygiene and aseptic aseptic technique to prevent introducing pathogens while while preserving beneficial flora. flora.

04

Monitoring

Evaluate patients for signs of dysbiosis such as such as diarrhea, yeast infections, or opportunistic opportunistic infections.

05

Nutritional Support

Encourage balanced diet and adequate hydration hydration to support healthy microbial communities.

Hygiene and Infection Control

Balancing Act

Proper hygiene is essential for preventing pathogen transmission, but excessive cleanliness can disrupt protective normal flora.

Best Practices:

- Regular, gentle hand washing with soap
- Avoid overuse of antibacterial products
- Maintain skin integrity





Case Study: Antibiotic-Induced Dysbiosis

Patient Scenario: A 65-year-old female is hospitalized for pneumonia and treated with broad-spectrum antibiotics for 10 days.

- 1** Day 1-3
Patient receives IV antibiotics. Pneumonia begins to improve with decreased fever and improved breathing.
- 2** Day 7
Patient develops watery diarrhea (8-10 episodes/day) with abdominal cramping. Stool test positive for *C. difficile* toxin.
- 3** Diagnosis
Clostridium difficile infection—antibiotics killed protective gut flora, allowing *C. diff* overgrowth.
- 4** Treatment
Discontinue broad-spectrum antibiotic. Start targeted *C. diff* treatment with oral vancomycin. Implement contact precautions.

? **Critical Thinking:** How could this complication have been prevented or detected earlier?



➤ Watch this short video, you will learn

5 Moments of Hand Hygiene

Important notes

Normal flora consists of beneficial beneficial microorganisms that permanently or temporarily colonize colonize specific body sites.

These microbes provide essential functions including pathogen exclusion, vitamin synthesis, and immune system development.

Disruption of normal flora through through antibiotics, stress, or immunosuppression can lead to opportunistic infections.

Nurses must balance infection control with preservation of preservation of beneficial microbial communities.

Understanding normal flora is fundamental to preventing and managing healthcare-associated infections.



Multiple Choice Questions

1. Which microorganism is part of normal skin flora?

- A. Mycobacterium tuberculosis
- B. Staphylococcus epidermidis
- C. Candida albicans
- D. None of the above

2. What type of symbiosis do E. coli bacteria have with humans in the colon?

- A. Mutualism
- B. Commensalism
- C. Parasitism
- D. Competition

3. Which body site is normally sterile in healthy individuals?

- A. Skin
- B. Large intestine
- C. Cerebrospinal fluid
- D. Mouth

4. Which vitamin is synthesized by intestinal bacteria?

- A. Vitamin C
- B. Vitamin K
- C. Vitamin D
- D. Vitamin A

5. What is the term for an imbalance in normal flora?

- A. Symbiosis
- B. Contamination
- C. Dysbiosis
- D. Mutualism

9. Opportunistic infections occur when:

- A. Pathogens are always present in the body
- B. Host defenses are weakened or microorganisms enter abnormal sites
- C. The immune system is functioning normally
- D. Normal flora are completely absent

Thank you for your attention! attention!

💡 **Remember:** Understanding normal flora is essential for providing safe, effective patient care and preventing healthcare-associated infections.

Continue exploring the fascinating world of microbiology and its applications in clinical nursing practice.