



Medical Surgical Nursing Department Clinical Adult Nursing II

Performance cast care Check List for Evaluation.

Nursing Action	Correctly Done (2)	Incorrectly Done (1)	Not Done (0)
1- Wash hands, Wear gloves and apron			
2- support body part to be casted, and position it in normal anatomical position			
3- Drape patient.			
4-Wash and dry part to be casted.			
5-Place knitted material over part to be casted, and allow additional material			
6-Wrap soft, nonwoven roll padding smoothly and evenly around part, use additional padding around bony prominences.			
7- Apply plaster or non-plaster casting material evenly on body part. 8- Choose appropriate width bandage. 9- Overlap preceding turn by half the width of the bandage. 10- Use additional casting material (splints) at joints and at points of anticipated cast stress.			
11- Finish cast: smooth edges, trim and reshape with cast cutter.			
12-Remove particles of casting materials from skin.			
13-Handle hardening casts with palms of hands, and support it on firm smooth surface. 14-Don't rest cast on hard surfaces or on sharp edges, and avoid pressure on it.			
15-Promote drying of cast. 16-Leave cast uncovered.			
17- Clean equipment that used, return to usual places and discard used materials			
18- Wash hands and documentation			
Total grade			

Student's Signature

Instructor's Signature



Medical Surgical Nursing Department

Clinical Adult Nursing II

Performance Skin Traction check list for Evaluation.

Nursing Action	Correctly Done (2)	Incorrectly Done (1)	Not Done (0)
1-Wash hands and explain procedure to client			
2-prepare client and area of body to be in traction			
3-Position client as requested by physician or apply halter, strips, elastic bandages while physician and other assistants hold client's tissues in desired positions			
4-Assist with application of equipment			
5-Assist with attachment of spreader bars, ropes, pulleys. Ropes are tied securely in knots and passed in grooves of pulleys to weights			
6-When all traction materials and spreader bars are in place, weights are placed on weight holder and hooked to loop in rope and lowered slowly and gently until rope is taught.			
7-Ask client how traction is affected by injured tissues and raise side rails as appropriate			
8-Gather unused materials and return to storage areas and wash hands			
Total grade			

Student's signature :

Instructor signature:



Medical Surgical Nursing Department

Clinical Adult Nursing II

Performance Burn Wound Care Check List for Evaluation

Nursing Action	Correctly Done (2)	Incorrectly Done (1)	Not Done (0)
1-Explain the procedure to the patient and the patient adequately premeditated with analgesics as ordered.			
2-The nurse perform wound care should be aware with principles of infection control			
3-All supplies will be set up using sterile technique. A- clean the wound B-Remove the old dressing if present			
4-Once the dressings are removed. The patient's burn wound are gently cleansed with gauze pad and then assessed.			
5-Any loose skin or eschar can be removed with gauze or debrided with scissors and forceps and the patient's unburned areas are then washed.			
6-Wound dressing after wound cleaning, the burned areas are patted dry and the prescribed topical agent is applied and circumferential dressings should be applied distally to proximally			
7-Record any observations in the patient's wound and ask the doctor about any abnormal findings			
Total grade			

Student's signature :

Instructor signature:



Medical Surgical Nursing Department

Clinical Adult Nursing II

Performance Renal Biopsy Check List for Evaluation

Nursing Action	Correctly Done (2)	Incorrectly Done (1)	Not Done (0)
1-Make sure that the informed consent is signed			
2-The patient lies in prone position on a firm table with cushioning under the abdomen for support			
3-The kidney is located with ultrasound or CT scan			
4-The biopsy site is marked over the right kidney.			
5-Local anesthetic is injected (1-2% Lignocaine from the skin down to the capsule)			
6-Once the area is numb, the nephrologist uses ultrasound to guide a spring-loaded biopsy needle into the kidney and the needle removes about 5 to 15 glomeruli. Two or three samples are usually required.			
7-After the core tissue samples are extracted, pressure is applied to the incision for 2-5 minutes to slow bleeding, and a bandage is placed on the wound.			
8-Document procedure (Date, time, site, observations and signature).			
Total grade			

Student's signature :

Instructor signature:

Medical Surgical Nursing Department

Clinical Adult Nursing II

Performance Lumbar Puncture check list for Evaluation.

Nursing Action	Correctly Done (2)	Incorrectly Done (1)	Not Done (0)
1-Ask the patient to empty urinary bladder and bowel and explain procedure to patient.			
2-Position the patient laterally with a small pillow under head and a pillow between legs and should be lying on a firm surface.			
3-Skin is prepared with antiseptic solution and the skin, and the subcutaneous spaces are infiltrated with local anesthetic agent.			
4-A spinal puncture needle is introduced at L3-L4 interspaces. After the needle enters the subarachnoid space, help the patient to slowly straighten legs.			
5-Instruct the patient to breathe quietly and ask the patient to not talk			
6-The initial pressure reading is obtained by measuring the level of the fluid column after it comes to rest and			
7-About 2-3 ml spinal fluid is placed in each of three test tubes			
8-Closing pressure after the cerebrospinal fluid specimen is collected			
9-After procedure, the patient is instructed to remain in prone position without pillow for about 2 hours.			
10- Ensure adequate hydration with oral or parental fluids.			
Total grade			

Student's signature :

Instructor signature



Medical Surgical Nursing Department

Clinical Adult Nursing II

Performance hemodialysis check list for Evaluation.

Nursing Action	Correctly Done (2)	Incorrectly Done (1)	Not Done (0)
1-Assess vital signs and lab investigations			
2-Check for holding any antihypertensive medications			
3- Observe site of access for any abnormalities as: ✓ Vascular swelling , or redness			
4- Prepare hemodialysis machine, dialyzer, dialysate and supplies			
5-Perform hand hygiene and gloves 6-Insert needles in vascular access			
7- Uncovered access during hemodialysis session 8- Don't sleep on access arm 9- Avoid using the patient's dialysis access arm for administering I.V. medications, taking BP, or blood sample			
10-Assess vital signs especially blood pressure every 30 minutes			
11-Post dialysis: assess vital signs, monitor lab investigations 12- Instruct patients to: ✓ Avoid wearing tight clothes at vascular access arm. ✓ Avoid carrying heavy weight at vascular access arm. ✓ Avoid measuring blood pressure from vascular access arm. ✓ Avoid laying on vascular access arm.			
13- Discard all equipment and supplies in special places			
Total grade			

Student's signature :

Instructor signature



Medical Surgical Nursing Department

Clinical Adult Nursing II

Performance Colostomy Care Check List for Evaluation.

Nursing Action	Correctly Done (2)	Incorrec tly Done (1)	Not Done (0)
1-Have patient assume a relaxed position and provide privacy.			
2-To remove pouching system: a. Wear non-sterile gloves. b. Push down gently on skin while lifting up on the wafer. c. Discard soiled pouch and wafer in odor proof plastic bag.			
3- To cleans skin: d. Use toilet tissue to remove feces from stoma and skin if needed. e. Cleans stoma and peristomal skin with soft cloth and water, soap optional. Clip or shave peristomal hair if appropriate. c. Rinse and dry skin thoroughly after cleansing.			
4-To apply wafer: a. Use measuring guide to determine stoma size. b. Trace correct size onto back of wafer and cut to stoma size. c. Apply a line of skin barrier paste around stoma or on tip of wafer opening. d. Remove paper backing from the wafer, center opening over stoma, and press wafer down onto peristomal skin			
5-Snap pouch on to the flange of water according to manufacturer's directions.			
6-Apply tail closure to pouch tail			
Total grade			



Medical Surgical Nursing Department

Clinical Adult Nursing II

Performance Colostomy irrigation Check List for Evaluation.

Step	Correctly Done (2)	Incorrectly Done (1)	Not Done (0)
1. Explain to procedure to the patient/caregiver.			
2. Assist the patient to sit on the toilet or in a chair facing the toilet.			
3. Fill the irrigation bag with 500 to 1000 ml of water.			
4. Flush the tubing with water to empty if of air.			
5. Hang up the irrigation bag on the bathroom door hook, or raise the bag 18 inches above the stoma before irrigation.			
6. Take off the pouch and the soiled colostomy bag.			
7. Remove any stool or drainage from the stoma with toilet paper.			
8. Gently clean with soap and warm water as needed.			
9. Place the ring and irrigation sleeve over the stoma and adjust the belt for a snug fit (a mild soapy solution may be squirted into the sleeve for easier cleaning).			
10. Put the bottom of the sleeve into the toilet.			
11. Attach the cone tip to the irrigation tubing and lubricate.			
12. Gently insert the cone into the stoma for a snug fit. If resistance is met, perform a lubricated digital exam to remove feces. NEVER use force because this could perforate the bowel.			
Total			



Medical Surgical Nursing Department

Clinical Adult Nursing II

Performance Paracentesis Check List for Evaluation.

Step	Correctly Done (2)	Incorrectly Done (1)	Not Done (0)
1- Check the patient's coagulation profile, vital signs			
2- Explain procedure to patient			
3- Instruct the patient to void.			
4- Place patient in upright position on edge of bed and drape him with exposing abdomen			
5- Preparing skin with antiseptic solution.			
6- Have collection bottle and tubing available			
7- Physician administers local anesthesia and introduces needle or trocar.			
8- After treatment : assist the patient to a comfortable position.			
9- Record amount and characteristics of fluid removed, number of specimens sent to laboratory, the patient's condition during treatment			
10- Check blood pressure and vital signs every 1/2 hour for 2 hours, every hour for 4 hours, and every 4 hours for 24 hours.			
Total			



Medical Surgical Nursing Department

Clinical Adult Nursing II

Performance liver dialysis Check List for Evaluation.

Step	Correctly Done (2)	Incorrectly Done (1)	Not Done (0)
1- Check the patient's coagulation profile, neurological, vascular access			
2- Prepare and check the main circuits (blood or patient, albumin, and dialysis) and explain the procedure to the patient			
3- Continuous Monitoring for blood pressure, heart rate, oxygen saturation, level of consciousness, and complications			
4- Post- dialysis : Reassess: vital signs mental status improvement catheter site			
5- Documentation: Duration of session Complications Patient response			
Total			



Medical Surgical Nursing Department

Clinical Adult Nursing II

Performance liver biopsy Check List for Evaluation.

Step	Correctly Done (2)	Incorrectly Done (1)	Not Done (0)
1- Check the patient's coagulation profile, vital signs			
2- Check for signed consent, explain procedure			
3- Expose the right side of the patient's upper abdomen (right hypochondria)			
4- Instruct the patient to inhale and exhale deeply several times, finally to exhale, and to hold breath at the end of expiration			
5- Immediately after the biopsy, assist the patient to be on the right side and place a pillow under the costal margin.			
6- Instruct the patient to avoid coughing or straining.			
7- Remeasuring vital signs			
8- Instruct the patient to avoid heavy lifting and strenuous activity for 1 week			
TOTAL			